Supporting Information:

Catalytic Asymmetric Protonation of Lithium Enolates Using Amino Acid Derivatives as Chiral Proton Sources

Kaori Mitsuhashi, Rie Ito, Takayoshi Arai, and Akira Yanagisawa*

Experimental Section

General Methods. Analytical TLC was done on precoated (0.25 mm) silica gel plates. Column chromatography was conducted with 70-230 mesh silica gel. ^1H NMR spectra were recorded on 400 MHz spectrometers. Chemical shifts of ^1H NMR spectra were reported relative to tetramethylsilane (δ 0) or chloroform (δ 7.26). Splitting patterns are indicated as s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br, broad. ^{13}C NMR spectra were recorded on 100 MHz spectrometers. Chemical shifts of ^{13}C NMR spectra were reported relative to CDCl_3 (δ 77.0). All experiments were carried out under an atmosphere of standard grade argon gas (oxygen <10 ppm). α,α -Alkylated ketones were prepared by treatment of the corresponding ketones with LDA at $-78\text{ }^\circ\text{C}$ for 1 h followed by addition of alkyl halide in the presence of DMI. The silyl enolate of 2-benzylcyclohexanone was prepared by treating the corresponding ketone with bromomagnesium diisopropylamide in THF followed by silylation (TMSCl , Et_3N , HMPA).¹ Other silyl enolates were prepared by treatment of corresponding ketones with LDA at $-78\text{ }^\circ\text{C}$ for 1 h followed by addition of trimethylsilyl chloride. N -L-Aspartyl-L-phenylalanine methyl ester (1) was purchased from Aldrich.

Typical Experimental Procedure for Catalytic Asymmetric Protonation of Lithium Enolates Generated from the Corresponding Trimethylsilyl Enolates: Synthesis of (*S*)-2-methyl-1-tetralone (entry 7 in Table 2, entry 1 in Table 3, entry 1 in Table 4). Under argon atmosphere, a solution of *n*-BuLi in hexane (1.59 N, 0.58 mL, 0.92 mmol) was added to the trimethylsilyl enolate of 2-methyl-1-tetralone (196.1 mg, 0.84 mmol) in THF (4 mL), and the mixture was stirred at $-78\text{ }^\circ\text{C}$ for 1 h. To the resulting lithium enolate was added a solution of 1 (25.8 mg, 0.084 mmol) in DMF (0.5 mL) at $-78\text{ }^\circ\text{C}$ and the mixture was stirred for 5 min. Then a solution of BHT (185 mg, 0.84 mmol) in THF (5 mL) was added over a period for 2 h at $-78\text{ }^\circ\text{C}$. After the

mixture was stirred for 1 h, the reaction was quenched by trimethylsilyl chloride (107 μ L, 0.84 mmol) for 30 min at -78 °C to exclude the unreacted enolate. The reaction mixture was extracted with ethyl acetate (20 mL) three times, and the combined organic layer was washed with brine (20 mL). After drying (Na₂SO₄), the solvent was evaporated to dryness. The residue, which was purified by column chromatography on silica gel (hexane/ethyl acetate = 20/1) to give (*S*)-2-methyl-1-tetralone (106.3 mg, 79%) as a colorless oil: TLC *R*_f 0.35 (1:10 ethyl acetate/hexane); IR (neat) 2931, 1681, 1601, 1454, 1226, 966, 737 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 1.28 (d, 3 H, *J* = 6.8 Hz, CH₃), 1.84-1.94 (m, 1 H, one proton of CH₂), 2.17-2.24 (m, 1 H, one proton of CH₂), 2.55-2.63 (m, 1 H, CH), 2.95-3.08 (m, 2 H, CH₂), 7.23 (d, 1 H, *J* = 7.5 Hz, aromatic), 7.30 (t, 1 H, *J* = 7.6 Hz, aromatic), 7.46 (dt, 1 H, *J* = 1.5, 7.5 Hz, aromatic), 8.04 (dd, 1 H, *J* = 1.2, 7.7 Hz, aromatic); ¹³C NMR (100 MHz, CDCl₃) δ 15.3, 28.7, 31.2, 42.4, 126.3, 127.2, 128.6, 132.2, 132.9, 144.0, 200.5; [α]_D -42 (*c* 2.6, dioxane). The optical purity was determined to be 80% ee by chiral HPLC analysis (DAICEL CHIRALCEL OD-H, Hexane/*i*-PrOH = 99/1, flow rate = 0.5 mL/min): *t*_{minor} = 14.3 min, *t*_{major} = 16.0 min. The spectral data (¹H NMR, ¹³C NMR) were identical to those of literatures.²

Typical Experimental Procedure for Catalytic Asymmetric Protonation of Lithium Enolates from Ketones: Synthesis of (*S*)-2-methyl-1-tetralone (entry 1 in Table 5). Under argon atmosphere, a solution of *n*-BuLi in hexane (1.66 *N*, 0.67 mL, 1.11 mmol) was added to diisopropylamine (169 μ L, 1.21 mmol) in THF (4 mL) at 0 °C, and the mixture was stirred at -78 °C for 1 h. Then, 2-methyl-1-tetralone (160 mg, 1.00 mmol) was added and stirred at -78 °C for another 1 h. To the resulting lithium enolate was added a solution of **1** (30.7 mg, 0.10 mmol) in DMF (0.5 mL) at -78 °C and the mixture was stirred for 5 min. Then a solution of BHT (220 mg, 1.00 mmol) in THF (5 mL) was added over a period for 2 h at -78 °C. After the mixture was stirred for 1 h, the reaction was quenched by trimethylsilyl chloride (127 μ L, 1.00 mmol) for 30 min at -78 °C to exclude the unreacted enolate. The reaction mixture was extracted with ethyl acetate (20 mL) three times, and the combined organic layer was washed with brine (20 mL). After drying (Na₂SO₄), the solvent was evaporated to dryness. The residue was purified by column chromatography on silica gel (hexane/ethyl acetate = 20/1) to give (*S*)-2-methyl-1-tetralone (104.0 mg, 65%, 81% ee) as a colorless oil. The spectral data (¹H NMR, ¹³C NMR) were identical to those of literatures.²

2-Methyl-5-methoxy-1-tetralone (entry 2 in Table 4 and entry 2 in Table 5)³. TLC *R_f* 0.27 (1:10 ethyl acetate/hexane); IR (neat) 2933, 1682, 1596, 1581, 1469, 1456, 1438, 1255, 1055, 979, 739 cm⁻¹; ¹H NMR (400MHz, CDCl₃) δ 1.25 (d, 3 H, *J* = 6.7 Hz, CH₃), 1.78-1.86 (m, 1 H, one proton of CH₂), 2.17-2.22 (m, 1 H, one proton of CH₂), 2.54-2.61 (m, 1 H, CH), 2.71-2.78 (m, 1 H, one proton of CH₂), 3.06-3.11 (dt, 1 H, *J* = 4.3, 18 Hz, CH₂), 3.86 (s, 3 H, OCH₃), 7.00 (d, 1 H, *J* = 8.2 Hz, aromatic), 7.26 (t, 1 H, *J* = 7.9 Hz, aromatic), 7.64 (d, 1 H, *J* = 7.9 Hz, aromatic); ¹³C NMR (100 MHz, CDCl₃) δ 15.2, 22.2, 30.5, 42.0, 55.6, 113.9, 118.9, 126.7, 133.1, 133.4, 156.7, 201.0; HRMS (FAB+) calcd for C₁₂H₁₄O₂ (M⁺) 190.0994: observed 190.0992; [α]_D -31.2 (*c* 1.8, dioxane). The optical purity was determined to be 64% ee by chiral HPLC analysis (DAICEL CHIRALCEL OB, Hexane/*i*-PrOH = 20/1, flow rate = 0.5 mL/min): *t*_{major} = 14.7 min, *t*_{minor} = 19.7 min.

2-Methyl-6-methoxy-1-tetralone (entry 3 in Table 4 and entry 3 in Table 5)^{3,4}. TLC *R_f* 0.19 (1:10 ethyl acetate/hexane); IR (neat) 2962, 2933, 1672, 1597, 1494, 1456, 1358, 1335, 1249, 1155, 1132, 1085, 1031, 970 cm⁻¹; ¹H NMR (400MHz, CDCl₃) δ 1.26 (d, 3 H, *J* = 6.7 Hz, CH₃), 1.82-1.90 (m, 1 H, one proton of CH₂), 2.15-2.20 (m, 1 H, one proton of CH₂), 2.49-2.59 (m, 1 H, CH), 2.89-3.04 (m, 2 H, CH₂), 3.85 (s, 3H, OCH₃), 6.68 (d, 1 H, *J* = 2.1 Hz, aromatic), 6.82 (dd, 1 H, *J* = 2.1, 8.5 Hz, aromatic), 8.01 (d, 1 H, *J* = 8.9 Hz, aromatic); ¹³C NMR (100 MHz, CDCl₃) δ 15.6, 29.3, 31.5, 42.4, 55.5, 112.6, 113.1, 126.1, 129.9, 146.7, 163.4, 199.7; HRMS (FAB+) calcd for C₁₂H₁₄O₂ (M⁺) 190.0994: observed 190.0985; [α]_D -30.8 (*c* 0.22, dioxane). The optical purity was determined to be 41% ee by chiral HPLC analysis (DAICEL CHIRALCEL OJ-H, Hexane/*i*-PrOH = 20/1, flow rate = 0.5 mL/min): *t*_{major} = 21.7 min, *t*_{minor} = 23.4 min.

2-Ethyl-1-tetralone (entry 4 in Table 4 and entry 4 in Table 5)^{1a} TLC *R_f* 0.46 (1:10 ethyl acetate/hexane); IR (neat) 2961, 2933, 1682, 1597, 1581, 1469, 1455, 1257, 1221, 1056, 980, 738 cm⁻¹; ¹H NMR (400MHz, CDCl₃) δ 1.01 (t, 3 H, *J* = 7.5 Hz, CH₃), 1.52-1.60 (m, 1 H, one proton of CH₂), 1.85-2.02 (m, 2 H, CH₂), 2.21-2.28 (m, 1 H, one proton of CH₂), 2.38-2.45 (m, 1 H, CH), 2.98-3.02 (m, 2 H, CH₂), 7.24 (d, 1 H, *J* = 7.0 Hz, aromatic), 7.30 (t, 1 H, *J* = 7.3 Hz, aromatic), 7.46 (dt, 1 H, *J* = 1.2, 7.3 Hz, aromatic), 8.03 (dd, 1 H, *J* = 1.2, 7.7 Hz, aromatic); ¹³C NMR (100 MHz, CDCl₃) δ 11.4, 22.4, 27.7, 28.3, 48.9, 126.4, 127.4, 128.6, 132.6, 133.0, 144.0, 200.3; [α]_D -20.6 (*c* 1.8, dioxane). The optical purity was determined to be 88% ee by chiral HPLC analysis (DAICEL CHIRALCEL OD-H, Hexane/*i*-PrOH = 150/1, flow rate = 0.25 mL/min): *t*_{major} = 28.0 min, *t*_{minor} = 29.8 min.

2-Propyl-1-tetralone (entry 5 in Table 4)^{1a}. TLC *R_f* 0.46 (1:10 ethyl acetate/hexane); IR (neat) 2961, 2933, 1597, 1581, 1469, 1455, 1257, 1221, 1056, 980, 738 cm⁻¹; ¹H NMR (400MHz, CDCl₃) δ 0.96 (t, 3 H, *J* = 7.2 Hz, CH₃) 1.33-1.53(m, 3 H, one proton of CH₂ and CH₂), 1.84-1.97 (m, 2 H, CH₂), 2.20-2.27 (m, 1 H, one proton of CH₂), 2.45-2.51 (m, 1 H, CH), 2.92-3.06 (m, 2 H, CH₂), 7.23 (d, 1 H, *J* = 7.5 Hz, aromatic), 7.29 (t, 1 H, *J* = 7.5 Hz, aromatic), 7.45 (dt, 1 H, *J* = 1.5, 7.5 Hz, aromatic), 8.03 (dd, 1 H, *J* = 1.2, 7.7 Hz, aromatic); ¹³C NMR (100 MHz, CDCl₃) δ 14.1, 20.1, 28.1, 28.2, 31.5, 47.1, 126.4, 127.3, 128.5, 132.4, 132.9, 143.8, 200.3; $[\alpha]_D^{25}$ -1.7 (*c* 2.37, chloroform). The optical purity was determined to be 11% ee by chiral HPLC analysis (DAICEL CHIRALCEL OJ-H, Hexane/*i*-PrOH = 99/1, flow rate = 0.5 mL/min): *t*_{minor} = 14.1 min, *t*_{major} = 15.1 min.

(S)-2-Benzyl-1-cyclohexanone (entry 6 in Table 4)². TLC *R_f* 0.38 (1:10 ethyl acetate/hexane); IR (neat) 2934, 2860, 1706, 1495, 1449, 1127, 731, 669 cm⁻¹; ¹H NMR (400MHz, CDCl₃) δ 1.30-1.41(m, 1 H, one proton of CH₂), 1.52-1.74 (m, 2 H, CH₂), 1.79-1.87 (m, 1 H, CH₂), 1.98-2.10 (m, 2 H, CH₂), 2.29-2.47 (m, 3 H, one proton of CH₂ and CH₂), 2.51-2.59 (m, 1 H, CH), 3.24(dd, 1H, *J* = 4.7, 13.9 Hz, CH₂), 7.12-7.21 (m, 3 H, aromatic), 7.25-7.29 (m, 2 H, aromatic); ¹³C NMR (100 MHz, CDCl₃) δ 25.0, 28.0, 33.4, 35.4, 42.1, 52.5, 125.9, 128.3 (2 C), 129.1 (2 C), 140.4, 212.6; HRMS (FAB+) calcd for C₁₃H₁₇O (*M*⁺+1) 189.1279; observed 189.1270; $[\alpha]_D^{25}$ -31.5 (*c* 2.02, methanol). The optical purity was determined to be 69% ee by chiral HPLC analysis (DAICEL CHIRALCEL OJ-H, Hexane/*i*-PrOH = 9/1, flow rate = 0.5 mL/min): *t*_{minor} = 13.5 min, *t*_{major} = 14.5 min.

3-Methyl-4-chromanone (entry 5 in Table 5)⁵. TLC *R_f* 0.32 (1:10 ethyl acetate/hexane); IR (neat) 2971, 2938, 2871, 1688, 1604, 1476, 1455, 1322, 1295, 1211, 1055, 1035, 1013, 755 cm⁻¹; ¹H NMR (400MHz, CDCl₃) δ 1.22(d, 3 H, *J* = 7.0, CH₃), 2.82-2.90 (m, 1 H, CH), 4.15(dd, 1 H, *J* = 9.7, 11.4, one proton of CH₂), 4.50 (dd, 1 H, *J* = 5.2, 11.3, one proton of CH₂), 6.95-7.04(m, 2 H, aromatic), 7.44-7.49 (m, 1 H, aromatic), 7.89-7.92 (m, 1 H, aromatic); ¹³C NMR (100 MHz, CDCl₃) δ 10.6, 40.7, 72.2, 117.6, 120.5, 121.3, 127.3, 135.7, 161.7, 194.8; $[\alpha]_D^{25}$ +7.2 (*c* 1.60, dioxane). The optical purity was determined to be 20% ee by chiral HPLC analysis (DAICEL CHIRALCEL OB, Hexane/*i*-PrOH = 99/1, flow rate = 0.5 mL/min): *t*_{minor} = 24.2 min, *t*_{major} = 29.2min.

References

- (1) Krafft, M. E.; Holton, R. A. *Tetrahedron Lett.* **1983**, *24*, 1345.

- (2) (a) Adameczyk, M.; Watt, D. S.; Netzel, D. A. *J. Org. Chem.* **1984**, *49*, 4226.
(b) Murakata, M.; Yasukata, T.; Aoki, T.; Nakajima, M.; Koga, K. *Tetrahedron* **1998**, *54*, 2449.
- (3) Kieboom, A. P. G.; Van Bekkum, H. *Synthesis* **1970**, 476.
- (4) Harrowven, D. C.; Dainty, R. F. *Tetrahedron* **1997**, *53*, 15771.
- (5) Barlocco, D.; Cignarella, G.; Curzu, M. M. *Synthesis* **1985**, 876.

¹H-NMR

Date : Fri May 13 20:45:23 2005

File Name : 050513mit_164.nmdata

Comment : 050513mit_164

Slcchistory : non

EXMODE : non

POINT : 32768 points

SAMPO : 32768 points

FREQ : 10000.0 Hz

FILTR : 5000 Hz

DELAY : 40.0 usec

DEADT : 57.2 usec

INVL : 100.0 usec

TIMES : 8 times

DUMY : 1 times

PD : 3.7232 sec

ACQIM : 3276.7996 msec

PREDL : 10.00000 msec

INWIT : 0.5000 msec

RESOL : 0.31 Hz

PW1 : 5.60 usec

OBNUC : ¹H

OBFRQ : 500.00 MHz

GBSET : 162160.00 Hz

RGAIN : 16

SCANS : 8 times

SLVNT : CDCL3

SPINNING : 10 Hz

TEMP : 25.0 C

Chemical structure: 1-methyl-2-phenylcyclohexan-1-one

CC1(C(=O)C2=CC=CC=C2)CCCCC1

¹³C-NMR

Date : Fri May 13 20:25:09 2005

File Name : 050513mit_bcm175.nmdata

Comment : bcm175

Slcchistory : bcm

EXMODE : bcm

POINT : 32768 points

SAMPD : 32768 points

FREQ : 33898.3 Hz

FILTR : 15950 Hz

DELAY : 11.8 usec

DEADT : 10.0 usec

Parameter	Value
INTVL	29.5 usec
TIMES	255 times
DUMMY	1 times
PD	2.0333 sec
ACQTM	966.6560 msec
PREDL	10.00000 msec
INITI	10.0000 msec
RESOL	1.03 Hz
PA1	7.25 usec
DBNUC	13C
DBFRQ	125.65 MHz
DBSET	127958.00 Hz
RGAIN	27
IRNUC	1H
IRFRQ	500.00 MHz
IRSET	162160.00 Hz
IRPW	45.0 usec
IRNS	0
SCANS	152 times
SLVNT	CDCL3
SPINNING	13 Hz
TEMP	26.2 C