Catalytic Asymmetric Reductive Michael Cyclization

Jung Woon Yang, Maria T. Hechavarria Fonseca, and Benjamin List*

Max-Planck-Institut für Kohlenforschung, D-45470 Mülheim an der Ruhr, Germany,

list@mpi-muelheim.mpg.de

Supporting Information

General: Reactions were monitored by thin layer chromatography using 0.25 mm E. Merck silica gel coated glass plates (60F-254) using UV light to visualize the course of reaction. Flash column chromatography was performed using E. Merck silica gel (60, particle size 0.040–0.063 mm). Chemical yields refer to pure isolated substances. ^1H and ^{13}C NMR spectra were obtained using either a Bruker DPX-300 or AV-400 spectrometer. Chemical shifts are reported in ppm from tetramethylsilane with the solvent resonance as the internal standard. The following abbreviations were used to designate chemical shift multiplicities: s = singlet, d = doublet, t = triplet, q = quartet, h = heptet, m = multiplet, br = broad. MS (EI): Finnigan MAT 8200 (70 eV), ESI-MS: Finnigan MAT 95, accurate mass determinations: Bruker APEX III FT-MS (7 T magnet).

Preparation of Catalysts

Catalysts **9e-f** were prepared according to the procedure of MacMillan *et al.*¹

^1H NMR (400 MHz, CDCl_3) δ 7.41-7.26 (m, 5H), 7.16 (d, J = 8.6 Hz, 2H), 6.89 (d, J = 8.6 Hz, 2H), 5.02 (s, 2H), 4.04 (s, 1H), 3.67-3.64 (m, 1H), 3.06 (dd, J = 14.0, 4.1 Hz, 1H), 2.92 (dd, J = 13.8, 7.0 Hz, 1H), 2.88 (s, 3H), 1.82 (br, 1H), 0.81 (s, 9H); ^{13}C

NMR (100 MHz, CDCl_3) δ 175.0, 157.3, 136.7, 130.3, 129.6, 128.2, 127.5, 127.0, 114.7, 82.1, 69.7, 59.1, 36.7, 34.6, 30.3, 25.0; HRMS (ESI) calcd for MNa^+ 375.204299, found 375.204124.

^1H NMR (300 MHz, CDCl_3) δ 7.07 (d, J = 8.4 Hz, 2H), 6.85 (d, J = 8.6 Hz, 2H), 3.98 (s, 1H), 3.63-3.60 (m, 1H), 2.98 (d, J = 5.3 Hz, 2H), 2.83 (s, 3H), 1.24 (s, 9H), 0.73 (s, 9H); ^{13}C NMR

¹ Paras, N. A.; MacMillan, D. W. C. *J. Am. Chem. Soc.* **2002**, 124, 7894 - 7895

(100 MHz, CDCl₃) δ 175.1, 153.8, 131.8, 129.8, 124.0, 82.1, 77.9, 58.9, 36.3, 34.5, 30.3, 28.4, 25.0; HRMS (ESI) calcd for MNa⁺ 341.219949, found 341.219842.

General procedure for the preparation of enal enones,² illustrated with compound 10

To a stirred solution of 2-bromobenzaldehyde (6.14 g, 33.17 mmol) in 40 mL of DMF were added acrolein diethyl acetal (8.64 g, 66.34 mmol), $n\text{Bu}_4\text{NOAc}$ (20.0 g, 66.34 mmol), K_2CO_3 (6.88 g, 49.76 mmol), KCl (2.47 g, 33.17 mmol), and $\text{Pd}(\text{OAc})_2$ (223.4 g, 0.995 mmol). The mixture was stirred at 90°C for 2 h. After cooling, the reaction mixture was quenched using saturated aqueous NaCl and extracted with diethyl ether. The combined organic layer were dried (Na_2SO_4), filtered, and concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (10% diethyl ether/hexane) to give 4.08 g (52% yield) of the unsaturated acetal.

^1H NMR (400 MHz, CDCl_3) δ 10.2 (s, 1H), 7.76 (d, $J = 0.7$ Hz, 1H), 7.49 (m, 2H), 7.36 (m, 1H), 6.07 (dd, $J = 15.9, 5.1$ Hz, 1H), 5.03 (m, 1H), 3.66 (m, 3H), 3.52 (m, 3H), 1.17 (t, $J = 5.0$ Hz, 6H); ^{13}C NMR (100 MHz, CDCl_3) δ 191.8, 134.2, 132.7, 131.8, 131.1, 128.7, 127.6, 127.3, 100.9, 61.0, 14.9.

To a solution of the aldehyde (1.60 g, 6.83 mmol) in 50 mL of CHCl_3 was added (benzoylmethylene)triphenylphosphorane (3.90 g, 10.25 mmol) at room temperature. The reaction mixture was stirred overnight under reflux condition. Then, the solvent was removed under reduced pressure and the crude material was purified by flash chromatography (10% ethyl acetate/hexane), providing the corresponding enone (1.93 g, 84%) as a brownish oil. Subsequently, to a stirred solution of the enone (1.80 g, 5.35 mmol) in 53.5 mL of acetone was added aqueous 4N HCl (5.35 mL) solution at room temperature. After 10 min, the reaction mixture was quenched using saturated aqueous NaCl and extracted with diethyl ether. The combined organic layer were dried (Na_2SO_4), filtered and concentrated *in vacuo*. The

² Battistuzzi, G.; Cacchi, S.; Fabrizi, G. *Org. Lett.*, **2003**, 5, 777-780.

residue was recrystallized from dichloromethane/diethylether to give 1.25 g (89%) of **10** as yellowish solid. Compound **12**, **14**, **16**, **18**, **20**, **22** (Knoevenagel instead of Wittig) were prepared analogously.

^1H NMR (300 MHz, CDCl_3) δ 9.71 (d, $J = 7.6$ Hz, 1H), 8.14 (d, $J = 15.4$ Hz, 1H), 7.97 (d, $J = 5.6$ Hz, 2H), 7.88 (d, $J = 15.4$ Hz, 1H), 7.71-7.39 (m, 8H), 6.62 (q, $J = 8.2$ Hz, 1H); ^{13}C NMR (75.5 MHz, CDCl_3) δ 193.8, 190.0, 149.2, 141.0, 138.1, 135.6, 134.4, 133.6, 131.9, 131.4, 130.8, 129.2, 129.0, 128.3, 128.1, 126.3; HRMS (CI, isobutane) calcd for MH^+ 263.107205, found 263.107547.

^1H NMR (400 MHz, CDCl_3) δ 9.78 (d, $J = 7.6$ Hz, 1H), 8.55 (s, 1H), 8.26 (d, $J = 15.4$ Hz, 1H), 8.12 (d, $J = 8.6$ Hz, 1H), 8.01-7.90 (m, 4H), 7.81 (d, $J = 7.6$ Hz, 1H), 7.69-7.47 (m, 6H), 6.70 (dd, $J = 15.8, 7.6$ Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 193.3,

189.4, 148.8, 140.6, 135.7, 135.3, 135.2, 134.1, 132.6, 131.5, 131.0, 130.4, 130.2, 129.6, 128.8, 128.7, 128.0, 127.9, 127.8, 127.0, 126.1, 124.4; HRMS (EI) calcd for MH^+ 312.115030, found 312.115074.

^1H NMR (300 MHz, CDCl_3) δ 9.70 (d, $J = 7.6$ Hz, 1H), 8.13 (d, $J = 15.3$ Hz, 1H), 7.85 (d, $J = 15.8$ Hz, 1H), 7.68-7.39 (m, 5H), 7.21-7.09 (m, 2H), 6.61 (dd, $J = 15.8, 7.6$ Hz, 1H); ^{13}C NMR (75.5 MHz, CDCl_3) δ 193.7, 180.6, 149.0, 147.1, 140.9, 135.1, 134.5, 132.6,

132.0, 131.9, 131.4, 131.0, 128.3, 128.2, 124.9, 124.0; HRMS (EI) calcd for MH^+ 345.966328, found 345.966600.

^1H NMR (300 MHz, CDCl_3) δ 9.78 (d, $J = 7.6$, 1H), 7.89 (dd, $J = 10.6, 15.8$ Hz, 2H), 7.63 (m, 2H), 7.46 (m, 2H), 6.66 (dd, $J = 15.8, 8.3$ Hz, 2H), 2.42 (s, 3H); ^{13}C NMR (75.5 MHz, CDCl_3) δ 197.5, 193.3, 148.6, 139.1, 134.6, 133.8, 131.6, 131.0, 130.5, 130.4, 127.9, 127.8, 28.4; HRMS (EI)

calcd for MH^+ 200.083730, found 200.083669.

^1H NMR (300 MHz, CDCl_3) δ 9.69 (d, $J = 7.6$ Hz, 1H), 8.08 (d, $J = 15.4$ Hz, 1H), 7.99-7.96 (m, 2H), 7.80 (d, $J = 15.8$ Hz, 1H), 7.63-7.53 (m, 2H), 7.49-7.44 (m, 3H), 7.40-7.34 (m, 1H), 7.12 (ddd, $J = 8.2, 8.2, 2.6$ Hz, 1H), 6.57 (dd, $J = 15.8, 7.6$ Hz, 1H); ^{13}C NMR

(75.5 MHz, CDCl_3) δ 193.6, 189.6, 166.1, 162.7, 147.9, 139.7, 137.9, 133.9, 131.6, 130.6,

130.4, 129.3, 129.0, 127.1, 118.3, 118.0, 114.9, 114.6; HRMS (ESI) calcd for MNa^+ 305.094828, found 305.094859.

^1H NMR (300 MHz, CDCl_3) δ 9.67 (d, $J = 7.6$ Hz, 1H), 8.14 (d, $J = 15.4$ Hz, 1H), 7.99-7.95 (m, 2H), 7.88 (d, $J = 15.7$ Hz, 1H), 7.57-7.43 (m, 3H), 7.33 (d, $J = 15.4$ Hz, 1H), 7.08 (d, $J = 22.0$ Hz, 1H), 6.56 (dd, $J = 15.6, 7.6$ Hz, 1H), 3.96 (s, 3H), 3.90 (s, 3H); ^{13}C NMR (75.5 MHz, CDCl_3) δ 193.9, 190.2, 152.0, 151.5, 148.6, 140.6, 138.3, 133.5, 129.9, 129.3, 129.1, 128.9, 128.0, 124.4, 109.7, 109.4, 56.6, 56.5; HRMS (ESI) calcd for MNa^+ 419.182895, found 419.182758.

^1H NMR (300 MHz, CDCl_3) δ 9.66 (d, $J = 7.6$ Hz, 1H), 7.96 (s, 1H), 7.62-7.56 (m, 2H), 7.40-7.30 (m, 2H), 6.59 (dd, $J = 15.8, 7.6$ Hz, 1H), 4.28 (q, $J = 7.1$ Hz, 2H), 4.09 (q, $J = 7.1$ Hz, 2H), 1.29 (t, $J = 7.1$ Hz, 3H), 1.03 (t, $J = 7.1$ Hz, 3H); ^{13}C NMR (75.5 MHz, CDCl_3) δ 193.8, 165.7, 163.8, 149.1, 141.2, 134.5, 133.4, 131.5, 131.2, 131.1, 130.4, 129.3, 127.8, 62.4, 62.0, 14.5, 14.1; HRMS (ESI) calcd for MNa^+ 327.120844, found 327.120691.

Substrates **24** and **26** were prepared according to the procedure of Roush *et al.*³

^1H NMR (400 MHz, CDCl_3) δ 9.46 (d, $J = 7.7$ Hz, 1H), 7.85 (d, $J = 7.6$ Hz, 2H), 7.51-7.38 (m, 3H), 6.99-6.75 (m, 3H), 6.14-6.08 (m, 1H), 2.51 (m, 4H); ^{13}C NMR (100 MHz, CDCl_3) δ 193.2, 189.9, 155.5, 146.2, 137.3, 133.3, 132.5, 128.2, 128.1, 126.6, 30.7, 30.5; HRMS (CI) calcd for MH^+ 215.107206, found 215.106931.

^1H NMR (300 MHz, CDCl_3) δ 9.53 (d, $J = 7.9$ Hz, 1H), 7.93 (d, $J = 8.2$ Hz, 2H), 7.57-7.44 (m, 3H), 7.09-6.80 (m, 3H), 6.19-6.11 (m, 1H), 2.45-2.35 (m, 4H), 1.82-1.72 (m, 2H); ^{13}C NMR (75.5 MHz, CDCl_3) δ 193.8, 190.5, 157.3, 148.1, 137.8, 133.5, 132.8, 128.6, 128.5, 126.6, 32.1, 26.3; HRMS (ESI) calcd for MNa^+ 251.104344, found 251.104252.

³ Frank, S. A.; Mergott, D. J.; Roush, W. R. *J. Am. Chem. Soc.* **2002**, *124*, 2404 -2405.

Screening of different Hantzsch esters

Representative experimental procedure for the catalytic asymmetric reductive Michael cyclization. To a solution of enal enones **10**, **12**, **14**, **16**, **18**, **20**, **22**, **24**, and **26** (0.5 mmol) and imidazolidinone-based chiral organocatalysts **9a-g** (20 mol%) in dry 1,4-dioxane (5 mL) was added Hantzsch ester **4a** (1.1 equiv). The reaction mixture was stirred at room temperature for 2-4 h, after which the solvent was removed and residue was chromatographed on silica gel to give the pure product (**11**, **13**, **15**, **17**, **19**, **21**, **23**, **25**, and **27**).

The synthesis of the racemic compounds were performed according to the above mentioned procedure at room temperature using dibenzylammonium trifluoroacetate (20 mol%) as the catalyst.

¹H NMR (300 MHz, CDCl₃) δ 9.91 (d, *J* = 2.5 Hz, 1H), 7.99-7.96 (m, 2H), 7.61-7.55 (m, 1H), 7.50-7.44 (m, 2H), 7.27-7.18 (m, 4H), 4.21-4.14 (m, 1H), 3.55 (dd, *J* = 17.9, 4.7 Hz, 1H), 3.37-3.25 (m, 2H), 3.15 (dd, *J* = 16.3, 8.9 Hz, 1H), 3.03-2.95 (m, 1H); ¹³C NMR (75.5 MHz, CDCl₃) δ 202.2, 198.5, 143.8, 141.1, 136.6, 133.4, 128.7, 128.1, 127.5, 127.1, 124.8, 123.9, 57.5, 44.3, 41.2, 31.8; HRMS (EI) calcd for MH⁺ 264.115030, found 264.115096; The enantiomeric excess was determined to be 96% by chiral HPLC (ChiralPak AD-H column, 5% *i*-PrOH/heptane, 1.0 mL/min, 220 nm, *t*_R (minor 16.2 min), *t*_R (major 17.6 min).

¹H NMR (300 MHz, CDCl₃) δ 9.85 (d, *J* = 1.7 Hz, 1H), 8.38 (s, 1H), 7.95 (dd, *J* = 8.6, 1.7 Hz, 1H), 7.85-7.76 (m, 3H), 7.53-7.42 (m, 2H), 7.18-7.10 (m, 4H), 4.17-4.10 (m, 1H), 3.58 (dt, *J* = 17.8, 4.8 Hz, 1H), 3.38-3.19 (m, 2H), 3.07 (dd, *J* = 16.1, 8.8 Hz, 1H), 2.97-2.90 (m, 1H); ¹³C NMR (75.5 MHz, CDCl₃) δ 202.6, 198.9, 144.3, 141.6, 136.1, 134.4, 132.9, 130.3, 130.0, 129.1, 128.2, 127.9, 127.5, 127.3, 125.3, 124.4, 124.1, 58.0, 44.8, 41.8,

32.2; HRMS (EI) calcd for MH^+ 312.115030, found 312.115074; The enantiomeric excess was determined to be 94% by chiral HPLC (ChiralPak AS-H column, 15% *i*-PrOH/heptane, 1.0 mL/min, 220 nm, t_R (minor 22.1 min), t_R (major 35.2 min).

1H NMR (300 MHz, $CDCl_3$) δ 9.78 (d, $J = 2.2$ Hz, 1H), 7.37 (d, $J = 4.1$ Hz, 1H), 7.20-7.11 (m, 4H), 7.03 (d, $J = 4.1$ Hz, 1H), 4.10-4.03 (m, 1H), 3.31 (dd, $J = 17.1, 5.1$ Hz, 1H), 3.23 (dd, $J = 16.2, 6.3$ Hz, 1H), 3.11 (dd, $J = 8.7, 4.6$ Hz, 1H), 3.06 (dd, $J = 8.7, 5.6$ Hz, 1H),

2.98-2.91 (m, 1H); ^{13}C NMR (75.5 MHz, $CDCl_3$) δ 202.3, 190.7, 145.7, 143.7, 141.3, 132.6, 131.8, 128.0, 127.5, 125.3, 124.3, 123.6, 57.7, 44.5, 41.6, 32.2; HRMS (ESI) calcd for MNa^+ 370.971747, found 370.971422; The enantiomeric excess was determined to be 92% by chiral HPLC (ChiralPak AD-H column, 5% *i*-PrOH/heptane, 1.0 mL/min, 220 nm, t_R (minor 20.7 min), t_R (major 24.9 min).

1H NMR (300 MHz, $CDCl_3$) δ 9.83 (t, $J = 1.6$ Hz, 1H), 7.26-7.14 (m, 4H), 4.01-3.94 (m, 1H), 3.26 (dd, $J = 16.3, 6.8$ Hz, 1H), 3.12 (dd, $J = 16.3, 8.8$ Hz, 1H), 3.02 (dd, $J = 17.8, 5.1$ Hz, 1H), 2.94-2.85 (m, 1H), 2.75 (dd, $J = 17.8, 8.7$ Hz, 1H), 2.21 (s, 3H); ^{13}C NMR (75.5 MHz, $CDCl_3$) δ 207.2, 202.2,

143.6, 140.9, 127.4, 127.1, 124.8, 123.8, 57.6, 49.0, 40.8, 31.9, 30.3; HRMS (EI) calcd for MH^+ 202.099380, found 202.099104; The enantiomeric excess was determined to be 91% by chiral HPLC (ChiralPak AD-H column, 5% *i*-PrOH/heptane, 1.0 mL/min, 220 nm, t_R (major 13.5 min), t_R (minor 14.4 min).

1H NMR (300 MHz, $CDCl_3$) δ 9.81 (d, $J = 2.1$ Hz, 1H), 7.91-7.86 (m, 2H), 7.53-7.47 (m, 1H), 7.41-7.35 (m, 2H), 7.11-7.06 (m, 1H), 6.83-6.76 (m, 2H), 4.09-4.03 (m, 1H), 3.42 (dd, $J = 18.0, 5.0$ Hz, 1H), 3.27-3.13 (m, 2H), 3.06-2.90 (m, 2H); ^{13}C NMR (75.5 MHz, $CDCl_3$)

δ 202.1, 198.6, 164.4, 161.2, 146.5, 136.8, 133.9, 129.2, 128.5, 126.1, 115.0, 114.7, 111.7, 111.4, 58.3, 44.4, 41.5, 31.4; HRMS (ESI) calcd for MNa^+ 305.094828, found 305.094859; The enantiomeric excess was determined to be 97% by chiral HPLC (ChiralPak AS-H column, 15% *i*-PrOH/heptane, 1.0 mL/min, 220 nm, t_R (minor 20.8 min), t_R (major 30.2 min).

1H NMR (300 MHz, $CDCl_3$) δ 9.81 (d, $J = 2.1$ Hz, 1H), 7.90-7.67 (m, 2H), 7.52-7.46 (m, 1H), 7.41-7.36 (m, 2H), 6.67 (d, $J = 8.8$ Hz, 2H), 4.06-4.02 (m, 1H), 3.77 (s, 3H), 3.74 (s, 3H), 3.42 (dd, $J =$

17.8, 4.9 Hz, 1H), 3.22 (dt, $J = 13.4, 8.9$ Hz, 1H), 3.15 (t, $J = 5.3$ Hz, 1H), 3.01 (dd, $J = 15.8, 8.7$ Hz, 1H), 2.94-2.87 (m, 1H); ^{13}C NMR (75.5 MHz, CDCl_3) δ 202.7, 199.1, 149.3, 149.0, 137.1, 135.7, 133.8, 133.1, 129.1, 128.4, 108.1, 107.5, 58.1, 56.5, 45.0, 41.9, 31.9; HRMS (ESI) calcd for MNa^+ 347.125381, found 347.125686; The enantiomeric excess was determined to be 96% by chiral HPLC (ChiralPak AD-H column, 10% *i*-PrOH/heptane, 1.0 mL/min, 220 nm, t_R (minor 23.4 min), t_R (major 29.7 min).

^1H NMR (300 MHz, CDCl_3) δ 9.72 (d, $J = 1.4$ Hz, 1H), 7.15-7.06 (m, 4H), 4.23 (dd, $J = 6.9, 4.2$ Hz, 1H), 4.14 (q, $J = 7.1$ Hz, 2H), 4.02 (q, $J = 7.1$ Hz, 2H), 3.72 (d, $J = 7.0$ Hz, 1H), 3.40-3.33 (m, 1H), 3.24-3.08 (m, 2H), 1.18 (t, $J = 7.1$ Hz, 3H), 1.04 (t, $J = 7.1$ Hz, 3H); ^{13}C NMR (75.5 MHz, CDCl_3) δ

200.4, 167.3, 167.0, 140.6, 139.6, 126.9, 125.9, 123.8, 123.6, 60.7, 60.5, 54.7, 52.7, 43.6, 30.5, 13.0, 12.8; HRMS (ESI) calcd for MNa^+ 327.120844, found 327.120691; The enantiomeric excess was determined to be 86% by chiral HPLC (ChiralPak AD-H column, 3% *i*-PrOH/heptane, 1.0 mL/min, 220 nm, t_R (major 13.3 min), t_R (minor 14.9 min).

All spectroscopic data (HRMS, ^1H NMR, ^{13}C NMR) matches data previously reported by List and Shabat *et al.*⁴ After derivatization of the aldehyde with trimethyl phosphonoacetate (4.0 equiv) and LiOH-mono hydrate (1.5 equiv) proceeds rapidly in THF to form an enone, the enantiomeric excess was

determined to be 72% by chiral HPLC (ChiralPak AD-H column, 5% *i*-PrOH/heptane, 1.0 mL/min, 220 nm, t_R (major 12.2 min), t_R (minor 15.1 min).

^1H NMR (400 MHz, CDCl_3) δ 9.50 (d, $J = 3.4$ Hz, 1H), 7.89-7.87 (m, 2H), 7.51-7.44 (m, 1H), 7.41-7.36 (m, 2H), 3.04 (dd, $J = 16.4, 4.3$ Hz, 1H), 2.69 (dd, $J = 16.4, 8.4$ Hz, 1H), 2.36-2.26 (m, 1H), 2.12-2.05 (m, 1H), 1.84-1.73 (m, 3H), 1.67-1.60 (m, 1H), 1.39-1.29 (m, 1H), 1.27-1.18 (m, 2H), 1.06-0.96 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) 203.5, 198.3,

136.0, 132.0, 127.6, 127.1, 54.5, 42.5, 31.6, 30.5, 25.1, 24.3, 23.7

After derivatization of the aldehyde with trimethyl phosphonoacetate (4.0 equiv) and LiOH-mono hydrate (1.5 equiv) in THF to form an enone, the enantiomeric excess was determined to be 95% by chiral GC (25 m, IVADEX1/PS086 G376, 0.5 bar H_2 as carrier gas): t_R (minor 154.0 min), t_R (major 154.8 min).

⁴ a) Hechavarria Fonseca, M. T.; List, B. *Angew. Chem., Int. Ed. Engl.* **2004**, 43, 3958-60. b) Weinstein, R.; Lerner, R. A.; Barbas, C. F., III; Shabat, D. *J. Am. Chem. Soc.* **2005**, 127, ASAP.

Determination of the relative configuration

The *trans*-configuration of compound **25** is known.⁴ The relative configuration of aldehyde **27** was deduced from NMR. The *trans*-configuration of **11**, **13**, **15**, **17**, **19**, **21**, and **23** was assigned analogously. In addition, treating aldehyde **11** with DBU (10 mol%) in CH₂Cl₂ did not result in any measurable isomerization (NMR) as would be expected for the *trans*-isomer.

Determination of the absolute configuration of 25

The absolute configuration of **25** was determined by comparing the optical rotation with the reported value.⁵ In addition, **25** was converted into known ester (-)-(*1R,2S*) as outlined in Scheme 1.

Scheme 1. Determination of the absolute stereochemistry of (*1R,2S*)-2-(2-oxo-2-phenylethyl)-cyclopentanecarboxylic acid methyl ester from **25**.

⁵ a) Enders, D.; Scherer, H. J.; Raabe, G. *Angew. Chem., Int. Ed. Engl.* **1991**, 30, 1664. b) Weinstein, R.; Lerner, R. A.; Barbas, C. F., III; Shabat, D. *J. Am. Chem. Soc.* **2005**, 127, ASAP.

13

Peak #	Ret. Time	Area %
1	17.432	1.955
2	19.419	0.060
3	22.603	2.992
4	35.501	94.994

Detector A Ch1 220nm		
Peak #	Ret. Time	Area %
1	17.182	2.017
2	19.198	0.427
3	20.469	4.043
4	24.409	93.512

Detector A Ch1 220nm		
Peak #	Ret. Time	Area %
1	9.847	3.291
2	10.443	1.611
3	13.374	90.708
4	14.337	4.389

Detector A Ch1 220nm		
Peak #	Ret. Time	Area %
1	17.200	4.394
2	20.604	2.690
3	28.943	92.916

Detector A Ch1 220nm		
Peak #	Ret. Time	Area %
1	23.698	4.187
2	24.666	4.120
3	30.025	91.693

Detector A Ch1 220nm		
Peak #	Ret. Time	Area %
1	10.307	0.085
2	10.922	1.102
3	12.350	91.516
4	13.839	7.298

Detector A Ch1 220nm		
Peak #	Ret. Time	Area %
1	9.669	5.682
2	10.307	0.798
3	11.772	80.643
4	14.436	12.878