Catalytic Enantioselective Alkylations of Tetrasubstituted Olefins. Synthesis of All-Carbon Quaternary Stereogenic Centers through Cu-Catalyzed Asymmetric Conjugate Additions of Alkylzinc Reagents to Enones

Alexander W. Hird and Amir H. Hoveyda*

Department of Chemistry, Merkert Chemistry Center, Boston College, Chestnut Hill, Massachusetts 02467

E-mail: amir.hoveyda@bc.edu

SUPPORTING INFORMATION

CONTENTS

General Information.....	Page S2
Preparation of substrates and reagents.....	Page S2 to S7
1	Page S2
2f	Page S3
8	Page S5
13	Page S2
17	Page S5
General Procedure for ECA.....	Page S7
Characterization Data and GLC traces	Pages S7 to S20
3	Page S7
6	Page S9
7	Page S10
9	Page S11
10	Page S12
11	Page S13
12	Page S14
14	Page S15
15	Page S16
16	Page S17
18	Page S18
19	Page S8
20a	Page S13
20b	Page S14
21	Page S20
22	Page S21
stereoproof	Page S17
Further functionalizations	Page S19 to S21
Ligand Screening Details.....	Pages S22 to S25
¹ H NMR spectra for all compounds.....	Pages S26 to S49

General. Infrared (IR) spectra were recorded on a Nicolet 210 spectrophotometer, $\nu_{\max}$ in cm^{-1} . Bands are characterized as broad (br), strong (s), medium (m) or weak (w). ^1H NMR spectra were recorded on a Varian Unity INOVA 400 (400 MHz) spectrometer. Chemical shifts are reported in ppm from tetramethylsilane with the solvent resonance resulting from incomplete deuteration as the internal standard (CHCl_3 : 7.26, C_6D_6 : 7.16).¹ Data are reported as follows: chemical shift, integration, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, br = broad, m = multiplet), coupling constants, and assignment. Many of the conjugate addition products were obtained as a mixture of keto diastereomers and enol tautomer; the ^1H NMR data is for the mixture, all peaks reported unless otherwise stated. In cases where it was difficult to discern whether a peak was due to a particular tautomer, or peaks overlapped, the integration value is given, where 1.0H corresponds to one proton from the major diastereomer. ^{13}C NMR spectra were recorded on a Varian Unity INOVA 400 (100 MHz) spectrometer with complete proton decoupling. Chemical shifts are reported in ppm from tetramethylsilane with the solvent resonance as the internal standard ($^{13}\text{CDCl}_3$: 77.16). Enantiomer ratios were determined by chiral GLC analysis: Alltech Associates Chiraldex GTA column (CDGTA, 30 m $\times$ 0.25 mm) or Supelco Betadex 120 column (β -dex, 30 m $\times$ 0.25 mm) in comparison with authentic racemic materials. Elemental analyses were performed by Robertson Microlit Laboratories (Madison, NJ). High resolution mass spectrometry was performed either by Mr. Chris Kowalczyk (Micromass LCT ESI-MS – positive mode at the mass spectrometry facility at Boston College, or by Dr Steven Mullen at the Mass Spectrometry Laboratory, University of Illinois at Urbana-Champaign. Optical rotation values were recorded on a Rudolph Research Analytical Autopol IV polarimeter.

Substrates: **1** was synthesized according to Paquette's procedure.² **13** was synthesized according to Taber's procedure.³

Reagents: Boc- and Fmoc- protected amino acids, EDC•HCl, HOBt•H₂O and amines for ligand synthesis were purchased from Advanced ChemTech, Nova Biochem or Aldrich. Piperidine, anthranilic acids, salicylic acids were purchased from Aldrich or Acros. Certain non-commercially available anthranilic acids were synthesized by known methods. $(\text{CuOTf})_2 \cdot \text{C}_6\text{H}_6$ was prepared according to a known method.⁴ All other Cu-salts were purchased from commercial sources and used without further purification. "Undistilled" toluene was purchased from Mallinkrodt Baker Inc and used as received (Assay 0.01% H₂O). "Dry" toluene was purified by a modified Advanced ChemTech purification system by passage through Cu and alumina columns.

(1) Gottlieb, H. E.; Kotlyar, V.; Nudelman, A. *J. Org. Chem.* **1997**, *62*, 7512-7515.

(2) Belmont, D. T.; Paquette, L. A. *J. Org. Chem.* **1985**, *50*, 4102-4107. Note: ^1H NMR data is incorrect. For correct ^1H NMR, see: Orban, J.; Turner, J. V. *Tetrahedron Lett.* **1983**, *24*, 2697-2700.

(3) Taber, D. F.; Amedio Jr., J. C.; Sherrill, R. G. *J. Org. Chem.* **1986**, *51*, 3382-3384.

(4) (a) Salomon, R. G.; Kochi, J. K. *J. Am. Chem. Soc.* **1973**, *95*, 1889-1897. (b) Salomon, R. G.; Kochi, J. K. *J. Am. Chem. Soc.* **1973**, *95*, 3300-3310.

CH₂Cl₂: purified through two alumina columns under a positive pressure of dry argon by a modified Advanced ChemTech purification system.

Et₂O: purified through two alumina columns under a positive pressure of dry argon by a modified Advanced ChemTech purification system.

THF: distilled under N₂ from sodium benzophenone ketyl.

DMSO: Aldrich

Acetic anhydride: Acros

***tert*-Butyl acetate:** distilled

Cyclopent-2-en-1-one: distilled

Chlorodiethylphosphate: Acros

Dibutylzinc (1M solution in heptane): Fluka

Diethylzinc: Aldrich

Diisopropylamine: distilled from NaOH

Diisopropylzinc (1M solution in toluene): Aldrich

Dimethylzinc: Strem

Ethylcyanoformate: distilled

HCl (4M in dioxane): Aldrich

Methyl lithium: Aldrich

Phosphorus Oxychloride: distilled under reduced pressure

Pyridine: distilled over CaH₂

Sodium Borohydride: Strem

Sodium hydride (60% in mineral oil): Strem

Sodium hydride (95%): Aldrich

Triethylborane: Strem

Trifluoromethanesulfonic anhydride: distilled over P₂O₅

Borane (dimethylsulfide complex): purified by vacuum transfer

Characterization Data

(-)-*N*-Methylantranilic-L-Leu-L-Trp-NMe₂. Boc-L-Trp-OH (500 mg, 1.64 mmol), dimethylamine hydrochloride (268 mg, 3.29 mmol), HOBt•H₂O (251 mg, 1.64 mmol) and EDC•HCl (315 mg, 1.64 mmol) were weighed in to a 100 mL flask. CH₂Cl₂ (50 mL) was added, followed by Et₃N (461 μL, 3.29 mmol). The solution was stirred at 23 °C for 24 h, then 50 mL of a 10% aqueous solution of citric acid was added. The layers were

separated. The lower organic layer was washed sequentially with 50 mL each of 10% aqueous citric acid, saturated aqueous NaHCO₃ and saturated NaCl. The organic layer was dried over Na₂SO₄ and concentrated in vacuo to yield a white solid which was identified only by ¹H NMR.

The white solid was cooled to 0 °C and HCl (4.90 mL, 19.7 mmol, 4M solution in dioxane) was added. The ice bath was removed and the solution was warmed to 23 °C, with swirling to ensure dissolution of all solids. After 30 min, the reaction was complete by TLC analysis (EtOAc, R_f starting material = 0.39, R_f product = 0). The solution was concentrated in vacuo and the solid residue was triturated with 10 mL Et₂O. The HCl salt was used directly in the next step. EDC•HCl (315 mg, 1.64 mmol), HOBt•H₂O (251 mg, 1.64 mmol) and Boc-L-Leu-OH (410 mg, 1.64 mmol), were added to a solution of the HCl salt and Et₃N (356 µL, 2.47 mmol) in 50 mL CH₂Cl₂. The solution was stirred rapidly to ensure dissolution of all solids, then for a further 16 h at 23 °C, at which point 50 mL of a 10% aqueous solution of citric acid was added. The layers were separated. The lower organic layer was washed sequentially with 50 mL each of 10% aqueous citric acid, saturated aqueous NaHCO₃ and saturated aqueous NaCl. The organic layer was dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by silica gel chromatography (gradient, 1:1 hexanes : EtOAc → EtOAc, R_f product = 0.39 in EtOAc) to yield an off-white solid (621 mg, 78%). The white solid was cooled to 0 °C and HCl (3.80 mL, 15.4 mmol, 4M solution in dioxane) was added. The ice bath was removed and the solution warmed to 23 °C, with swirling to ensure dissolution of all solids. After 30 min, the reaction was complete by TLC analysis (EtOAc, R_f starting material = 0.39, R_f product = 0). The solution was concentrated in vacuo and the solid residue was triturated with 10 mL Et₂O. The HCl salt was used directly in the next step. EDC•HCl (246 mg, 1.28 mmol), HOBt•H₂O (196 mg, 1.28 mmol) and *N*-methylantranilic acid (194 mg, 1.28 mmol) were added to a solution of the HCl salt and Et₃N (270 µL, 1.92 mmol) in 50 mL CH₂Cl₂. The solution was stirred rapidly to ensure dissolution of all solids, then for a further 16 h at 23 °C at which point 50 mL of a 10% aqueous solution of citric acid was added. The layers were separated. The lower organic layer was washed sequentially with 50 mL each of 10% aqueous citric acid, saturated aqueous NaHCO₃ and saturated NaCl. The organic layer was dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by silica gel chromatography (gradient, 1:1 → 1:2 hexanes : EtOAc, R_f product = 0.07 in 1:1 hexanes : EtOAc) to yield a white solid (421 mg, 69%, or 54% over 5 steps). mp 118 °C. IR (solid film): 3304 (br), 2956 (w), 2932 (w), 2969 (w), 2817 (w), 1626 (s), 1576 (w), 1520 (m); ¹H NMR (400 MHz, CDCl₃), *note*: chemical shifts are dependent upon concentration: δ 8.16 (1H, s), 7.58 (1H, d, J = 7.7 Hz), 7.48 (1H, br s), 7.36-7.24 (3H, m), 7.17-7.04 (3H, m), 6.99 (1H, d, J = 2.2 Hz), 6.67 (1H, d, J = 8.3 Hz), 6.57 (1H, dd, J = 7.5, 7.5 Hz), 6.46 (1H, d, J = 8.3 Hz), 5.20 (1H, ddd, J = 7.7, 7.2, 6.7 Hz), 4.67 (1H, ddd, J = 8.7, 8.7, 4.9 Hz), 3.16 (1H, A of ABX, J_{AB} = 14.4, J_{AX} = 7.4 Hz), 3.14 (1H, B of ABX, J_{AB} = 14.4, J_{BX} = 6.3 Hz), 2.85 (3H, s), 2.82 (3H, s), 2.65 (3H, s), 1.80-1.50 (3H, m), 0.93 (3H, d, J = 5.8 Hz), 0.92 (3H, d, J = 6.0 Hz); ¹³C NMR (100 MHz, CDCl₃): δ 171.6, 171.3, 169.5, 150.6, 136.0, 133.1, 127.6, 127.5, 123.2, 122.0, 119.5, 118.5, 114.6, 114.5, 111.1, 111.1, 110.2, 51.6, 49.8, 41.6, 36.9, 35.7, 29.7, 29.1, 24.8, 23.0, 22.0; HRMS Calcd for C₂₇H₃₅N₅O₃Na (M+Na⁺): 500.2638, Found: 500.2644; [α]_D²⁰ = -59.6 ° (c = 1.24, CHCl₃).

tert-Butyl 2-methyl-6-oxocyclohex-1-enecarboxylate. Prepared in a similar manner to the procedure used by Paquette to synthesize **1**.² To a stirred solution of BuLi (42.0 mL of a 1.62 M solution in hexanes, 68.4 mmol) in THF (95 mL) was added diisopropylamine (9.59 mL, 68.4 mmol) dropwise at -20 °C over 20 min. The pale yellow solution of LDA was cooled to -78 °C. After 10 min, *t*-BuOAc (9.20 mL, 68.4 mmol) was added dropwise over 20 min. After 45 min at -78 °C, a solution of 6-methyl-3,4-dihydropyran-2-one (**1**) in THF (10 mL) was added dropwise. The solution was stirred at -78 °C for 4 h, then allowed to warm slowly to room temperature over 14 h. The reaction mixture was poured in to a mixture of ice and pH 7 buffer. The layers were separated and the aqueous layer was washed with Et₂O (3 × 100 mL). The combined organic extracts were washed with a saturated aqueous solution of NaCl (100 mL), dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by silica gel chromatography (5:1 hexanes : EtOAc; *R_f* = 0.18) to give the product as a pale yellow oil (2.33 g, 11.1 mmol, 21%). This oil was also purified by Kugelrohr distillation prior to use (100 °C, ~0.25 mm Hg) to yield a colorless oil. IR (neat): 2979 (m), 2932 (m), 2872 (w), 1725 (s), 1672 (s), 1634 (m), 1368 (s), 1258 (s), 1169 (s), 1078 (s); ¹H NMR (400 MHz, CDCl₃), δ 2.39 (2H, dd, *J* = 7.0, 6.5 Hz, CH₂), 2.35 (2H, dd, *J* = 6.1, 6.1 Hz, CH₂), 2.03-1.97 (2H, m, CH₂CH₂CH₂), 1.98 (3H, s, CH₃), 1.53 (9H, s, C(CH₃)₃); ¹³C NMR (100 MHz, CDCl₃): δ 195.2, 166.4, 158.2, 134.5, 82.1, 37.1, 31.6, 28.3, 21.9, 21.9; Anal Calcd for C₁₂H₁₈O₃: C, 68.54; H, 8.63; Found C, 68.33; H, 8.66; HRMS Calcd for C₁₂H₁₈O₃ (M⁺): 210.1256, Found: 210.1259.

Ethyl 2-butyl-5-oxocyclopent-1-enecarboxylate. To a stirred suspension of CuI (1.36 g, 7.16 mmol) in Et₂O (50 mL) was added BuLi (8.70 mL of a 1.64 M in hexanes, 14.3 mmol) at 0 °C. The solution was stirred for 30 min and then cooled to -78 °C. 2-Cyclopentenone (500 μL, 5.97 mmol) was added and the solution was allowed to warm slowly to 23 °C. Consumption of starting material was confirmed by TLC and the solution was recooled to -78 °C. Ethylcyanoformate (649 μL, 6.57 mmol) was added and the reaction mixture was slowly warmed to 0 °C and stirred for 16 h. The reaction was quenched by the addition of a saturated aqueous solution of NH₄Cl (30 mL). After stirring for 30 min, the layers were separated and the aqueous layer was washed with Et₂O (3 × 40 mL). The combined organic layers were washed with saturated aqueous NaCl (50 mL), dried over MgSO₄ and concentrated in vacuo. The residue was

purified by silica gel chromatography (gradient, 10:1 → 5:1 hexanes : EtOAc, product R_f = 0.26 in 3:1, I_2 stain) to yield **II** as a pale yellow oil (920 mg, 4.33 mmol, 73%).

PhSe(O)Cl was prepared according to the method of Reich⁵ and used according to Smith's procedure.⁶ To a suspension of NaH (41.0 mg, 1.63 mmol) in THF (25 mL) at 23 °C was added **II** (288 mg, 1.35 mmol in 5 mL THF). After 1 h, the yellow solution was allowed to settle and cannulated in to a N_2 -filled 100 mL flask. The solution was cooled to -78 °C and PhSe(O)Cl (338 mg, 1.63 mmol in 5 mL THF) was added dropwise over 2 min. After 45 min at -78 °C, TLC analysis indicated complete consumption of starting material (R_f product = 0.17 in 3:1 hexanes : EtOAc). The solution was allowed to warm to room temperature and added to Et₂O (100 mL) and 1 M aqueous HCl (30 mL). The layers were separated and the aqueous layer was washed with Et₂O (2 × 30 mL). The combined organic layers were dried over MgSO₄ and concentrated in vacuo. The yellow residue was purified by silica gel chromatography (gradient, 6:1 → 3:1 hexanes : EtOAc) to yield **17** as a pale yellow oil (195 mg, 0.927 mmol, 68%). IR (neat): 2960 (s), 2932 (s), 2873 (m), 1745 (s), 1711 (s), 1623 (m); ¹H NMR (400 MHz, CDCl₃), δ 4.31 (2H, q, J = 7.1 Hz, OCH₂CH₃), 2.74 (2H, m, C=CCH₂CH₂CH₂CH₃), 2.68-2.64 (2H, m, aliphatic CH₂), 2.49-2.45 (2H, m, aliphatic CH₂), 1.62-1.52 (2H, m, aliphatic CH₂), 1.40 (2H, apparent sextet, J = 7.3 Hz, CH₂CH₂CH₃), 1.34 (3H, t, J = 7.1 Hz, OCH₂CH₃), 0.94 (3H, t, J = 7.3 Hz, CH₂CH₂CH₃); ¹³C NMR (100 MHz, CDCl₃): δ 203.9, 187.5, 163.6, 132.9, 60.9, 35.1, 32.5, 30.4, 29.9, 22.9, 14.4, 13.9; Anal Calcd for C₁₂H₁₈O₃: C, 68.54; H, 8.63; Found C, 68.29; H, 8.43; HRMS Calcd for C₁₂H₁₈O₃ (M⁺): 210.1256, Found: 210.1260.

Bis(4-acetoxybutyl)zinc.⁷ **Caution:** Dialkylzinc reagents are pyrophoric! 3-buten-1-ol (3.00 g, 41.6 mmol), Et₃N (8.76 mL, 62.4 mmol), and DMAP (254 mg, 2.08 mmol) were dissolved in dry CH₂Cl₂ (83 mL). Ac₂O (5.90 mL, 62.4 mmol) was added and the reaction was stirred for 3 h at 23 °C. After this time, HCl (120 mL, 1 M aqueous) was added and the layers were separated. Saturated aqueous NaHCO₃ (20 mL) was added and the mixture was stirred vigorously for 30 min to destroy excess Ac₂O. After this time, the layers were separated. The organic layer was dried over Na₂SO₄ and concentrated in vacuo. The residue was distilled under a nitrogen atmosphere to yield but-3-enyl acetate as a colorless oil (2.88 g, 25.27 mmol, 61%). Triethylborane (5.79 mL, 40.0 mmol) was added to a 25 mL pear-shaped flask and cooled to 0 °C. BH₃•SMe₂⁸ (2.40 mL, 20.0 mmol) was added via syringe. The reaction vessel was allowed to warm to 23 °C and stirred for an additional 1 h to produce HB(Et)₂• $\frac{1}{3}$ SMe₂. But-3-enyl acetate (2.88 g, 25.3 mmol) was placed in a 50 mL flask equipped with a 3-way vacuum adapter. The

(5) Reich, H. J.; Renga, J. M.; Reich, I. L. *J. Am. Chem. Soc.* **1975**, *97*, 5434-5447.

(6) Smith III, A. B.; Sestelo, J. P.; Dormer, P. G. *J. Am. Chem. Soc.* **1995**, *117*, 10755-10756. Note: the simpler procedure using PhSeCl and subsequent oxidation by H₂O₂ gave much lower yields with a related substrate (maximum of 20% yield was obtained when exactly 2 equiv of H₂O₂ were employed).

(7) For a general reference for the preparation and reactions of organozinc compounds, see: Knochel, P.; Singer, R. D. *Chem. Rev.* **1993**, *93*, 2117-2188.

(8) Borane dimethylsulfide complex was purified by vacuum transfer prior to use.

flask was placed under nitrogen and cooled to -10 °C. $\text{HB}(\text{Et})_2 \cdot \frac{1}{3} \text{SMe}_2$ (3.11 mL, 25.3 mmol) was added dropwise over 5 min. The reaction was stirred at 23 °C for 3 h at which time unreacted reagents and excess dimethylsulfide were removed under vacuum. The flask was then placed under nitrogen and cooled to 0 °C. Et_2Zn (5.29 mL, 50.5 mmol) was added dropwise. The boron-zinc exchange reaction was allowed to proceed at 23 °C for 14 h. Triethylborane and excess Et_2Zn were removed under vacuum (caution – pyrophoric). The residue was brought in to a glovebox and filtered through a syringe filter (Fisherbrand, PTFE, 0.45 μm) to yield a colorless solution (3.00 g, 20.3 mmol, 80%). ^1H NMR (400 MHz, C_6D_6) analysis showed the desired product was present, along with ca. 10% Et_2Zn : δ 4.06 (2H, t, J = 6.3 Hz, CH_2OAc), 1.68 (3H, s, CH_3), 1.56-1.49 (4H, m, aliphatic CH_2), 0.22 (2H, t, J = 7.5 Hz, CH_2Zn).

General Procedure:⁹ **(2S)-(+)-Methyl 2-ethyl-2-methyl-6-oxo-cyclohexane -3-carboxylate**. CuCN (0.3 mg, 3 μmol) and **2f** (1.4 mg, 3.0 μmol) were weighed into a 13 $\times$ 100 mm test tube. Toluene (1 mL) was added and the tube was sealed with a rubber septum and purged with nitrogen. The suspension was allowed to stir at 23 °C for 3 h and then cooled to 0 °C. Et_2Zn (24 μL , 0.23 mmol) was added

and the pale yellow solution was stirred at 0 °C for 5 min. A solution of **1** (24.8 mg, 0.148 mmol) in toluene (0.5 mL) was added via cannula. The septum was wrapped with Teflon tape and stirred for a further 11 h at 0 °C. The reaction solution was cooled to -78 °C, diluted with EtOAc (1 mL), and quenched with aqueous 1M HCl (1 mL). The upper organic layer was separated and the aqueous layer was washed with EtOAc (3 $\times$ 1 mL). The combined organics were concentrated in vacuo. The residue was purified by silica gel chromatography (6:1 hexanes: Et_2O) to yield **3** as a pale yellow oil (26.0 mg, 0.131 mmol, 89%). IR (neat): 2963 (s), 2879 (m), 1753 (m), 1732 (m), 1713 (s), 1636 (w), 1600 (w), 1462 (m), 1436 (m); ^1H NMR (400 MHz, CDCl_3), reported for a 2.8 : 1.9 : 1 ratio of two keto diastereomers and the enol tautomer: δ 3.77 (3H, s, $\text{OCH}_3_{\text{enol}}$), 3.69 (3H, s, $\text{OCH}_3_{\text{diast 2}}$), 3.68 (3H, s, $\text{OCH}_3_{\text{diast 1}}$), 3.25 (1H, br s, $\text{C}(\text{O})\text{CHC}(\text{O})_{\text{diast 2}}$), 3.21 (1H, dd, J = 1.2, 1.2 Hz, $\text{C}(\text{O})\text{CHC}(\text{O})_{\text{diast 1}}$), 2.77 (1H, ddd, J = 14.2, 10.0, 6.8 Hz, $\text{CHHC}(\text{O})_{\text{diast 1}}$), 2.70-2.62 (1H, m, $\text{CHHC}(\text{O})_{\text{diast 2}}$), 2.35-2.25 (2H, m, $\text{CHHC}(\text{O})_{\text{diast 1\&2}}$ and $\text{CH}_2\text{C}(\text{OH})_{\text{enol}}$), 2.07 (1H, ddd, J = 14.7, 10.5, 4.6 Hz, aliphatic CH), 1.98-1.75 (4.3H, m, aliphatic CH), 1.69-1.54 (2.5H, m, aliphatic CH), 1.47-1.24 (4.8H, m, aliphatic CH), 1.15 (3H, s, $\text{C}(\text{Et})\text{CH}_3_{\text{enol}}$), 1.02 (3H, s, $\text{C}(\text{Et})\text{CH}_3_{\text{diast 2}}$), 0.93 (3H, s, $\text{C}(\text{Et})\text{CH}_3_{\text{diast 1}}$), 0.85 (3H, t, J = 7.5 Hz, $\text{CH}_2\text{CH}_3_{\text{diast 2}}$), 0.84 (3H, t, J = 7.5 Hz, $\text{CH}_2\text{CH}_3_{\text{diast 1}}$), 0.70 (3H, t, J = 7.5 Hz, $\text{CH}_2\text{CH}_3_{\text{enol}}$); ^{13}C NMR (100 MHz, CDCl_3): δ 207.2, 207.2, 174.5, 174.4, 169.5, 169.4, 104.8, 67.4, 66.2, 52.0, 51.9, 51.2, 41.9, 41.6, 39.7, 39.3, 36.1, 35.4, 33.0, 32.7, 32.1, 30.8, 30.8, 27.1, 23.2, 22.0, 22.0, 21.8, 21.6, 17.6, 9.0, 7.7, 7.6; Anal Calcd for $\text{C}_{11}\text{H}_{18}\text{O}_3$: C, 66.64; H, 9.15; Found C, 66.43; H, 9.35; HRMS Calcd for $\text{C}_{11}\text{H}_{18}\text{O}_3\text{Na}$ ($\text{M}+\text{Na}^+$): 221.1154, Found: 221.1147;

(9) Note: CuCN is highly toxic and contact should be avoided. Acidic workup of CuCN containing solutions will result in evolution of HCN . For small scale reactions, the quantity of HCN is negligible. However, for larger scale reactions, aqueous NH_4Cl should be used for workup instead of 1N HCl . This has no effect upon the enantioselectivity or yield of the reaction.

$[\alpha]_{\text{D}}^{20} +28.8^{\circ}$ ($c = 0.883$, CHCl_3) for a sample of 84% ee. Optical purity was determined by chiral GLC analysis (β -dex, 120 $^{\circ}\text{C}$, 15 psi).

Peak #	RetTime [min]	Type	Width [min]	Area [pA*s]	Height [pA]	Area %
1	44.961	MM	0.3192	6.12454	3.19751e-1	49.72508
2	45.931	MM	0.3354	6.19227	3.07671e-1	50.27492

authentic racemic

Peak #	RetTime [min]	Type	Width [min]	Area [pA*s]	Height [pA]	Area %
1	45.000	MM	0.3346	26.17152	1.30344	8.91025
2	46.000	MM	0.3328	267.55228	13.39878	91.08975

82% ee

(S)-(-)-3-Ethyl-3-methylcyclohexanone.¹⁰ Using a microscale kit: β -Keto-ester **3** (81 mg, 0.41 mmol) was dissolved in DMSO (272 μL) and H_2O (22 μL , 1.2 mmol). NaCl (26 mg, 0.45 mmol) was added and the vial was topped with a condenser and heated to 150 $^{\circ}\text{C}$ by an oil bath. After 3 h, the reaction was allowed to cool and diluted with H_2O (2 mL). The aqueous mixture was washed with pentane (4×5 mL). The combined organics were dried over Na_2SO_4 and concentrated in vacuo (*care*: volatile product) to yield a pale yellow oil (53.0 mg, 0.378 mmol, 93%). IR (neat): 2962 (s), 2938 (s), 2879 (m), 2852 (w), 1711 (s), 1461 (m); ^1H NMR (400 MHz, CDCl_3): δ 2.30-2.24 (2H, m, aliphatic $\text{C}(=\text{O})\text{CH}_2\text{CH}_2$), 2.17 (1H, AB, $J = 13.5$ Hz, $\text{C}(=\text{O})\text{CHHC}(\text{CH}_3)\text{Et}$), 2.09 (1H, AB, $J = 13.5$ Hz, $\text{C}(=\text{O})\text{CHHC}(\text{CH}_3)\text{Et}$), 1.89-1.82 (2H, m, aliphatic CH_2), 1.66-1.48 (2H, m, aliphatic CH_2), 1.34-1.24 (2H, m, aliphatic CH_2), 0.89 (3H, s, $\text{C}(\text{Et})\text{CH}_3$), 0.84 (3H, dd, $J = 7.5, 7.5$ Hz, CH_2CH_3); ^{13}C NMR (100 MHz, CDCl_3): δ 212.7, 53.5, 41.2, 38.8, 35.5, 34.1, 24.6, 22.3, 7.9; HRMS Calcd for $\text{C}_9\text{H}_{16}\text{ONa}$ ($\text{M}+\text{Na}^+$): 163.1099, Found: 163.1098; $[\alpha]_{\text{D}}^{20} -4.3^{\circ}$ ($c = 1.6$, CHCl_3) for a sample of 81% ee.¹¹

Optical purity was determined by chiral GLC analysis (β -dex, 100 $^{\circ}\text{C}$, 15 psi): for GC retention times, see trace for dealkoxycarbonylated product of **9** (major enantiomer is same in both cases).

(10) For a review of dealkoxycarbonylations of 1,3-keto methyl esters, see: Krapcho, A. P. *Synthesis*, **1982**, 893-914.

(11) According to Alexakis' inferred stereochemistry, this corresponds to the (*S*) enantiomer: d' Augustin, M.; Palais, L.; Alexakis, A. *Angew. Chem. Int. Ed.* **2005**, *44*, 1376-1378. This is also consistent with our stereoproof: see ref (12).

(+)-Methyl 2-butyl-2-methyl-6-oxocyclohexanecarboxylate. IR (neat): 2955 (s), 2933 (s), 2872 (m), 2861 (m), 1753 (m), 1731 (s), 1709 (s), 1637 (w), 1599 (w), 1460 (m), 1435 (m); ^1H NMR (400 MHz, CDCl_3), reported for a 3.4 : 1.9 : 1 ratio of two keto diastereomers and enol tautomer: δ 3.76 (3H, s, $\text{OCH}_{3\text{enol}}$), 3.69 (3H, s, $\text{OCH}_{3\text{diast2}}$), 3.68 (3H, s, $\text{OCH}_{3\text{diast1}}$), 3.25 (1H, s, $\text{C(=O)CHCO}_2\text{CH}_{3\text{diast2}}$), 3.20 (1H, s, $\text{C(=O)CHCO}_2\text{CH}_{3\text{diast1}}$), 2.76 (1H, ddd, $J = 14.3, 9.8, 6.8$ Hz, $\text{CHHC(=O)}_{\text{diast1}}$), 2.70-2.61 (0.7H, m, aliphatic CH), 2.34-2.23 (2.5H, m, aliphatic CH), 2.08 (1H, ddd, $J = 14.3, 10.3, 4.5$ Hz, $\text{CHHC(=O)}_{\text{diast1}}$), 1.96-1.75 (4.7H, m, aliphatic CH), 1.69-1.49 (3.8H, m, aliphatic CH), 1.41 (1H, ddd, $J = 13.5, 5.0, 5.0$ Hz, aliphatic $\text{CH}_{\text{diast1}}$), 1.35-1.15 (12.2H, m, aliphatic CH), 1.15 (3H, s, $\text{CCH}_{3\text{enol}}$), 1.03 (3H, s, $\text{CCH}_{3\text{diast2}}$), 0.94 (3H, s, $\text{CCH}_{3\text{diast1}}$), 0.91-0.84 (6.9H, m, aliphatic CH); ^{13}C NMR (100 MHz, CDCl_3): δ 207.2, 207.2, 174.3, 169.5, 169.4, 174.4, 105.1, 67.6, 66.5, 51.9, 51.8, 51.1, 41.7, 41.5, 40.5, 40.0, 39.7, 39.3, 38.1, 36.9, 35.1, 33.5, 32.7, 30.7, 27.5, 27.0, 25.5, 25.3, 23.9, 23.5, 23.5, 23.4, 22.5, 21.8, 21.6, 17.7, 14.3, 14.2, 14.2; Anal Calcd for $\text{C}_{13}\text{H}_{22}\text{O}_3$: C, 68.99; H, 9.80; Found C, 69.01; H, 9.84; HRMS Calcd for $\text{C}_{13}\text{H}_{22}\text{O}_3\text{Na}$ ($\text{M}+\text{Na}^+$): 249.1467, Found: 249.1464; $[\alpha]_{\text{D}}^{20} +28.5^\circ$ ($c = 1.05$, CHCl_3) for a sample of 89% ee.

Optical purity determined by chiral GLC analysis of dealkoxycarbonylated product (1 equiv NaCl , 3 equiv H_2O , 1.5 M in DMSO, 150°C , 3 h) in comparison with authentic racemic material. Conditions: β -dex, 120°C , 15 psi.

Peak #	RetTime [min]	Type	Width [min]	Area [pA*s]	Height [pA]	Area %
1	26.058	MM	0.1889	166.10118	14.65485	50.05075
2	26.730	MM	0.2025	165.76436	13.64158	49.94925

authentic racemic

Peak #	RetTime [min]	Type	Width [min]	Area [pA*s]	Height [pA]	Area %
1	25.927	MM	0.1649	77.84735	7.86863	4.09278
2	26.316	MM	0.2842	1824.21936	106.96761	95.90722

92% ee

(+)-Methyl 2-isopropyl-2-methyl-6-oxocyclohexanecarboxylate. IR (neat): 2955 (s), 2928 (s), 2856 (s), 1743 (s), 1471 (m), 1462 (m), 1361 (w), 1255 (s); ^1H NMR (400 MHz, CDCl_3), ~15:1 enol : keto; reported peaks for enol tautomer: δ 3.77 (3H, s, OCH_3), 2.38 (1H, qq, $J = 6.9, 6.9$ Hz, $\text{CH}(\text{CH}_3)_2$), 2.27-2.20 (2H, m, aliphatic CH), 1.75-1.52 (2H, m, aliphatic CH), 1.43 (1H, ddd, $J = 13.0, 13.0, 3.4$ Hz, aliphatic CH), 1.33 (1H, ddd, $J = 13.2, 3.9, 3.9$ Hz, aliphatic CH), 1.21 (3H, s, $\text{C}(\text{i-Pr})\text{CH}_3$), 0.85 (3H, d, $J = 6.9$ Hz, $\text{CH}(\text{CH}_3)\text{CH}_3$), 0.66 (3H, d, $J = 6.9$ Hz, $\text{CH}(\text{CH}_3)\text{CH}_3$); ^{13}C NMR (100 MHz, CDCl_3), reported peaks for two keto diastereomers and enol tautomer: δ 208.0, 207.6, 174.5, 174.1, 169.6, 169.5, 105.6, 66.8, 64.6, 52.1, 51.8, 51.2, 44.8, 44.1, 40.0, 38.3, 38.1, 34.9, 33.9, 33.7, 30.9, 30.6, 30.0, 29.9, 25.5, 22.0, 21.2, 19.7, 18.2, 17.6, 17.5, 17.3, 17.0, 16.8, 16.7, 16.5; HRMS Calcd for $\text{C}_{12}\text{H}_{20}\text{O}_3\text{Na}$ ($\text{M}+\text{Na}^+$): 235.1310, Found: 235.1304; $[\alpha]_D^{20} +53.6^\circ$ ($c = 1.17$, CHCl_3) for a sample of 86% ee.

Optical purity determined by chiral GLC analysis of dealkoxycarbonylated product (1 equiv NaCl , 3 equiv H_2O , 1.5 M in DMSO, 150°C , 3 h) in comparison with authentic racemic material. Conditions: CDGTA, 105°C , 15 psi.

Peak #	RetTime [min]	Type	Width [min]	Area counts*s	Height [counts]	Area %
1	16.850	MM	0.2628	1.15428e4	731.97931	49.64193
2	19.449	MM	0.3180	1.17093e4	613.78320	50.35807

authentic racemic

Peak #	RetTime [min]	Type	Width [min]	Area counts*s	Height [counts]	Area %
1	16.844	MM	0.2449	9368.16309	637.52991	7.08114
2	19.167	MM	0.4594	1.22929e5	4459.52979	92.91886

86% ee

(+)-(S)-tert-Butyl 2-ethyl-2-methyl-6-oxocyclohexanecarboxylate. IR (neat): 2969 (s), 2937 (s), 2877 (s), 1743 (s), 1708 (s), 1628 (m), 1599 (m), 1460 (m), 1392 (m); ^1H NMR (400 MHz, CDCl_3), reported for 1.5 : 1 ratio of two keto diastereomers: δ 3.11 (1H, s, $\text{C}(=\text{O})\text{CHCO}_2t\text{-Bu}_{\text{diast 1}}$), 3.08 (1H, s, $\text{C}(=\text{O})\text{CHCO}_2t\text{-Bu}_{\text{diast 2}}$), 2.81 (1H, ddd, $J = 14.2, 11.1, 7.1$ Hz, $\text{CHHC}(=\text{O})_{\text{diast 2}}$), 2.67 (1H, ddd, $J = 16.9, 8.4, 6.2$ Hz, $\text{CHHC}(=\text{O})_{\text{diast 1}}$), 2.40-2.22 (2.7H, m, aliphatic CH), 2.09 (1H, ddd, $J = 13.7, 11.1, 4.5$ Hz, aliphatic $\text{CH}_{\text{diast 2}}$), 2.00-1.75 (4.7H, m, aliphatic CH), 1.70-1.25 (9.1H, m, aliphatic CH), 1.46 (9H, s, $\text{C}(\text{CH}_3)_3_{\text{diast 1}}$), 1.45 (9H, s, $\text{C}(\text{CH}_3)_3_{\text{diast 2}}$), 1.03 (3H, s, $\text{C}(\text{Et})\text{CH}_3_{\text{diast 1}}$), 0.89 (3H, s, $\text{C}(\text{Et})\text{CH}_3_{\text{diast 2}}$), 0.87 (3H, dd, $J = 7.5, 7.5$ Hz, $\text{CH}_2\text{CH}_3_{\text{diast 2}}$), 0.85 (3H, dd, $J = 7.5, 7.4$ Hz, $\text{CH}_2\text{CH}_3_{\text{diast 1}}$); ^{13}C NMR (100 MHz, CDCl_3), reported for mix of two keto diastereomers and enol: δ 208.1, 207.9, 173.9, 173.6, 168.2, 168.2, 105.6, 81.7, 81.5, 81.5, 68.0, 67.2, 41.7, 41.4, 39.6, 39.1, 36.4, 35.5, 33.1, 32.6, 32.4, 32.2, 31.5, 30.8, 28.6, 28.2, 28.1, 27.3, 22.9, 22.3, 21.8, 21.5, 17.7, 9.1, 7.7, 7.7; HRMS Calcd for $\text{C}_{14}\text{H}_{24}\text{O}_3\text{Na}$ ($\text{M}+\text{Na}^+$): 263.1623, Found: 263.1621; $[\alpha]_{\text{D}}^{20} +32.5^\circ$ ($c = 1.08$, CHCl_3) for a sample of 84% ee.

Optical purity determined by chiral GLC analysis of dealkoxycarbonylated product (5:1 CH_2Cl_2 : TFA, 24 h) in comparison with authentic racemic material. Conditions: β -dex, 100 $^\circ\text{C}$, 15 psi.

Peak #	RetTime [min]	Type	Width [min]	Area [pA*s]	Height [pA]	Area %
1	26.058	MF	0.3346	991.75940	49.39467	50.14263
2	27.008	FM	0.3981	986.11725	41.28365	49.85737

authentic racemic

Peak #	RetTime [min]	Type	Width [min]	Area [pA*s]	Height [pA]	Area %
1	26.667	MM	0.2103	11.49217	9.10909e-1	6.38476
2	27.452	MM	0.2487	168.50165	11.29000	93.61524

87% ee

(+)-tert-Butyl 2-butyl-2-methyl-6-oxocyclohexanecarboxylate. IR (neat): 2958 (s), 2933 (s), 2873 (m), 2860 (m), 1745 (m), 1708 (s), 1627 (m), 1602 (m), 1458 (m), 1393 (m), 1369 (m), 1330 (m), 1249 (m); ^1H NMR (400 MHz, CDCl_3), reported for a 1.25 : 1 mix of keto diastereomers: δ 3.11 (1H, s, $\text{C(=O)CHCO}_2t\text{-Bu}_{\text{diast 2}}$), 3.08 (1H, s, $\text{C(=O)CHCO}_2t\text{-Bu}_{\text{diast 1}}$), 2.80 (1H, ddd, $J = 14.2, 10.8, 7.0$ Hz, $\text{CHHC(=O)}_{\text{diast 1}}$), 2.71-2.62 (1H, m, $\text{CHHC(=O)}_{\text{diast 2}}$), 2.42-2.20 (2.4H, m, aliphatic **CH**), 2.10 (1H, ddd, $J = 13.8, 11.3, 4.6$ Hz, aliphatic **CH**), 2.04-1.74 (5H, m, aliphatic **CH**), 1.70-1.50 (4.6H, m, aliphatic **CH**), 1.45 (9H, s, $\text{C(CH}_3)_3_{\text{diast 2}}$), 1.44 (9H, s, $\text{C(CH}_3)_3_{\text{diast 1}}$), 1.40-1.12 (13.7H, m, aliphatic **CH**), 1.03 (3H, s, $\text{C(Bu)CH}_3_{\text{diast 2}}$), 0.92-0.86 (5.4H, m, aliphatic **CH**), 0.91 (3H, s, $\text{C(Bu)CH}_3_{\text{diast 1}}$); ^{13}C NMR (100 MHz, CDCl_3), reported for mix of two keto diastereomers and enol: δ 208.1, 208.0, 173.7, 173.6, 168.2, 168.2, 105.8, 81.7, 81.5, 81.5, 68.4, 67.4, 41.6, 41.3, 40.9, 39.8, 39.6, 39.1, 38.9, 37.2, 35.2, 33.2, 32.6, 30.8, 28.6, 28.2, 28.2, 27.7, 27.1, 25.5, 25.4, 23.7, 23.7, 23.5, 23.4, 22.9, 21.8, 21.6, 17.8, 14.4, 14.3, 14.2; Anal Calcd for $\text{C}_{16}\text{H}_{28}\text{O}_3$: C, 71.60; H, 10.52; Found C, 71.40; H, 10.70; HRMS Calcd for $\text{C}_{16}\text{H}_{28}\text{O}_3\text{Na}$ ($\text{M}+\text{Na}^+$): 291.1936, Found: 291.1937; $[\alpha]_{\text{D}}^{20} +29.2^\circ$ ($c = 1.05$, CHCl_3) for a sample of 95% ee.

Optical purity determined by chiral GLC analysis of dealkoxycarbonylated product (5:1 CH_2Cl_2 :

Peak #	RetTime [min]	Type	Width [min]	Area [pA*s]	Height [pA]	Area %
1	26.058	MM	0.1889	166.10118	14.65485	50.05075
2	26.730	MM	0.2025	165.76436	13.64158	49.94925

Peak #	RetTime [min]	Type	Width [min]	Area [pA*s]	Height [pA]	Area %
1	25.978	MM	0.1799	20.73734	1.92091	2.53965
2	26.475	MM	0.2240	795.80585	59.19953	97.46035

TFA, 24 h) in comparison with authentic racemic material. Conditions: β -dex, 100 $^\circ\text{C}$, 15 psi.

authentic racemic

95% ee

(+)-tert-Butyl 2-hydroxy-6-isopropyl-6-methylcyclohex-1-enecarboxylate. IR (neat): 2964 (s, br), 2877 (s), 1745 (m), 1709 (s), 1628 (s), 1599 (s), 1459 (m), 1391 (s), 1368 (s), 1302 (s), 1243 (s), 1217 (m), 1158 (s); ^1H NMR (400 MHz, CDCl_3), reported for enol (>15:1 enol : keto): δ 2.43 (1H, qq, $J = 7.0, 7.0$ Hz, $\text{CH}(\text{CH}_3)_2$), 2.23-2.17 (2H, m, $\text{C}=\text{C}(\text{OH})\text{CH}_2$), 1.70-1.54 (2H, m, aliphatic CH), 1.51 (9H, s, $\text{C}(\text{CH}_3)_3$), 1.44-1.28 (2H, m, aliphatic CH), 1.20 (3H, s, $\text{C}(i\text{-Pr})\text{CH}_3$), 0.83 (3H, d, $J = 7.0$ Hz, $\text{CH}(\text{CH}_3)(\text{CH}_3)$), 0.68 (3H, d, $J = 7.0$ Hz, $\text{CH}(\text{CH}_3)\text{CH}_3$); ^{13}C NMR (100 MHz, CDCl_3), reported for enol (>15:1 enol : keto): δ 173.7, 173.3, 106.6, 81.6, 38.3, 33.9, 30.9, 30.8, 28.6 (3), 25.4, 18.2, 17.6, 17.3; Anal Calcd for $\text{C}_{15}\text{H}_{26}\text{O}_3$: C, 70.83; H, 10.30; Found C, 70.59; H, 10.19; HRMS Calcd for $\text{C}_{15}\text{H}_{26}\text{O}_3\text{Na}$ ($\text{M}+\text{Na}^+$): 277.1780, Found: 277.1779; $[\alpha]_{\text{D}}^{20} +88.4^\circ$ ($c = 0.993$, CHCl_3) for a sample of 90% ee (ee determined after dealkoxycarbonylation – see below).

(-)-3-isopropyl-3-methylcyclohexanone. A sample of **11** (30.5 mg, 0.120 mmol) was dissolved in 5:1 CH_2Cl_2 : TFA (250:50 μL) and stirred under nitrogen for 20 h. The solution was diluted with 1 mL CH_2Cl_2 and saturated aqueous NaHCO_3 (1 mL) was added. After gas evolution had ceased, the layers were separated and the aqueous layer was washed with additional CH_2Cl_2 (2×1 mL). The combined organic layers were dried over Na_2SO_4 and concentrated in vacuo (*care*: volatile product) to yield the title compound as a pale yellow oil (16.4 mg, 0.106 mmol, 89%). IR (neat): 2961 (s), 2877 (m), 1711 (s); ^1H NMR (400 MHz, CDCl_3), δ 2.36-2.20 (3H, m, aliphatic CH), 2.10 (1H, ddd, $J = 13.3, 1.6, 1.6$ Hz, aliphatic CH), 1.98-1.75 (2H, m, aliphatic CH), 1.69-1.42 (3H, m, aliphatic CH), 0.86 (3H, d, $J = 6.9$ Hz, $\text{CH}(\text{CH}_3)\text{CH}_3$), 0.86 (3H, d, $J = 6.9$ Hz, $\text{CH}(\text{CH}_3)\text{CH}_3$), 0.81 (3H, s, $\text{C}(i\text{-Pr})\text{CH}_3$); ^{13}C NMR (100 MHz, CDCl_3): δ 213.1, 52.0, 41.3, 41.2, 36.4, 34.2, 22.1, 20.0, 17.1, 17.0; HRMS Calcd for $\text{C}_{10}\text{H}_{18}\text{ONa}$ ($\text{M}+\text{Na}^+$): 177.1255, Found: 177.1256; $[\alpha]_{\text{D}}^{20} -15.3^\circ$ ($c = 1.00$, CHCl_3) for a sample of 88% ee.

Optical purity determined by chiral GLC analysis in comparison with authentic racemic material. Conditions: CDGTA, 105 $^\circ\text{C}$, 15 psi.

Peak #	RetTime [min]	Type	Width [min]	Area counts*s	Height [counts]	Area %
1	16.850	MM	0.2628	1.15428e4	731.97931	49.64193
2	19.449	MM	0.3180	1.17093e4	613.78320	50.35807

authentic racemic

Peak #	RetTime [min]	Type	Width [min]	Area counts*s	Height [counts]	Area %
1	16.800	MM	0.2084	2083.76050	166.64851	5.16497
2	19.205	MM	0.3508	3.82603e4	1817.76514	94.83503

90% ee

(+)-tert-Butyl 2-(4-acetoxybutyl)-2-methyl-6-oxocyclohexanecarboxylate. IR (neat): 2947 (m), 2874 (w), 1741 (s), 1708 (s), 1460 (w), 1367 (m), 1242 (s), 1151 (m); ^1H NMR (400 MHz, CDCl_3), 1:1:0.4 mixture of two keto diastereomers and enol tautomer; NMR data reported only for keto diastereomers: δ 4.07-4.02 (2H, m, $\text{CH}_2\text{OAc}_{\text{diast 1,2 and enol}}$), 3.11 (1H, s, $\text{C(=O)CHCO}_2t\text{-Bu}$), 3.08 (1H, s, $\text{C(=O)CHCO}_2t\text{-Bu}$), 2.79 (1H, ddd,

$J = 14.3, 10.6, 7.1$ Hz, aliphatic CH), 2.66 (1H, ddd, $J = 14.3, 7.8, 6.2$ Hz, aliphatic CH), 2.34-2.20 (2H, m, aliphatic CH), 2.09 (1H, ddd, $J = 14.0, 11.0, 4.5$ Hz, aliphatic CH), 2.04 and 2.04 (3H, s, C(=O)CH_3), 2.04 (3H, s, C(=O)CH_3), 1.96-1.74 (4.4H, m, aliphatic CH), 1.66-1.52 (6.5H, m, aliphatic CH), 1.45 (9H, s, $\text{OC(CH}_3)_3$), 1.44 (9H, s, $\text{OC(CH}_3)_3$), 1.42-1.20 (8.6H, m, aliphatic CH), 1.04 (3H, s, CH_2CCH_3), 0.92 (3H, s, CH_2CCH_3); ^{13}C NMR (100 MHz, CDCl_3) δ 207.7, 207.6, 171.2 (2), 168.1, 168.0, 81.8, 81.7, 68.2, 67.2, 64.5, 64.4, 41.5, 41.3, 39.8, 39.6, 39.1, 38.7, 33.3, 32.6, 29.4, 29.3, 28.2 (3), 28.2 (3), 23.6, 22.7, 21.7, 21.5, 21.1 (2), 19.9, 19.8; Anal Calcd for $\text{C}_{18}\text{H}_{30}\text{O}_5$: C, 66.23; H, 9.26; Found C, 66.28; H, 9.47; HRMS Calcd for $\text{C}_{18}\text{H}_{30}\text{O}_5$ (M): 327.2171, Found: 327.2169; $[\alpha]_{\text{D}}^{20} +4.6^\circ$ ($c = 1.0$, CHCl_3) for a sample of 72% ee (for ee determination, see below).

(+)-4-(1-Methyl-3-oxocyclohexyl)butylacetate. A sample of **12** (9.6 mg, 0.029 mmol) was sealed in a vial and heated to 150 $^\circ\text{C}$ in an oil bath. After 3 h, the vial was cooled to room temperature and the contents were purified by silica gel chromatography (gradient, 3:1 $\rightarrow$ 1:1 pentanes : Et_2O) to yield the product as a pale yellow oil (5.6 mg, 0.024 mmol, 84%). IR (neat): 2937 (w), 1737 (s), 1712 (w), 1242 (s); ^1H NMR (400 MHz, CDCl_3): 4.05 (2H,

apparent t, $J = 6.7$ Hz, CH_2OAc), 2.30-2.24 (2H, m, $\text{C(=O)CH}_2\text{CH}_2$), 2.18 (1H, AB, $J_{\text{AB}} = 13.5$ Hz, $\text{C(=O)CHHC(CH}_3)$), 2.10 (1H, AB, $J_{\text{AB}} = 13.5$ Hz, $\text{C(=O)CHHC(CH}_3)$), 2.04 (3H, s, C(=O)CH_3), 1.90-1.81 (2H, m, aliphatic CH), 1.66-1.53 (4H, m, aliphatic CH), 1.37-1.22 (4H, m, aliphatic CH), 0.91 (3H, s, CH_2CCH_3); ^{13}C NMR (100 MHz, CDCl_3): δ 212.2, 171.3, 64.5, 53.8, 41.4, 41.1, 38.7, 36.0, 29.4, 25.1, 22.3, 21.1, 20.0; HRMS Calcd for $\text{C}_{13}\text{H}_{22}\text{O}_3$ (M): 227.1647, Found: 227.1649; $[\alpha]_{\text{D}}^{20} +3.0^\circ$ ($c = 1.00$, CHCl_3) for a sample of 72% ee. Optical purity determined by chiral GLC analysis in comparison with authentic racemic material. Conditions: CDGTA, 130 $^\circ\text{C}$, 15 psi.

Peak #	RetTime [min]	Type	Width [min]	Area [pA*s]	Height [pA]	Area %
1	81.920	MM	0.6770	347.93915	8.56600	49.99990
2	90.439	MM	0.8523	347.94052	6.80423	50.00010

authentic racemic

Peak #	RetTime [min]	Type	Width [min]	Area [pA*s]	Height [pA]	Area %
1	82.522	MM	0.5700	38.53152	1.12674	6.27480
2	90.189	MM	1.0104	575.53632	9.49331	93.72520

87% ee

(–)-Methyl 2-ethyl-2-methyl-5-oxocyclopentanecarboxylate. IR (neat): 2961 (s), 2877 (m), 1734 (w), 1712 (m), 1636 (s), 1598 (s), 1463 (m), 1439 (s), 1383 (s), 1351 (m); ^1H NMR (400 MHz, CDCl_3), reported as a 2.3 : 1.1 : 1 mix of two keto diastereomers and enol tautomer: δ 10.83 (1H, s, OH_{enol}), 3.75 (3H, s, $\text{OCH}_{3\text{enol}}$), 3.71 (3H, s, $\text{OCH}_{3\text{diast 1}}$), 3.70 (3H, s, $\text{OCH}_{3\text{diast 2}}$), 2.98 (1H, s, $\text{C(=O)CHCO}_2\text{CH}_{3\text{diast 1}}$), 2.92 (1H, s, $\text{C(=O)CHCO}_2\text{CH}_{3\text{diast 2}}$), 2.55-2.28 (3.8H, m, aliphatic CH), 2.13 (0.5H, ddd, $J = 12.9, 9.6, 7.8$ Hz, aliphatic CH), 1.90 (1H, ddd, $J = 12.9, 9.1, 5.0$ Hz, aliphatic $\text{CH}_{\text{diast 1}}$), 1.85-1.75 (1.4H, m, aliphatic CH), 1.70 (0.6H, dddd, $J = 13.0, 8.8, 5.3, 0.7$ Hz, aliphatic CH), 1.65-1.35 (4.4H, m, aliphatic CH), 1.15 (3H, s, C(=O)CHCH_3), 1.09 (3H, s, C(=O)CHCH_3), 1.05 (3H, s, C(=O)CHCH_3), 0.92 (3H, dd, $J = 7.5, 7.5$ Hz, $\text{CH}_2\text{CH}_{3\text{diast 1}}$), 0.91 (3H, dd, $J = 7.5, 7.5$ Hz, $\text{CH}_2\text{CH}_{3\text{diast 2}}$), 0.76 (3H, dd, $J = 7.5, 7.5$ Hz, $\text{CH}_2\text{CH}_{3\text{enol}}$); ^{13}C NMR (100 MHz, CDCl_3), peaks reported for both keto diastereomers and enol tautomer: δ 213.5, 213.0, 176.9, 171.0, 169.5, 169.3, 106.9, 65.9, 64.6, 52.0, 52.0, 50.9, 45.5, 44.4, 44.3, 36.3 (2), 34.1, 33.5, 32.9, 32.8, 32.8, 31.1, 29.9, 27.0, 25.0, 21.0, 9.3, 8.8, 8.6; HRMS Calcd for $\text{C}_{10}\text{H}_{16}\text{O}_3\text{Na}$ ($\text{M}+\text{Na}^+$): 207.0997, Found: 207.0997; $[\alpha]_{\text{D}}^{20} -16.4^\circ$ ($c = 1.06$, CHCl_3) for a sample of 77% ee.

Optical purity determined by chiral GLC analysis of dealkoxycarbonylated product (1 equiv NaCl , 3 equiv H_2O , 1.5 M in DMSO, 150°C , 3 h) in comparison with authentic racemic material. Conditions: CDGTA, 100°C , 15 psi.

Peak #	RetTime [min]	Type	Width [min]	Area counts*s	Height [counts]	Area %
1	6.772	MM	0.0864	2.78002e4	5365.13672	49.92948
2	7.086	MM	0.0961	2.78788e4	4834.10498	50.07052

authentic racemic

Peak #	RetTime [min]	Type	Width [min]	Area counts*s	Height [counts]	Area %
1	6.920	MM	0.0754	489.71146	108.26046	12.00351
2	7.202	MM	0.0904	3590.02393	661.79437	87.99649

76% ee

(–)-Methyl 2-butyl-2-methyl-5-oxocyclopentanecarboxylate. IR (neat): 2957 (s), 2932 (s), 2872 (m), 2862 (m), 1757 (s), 1729 (s), 1654 (s), 1614 (m), 1458 (m), 1441 (m); ^1H NMR (400 MHz, CDCl_3), reported for a 2 : 1 : 1 mix of two keto diastereomers and enol tautomer: δ 10.81 (1H, s, OH_{enol}), 3.76 (3H, s, $\text{OCH}_{3\text{enol}}$), 3.71 (3H, s, $\text{OCH}_{3\text{diast 1}}$), 3.70 (3H, s, $\text{OCH}_{3\text{diast 2}}$), 2.99 (1H, s, $\text{C(=O)CHCO}_2\text{CH}_{3\text{diast 1}}$), 2.92 (1H, s, $\text{C(=O)CHCO}_2\text{CH}_{3\text{diast 2}}$), 2.52-2.29 (3.7H, m, aliphatic CH), 2.14 (1H, ddd, $J = 12.9, 9.6, 7.9$ Hz, aliphatic $\text{CH}_{\text{diast 2}}$), 1.91 (1H, ddd, $J = 14.0, 9.1, 5.0$ Hz, aliphatic $\text{CH}_{\text{diast 1}}$), 1.85-1.75 (1.4H, m, aliphatic CH), 1.71 (1H, dddd, $J = 12.9, 8.8, 5.0, 0.8$ Hz, aliphatic $\text{CH}_{\text{diast 2}}$), 1.61-1.18 (11.6H, m, aliphatic CH), 1.15 (3H, s, $\text{C(Bu)CH}_{3\text{enol}}$), 1.10 (3H, s, $\text{C(Bu)CH}_{3\text{diast 2}}$), 1.06 (3H, s, $\text{C(Bu)CH}_{3\text{diast 1}}$), 0.91 (3H, dd, $J = 6.9, 6.8$ Hz, $\text{CH}_2\text{CH}_{3\text{diast 1}}$), 0.89 (3H, dd, $J = 5.6, 5.6$ Hz, $\text{CH}_2\text{CH}_{3\text{diast 2}}$), 0.87 (3H, dd, $J = 7.3, 7.3$ Hz, $\text{CH}_2\text{CH}_{3\text{enol}}$); ^{13}C NMR (100 MHz, CDCl_3): δ 213.6, 213.0, 176.7, 171.0, 169.5, 169.2, 107.2, 66.0, 64.9, 52.0 (2), 50.9, 45.1, 44.1, 44.0, 41.6, 40.2, 37.2, 36.4, 36.3, 34.0, 33.5, 33.4, 31.1, 27.4, 27.3, 26.7, 26.5, 25.6, 23.5, 23.4, 23.3, 21.4, 14.3, 14.1, 14.1; Anal Calcd for $\text{C}_{12}\text{H}_{20}\text{O}_3$: C, 67.89; H, 9.50; Found C, 68.12; H, 9.58; HRMS Calcd for $\text{C}_{12}\text{H}_{20}\text{O}_3\text{Na}$ ($\text{M} + \text{Na}^+$): 235.1310, Found: 235.1304; $[\alpha]_{\text{D}}^{20} -12.2^\circ$ ($c = 1.23$, CHCl_3) for a sample of 65% ee.

Optical purity determined by chiral GLC analysis of dealkoxycarbonylated product (1 equiv NaCl, 3 equiv H_2O , 1.5 M in DMSO, 150°C , 3 h) in comparison with authentic racemic material. Conditions: β -dex, 80°C , 15 psi.

Peak #	RetTime [min]	Type	Width [min]	Area [pA*s]	Height [pA]	Area %
1	115.034	MF	1.2709	418.25970	5.48513	50.12166
2	118.172	FM	1.5362	416.22922	4.51592	49.87834

authentic racemic

Peak #	RetTime [min]	Type	Width [min]	Area [pA*s]	Height [pA]	Area %
1	116.952	MM	0.7754	18.88666	4.05980e-1	7.20226
2	118.808	MM	1.2094	243.34563	3.35363	92.79774

86% ee

(-)-(2R)-Methyl 2-isopropyl-2-methyl-5-oxocyclopentanecarboxylate. IR (neat): 2964 (s), 2882 (m), 1759 (s), 1729 (s), 1654 (m), 1614 (w), 1462 (m), 1441 (m), 1350 (m); ^1H NMR (400 MHz, CDCl_3), reported for a 1.7 : 1 : 1.2 ratio of two keto diastereomers and enol tautomer: δ 10.89 (1H, s, OH_{enol}), 3.75 (3H, s, $\text{OCH}_{3\text{enol}}$), 3.71 (3H, s, $\text{OCH}_{3\text{diast 1}}$), 3.69 (3H, s, $\text{OCH}_{3\text{diast 2}}$), 3.12 (1H, s, $\text{C(=O)CHCO}_2\text{CH}_{3\text{diast 1}}$), 2.95 (1H, s, $\text{C(=O)CHCO}_2\text{CH}_{3\text{diast 2}}$), 2.60-2.28 (4.6H, m, aliphatic CH), 2.22-2.02 (1.2H, m, aliphatic CH), 1.89-1.66 (5H, m, aliphatic CH), 1.35 (0.7H, ddd, $J = 13.2, 9.6, 5.4$ Hz, aliphatic CH), 1.18 (3H, s, $\text{C}(i\text{-Pr})\text{CH}_{3\text{enol}}$), 1.04 (3H, s, $\text{C}(i\text{-Pr})\text{CH}_{3\text{diast 1}}$), 0.94 (3H, d, $J = 6.8$ Hz, $\text{CH}(\text{CH}_3)(\text{CH}_3)_{\text{diast 1}}$), 0.92 (3H, d, $J = 6.8$ Hz, $\text{CH}(\text{CH}_3)(\text{CH}_3)_{\text{diast 1}}$), 0.91 (3H, d, $J = 6.8$ Hz, $\text{CH}(\text{CH}_3)(\text{CH}_3)_{\text{diast 2}}$), 0.90 (3H, d, $J = 6.8$ Hz, $\text{CH}(\text{CH}_3)(\text{CH}_3)_{\text{diast 2}}$), 0.89 (3H, s, $\text{C}(i\text{-Pr})\text{CH}_{3\text{diast 2}}$), 0.85 (3H, d, $J = 6.9$ Hz, $\text{CH}(\text{CH}_3)(\text{CH}_3)_{\text{enol}}$), 0.72 (3H, d, $J = 6.9$ Hz, $\text{CH}(\text{CH}_3)(\text{CH}_3)_{\text{enol}}$); ^{13}C NMR (100 MHz, CDCl_3): δ 214.3, 213.0, 176.5, 171.1, 170.1, 169.2, 107.6, 66.0, 64.5, 52.1, 52.0, 51.0, 48.8, 48.0, 47.6, 37.3, 36.5, 36.1, 34.1, 34.0, 33.4, 32.7, 31.3, 28.0, 26.1, 18.8, 18.7, 18.6, 18.1, 17.6, 17.5, 17.4, 17.1; HRMS Calcd for $\text{C}_{11}\text{H}_{18}\text{O}_3\text{Na}$ ($\text{M}+\text{Na}^+$): 221.1154, Found: 221.1152; $[\alpha]_{\text{D}}^{20} -28.9^\circ$ ($c = 1.17$, CHCl_3) for a sample of 61% ee.

(S)-(-)-3-Isopropyl-3-methylcyclopentanone. In a 300 μL vial was added **16** (30 mg, 0.15 mmol, 60% ee), NaCl (8.8 mg, 0.15 mmol), DMSO (100 μL), and H_2O (8.1 μL , 0.45 mmol). A condenser was attached to the vial and the mixture was heated at 150 $^\circ\text{C}$ for 3 h. After cooling to 23 $^\circ\text{C}$, water (0.5 mL) and pentane (0.5 mL) were added. The layers were separated and the aqueous layer was washed with pentane (3×1 mL). The combined organics were dried over Na_2SO_4 , filtered and concentrated in vacuo (*care*: volatile product) to yield a pale yellow oil (10 mg, 0.071 mmol, 48%). IR (neat): 2961 (m), 2875 (w), 1742 (s); ^1H NMR (400 MHz, CDCl_3): δ 2.38-2.22 (2H, m, $\text{C(O)CH}_2\text{CH}_2$), 2.05 (2H, s, $\text{C(O)CH}_2\text{C}(\text{CH}_3)(i\text{-Pr})$), 1.89-1.68 (2H, m, $\text{C(O)CH}_2\text{CH}_2$), 1.61 (1H, qq, $J = 6.9, 6.9$ Hz, $\text{CH}(\text{CH}_3)_2$), 0.91 (3H, d, $J = 6.9$ Hz, $\text{CH}(\text{CH}_3)\text{CH}_3$), 0.90 (3H, s, $\text{C}(i\text{-Pr})\text{CH}_3$), 0.89 (3H, d, $J = 6.9$ Hz, $\text{CH}(\text{CH}_3)\text{CH}_3$); ^{13}C NMR (100 MHz, CDCl_3): δ 220.4, 52.2, 43.0, 38.4, 37.1, 34.7, 19.2, 18.6, 17.4; HRMS Calcd for $\text{C}_9\text{H}_{16}\text{O}$ (M^+): 140.1201, Found: 140.1200; $[\alpha]_{\text{D}}^{20} -92.4^\circ$ ($c = 1.64$, CHCl_3) for a sample of 60% e.e, indicating the *S*-isomer.¹²

Optical purity determined by chiral GLC analysis in comparison with authentic racemic material. Conditions: CDGTA, 100 $^\circ\text{C}$, 15 psi.

(12) Norin, T. *Acta. Chem. Scand.* **1965**, *19*, 1289-1292; Reported: $[\alpha]_{\text{D}}^{23} +162^\circ$ ($c = 1.63$, CHCl_3) for (*R*)-3-isopropyl-3-methylcyclopentanone.

Peak #	RetTime [min]	Type	Width [min]	Area [pA*s]	Height [pA]	Area %
1	9.752	MM	0.1436	986.66125	114.49625	49.99956
2	10.704	MM	0.1821	986.67847	90.31216	50.00044

authentic racemic

Peak #	RetTime [min]	Type	Width [min]	Area [pA*s]	Height [pA]	Area %
1	10.049	MM	0.0799	22.01718	4.59456	17.17379
2	11.012	MM	0.0976	106.18504	18.13884	82.82621

66% ee

(+)-Ethyl 2-butyl-2-methyl-5-oxocyclopentanecarboxylate. IR (neat): 2959

(s), 2932 (s), 2873 (m), 2862 (m), 1757 (s), 1726 (s), 1651 (s), 1614 (m), 1467 (m); ¹H NMR (400 MHz, CDCl₃), reported for a 1.0 : 0.62 : 0.78 ratio of two keto diastereomers and enol tautomer: δ 10.89 (1H, s, OH_{enol}), 4.30-4.09 (2H, m, OCH₂CH₃_{3diast 1,2 and enol}), 2.96 (1H, br s, C(=O)CHCO₂Et_{diast 1}), 2.89 (1H, s, C(=O)CHCO₂Et_{diast 2}), 2.55-2.26 (4.50H, m, aliphatic CH), 2.15 (0.5H, ddd, *J* = 12.8, 9.6, 8.2 Hz, aliphatic CH), 1.92 (0.9H, ddd, *J* = 12.9, 9.1, 5.1 Hz, aliphatic CH), 1.85-1.76 (1.7H, m, aliphatic CH), 1.71 (0.6H, dddd, *J* = 12.8, 8.8, 4.7, 0.8 Hz, aliphatic CH), 1.65-1.35 (6.2H, m, aliphatic CH), 1.35-1.20 (15.2H, m, aliphatic CH), 1.16 (3H, s, C(Bu)CH₃_{enol}), 1.09 (3H, s, C(Bu)CH₃_{diast 2}), 1.07 (3H, s, C(Bu)CH₃_{diast 1}), 0.95-0.84 (3H, m, CH₂CH₂CH₃_{diast 1,2 and enol}); ¹³C

NMR (100 MHz, CDCl₃): δ 213.9, 213.2, 176.6, 170.6, 169.1, 168.8, 107.3, 66.0, 65.0, 61.0, 60.9, 59.7, 45.2, 44.2, 44.0, 41.6, 40.2, 37.4, 36.4, 36.3, 34.0, 33.6, 33.4, 31.1, 27.6, 27.4, 26.7, 26.5, 25.6, 23.5, 23.5, 23.4, 21.5, 14.4, 14.4, 14.3, 14.3, 14.1, 14.1; HRMS Calcd for C₁₃H₂₂O₃ (M): 226.1569, Found: 226.1574; [α]_D²⁰ +19.2° (*c* = 1.00, CHCl₃) for a sample of 82% ee.

Optical purity determined by chiral GLC analysis of dealkoxycarbonylated product (1 equiv NaCl, 3 equiv H₂O, 1.5 M in DMSO, 150 °C, 3 h) in comparison with authentic racemic material. Conditions: β-dex, 80 °C, 15 psi.

Peak #	RetTime [min]	Type	Width [min]	Area [pA*s]	Height [pA]	Area %
1	115.034	MF	1.2709	418.25970	5.48513	50.12166
2	118.172	FM	1.5362	416.22922	4.51592	49.87834

authentic racemic

Peak #	RetTime [min]	Type	Width [min]	Area [pA*s]	Height [pA]	Area %
1	114.387	MM	0.9462	108.01502	1.90254	91.18168
2	117.802	MM	0.8500	10.44630	2.04818e-1	8.81832

82% ee

(+)-(S)- Methyl 2- (diethoxyphosphoryloxy) -6-ethyl-6-methylcyclohex-1-enecarboxylate.

Sodium hydride (60% in mineral oil, 3.9 mg, 0.098 mmol) was washed with pentane (0.5 mL) and Et₂O (1.0 mL) was added. The resulting suspension was cooled to 0 °C and a solution of conjugate addition adducts (16 mg, 0.082 mmol in 0.3 mL Et₂O) was added via cannula, and rinsed with an additional 0.2 mL Et₂O. The solution was warmed to room temperature over 30 min and then re-cooled to 0 °C. Diethylchlorophosphate (14 µL, 0.098 mmol) was added and the solution was stirred at room temperature for 16 h. Excess NaH and diethylchlorophosphate were quenched by addition of a saturated aqueous solution of NH₄Cl (1 mL). The layers were separated and the aqueous layer was washed with EtOAc (3 × 1 mL). The combined organics were washed with brine (2 mL), dried over MgSO₄ and concentrated in vacuo. The residue was purified by silica gel chromatography (3:1 hexanes : EtOAc → 1:1 hexanes : EtOAc) to yield a pale yellow oil (22 mg, 0.066 mmol, 81%). IR (neat): 2966 (m), 2943 (m), 2879 (w), 1727 (s), 1673 (m), 1457 (m), 1436 (m); ¹H NMR (400 MHz, CDCl₃): δ 4.18-4.08 (4H, m, OCH₂CH₃), 3.74 (3H, s, OCH₃), 2.44-2.38 (2H, m, OC(=C)CH₂), 1.79-1.70 (2H, m, aliphatic CH₂), 1.63-1.53 (2H, m, aliphatic CH₂), 1.48 (1H, q, *J* = 7.5 Hz, CCHHCH₃), 1.47 (1H, q, *J* = 7.5 Hz, CCHHCH₃), 1.36-1.31 (6H, m, OCH₂CH₃), 1.13 (3H, s, C(Et)CH₃), 0.83 (3H, dd, *J* = 7.5, 7.5 Hz, CCH₂CH₃); ¹³C NMR (100 MHz, CDCl₃): δ 168.1, 147.9, 125.7 (d, *J*_{C-P} = 8.4 Hz), 64.5 (2C, d, *J*_{C-P} = 3.5 Hz), 51.5, 37.3, 32.9, 32.7, 27.7, 25.8, 18.5, 16.2, 16.2, 8.5; HRMS Calcd for C₁₅H₂₇O₆PNa (M+Na⁺): 357.1443, Found: 357.1447; [α]_D²⁰ = +21.0 ° (*c* = 1.00, CHCl₃) for a sample of 80% ee.

(+)-Methyl 2-(diethoxyphosphoryloxy)-6-isopropyl-6-methylcyclohex-1-enecarboxylate.

Prepared from **7** (71% ee sample) as described above for **VI**. IR (neat): 2949 (br s), 2876 (m), 2842 (w), 1727 (s), 1673 (m), 1456 (m), 1434 (m), 1362 (m); ^1H NMR (400 MHz, CDCl_3): δ 4.20-4.05 (4H, m, $\text{P}(\text{OCH}_2\text{CH}_3)_2$), 3.72 (3H, s, CO_2CH_3), 2.46-2.29 (2H, m, aliphatic CH), 1.81-1.62 (3H, m, aliphatic CH), 1.49 (1H, ddd, $J = 12.9, 12.5, 4.0$ Hz, aliphatic CH), 1.35-1.28 (1H, m, aliphatic CH), 1.32 (3H, qd, $J_q = 7.0$ Hz, $J_{\text{H-P}} = 1.1$ Hz, OCH_2CH_3), 1.31 (3H, qd, $J_q = 7.0$ Hz, $J_{\text{H-P}} = 1.1$ Hz, OCH_2CH_3), 1.22 (3H, s, $\text{CCH}_3(i\text{-Pr})$), 0.87 (3H, d, $J = 6.8$ Hz, $\text{CH}(\text{CH}_3)\text{CH}_3$), 0.85 (3H, d, $J = 6.8$ Hz, $\text{CH}(\text{CH}_3)\text{CH}_3$); ^{13}C NMR (100 MHz, CDCl_3): δ 168.2, 148.0 (d, $J_{\text{C-P}} = 7.5$ Hz), 125.9 (d, $J_{\text{C-P}} = 8.3$ Hz), 64.4 (d, $J_{\text{C-P}} = 6.6$ Hz), 64.4 (d, $J_{\text{C-P}} = 6.5$ Hz), 51.5, 40.7, 35.4, 28.1, 28.0, 26.2, 18.6, 18.5, 16.8, 16.2, 16.2; HRMS Calcd for $\text{C}_{16}\text{H}_{29}\text{O}_6\text{P}$ (M^+): 348.1702, Found: 348.1701; $[\alpha]_{\text{D}}^{20} +41.9^\circ$ ($c = 1.00$, CHCl_3).

(+)-Methyl 6-isopropyl-2,6-dimethylcyclohex-1-enecarboxylate.

CuI (49 mg, 0.26 mmol), was suspended in Et_2O (1 mL) and cooled to 0°C . MeLi (330 μL , 0.52 mmol of a 1.56 M solution in Et_2O) was added dropwise. The resulting clear colorless solution of Gilman's reagent was cooled to -78°C and enol phosphate **VII** (45 mg, 0.13 mmol, 71% ee) was added dropwise. The reaction was allowed to warm slowly to 0°C , after which it was quenched by addition of 1 M aqueous HCl (1 mL). The layers were separated and the aqueous layer was washed with Et_2O (3×1.5 mL). The combined organic extracts were washed sequentially with saturated aqueous NaHCO_3 and saturated aqueous NaCl , dried over MgSO_4 and concentrated in vacuo. The residue was purified by silica gel chromatography (20:1 hexanes : Et_2O) to yield a pale yellow oil (19 mg, 0.090 mmol, 70%). IR (neat): 2942 (m), 2873 (w), 2831 (w), 1722 (s), 1454 (w), 1429 (w), 1268 (m), 1222 (m); ^1H NMR (400 MHz, CDCl_3): δ 3.72 (3H, s, CO_2CH_3), 2.09-1.87 (2H, m, aliphatic CH), 1.71-1.54 (3H, m, aliphatic CH), 1.66 (3H, s, $\text{C}(\text{=C})\text{CH}_3$), 1.48 (1H, ddd, $J = 12.3, 12.3, 4.1$ Hz, aliphatic CH), 1.32 (1H, dddd, $J = 12.7, 4.3, 3.1, 1.3$ Hz, aliphatic CH), 1.18 (3H, s, $\text{C}(i\text{-Pr})\text{CH}_3$), 0.85 (3H, d, $J = 6.8$ Hz, $\text{CH}(\text{CH}_3)\text{CH}_3$), 0.85 (3H, d, $J = 6.8$ Hz, $\text{CH}(\text{CH}_3)\text{CH}_3$); ^{13}C NMR (100 MHz, CDCl_3): δ 171.5, 135.7, 135.4, 51.1, 39.9, 35.6, 31.7, 28.7, 26.5, 21.8, 18.8, 18.6, 16.9; HRMS Calcd for $\text{C}_{13}\text{H}_{22}\text{O}_2$ (M^+): 211.1698, Found: 211.1701; $[\alpha]_{\text{D}}^{20} +82.8^\circ$ ($c = 1.00$, CHCl_3).

(+)-(S)-tert-Butyl 6-ethyl-6-methyl-2-(trifluoromethylsulfonyloxy)cyclohex-1-enecarboxylate.

^{13}NaH (33 mg, 0.83 mmol, 60% in mineral oil) was washed with pentane (1 mL) under N_2 . **9** (100 mg, 0.416 mmol, 87% ee) in Et_2O (0.8 mL) was added at 0°C . The suspension was allowed to stir at 0°C for 30 min, then at 23°C for 30 min. The suspension was then recooled to 0°C and Tf_2O (distilled over P_2O_5 , 105 μL , 0.624 mmol) was

added dropwise. The suspension was allowed to stir at 0 °C for 1 h, then at 23 °C for 1 h before cooling again to 0 °C, diluting with Et₂O (2 mL) and quenching excess Tf₂O by slow addition of H₂O (1 mL). The layers were separated and the organic layer was washed with Et₂O (3 × 2 mL). The combined organics were washed with saturated aqueous NaHCO₃ (1 mL), saturated aqueous NaCl (1 mL), dried over MgSO₄ and concentrated in vacuo to yield **VIII** as a pale yellow oil (143 mg, 0.384 mmol, 92%). IR (neat): 2971 (m), 2945 (m), 2881 (w), 1722 (s), 1422 (s), 1211 (s); ¹H NMR (400 MHz, CDCl₃): δ 2.38 (1H, A of ABXY, *J*_{AB} = 17.5, *J*_{AX} = 7.4, *J*_{AY} = 7.4 Hz, C(OTf)CHH), 2.30 (1H, B of ABXY, *J*_{AB} = 17.5, *J*_{BX} = 5.4, *J*_{BY} = 5.4 Hz, C(OTf)CHH), 1.97-1.71 (2H, m, aliphatic CH), 1.68-1.44 (3H, m, aliphatic CH), 1.52 (9H, s, OC(CH₃)₃), 1.32 (1H, ddd, *J* = 13.3, 6.4, 3.3 Hz, aliphatic CH), 1.18 (3H, s, C(Et)CH₃), 0.88 (3H, dd, *J* = 7.5, 7.5 Hz, CH₂CH₃); ¹³C NMR (100 MHz, CDCl₃): δ 164.6, 146.1, 134.0, 118.3 (q, *J*_{C-F} = 319 Hz), 83.4, 39.0, 32.6, 32.5, 28.2, 27.1, 26.0, 18.9, 8.6; Anal Calcd for C₁₅H₂₃F₃O₅S: C, 48.38; H, 6.23; Found C, 48.59; H, 6.47; [α]_D²⁰ +10.5° (*c* = 1.00, CHCl₃).

(+)-(S)-tert-Butyl 6-ethyl-6-methylcyclohex-1-enecarboxylate. To a solution of **9** (50 mg, 0.21 mmol, 87% ee) in MeOH (700 μ L) at 0 °C was added NaBH₄ (24 mg, 0.62 mmol). The solution was warmed to 23 °C and allowed to stir for 1 h. At this time, the reaction was complete by TLC analysis (*R*_f starting materials = 0.67 and 0.26, *R*_f product = 0.12, 10:1 hexanes : Et₂O). The excess

NaBH₄ was quenched by addition of H₂O (0.5 mL) and 1M aqueous HCl (0.5 mL). The solution was stirred vigorously for 5 min, then washed with EtOAc (3 × 3 mL). The combined organics were washed with saturated aqueous NaCl (3 mL), dried over MgSO₄ and concentrated in vacuo to yield the alcohol as a colorless oil. The alcohol was identified only by ¹H NMR (approx 6:3:2:1 mixture of diastereomers) and then used directly in the next step. The alcohol was dissolved in pyridine (700 μ L) and cooled to 0 °C. Freshly distilled POCl₃ (78 μ L, 0.83 mmol) was added dropwise with stirring. The ice bath was removed and the white suspension was allowed to warm to 23 °C and stirred for a further 16 h. At this point, the reaction mixture was diluted with Et₂O (2 mL) and added to ice and water (0.5 mL). After the ice had melted, the layers were separated and the aqueous layer was washed with EtOAc (3 × 3 mL). The combined organics were washed with 1M aqueous HCl (3 × 2 mL), saturated aqueous NaCl (3 mL), dried over MgSO₄ and concentrated in vacuo. The residue was purified by silica gel chromatography (25:1 hexanes : Et₂O, *R*_f product = 0.58 in 10:1 hexanes : Et₂O) to yield a colorless oil (33 mg, 0.15 mmol, 71%). IR (neat): 2965 (s), 2935 (s), 2867 (m), 1706 (s), 1456 (m), 1367 (s), 1269 (s), 1253 (s), 1166 (s); ¹H NMR (400 MHz, CDCl₃): δ 6.74 (1H, dd, *J* = 4.4, 3.6 Hz, C=CH), 2.19-2.01 (2H, m, C=CCH₂), 1.81 (1H, dddd, *J* = 13.7, 7.6, 7.6, 7.4 Hz, aliphatic CH), 1.67-1.55 (3H, m, aliphatic CH), 1.48 (9H, s, C(CH₃)₃), 1.48-1.40 (1H, m, aliphatic CH), 1.29-1.24 (1H, m, aliphatic CH), 1.19 (3H, s, C(Et)CH₃), 0.75 (3H, dd, *J* = 7.5, 7.5 Hz, CH₂CH₃); ¹³C NMR (100 MHz, CDCl₃): δ 167.5, 139.5, 138.8, 79.9, 36.3, 35.2, 32.3, 28.3 (3), 26.6, 26.3, 18.3, 8.8; HRMS Calcd for C₁₄H₂₅O₂ (M+H⁺): 225.1855, Found: 225.1866; [α]_D²⁰ +40.3° (*c* = 1.00, CHCl₃).

Ligand Screening Details

Experimental procedure for ligand screening reactions

Cu salt (6.0 μmol) and ligand (7.5 μmol) are weighed in to a 13 $\times$ 100 mm test tube in a N_2 -filled glovebox. The tube was capped with a septum and taped with Parafilm. A solution of substrate (0.15 mmol) in dry solvent (1.5 mL) with decane (20 μL , internal standard to calculate conversion) was added. The mixture was cooled to 0 $^\circ\text{C}$ and Et_2Zn (0.45 mmol) was added. After 24 h, the reaction mixture was cooled to -78 $^\circ\text{C}$. Et_2O (1 mL) was added, followed by 1 mL 1 N aqueous HCl. The solution was stirred rapidly. The upper organic layer was removed and passed through a plug of silica gel. The aqueous layer was washed with EtOAc (3 $\times$ 1.5 mL) and passed through the silica plug. The crude organic filtrate was analyzed by chiral GLC.

Initial ligand and Cu salt screening results are depicted in Table 1 below.

Table 1: Screening combinations of Cu-salt, ligand and solvent

	2a			4			5		
conv (%); e.e. (%)	THF	toluene	CH_2Cl_2	THF	toluene	CH_2Cl_2	THF	toluene	CH_2Cl_2
(CuOTf) $_2$ ·C $_6$ H $_6$	<10; n.d.	<10; n.d.	<10; n.d.	<10; n.d.	<10; n.d.	<10; n.d.	10; 18	18; 5	21; -6
Cu(OTf) $_2$	<10; n.d.	51; 4*	10; n.d.	<10; n.d.	78; <5	<10; n.d.	<10; n.d.	>95; -19	<10; n.d.
CuCN	23; 13	>95; 42	>95; 22	<10; n.d.	>95; <5	88; 7	12; 13	>95; <5	>95; -6
CuOAc	<10; n.d.	<10; n.d.	<10; n.d.	<10; n.d.	<10; n.d.	<10; n.d.	<10; n.d.	<10; n.d.	11; <5
Cu(OAc) $_2$ ·H $_2$ O	<10; n.d.	<10; n.d.	<10; n.d.	<10; n.d.	<10; n.d.	<10; n.d.	<10; n.d.	<10; n.d.	15; <5
CuCl	<10; n.d.	<10; n.d.	<10; n.d.	11; <5	27; 6	<10; n.d.	<10; n.d.	<10; n.d.	66; <5
CuBr·SMe $_2$	<10; n.d.	15; 13	<10; n.d.	<10; n.d.	31; <5	<10; n.d.	<10; n.d.	34; 4	<10; n.d.
CuI	13; <5	<10; n.d.	<10; n.d.	<10; n.d.	17; <5	<10; n.d.	14; <5	<10; n.d.	<10; n.d.

n.d. = not determined; * substantial amounts of unidentified byproduct

Amino acid screening was conducted on ligand **2a** as shown in Chart 1 below. *tert*-Leucine proved optimal as AA-1. However, with only a modest increase in enantioselectivity, we were encouraged to investigate the other ligand modules before returning to the AA-1 screen.

An *N*-terminus screen was conducted next. The first round (Figure 1) indicated that all variations of a salicyl terminus were inferior to unsubstituted salicylic acid. However, anthranilic acid gave higher selectivity at 47% ee.

Further screening of additional anthranilic acids (Figure 2) indicated *N*-methyl anthranilic acid as the most promising candidate.

Figure 2: Second round screening of *N*-terminus: substituted anthranilic acids.

C-terminus screening was carried out next. As shown in Figure 3, the dimethyl amide bearing ligand gave the highest enantioselectivity.

Figure 3: *C*-terminus screening

With the optimal two termini in hand, we again turned our attention to screening of the amino acid modules. The screening results are summarized in Charts 2 and 3 to give the optimal ligand **2f**.

Chart 2: Ligand screening data: AA-1 screen

Chart 3: Ligand screening data: AA-2 screen

ASH-1U-071-02-F1
Pulse Sequence: zgpg30

400-MV-LB-C
Pulse Sequence: zgpg1

APM-10-19-B-C
 Pulse Sequence: zgpg30