Supporting Information

Catalytic Enantioselective Mannich-Type Reactions of Ketoimines

Yutaka Suto, Motomu Kanai,* and Masakatsu Shibasaki*

Graduate School of Pharmaceutical Sciences, The University of Tokyo

Hongo, Bunkyo-ku, Tokyo 113-0033, Japan

General: NMR spectra were recorded on a JEOL JNM-LA500 spectrometer, operating at 500 MHz for ^1H NMR, 125.65 MHz for ^{13}C NMR and 202.35 MHz for ^{31}P NMR. Chemical shifts were reported downfield from TMS ($= 0$) for ^1H NMR. For ^{13}C NMR, chemical shifts were reported in the scale relative to the solvent used as an internal reference. ^{31}P NMR were carried out with H_3PO_4 ($= 0$ ppm) as an external standard. Optical rotations were measured on a JASCO P-1010 polarimeter. ESI-Q mass spectra were measured on Waters-ZQ4000. Column chromatographies were performed with silica gel Merck 60 (230-400 mesh ASTM). The enantiomeric excess (ee) was determined by HPLC analysis. HPLC analysis was performed on JASCO HPLC systems consisting of the following: pump, 880-PU or PU-980; detector, 875-UV or UV-970, measured at 254 nm; mobile phase, hexane-2-propanol. In general, reactions were carried out in dry solvents under an argon atmosphere, unless noted otherwise. Normal glassware can be used to conduct the reactions in this manuscript. (*R*)-DTBM-SEGPBOS was kindly provided from Takasago International Co. (Fax: +81-463-25-2084).

Preparation of *N*-di(3,5-xylyl)phosphinoylimine (**2**).

To a degassed solution of the corresponding ketoxime (3.6 mmol) and triethylamine (1.0 mL, 7.2 mmol) in hexane: $\text{CH}_2\text{Cl}_2 = 1:1$ (30 mL), chlorodi(3,5-xylyl)phosphine (1.0 mL, 3.6 mmol) was slowly added over 30 min at -40 °C. After the addition was complete, the temperature was gradually increased to room temperature and the mixture was allowed to stir overnight at room temperature. Filtration through celite, evaporation and purification by silica gel column chromatography ($\text{AcOEt}/\text{CH}_2\text{Cl}_2 = 1/10-1/5$) gave the product. The product was further purified by recrystallization from hexane/ CH_2Cl_2 (**2a–2h**, **2j**), or used without further purification (**2i**, **2k**, **2l**). ^{13}C NMR peaks assigned to the aromatic carbons were complicated due to the coupling to the phosphorus nucleus. *N*-Diphenylphosphinoylimine **1a** is a known compound.¹

N-(1-Phenylethylidene)-di(3,5-xylyl)phosphinamide (**2a**)

^1H NMR (C_6D_6) δ 2.02(s, 12H), 2.83 (d, $J = 1.3$ Hz, 3H), 6.75 (s, 2H), 7.04-7.11 (m, 3H), 7.87-7.90 (m, 2H), 8.06 (d, $J = 12.1$ Hz, 4H); ^{13}C NMR (C_6D_6) δ 21.18, 22.61, 22.70, 128.47, 129.85, 129.92, 131.89, 133.12, 133.14, 135.94, 136.96, 138.13, 138.23, 140.21, 140.40, 181.11 (d, $J_{\text{C-P}} = 8.4$ Hz); ^{31}P NMR (C_6D_6) δ 18.8; MS m/z 398 ($\text{M}+\text{Na}^+$); HRMS Calcd for $\text{C}_{24}\text{H}_{26}\text{NOP}$ (M^+) 375.1747, Found 375.1745; IR (KBr) 1637 cm^{-1} .

N-(1-*p*-Chlorophenylethylidene)-di(3,5-xylyl)phosphinamide (**2b**)

^1H NMR (C_6D_6) δ 2.04(s, 12H), 2.72 (d, $J = 1.8$ Hz, 1H), 6.77 (s, 2H), 6.99-7.05 (m, 2H), 7.58-7.63 (m, 2H), 8.01 (d, $J = 11.9$ Hz, 4H); ^{13}C NMR (C_6D_6) δ 21.18, 22.39, 22.49, 128.68, 129.45, 129.79, 129.87, 133.22,

133.23, 135.72, 136.74, 138.17, 138.22, 138.27, 138.49, 138.68, 179.65 (d, $J_{C-P} = 7.2$ Hz); ^{31}P NMR (C_6D_6) δ 17.1; MS m/z 432 ($\text{M}+\text{Na}^+$); HRMS Calcd for $\text{C}_{24}\text{H}_{25}\text{ClNOP}^+$ (M^+) 409.1357, Found 409.1356; IR (KBr) 1636 cm^{-1} .

***N*-(1-*p*-Methoxyphenylethylidene)-di(3,5-xylyl)phosphinamide (2c)**

^1H NMR (C_6D_6) δ 2.04(s, 12H), 2.85 (d, $J = 1.8$ Hz, 3H), 3.23 (s, 3H), 6.63-6.67 (m, 2H), 6.77 (s, 2H), 7.90-7.95 (m, 2H), 8.07 (d, $J = 11.9$ Hz, 4H); ^{13}C NMR (C_6D_6) δ 21.21, 22.44, 22.54, 54.86, 113.84, 129.86, 129.93, 130.23, 132.99, 133.00, 133.19, 136.41, 137.43, 138.04, 138.15, 163.20, 180.12, 180.18; ^{31}P NMR (C_6D_6) δ 16.5; MS m/z 428 ($\text{M}+\text{Na}^+$); HRMS Calcd for $\text{C}_{24}\text{H}_{25}\text{NO}_2\text{P}^+$ ($\text{M}-\text{Me}^+$) 390.1617, Found 390.1614 IR (KBr) 1699 cm^{-1} .

***N*-(1-Naphthalene-2-ylidene)-di(3,5-xylyl)phosphinamide (2d)**

^1H NMR (C_6D_6) δ 2.04(s, 12H), 2.95 (d, $J = 1.9$ Hz, 3H), 6.77 (s, 2H), 7.19-7.26 (m, 2H), 7.51-7.58 (m, 2H), 7.60 (d, $J = 8.6$ Hz 1H), 8.08 (s, 1H), 8.13 (d, $J = 11.7$, 4H), 8.40 (dd, $J = 1.9$ Hz, 8.9 Hz, 1H); ^{13}C NMR (C_6D_6) δ 21.21, 22.59, 22.68, 124.55, 126.57, 129.62, 129.92, 129.99, 133.14, 135.49, 136.05, 137.07, 137.73, 137.90, 138.16, 138.26, 180.96 (d, $J_{C-P} = 8.3$ Hz); ^{31}P NMR (C_6D_6) δ 17.1; MS m/z 448 ($\text{M}+\text{Na}^+$); HRMS Calcd for $\text{C}_{28}\text{H}_{28}\text{NOP}^+$ (M^+) 425.1903, Found 425.1908; IR (KBr) 1619 cm^{-1} .

***N*-[1-(2-furanyl)ethylidene]-di(3,5-xylyl)phosphinamide (2e)**

^1H NMR (C_6D_6) δ 2.03(s, 12H), 2.70 (d, $J = 2.2$ Hz, 3H), 5.88 (dd, $J = 1.9$ Hz, $J = 3.7$ Hz, 1H), 6.69 (d, $J = 3.7$ Hz, 1H), 6.75 (s, 2H), 6.91 (d, $J = 1.9$ Hz, 1H), 8.07 (d, $J = 12.2$ Hz, 4H); ^{13}C NMR (C_6D_6) δ 21.16, 21.85, 21.95, 112.27, 116.04, 129.91, 129.97, 133.08, 133.11, 135.93, 136.95, 138.06, 138.16, 145.92, 154.86, 155.09, 169.64 (d, $J_{C-P} = 6.3$ Hz); ^{31}P NMR (C_6D_6) δ 17.7; MS m/z 388 ($\text{M}+\text{Na}^+$); HRMS Calcd for $\text{C}_{22}\text{H}_{24}\text{NO}_2\text{P}^+$ (M^+) 365.1539, Found 365.1535; IR (KBr) 1627 cm^{-1} .

***N*-[1-(3-Thienyl)ethylidene]-di(3,5-xylyl)phosphinamide (2f)**

^1H NMR (C_6D_6) δ 2.03(s, 12H), 2.70 (d, $J = 2.1$ Hz, 3H), 6.68 (dd, $J = 2.9$ Hz, $J = 5.1$ Hz, 1H), 6.76 (s, 2H), 7.16 (dd, $J = 1.2$ Hz, $J = 2.9$ Hz, 1H), 7.68 (dd, $J = 1.2$ Hz, 5.1 Hz, 1H), 8.03 (d, $J = 11.9$ Hz, 4H); ^{13}C NMR (C_6D_6) δ 21.18, 23.28, 23.37, 125.84, 127.29, 129.83, 129.91, 130.47, 131.26, 133.05, 136.08, 137.10, 138.07, 138.18, 145.109, 175.62, 175.68; ^{31}P NMR (C_6D_6) δ 15.6; MS m/z 404 ($\text{M}+\text{Na}^+$); HRMS Calcd for $\text{C}_{22}\text{H}_{26}\text{NOPS}^+$ ($\text{M}+\text{H}^+$) 382.1394, Found 382.1400; IR (KBr) 1616 cm^{-1} .

***N*-(1-Phenylpropylidene)-di(3,5-xylyl)phosphinamide (2g)**

^1H NMR (C_6D_6) δ 1.16 (t, $J = 8.1$ Hz, 3H), 2.01 (s, 12H), 3.47 (dt, $J = 1.8$ Hz, $J = 8.1$ Hz, 2H), 6.75 (s, 2H), 7.07-7.13 (m, 3H), 7.92-7.97 (m, 2H), 8.08 (d, $J = 12.0$ Hz, 4H); ^{13}C NMR (C_6D_6) δ 13.47, 13.48, 21.17, 28.91, 29.00, 128.44, 128.69, 129.85, 129.92, 131.84, 133.08, 133.09, 136.14, 137.16, 138.11, 138.22, 138.90, 139.09, 186.17 (d, $J_{C-P} = 7.2$ Hz); ^{31}P NMR (C_6D_6) δ 17.9; MS m/z 412 ($\text{M}+\text{Na}^+$); HRMS Calcd for $\text{C}_{25}\text{H}_{28}\text{NOP}^+$ (M^+) 389.1903, Found 389.1907; IR (KBr) 1630 cm^{-1} .

***N*-(1-Cyclohex-1-enylethylidene)-di(3,5-xylyl)phosphinamide (2h)**

^1H NMR (C_6D_6) δ 1.26-1.32 (m, 2H), 1.43-1.50 (m, 2H), 1.74-1.80 (m, 2H), 2.04 (s, 12H), 2.53-2.59 (m, 2H), 2.62 (d, $J = 1.9$ Hz, 3H), 6.17-6.21 (m, 1H), 6.76 (s, 2H), 8.02 (d, $J = 11.6$ Hz, 4H); ^{13}C NMR (C_6D_6) δ 21.10,

21.21, 21.78, 22.65, 25.00, 26.36, 26.38, 129.89, 129.97, 132.92, 132.94, 136.32, 137.34, 137.98, 138.08, 138.31, 140.72, 140.91, 181.51 (d, J_{C-P} = 7.2 Hz); ^{31}P NMR (C_6D_6) δ 16.2; MS m/z 402 ($\text{M}+\text{Na}^+$); HRMS Calcd for $\text{C}_{24}\text{H}_{30}\text{NOP}^+$ (M^+) 379.2060, Found 379.2064; IR (KBr) 2926, 1617 cm^{-1} .

***N*-(1-Methylhex-2-enylidene)-di(3,5-xylyl)phosphinamide (2i)**

^1H NMR (C_6D_6) δ 0.69 (t, J = 7.3 Hz, 3H), 1.10-1.18 (m, 2H), 1.76-1.82 (m, 2H), 2.40 (s, 12H), 6.13 (dt, J = 7.0 Hz, J = 15.9 Hz, 1H), 6.74 (s, 2H), 6.92 (brs, 1H), 8.03 (d, J = 11.3 Hz, 4H); ^{13}C NMR (C_6D_6) δ 13.72, 14.29, 21.16, 21.76, 22.99, 31.90, 34.83, 129.93, 130.01, 132.95, 136.07, 137.08, 137.95, 138.04, 145.22, 180.68 (d, J_{C-P} = 8.3 Hz); ^{31}P NMR (C_6D_6) δ 16.5; MS m/z 390 ($\text{M}+\text{Na}^+$); HRMS Calcd for $\text{C}_{23}\text{H}_{30}\text{NOP}^+$ (M^+) 367.2060, Found 367.2061; IR (KBr) 2958, 1613 cm^{-1} .

***N*-(1-Cyclohexylethylidene)-di(3,5-xylyl)phosphinamide (2j)**

^1H NMR (C_6D_6) δ 0.97-1.12 (m, 3H), 1.24-1.34 (m, 2H), 1.46-1.53 (m, 1H), 1.57-1.64 (m, 2H), 1.67-1.74 (m, 2H), 2.01-2.08 (m, 1H), 2.04 (s, 12H), 2.39 (d, J = 2.2 Hz, 3H), 6.75 (s, 2H), 8.00 (d, J = 12.2 Hz, 4H); ^{13}C NMR (C_6D_6) δ 21.20, 26.13, 26.25, 30.27, 51.78, 51.94, 129.87, 129.94, 132.93, 136.21, 137.22, 137.95, 138.05, 194.18 (d, J_{C-P} = 11.3 Hz); ^{31}P NMR (C_6D_6) δ 15.6; MS m/z 404 ($\text{M}+\text{Na}^+$); HRMS Calcd for $\text{C}_{24}\text{H}_{32}\text{NOP}^+$ (M^+) 381.2216, Found 381.2218; IR (KBr) 2927, 1656 cm^{-1} .

General Procedure for Catalytic Enantioselective Mannich Reaction of Ketimines (Table 3, Entry 1).

(5a): Imine **2a** (75.1 mg, 0.2 mmol) was weighed in the open air into a Schlenk tube equipped with a three-way stopper, and dried under vacuum for 2 h at room temperature. This reaction vessel was put into a glovebox. After (*R*)-DTBM-SEGPBOS (21.6 mg, 0.02 mmol) and CuOAc (2.5 mg, 0.02 mmol) were added to the vessel in a glovebox, the stopper was closed and the reaction vessel was taken out of the glovebox. Dry THF (0.5 mL) was added to the mixture under Ar atmosphere, which was stirred for 10 min at room temperature. To the solution, diacetoxydiethoxysilane (51 μL , 0.02 mmol) and ketene silyl acetal **3** (82 μL , 0.4 mmol) were added at room temperature. The three-way stopper was closed, and the mixture was stirred for 20 h at 40 $^\circ\text{C}$. After cooling, the reaction was quenched with aqueous NaCl, extracted with AcOEt, and the combined organic layer was dried over Na_2SO_4 . Filtration, evaporation, and purification by silica gel column chromatography (AcOEt/ CH_2Cl_2 = 1/10–1/5) gave the product **5a** in 81% yield (80.1 mg).

***N*-[2-butoxycarbonyl-1-methyl-1-(2-naphthyl)ethyl]-diphenylphosphinamide (4d)**

^1H NMR (CDCl_3) δ 0.9 (t, J = 7.3 Hz, 3H), 1.09-1.17 (m, 2H), 1.33-1.40 (m, 2H), 1.62 (s, 3H), 3.23 (d, J = 16.2 Hz, 1H), 3.45 (d, J = 16.2 Hz, 1H), 3.84-3.96 (m, 2H), 5.18 (d, J = 5.2 Hz, 1H), 7.38-7.52 (m, 8H), 7.71 (dd, J = 2.2 Hz, J = 8.6 Hz, 1H), 7.75-7.85 (m, 3H), 7.88-8.04 (m, 5H); ^{13}C NMR (CDCl_3) δ 13.47, 18.92, 29.55 (d, J_{C-P} = 3.1 Hz), 30.39, 47.59 (d, J_{C-P} = 2.1 Hz), 58.95 (d, J_{C-P} = 2.0 Hz), 64.30, 123.60, 123.71, 125.88, 126.05, 127.365, 128.20, 128.24, 128.37, 128.47, 128.50, 128.60, 131.17, 131.25, 131.46, 131.48, 131.50, 131.52, 132.01, 132.08, 132.29, 133.08, 134.23, 134.58, 135.27, 135.59, 144.08, 144.14, 172.20; ^{31}P NMR (CDCl_3) δ 19.4; $[\alpha]_D^{23}$ -45.4 (c = 1.11, CHCl_3) (91% ee); HPLC DAICEL CHIRALCEL AD-H, hexane/2-propanol = 4/1, 1.0 mL/min, t_R 11.3 min (major), 18.7 min (minor); MS m/z 508 ($\text{M}+\text{Na}^+$); HRMS

Calcd for $C_{30}H_{32}NO_3P^+$ (M^+) 485.2114, Found 485.2112; IR (neat) 1720, 1198 cm^{-1} .

***N*-(2-butoxycarbonyl-1-methyl-1-phenylethyl)-di(3,5-xylyl)phosphinamide (5a)**

1H NMR ($CDCl_3$) δ 0.81 (t, $J = 7.3$ Hz, 3H), 1.14-1.23 (m, 2H), 1.37-1.44 (m, 2H), 1.52 (s, 3H), 2.31 (s, 6H), 2.33 (s, 6H), 3.06 (d, $J = 16.2$ Hz, 1H), 3.38 (d, $J = 16.2$ Hz, 1H), 3.86-3.96 (m, 2H), 4.91 (d, $J = 5.2$ Hz, 1H), 7.07 (s, 1H), 7.08 (s, 1H), 7.19-7.23 (m, 1H), 7.30-7.35 (m, 2H), 7.48 (d, $J = 12.2$ Hz, 2H), 7.54-7.59 (m, 4H); ^{13}C NMR ($CDCl_3$) δ 13.54, 18.94, 21.27, 21.30, 29.73 (d, $J_{C-P} = 4.1$ Hz), 30.40, 47.71 (d, $J_{C-P} = 2.1$ Hz), 58.70 (d, $J_{C-P} = 3.0$ Hz), 64.19, 125.00, 126.70, 128.29, 128.69, 128.76, 129.56, 129.64, 133.12, 133.13, 133.16, 133.19, 134.14, 134.46, 135.17, 135.46, 137.87, 137.98, 138.03, 138.13, 147.08, 147.14, 172.07; ^{31}P NMR ($CDCl_3$) δ 20.3; $[\alpha]^{26}_D -17.0$ ($c = 1.12$, $CHCl_3$) (95% ee); HPLC DAICEL CHIRALCEL AD-H, hexane/2-propanol = 20/1, 1.0 mL/min, t_R 16.1 min (major), 29.4 min (minor); MS m/z 514 ($M+Na^+$); HRMS Calcd for $C_{30}H_{38}NO_3P^+$ (M^+) 491.2584, Found 491.2591; IR (neat) 1723, 1199 cm^{-1} .

***N*-(2-butoxycarbonyl-1-*p*-chlorophenyl-1-methylethyl)-di(3,5-xylyl)phosphinamide (5b)**

1H NMR ($CDCl_3$) δ 0.81 (t, $J = 7.4$ Hz, 3H), 1.18-1.24 (m, 2H), 1.40-1.44 (m, 2H), 1.51 (s, 3H), 2.31 (s, 6H), 2.33 (s, 6H), 3.05 (d, $J = 16.2$ Hz, 1H), 3.30 (d, $J = 16.2$ Hz, 1H), 3.90-3.96 (m, 2H), 4.90 (d, $J = 5.5$ Hz, 1H), 7.08 (s, 1H), 7.09 (s, 1H), 7.25-7.29 (m, 2H), 7.46-7.52 (m, 6H); ^{13}C NMR ($CDCl_3$) δ 13.54, 18.97, 21.29, 21.31, 29.58 (d, $J_{C-P} = 4.1$ Hz), 30.43, 47.59, 58.36, 64.38, 126.74, 128.39, 128.75, 128.83, 129.40, 129.47, 132.61, 133.189, 133.21, 133.23, 133.26, 133.91, 133.424, 134.94, 135.25, 137.96, 138.08, 138.18, 145.49, 145.54, 171.94; ^{31}P NMR ($CDCl_3$) δ 20.1; $[\alpha]^{26}_D -24.9$ ($c = 0.80$, $CHCl_3$) (97% ee); HPLC DAICEL CHIRALCEL AD-H, hexane/2-propanol = 20/1, 1.0 mL/min, t_R 19.0 min (major), 26.9 min (minor); MS m/z 548 ($M+Na^+$); HRMS Calcd for $C_{30}H_{37}ClNO_3P^+$ (M^+) 525.2194, Found 525.2194; IR (neat) 1722, 1199 cm^{-1} .

***N*-(2-butoxycarbonyl-1-*p*-methoxyphenyl-1-methylethyl)-di(3,5-xylyl)phosphinamide (5c)**

1H NMR ($CDCl_3$) δ 0.83 (t, $J = 7.4$ Hz, 3H), 1.16-1.25 (m, 2H), 1.39-1.43 (m, 2H), 1.52 (s, 3H), 2.31 (s, 6H), 2.32 (s, 6H), 3.06 (d, $J = 16.2$ Hz, 1H), 3.30 (d, $J = 16.2$ Hz, 1H), 3.78 (s, 3H), 3.89-3.97 (m, 2H), 4.86 (d, $J = 5.5$ Hz, 1H), 6.81-6.86 (m, 2H), 7.06 (s, 1H), 7.08 (s, 1H), 7.44-7.49 (m, 4H), 7.54 (d, $J = 12.2$ Hz, 2H); ^{13}C NMR ($CDCl_3$) δ 13.56, 18.98, 21.28, 21.30, 29.80 (d, $J_{C-P} = 4.1$ Hz), 30.45, 47.77, 55.17, 58.26 (d, $J_{C-P} = 2.0$ Hz), 64.19, 113.40, 126.33, 128.75, 128.82, 129.51, 129.59, 133.06, 133.07, 133.11, 133.13, 134.17, 134.47, 135.20, 135.47, 137.84, 137.94, 137.99, 138.08, 139.00, 139.05, 158.24, 172.13; ^{31}P NMR ($CDCl_3$) δ 19.9; $[\alpha]^{25}_D -20.4$ ($c = 0.91$, $CHCl_3$) (97% ee); HPLC DAICEL CHIRALCEL AD-H, hexane/2-propanol = 20/1, 1.0 mL/min, t_R 25.7 min (major), 47.8 min (minor); MS m/z 544 ($M+Na^+$); HRMS Calcd for $C_{31}H_{40}NO_4P^+$ (M^+) 521.2689, Found 521.2694; IR (neat) 1720, 1185 cm^{-1} .

***N*-[2-butoxycarbonyl-1-methyl-1-(2-naphthyl)ethyl]-di(3,5-xylyl)phosphinamide (5d)**

1H NMR ($CDCl_3$) δ 0.81 (t, $J = 7.7$ Hz, 3H), 1.11-1.19 (m, 2H), 1.34-1.42 (m, 2H), 1.64 (s, 3H), 2.28 (s, 6H), 2.33 (s, 6H), 3.23 (d, $J = 16.2$ Hz, 1H), 3.41 (d, $J = 16.2$ Hz, 1H), 3.88-3.94 (m, 2H), 5.02 (d, $J = 5.5$ Hz, 1H), 7.04 (s, 1H), 7.09 (s, 1H), 7.42-7.52 (m, 4H), 7.57 (d, $J = 12.5$ Hz, 2H), 7.70 (dd, $J = 2.2$ Hz, $J = 8.9$ Hz, 1H), 7.77-7.84 (m, 3H), 7.95 (d, $J = 1.9$ Hz, 1H); ^{13}C NMR ($CDCl_3$) δ 13.47, 18.94, 21.26, 21.32, 29.55 (d, $J_{C-P} = 4.1$ Hz), 30.43, 47.63 (d, $J_{C-P} = 2.1$ Hz), 58.77 (d, $J_{C-P} = 3.0$ Hz), 64.26, 123.75, 123.85, 125.83, 125.98,

127.35, 128.03, 120.20, 128.77, 128.85, 129.58, 129.65, 132.28, 133.09, 133.12, 133.17, 133.20, 134.04, 134.40, 135.07, 135.36, 135.40, 137.86, 137.97, 138.04, 138.15, 144.21, 144.26, 172.10; ^{31}P NMR (CDCl_3) δ 20.1; $[\alpha]_D^{27}$ -50.8 ($c = 0.83$, CHCl_3) (96% ee); HPLC DAICEL CHIRALCEL AD-H, hexane/2-propanol = 20/1, 1.0 mL/min, t_R 17.2 min (major), 26.8 min (minor); MS m/z 564 ($\text{M}+\text{Na}^+$); HRMS Calcd for $\text{C}_{34}\text{H}_{40}\text{NO}_3\text{P}^+$ (M^+) 541.2740, Found 541.2742; IR (neat) 1723, 1197 cm^{-1} .

***N*-[2-butoxycarbonyl-1-(2-furanyl)-1-methylethyl]-di(3,5-xylyl)phosphinamide (5e)**

^1H NMR (CDCl_3) δ 0.87 (t, $J = 7.3$ Hz, 3H), 1.24-1.32 (m, 2H), 1.48-1.54 (m, 2H), 1.64 (s, 3H), 2.29 (s, 6H), 2.31 (s, 6H), 3.01 (d, $J = 15.3$ Hz, 1H), 3.07 (d, $J = 15.3$ Hz, 1H), 3.98-4.06 (m, 2H), 4.74 (d, $J = 6.4$ Hz, 1H), 6.19 (dd, $J = 1.8$ Hz, 3.4 Hz, 1H), 6.24 (dd, $J = 0.9$ Hz, 3.4 Hz, 1H), 7.04 (s, 1H), 7.05 (s, 1H), 7.19 (dd, $J = 0.9$ Hz, 1.8 Hz, 1H), 7.40 (d, $J = 13.1$ Hz, 2H), 7.43 (d, $J = 13.4$ Hz, 2H); ^{13}C NMR (CDCl_3) δ 13.58, 19.00, 19.02, 21.26, 26.31 (d, $J_{C-P} = 4.1$ Hz), 30.46, 46.56 (d, $J_{C-P} = 3.0$ Hz), 54.67 (d, $J_{C-P} = 2.1$ Hz), 64.39, 105.76, 105.77, 110.11, 128.76, 128.84, 129.50, 129.57, 133.02, 133.04, 133.11, 133.13, 133.53, 133.89, 134.54, 134.89, 137.70, 137.80, 137.88, 137.99, 141.14, 157.66, 157.72, 171.55; ^{31}P NMR (CDCl_3) δ 19.9; $[\alpha]_D^{27}$ -10.9 ($c = 0.76$, CHCl_3) (95% ee); HPLC DAICEL CHIRALCEL AD-H, hexane/2-propanol = 20/1, 1.0 mL/min, t_R 15.5 min (major), 21.9 min (minor); MS m/z 504 ($\text{M}+\text{Na}^+$); HRMS Calcd for $\text{C}_{28}\text{H}_{36}\text{NO}_4\text{P}^+$ (M^+) 481.2376, Found 481.2379; IR (neat) 1726, 1194 cm^{-1} .

***N*-[2-butoxycarbonyl-1-methyl-1-(3-thienyl)phenylethyl]-di(3,5-xylyl)phosphinamide (5f)**

^1H NMR (CDCl_3) δ 0.84 (t, $J = 7.3$ Hz, 3H), 1.19-1.27 (m, 2H), 1.43-1.49 (m, 2H), 1.57 (s, 3H), 2.30 (s, 6H), 2.32 (s, 6H), 3.00 (d, $J = 15.9$ Hz, 1H), 3.21 (d, $J = 15.9$ Hz, 1H), 3.93-4.01 (m, 2H), 4.91 (d, $J = 5.5$ Hz, 1H), 7.07 (bs, 2H), 7.13 (dd, $J = 1.3$ Hz, $J = 5.2$ Hz, 1H), 7.20 (dd, $J = 3.1$ Hz, $J = 5.2$ Hz, 1H), 7.25 (dd, $J = 1.3$ Hz, $J = 3.1$ Hz, 1H), 7.44 (d, $J = 11.8$ Hz, 2H), 7.50 (d, $J = 12.2$ Hz, 2H); ^{13}C NMR (CDCl_3) δ 13.57, 13.59, 18.99, 19.00, 21.29, 29.22 (d, $J_{C-P} = 5.2$ Hz), 30.44 (d, $J_{C-P} = 2.0$ Hz), 48.14, 56.71, 56.72, 56.74, 64.30, 120.34, 125.60, 125.98, 128.81, 128.89, 129.43, 129.50, 133.11, 134.29, 135.06, 135.29, 137.84, 137.94, 138.05, 148.54, 148.58, 171.92; ^{31}P NMR (CDCl_3) δ 19.6; $[\alpha]_D^{27}$ -19.5 ($c = 1.09$, CHCl_3) (96% ee); HPLC DAICEL CHIRALCEL AD-H, hexane/2-propanol = 20/1, 1.0 mL/min, t_R 8.5 min (major), 14.2 min (minor); MS m/z 520 ($\text{M}+\text{Na}^+$); HRMS Calcd for $\text{C}_{28}\text{H}_{36}\text{NO}_3\text{PS}^+$ (M^+) 497.2148, Found 497.2154; IR (neat) 1723, 1197 cm^{-1} .

***N*-(2-butoxycarbonyl-1-ethyl-1-phenylethyl)-di(3,5-xylyl)phosphinamide (5g)**

^1H NMR (CDCl_3) δ 0.46 (t, $J = 7.3$ Hz, 3H), 0.86 (t, $J = 7.4$ Hz, 3H), 1.22-1.31 (m, 2H), 1.45-1.53 (m, 2H), 1.88-2.04 (m, 2H), 2.31 (s, 6H), 2.34 (s, 6H), 3.26 (d, $J = 16.8$ Hz, 1H), 3.31 (d, $J = 16.8$ Hz, 1H), 3.96-4.05 (m, 2H), 4.82 (d, $J = 5.2$ Hz, 1H), 7.06 (s, 1H), 7.09 (s, 1H), 7.19 (t, $J = 7.6$ Hz, 1H), 7.30 (t, $J = 7.6$ Hz, 2H), 7.44 (d, $J = 12.2$ Hz, 2H), 7.51 (d, $J = 7.6$ Hz, 2H), 7.58 (d, $J = 12.2$ Hz, 2H); ^{13}C NMR (CDCl_3) δ 8.81, 13.57, 19.02, 21.28, 21.31, 30.48, 34.02 (d, $J_{C-P} = 4.1$ Hz), 42.67 (d, $J_{C-P} = 2.1$ Hz), 62.76 (d, $J_{C-P} = 3.1$ Hz), 64.21, 126.35, 126.74, 127.96, 128.76, 128.83, 129.39, 129.46, 133.06, 134.41, 134.60, 135.44, 135.58, 137.80, 137.91, 138.01, 144.35, 144.41, 172.50; ^{31}P NMR (CDCl_3) δ 20.0; $[\alpha]_D^{22}$ +6.3 ($c = 0.98$, CHCl_3) (91% ee); HPLC DAICEL CHIRALCEL AD-H, hexane/2-propanol = 20/1, 1.0 mL/min, t_R 14.4 min (major),

21.8 min (minor); MS m/z 528 ($M+Na^+$); HRMS Calcd for $C_{31}H_{40}NO_3P^+$ (M^+) 505.2740, Found 505.2736; IR (neat) 2960, 1725, 1198 cm^{-1} .

***N*-(2-butoxycarbonyl-1-cyclohex-1-enyl-1-methylethyl)-di(3,5-xylyl)phosphinamide (5h)**

1H NMR ($CDCl_3$) δ 0.88(t, $J = 7.3$ Hz, 3H), 1.28-1.35 (m, 5H), 1.44-1.58 (m, 6H), 1.96-2.08 (m, 4H), 2.31 (s, 6H), 2.34 (s, 6H), 2.76 (d, $J = 15.3$ Hz, 1H), 2.97 (d, $J = 15.3$ Hz, 1H), 3.99-4.08 (m, 2H), 4.37 (d, $J = 5.8$ Hz, 1H), 5.85-5.90 (m, 1H), 7.05 (s, 1H), 7.06 (s, 1H), 7.44 (d, $J = 12.2$ Hz, 2H), 7.52 (d, $J = 12.2$ Hz, 2H); ^{13}C NMR ($CDCl_3$) δ 13.60, 19.08, 21.28, 22.03, 22.94, 24.71, 25.29, 26.03 (d, $J_{C-P} = 4.1$ Hz), 30.58, 45.84 (d, $J_{C-P} = 2.1$), 59.33 (d, $J_{C-P} = 2.1$ Hz), 64.10, 121.62, 128.77, 128.84, 129.51, 129.59, 132.92, 132.94, 139.96, 132.98, 134.44, 134.71, 135.47, 135.71, 137.71, 137.81, 137.86, 137.97, 141.01, 141.06, 172.29; ^{31}P NMR ($CDCl_3$) δ 19.3; $[\alpha]_D^{27} +2.1$ ($c = 1.06$, $CHCl_3$) (81% ee, using (*R,R*)-**8**); HPLC DAICEL CHIRALCEL AD-H, hexane/2-propanol 20/1, 1.0 mL/min, t_R 10.1 min (minor), 15.5 min (major); MS m/z 518 ($M+Na^+$); HRMS Calcd for $C_{30}H_{42}NO_3P^+$ (M^+) 495.2897, Found 495.2903; IR (neat) 2929, 1725, 1199 cm^{-1} .

***N*-[1-(2-butoxycarbonyl)ethyl-1-methyl-2-heptenyl]-di(3,5-xylyl)phosphinamide (5i)**

1H NMR ($CDCl_3$) δ 0.81(t, $J = 7.3$ Hz, 3H), 0.88 (t, $J = 7.4$ Hz, 3H), 1.20-1.35 (m, 4H), 1.36 (s, 3H), 1.51-1.59 (m, 2H), 1.81-1.88 (m, 2H), 2.30 (s, 6H), 2.31 (s, 6H), 2.67 (d, $J = 15.3$ Hz, 1H), 2.78 (d, $J = 15.3$ Hz, 1H), 4.01- 4.10 (m, 2H), 4.44 (d, $J = 6.1$ Hz, 1H), 5.51-5.60 (m, 2H), 7.45 (s, 2H), 7.44 (d, $J = 12.2$ Hz, 2H), 7.49 (d, $J = 12.2$ Hz, 2H); ^{13}C NMR ($CDCl_3$) δ 13.53, 13.60, 19.09, 21.25, 22.09, 27.31 (d, $J_{C-P} = 4.1$ Hz), 30.53, 34.06, 47.30 (d, $J_{C-P} = 3.1$ Hz), 55.99 (d, $J_{C-P} = 3.1$ Hz), 64.23, 128.88, 128.98, 129.05, 129.45, 129.52, 132.89, 132.90, 132.93, 134.43, 134.55, 135.46, 135.56, 135.60, 135.64, 137.70, 137.76, 137.80, 137.87, 172.13; ^{31}P NMR ($CDCl_3$) δ 19.5; $[\alpha]_D^{24} +4.7$ ($c = 1.00$, $CHCl_3$) (77% ee, using (*R,R*)-**8**); HPLC DAICEL CHIRALCEL AD-H, hexane/2-propanol 50/1, 1.0 mL/min, t_R 31.4 min (minor), 33.6 min (major); MS m/z 506 ($M+Na^+$); HRMS Calcd for $C_{29}H_{42}NO_3P^+$ (M^+) 483.2897, Found 483.2895; IR (neat) 2958, 1725, 1197 cm^{-1} .

***N*-[2-butoxycarbonyl-1-cyclohexyl-1-methylethyl]-di(3,5-xylyl)phosphinamide (5j)**

1H NMR ($CDCl_3$) δ 0.90(t, $J = 7.4$, 3H), 0.92-1.25 (m, 5H), 1.08 (s, 3H), 1.30-1.39 (m, 2H), 1.53-1.85 (m, 7H), 2.14 (d, $J = 12.5$, 1H), 2.30 (s, 6H), 2.32 (s, 6H), 2.56 (d, $J = 15.6$ Hz, 1H), 2.94 (1H, $J = 15.6$ Hz, 1H), 3.98 (d, $J = 5.2$ Hz, 1H), 4.02-4.15 (m, 2H), 7.05 (s, 1H), 7.07 (s, 1H), 7.43 (d, $J = 12.2$ Hz, 2H), 7.52 (d, $J = 12.2$ Hz, 2H); ^{13}C NMR ($CDCl_3$) δ 13.60, 19.10, 21.25, 21.63, 21.66, 26.41, 26.65, 26.77, 27.33, 27.59, 30.58, 44.68 (d, $J_{C-P} = 2.1$ Hz), 47.06 (d, $J_{C-P} = 6.3$ Hz), 58.98 (d, $J_{C-P} = 2.1$ Hz), 64.05, 128.76, 128.84, 129.47, 129.55, 132.89, 132.93, 132.95, 134.52, 134.83, 135.55, 135.82, 137.68, 137.79, 137.83, 137.94, 172.52; ^{31}P NMR ($CDCl_3$) δ 19.4; $[\alpha]_D^{25} +12.6$ ($c = 1.14$, $CHCl_3$) (74% ee, using (*S,S*)-*i*-Pr-Duphos **7**); HPLC DAICEL CHIRALCEL AD-H, hexane/2-propanol 20/1, 1.0 mL/min, t_R 10.3 min (major), 12.8 min (minor); MS m/z 520 ($M+Na^+$); HRMS Calcd for $C_{29}H_{41}NO_3P$ ($M-Me^+$) 482.2819, Found 482.2823; IR (neat) 2927, 1720, 1192 cm^{-1} .

Preparation of Diacetoxidiethoxysilane.

Tetraacetoxysilane (20 g, 76 mmol) was dried under vacuum and suspended in THF (50 mL). To the suspension, EtOH (8.4 mL, 144 mmol) was slowly added over 3 h, and the mixture was stirred further at room temperature overnight. THF and AcOH were removed under reduced pressure (80 mmHg, 1 h and 12 mmHg, 1 h). Purification by distillation under reduced pressure (13 mmHg, 85-92 °C) gave 6.4 g of product ((EtO)₂Si(OAc)₂:(EtO)₃SiOAc = 3:1). ¹H NMR (C₆D₆) δ 1.15 (t, *J* = 7.0 Hz, 6H), 1.67 (s, 6H), 4.02 (q, *J* = 7.0 Hz, 4H); ¹³C NMR (C₆D₆) δ 17.90, 21.78, 60.84, 168.89.

Preparation of DuPHOS Derivative (8)

DuPHOS derivative **8** was synthesized following the reported procedures.²

¹H NMR (CDCl₃) δ 0.52-0.70 (m, 5H), 0.73 (s, 18H), 0.81 (s, 18H), 0.85-1.10 (m, 16H), 1.21-1.28 (m, 2H), 1.40-1.85 (m, 19H), 1.96-2.08 (m, 5H), 2.17-2.28 (m, 2H), 2.36-2.47 (m, 2H), 7.24-7.28 (m, 2H), 7.45-7.55 (m, 2H); ¹³C NMR (CDCl₃) δ 27.22, 27.41, 27.51, 27.56, 27.69, 29.62, 30.62, 30.95, 31.00, 31.04, 31.96, 32.24, 32.34, 34.44, 34.49, 34.54, 35.73, 35.78, 35.83, 38.22, 42.43, 42.54, 42.65, 48.05, 48.29, 48.34, 49.32, 49.38, 49.43, 127.46, 133.66, 145.84; ³¹P NMR (CDCl₃) δ -12.5; [α]²³_D +65.9 (c = 0.99, CHCl₃) MS *m/z* 803 (M+H⁺); HRMS Calcd for C₅₄H₉₃P₂ (M+H⁺) 803.6747, Found 803.6730; IR 2934, 1364, 1098 cm⁻¹.

Conversion to β-Amino Acid (9) (Scheme 1)

The Mannich adduct **5a** (71 mg, 0.14 mmol, 95% ee) was treated with 3 N HCl (THF:H₂O = 3:1, 1 mL) at 40 °C for 2.5 h. After cooling, volatiles were evaporated and dried under vacuum. The crude mixture was treated with 0.5 N NaOH (2-propanol:H₂O = 1:1, 2 mL) at room temperature for 20 h. The reaction mixture was neutralized with aqueous 1 N HCl and volatiles were evaporated. The residue was purified through ion exchange chromatography [Dowex 50Wx8-100 (acidic) (2 g), loading and eluting with MeOH to get rid of Xy₂P(O)OH, and then eluting with 25% NH₃ aq.] to give β-amino acid **9** as a white solid in 81% yield (20.1 mg). Based on the sign of the optical rotation, the absolute configuration was determined to be (*S*). [α]²⁴_D = +14.7 (c = 1.00, CH₃OH), lit. [α]²²_D = -10.2 (c = 0.78, CH₃OH) for (*R*) isomer (100% ee).³ ¹H NMR (CD₃OD) δ 1.69 (s, 3H), 2.72 (d, *J* = 16.5 Hz, 1H), 2.82 (d, *J* = 16.5 Hz, 1H), 7.31-7.35 (m, 1H), 7.39-7.48 (m, 4H); ¹³C NMR (CD₃OD) δ 27.15, 46.24, 58.07, 125.64, 129.18, 130.02, 142.98, 177.21; MS *m/z* 202 (M+Na⁺); HRMS Calcd for C₁₀H₁₃NO₂ (M⁺) C₁₀H₁₃NO₂ 179.0941, Found 179.0940; IR (KBr) 3398, 2980, 1573, 1473, 1390 cm⁻¹.

Preliminary Ligand Screening (Table S-1)

Table S-1. Effects of Chiral Ligand and Cu(I) Sources

entry	Cu(I)	ligand	yield (%) ^a	ee (%) ^b
1	CuOAc	(<i>R</i>)-SEGPPOS	36	46
2	CuOAc	(<i>R</i>)-DTBM-SEGPPOS	54	94
3	CuOAc	(<i>S,S</i>)- <i>i</i> Pr-Duphos	97	66
4	CuOAc	Taniaphos-Ph	11	N.D.
5	CuOAc	Taniaphos-Cy	11	N.D.
6	CuF•3PPh ₃ •2EtOH	(<i>R</i>)-DTBM-SEGPPOS	60	56
7	CuF ^c	(<i>R</i>)-DTBM-SEGPPOS	30	6

^a Determined from ¹H NMR. ^b Determined by chiral HPLC

^c Prepared by reducing CuF₂•H₂O with 2 equiv of chiral phosphine to Cu in situ.

(*R*)-SEGPPOS; R¹ = X = H

(*R*)-DTBM-SEGPPOS; R¹ = *t*Bu, X = OMe

(*S,S*)-*i*Pr-Duphos

Taniaphos-Ph; R² = Ph
Taniaphos-Cy; R² = Cy

Comparison between *N*-Diphenylphosphinoylimines and *N*-Di(3,5-xylyl)phosphinoylimines (Table S-2)

Table S-2. Effects of *N*-Phosphinoyl Moiety

entry	R	substrate Ar		yield ^a (%)	ee ^b (%)
1		Ph	1a	69	94
2		3,5-Xy	2a	81	95
3		Ph	1b	73	93
4		3,5-Xy	2b	82	97
5 ^c		Ph	1c	59	94
6 ^c		3,5-Xy	2c	87	97
7		Ph	1d	82	92
8		3,5-Xy	2d	74	96
9		Ph	1f	69	93
10		3,5-Xy	2f	92	97

^a Isolated yield. ^b Determined by chiral HPLC.

^c 4 equiv of **3** was used.

Proposed Catalytic Cycle (Scheme S-1)

Scheme S-1. Proposed Catalytic Cycle

On the basis of the reaction mechanism of the catalytic asymmetric aldol reaction of ketones,⁴ we proposed the catalytic cycle of the present reaction as shown in Scheme S-1. The additive $[(\text{EtO})_2\text{Si}(\text{OAc})_2]$ in the aromatic ketoimine case and $(\text{EtO})_3\text{SiF}$ in the aliphatic ketoimine case] should trap the intermediate copper amide (**A**), and facilitate the presumable turnover-limiting catalyst regeneration step.

References

- ¹ Krzyzanowsk, B.; Stec, W. J. *Synthesis*, **1982**, 270.
- ² Wada, R.; Shibuguchi, T.; Makino, S.; Oisaki, K.; Kanai, M.; Shibasaki, M. *J. Am. Chem. Soc.* **2006**, 128, 7687.
- ³ Hua, D. H.; Miao, S. W.; Chen, J. S.; Iguchi, S. *J. Org. Chem.* **1991**, 56, 4.
- ⁴ Oisaki, K.; Zhao, D.; Kanai, M.; Shibasaki, M. *J. Am. Chem. Soc.* **2006**, 128, 7164.