Catalytic Enantioselective Negishi Reactions of Racemic Secondary Benzylic Halides

Forrest O. Arp and Gregory C. Fu*

Department of Chemistry, Massachusetts Institute of Technology,
Cambridge, Massachusetts 02139

Supporting Information

I. General

N,N-Dimethylacetamide (DMA; anhydrous in a Sure-Seal[®] bottle; Fluka), NiBr₂·diglyme (Aldrich), (*R*)-(*i*-Pr)-Pybox (Aldrich), (*S*)-(*i*-Pr)-Pybox (Fluka), and pyridine (anhydrous in a Sure-Seal[®] bottle; Fluka) were used as received. CH₂Cl₂ was dried by passage through a neutral alumina column under argon pressure prior to use. All other chemicals were reagent-grade and used as received.

Note: Although NiBr₂·diglyme and (*i*-Pr)-Pybox can be left in the air without any impact on cross-coupling efficiency, we recommend that, for long-term storage, they be kept under an inert atmosphere.

HPLC analyses were carried out on an Agilent 1100 Series system with Daicel Chiralpak[®] columns in hexanes/iso-propanol mixtures. GC analyses were performed on a Hewlett-Packard HP 6850 Series apparatus with a Chrompack capillary column 'CP Chirasil-Dex CB' (25 m x 0.25 mm x 0.25 mm), unless otherwise noted.

II. Preparation of Substrates

The yields have not been optimized.

NaBH₄ Reduction Procedure. A 250-mL round-bottomed flask was charged with the indanone and a mixture of 1,2-dichloroethane/methanol (1:1; 50 mL). NaBH₄ (2 equiv) was then added to the stirred mixture, and the resulting clear, colorless solution was allowed to stir open to the air for 30 min. Then, the reaction mixture was transferred to a separatory funnel with the aid of water (50 mL), and the aqueous layer was extracted with CH₂Cl₂ (3 x 50 mL). The combined organic layers were dried (Na₂SO₄) and concentrated under vacuum to furnish the product as either a white solid or a clear, colorless oil.

5-Chloro-1-indanol [33781-38-3]. This compound was reduced according to the

Reduction Procedure using 5-chloro-1-indanone (2.08 g, 12.5 mmol) and NaBH₄ (946 mg, 25.0 mmol). The product was obtained as a white solid (2.01 g, 95%).

¹H NMR (CDCl₃, 400 MHz): δ 7.33-7.20 (m, 1H), 7.23-7.20 (m, 2H), 5.20 (t, *J* = 6.0 Hz, 1H), 3.07-2.99 (m, 1H), 2.84-2.76 (m, 1H), 2.54-2.46 (m, 1H), 2.00-1.91 (m, 2H);

¹³C NMR (CDCl₃, 100 MHz): δ 145.3, 143.4, 134.0, 126.9, 125.3, 125.0, 75.7, 36.1, 29.6.

6-Methyl-1-indanol [200425-63-4]. This compound was reduced according to the Reduction Procedure using 6-methyl-1-indanone (0.989 g, 6.77 mmol) and NaBH₄ (500 mg, 13 mmol). The product was obtained as a clear, colorless oil (0.947 mg, 94%).

¹H NMR (CDCl₃, 400 MHz): δ 7.25 (s, 1H), 7.17-7.15 (m, 1H), 7.11-7.09 (m, 1H), 5.24-5.19 (m, 1H), 3.06-2.99 (m, 1H), 2.83-2.75 (m, 1H), 2.53-2.45 (m, 1H), 2.38 (s, 3H), 1.99-1.89 (m, 2H);

¹³C NMR (CDCl₃, 100 MHz): δ 145.1, 140.2, 136.3, 129.1, 124.7, 124.6, 76.4, 36.2, 29.3, 21.2.

5-Cyano-1-indanol [125114-88-7]. This compound was reduced according to the Reduction Procedure using 5-cyano-1-indanone¹ (540 mg, 3.44 mmol) and NaBH₄ (260 mg, 6.87 mmol). The product was obtained as a white solid (498 mg, 91%).

¹H NMR (CDCl₃, 500 MHz): δ 7.55-7.51 (m, 3H), 5.31-5.26 (m, 1H), 3.10-3.04 (m, 1H), 2.89-2.83 (m, 1H), 2.60-2.53 (m, 1H), 2.09 (br s, 1H), 2.02-1.95 (m, 1H);

¹³C NMR (CDCl₃, 125 MHz): δ 150.3, 144.1, 130.7, 128.4, 124.9, 119.1, 111.3, 75.5, 35.5, 29.4.

1-Benz[flindanol [123332-18-3]. This compound was reduced according to the Reduction Procedure using benz[flindanone² (499 mg, 2.74 mmol) and NaBH₄ (200 mg, 5.3 mmol). The product was obtained as a white solid (503 mg, 100%).

¹H NMR (CDCl₃, 500 MHz): δ 7.87-7.84 (m, 2H), 7.81-7.79 (m, 1H), 7.70 (s, 1H), 7.48-7.41 (m, 2H), 5.39 (dd, *J* = 6.0, 6.1 Hz, 1H), 3.26-3.19 (m, 1H), 3.02-2.94 (m, 1H), 2.60-2.52 (m, 1H), 2.09-2.01 (m, 1H), 1.81 (d, *J* = 6.6 Hz, 1H);

¹³C NMR (CDCl₃, 125 MHz): δ 144.2, 141.5, 133.9, 132.9, 128.1, 127.6, 125.8, 125.2,

122.9, 122.6, 75.9, 36.7, 29.3.

6-Methoxy-1-indanol [3469-09-8]. This compound was reduced according to the Reduction Procedure using 6-methoxy-1-indanone (2.23 g, 13.7 mmol) and NaBH₄ (1.04 g, 27.5 mmol). The product was obtained as a white solid (2.18 g, 97%).

¹H NMR (CDCl₃, 500 MHz): δ 7.15 (d, *J* = 8.2 Hz, 1H), 6.97 (d, *J* = 2.4 Hz, 1H), 6.83 (dd, *J* = 2.5, 8.3 Hz, 1H), 5.22-5.21 (m, 1H), 3.82 (s, 3H), 3.02-2.95 (m, 1H), 2.80-2.72 (m, 1H), 2.57-2.48 (m, 1H), 1.99-1.91 (m, 1H), 1.81 (br s, 1H).

¹³C NMR (CDCl₃, 125 MHz): δ 159.0, 146.3, 135.0, 125.5, 115.0, 108.7, 76.6, 55.5, 36.6, 28.9.

PBr₃/PCl₃ Halogenation Procedure. A dry 250-mL round-bottomed flask was charged with the benzylic alcohol and then placed under a nitrogen atmosphere. Dry CH₂Cl₂ (~50 mL) and dry pyridine (0.1 mL) were added, and then the reaction mixture was cooled to -10 °C in an *iso*-propanol/ice bath. PBr₃ (0.5 equiv) was dissolved separately in dry CH₂Cl₂ (5 mL), and the resulting solution was added portion-wise over a period of 10 min to the solution of the alcohol. The mixture was allowed to stir for an additional 20 min, and then the solution was transferred to a separatory funnel with the aid of 5% aqueous Na₂CO₃ (50 mL) and extracted with CH₂Cl₂ (3 × 50 mL). The combined organic layers were dried (Na₂SO₄) and concentrated under vacuum to give a yellow residue that was taken up in ether (10 mL), passed through an Acrodisc[®], and then concentrated, furnishing the desired benzylic halide as either a clear, colorless oil or a white solid.

1-Bromo-5-chloroindane [192702-71-9]. This compound was prepared according to the Halogenation Procedure using 5-chloro-1-indanol (2.01 g, 11.9 mmol), PBr₃ (1.61 g, 5.95 mmol), and dry pyridine (0.1 mL). The product was obtained as a clear, light-pink oil (1.57 g, 57%).

¹H NMR (CDCl₃, 500 MHz): δ 7.36-7.34 (m, 1H), 7.26 (s, 1H), 7.23-7.21 (m, 1H), 5.54 (dd, *J* = 2.2, 6.2 Hz, 1H), 3.25-3.23 (m, 1H), 2.88 (ddd, *J* = 2.6, 7.6, 16.3 Hz, 1H), 2.67-2.59 (m, 1H), 2.58-2.50 (m, 1H);

¹³C NMR (CDCl₃, 125 MHz): δ 145.5, 142.4, 134.7, 127.4, 126.1, 125.0, 53.4, 37.8, 30.5.

6-Methyl-1-bromoindane. This compound was prepared according to the Halogenation Procedure using 6-methyl-1-indanol (0.920 g, 6.21 mmol), PBr₃ (870 mg, 3.2 mmol), and dry pyridine (0.1 mL). The product was obtained as a clear, colorless oil (1.03 g, 79%).

¹H NMR (CDCl₃, 400 MHz): δ 7.26 (s, 1H), 7.18-7.16 (m, 1H), 7.10-7.08 (m, 1H), 5.58 (dd, *J* = 2.3, 6.0 Hz, 1H), 3.20-3.12 (m, 1H), 2.89-2.82 (m, 1H), 2.64-2.50 (m, 2H), 2.37 (s, 3H);

¹³C NMR (CDCl₃, 100 MHz): δ 143.9, 140.6, 136.8, 129.9, 125.5, 124.5, 55.2, 37.9, 30.6, 21.2;

IR (film) 3011, 2976, 2945, 2916, 2846, 1493, 1442, 1184, 1162, 861, 812 cm⁻¹;

EIMS (70 eV) *m/z*: M⁺ 210, 208, 130, 115, 102, 89.

1-Bromoindane [24373-98-6]. This compound was prepared according to the Halogenation Procedure using 1-indanol (4.00 g, 29.8 mmol), PBr₃ (4.03 g, 14.9 mmol), and dry pyridine (0.1 mL). The product was obtained as a clear, colorless oil (5.47 g, 93%) that contained ~5% 1-indene according to ¹H NMR.

¹H NMR (CDCl₃, 500 MHz): δ 7.48-7.46 (m, 1H), 7.31-7.25 (m, 3H), 5.63 (dd, *J* = 2.4, 6.1 Hz, 1H), 3.27-3.19 (m, 1H), 2.96-2.89 (m, 1H), 2.67-2.53 (m, 2H);

¹³C NMR (CDCl₃, 125 MHz): δ 143.8, 143.6, 128.8, 127.1, 125.1, 124.8, 54.8, 37.7, 30.6.

1-Bromo-5-cyanoindane [475475-78-6]. This compound was prepared according to the Halogenation Procedure using 5-cyano-1-indanol (498 mg, 3.13 mmol), PBr₃ (425 mg, 1.57 mmol), and dry pyridine (0.1 mL). The product was obtained as a white solid (472 mg, 68%).

¹H NMR (CDCl₃, 500 MHz): δ 7.56-7.51 (m, 3H), 5.51 (dd, *J* = 2.5, 6.5 Hz, 1H), 3.25-3.17 (m, 1H), 2.98-2.92 (m, 1H), 2.70-2.60 (m, 1H), 2.58-2.51 (m, 1H);

¹³C NMR (CDCl₃, 125 MHz): δ 148.8, 144.3, 131.0, 128.4, 125.7, 118.7, 112.1, 51.8, 37.1, 30.2.

1-Bromoacenaphthene [24171-73-1]. This compound was prepared according to the Halogenation Procedure using 1-acenaphthenol (1.00 g, 5.88 mmol), PBr_3 (796 mg, 2.94 mmol), and dry pyridine (0.1 mL). The resulting yellow solid was further purified by recrystallization from benzene/hexanes (1:5; 6 mL) to give the desired product as a lustrous, white crystalline solid (0.925 g, 68%).

^1H NMR (CDCl_3 , 500 MHz): δ 7.77-7.75 (m, 1H), 7.70-7.69 (m, 1H), 7.60-7.56 (m, 2H), 7.55-7.52 (m, 1H), 7.36-7.34 (m, 1H), 5.95 (dd, J = 2.1, 7.3 Hz, 1H), 4.14 (dd, J = 7.3, 18.4 Hz, 1H), 3.84 (d, J = 18.5 Hz, 1H);

^{13}C NMR (CDCl_3 , 125 MHz): δ 144.7, 140.6, 136.5, 131.1, 128.3, 128.2, 125.2, 123.1, 121.6, 120.0, 47.6, 43.6.

1-Bromobenz[f]indane [146000-51-3]. This compound was prepared according to the Halogenation Procedure using 1-benz[f]indanol (503 mg, 2.73 mmol), PBr_3 (371 mg, 1.37 mmol), and dry pyridine (0.1 mL). The product was obtained as a white solid (598 mg, 89%).

^1H NMR (CDCl_3 , 500 MHz): δ 7.92 (s, 1H), 7.85-7.79 (m, 2H), 7.72 (s, 1H), 7.49-7.42 (m, 2H), 5.74 (dd, J = 3.1, 5.1 Hz, 1H), 3.42-3.33 (m, 1H), 3.10-3.03 (m, 1H), 2.65-2.59 (m, 2H);

^{13}C NMR (CDCl_3 , 125 MHz): δ 142.8, 141.3, 134.0, 132.9, 128.3, 127.6, 126.3, 125.4, 124.0, 123.0, 53.9, 38.4, 30.1.

1-(1-Bromoethyl)-4-methylbenzene [69150-78-3]. This compound was prepared according to the Halogenation Procedure using 1-(4-methylphenyl)ethanol (1.50 g, 11.0 mmol), PBr_3 (1.49 g, 5.50 mmol), and dry pyridine (0.1 mL). The product was obtained as a clear, colorless oil (1.93 g, 88%).

^1H NMR (CDCl_3 , 400 MHz): δ 7.35 (d, J = 8.2 Hz, 2H), 7.16 (d, J = 7.9 Hz, 2H), 5.23 (q, J = 6.9 Hz, 1H), 2.35 (s, 3H), 2.05 (d, J = 6.9 Hz, 3H);

^{13}C NMR (CDCl_3 , 100 MHz): δ 140.3, 138.3, 129.3, 126.7, 49.8, 26.8, 21.2.

1-Chloroindane [35275-62-8]. This compound was prepared according to the Halogenation Procedure using 1-indanol (1.00 g, 7.45 mmol), PCl_3 (512 mg, 3.73 mmol), and dry pyridine (0.1 mL). The product was obtained as a clear, colorless oil (0.866 g, 76%).

^1H NMR (CDCl_3 , 500 MHz): δ 7.44-7.43 (m, 1H), 7.29-7.24 (m, 3H), 5.44 (dd, $J = 3.3$, 6.7 Hz, 1H), 3.23-3.16 (m, 1H), 2.93-2.87 (m, 1H), 2.64-2.57 (m, 1H), 2.42-2.36 (m, 1H);

^{13}C NMR (CDCl_3 , 125 MHz): δ 143.3, 143.0, 128.8, 127.0, 124.9, 124.7, 63.2, 36.9, 30.3.

1-Chloro-6-methoxyindane [128226-43-7]. This compound was prepared according to the Halogenation Procedure using 6-methoxy-1-indanol (651 mg, 3.96 mmol), PCl_3 (272 mg, 1.98 mmol), and dry pyridine (0.1 mL). The product was obtained as a clear, colorless oil (510 mg, 70%).

^1H NMR (CDCl_3 , 500 MHz): δ 7.18-7.16 (m, 1H), 6.97-6.96 (m, 1H), 6.86-6.84 (m, 1H), 5.40 (dd, $J = 3.4$, 6.7 Hz, 1H), 3.82 (s, 3H), 3.15-3.09 (m, 1H), 2.87-2.81 (m, 1H), 2.67-2.59 (m, 1H), 2.42-2.37 (m, 1H);

^{13}C NMR (CDCl_3 , 125 MHz): δ 159.1, 144.3, 135.2, 125.4, 115.8, 109.3, 63.4, 55.5, 37.4, 29.5.

Preparation of the Organozinc Reagents.³ A 25-mL Schlenk tube was charged with zinc powder (1.47 g, 22.5 mmol) and heated to 70 °C under high vacuum for 30 min. After back-filling with argon, iodine (0.19 g, 0.75 mmol) and DMA (to give a total volume of 10 mL) were added, and the resulting heterogeneous red mixture was allowed to stir until the red color of the iodine had faded (typically 1-2 min). Then, the alkyl halide (16 mmol; freshly distilled) was added. The colorless reaction mixture was allowed to stir for 12 h at 70 °C (exceptions: For MeI, the reaction was *not* heated; for cyclohexylmethylbromide, the reaction was heated for 24 h at 85 °C; for EtBr the reaction was *not* heated and it was allowed to stir for 48 h), then the mixture was allowed to cool to room temperature (the disappearance of the starting material and the formation of the organozinc reagent can readily be monitored by no-D NMR). The gray solution (~1.6 M) was passed through an Acrodisc[®] and stored under argon in a dry container with a pierceable septum.

These organozinc solutions can be stored at room temperature for several weeks without deterioration.

III. Negishi Cross-Coupling Reactions

General Procedure. In the air (no special precautions are necessary), a 4-mL glass vial was charged with NiBr₂·diglyme (35.3 mg, 0.100 mmol), (*R*)-(*i*-Pr)-Pybox (39.2 mg, 0.130 mmol), and the benzylic halide (1.00 mmol). The vial was fitted with a septum cap and purged with argon for 15 min. DMA (1.75 mL) was then added, and the resulting orange mixture was allowed to equilibrate for 15 min under argon in an isopropanol bath cooled to 0 °C. The organozinc solution (~1.6 M in DMA; 1.0 mL, 1.6 mmol) was then added in a single portion to the heterogeneous mixture, which rapidly became a clear, red-brown solution. The reaction mixture was allowed to stir for 24 h at 0 °C. Then, the excess organozinc reagent was quenched by the addition of ethanol (0.3 mL), and the resulting orange liquid was purified directly by flash chromatography.

The ee was determined on Chirasil[®] DEX-CB or Daicel Chiralpak[®] columns.

The second runs were conducted with (*S*)-(*i*-Pr)-Pybox.

2-[2-(5-Chloroindan-1-yl)ethyl]-[1,3]dioxolane (eq 2; Table 1, entry 2). This compound was prepared according to the General Procedure using (±)-1-bromo-5-chloroindane (232 mg, 1.00 mmol) and the appropriate organozinc reagent (~1.6 M in DMA; 1.0 mL, 1.6 mmol). After purification by column chromatography (8% EtOAc in hexanes), the title compound was isolated as a clear, colorless oil.

1st run: 210 mg (83%; 91% ee); 2nd run: 204 mg (81%; 91% ee).

The ee's were determined via HPLC on an OD-H column (eluent: 0.5% isopropanol in hexanes; flow rate: 1.0 mL/min) with t_r (major): 16.3 min, t_r (minor): 15.3 min.

[α]_D²³ = -2.0 (c = 0.83, CHCl₃);

¹H NMR (CDCl₃, 500 MHz): δ 7.18 (s, 1H), 7.12 (s, 2H), 4.90 (t, *J* = 4.7 Hz, 1H), 4.00-3.98 (m, 2H), 3.89-3.86 (m, 2H), 3.13-3.07 (m, 1H), 2.91 (ddd, *J* = 4.8, 8.6, 15.9 Hz, 1H), 2.85-2.78 (m, 1H), 2.34-2.27 (m, 1H), 1.98-1.92 (m, 1H), 1.82-1.67 (m, 3H), 1.58-1.49 (m, 1H);

¹³C NMR (CDCl₃, 125 MHz): δ 146.0, 145.6, 131.9, 126.1, 124.6, 124.6, 104.5, 64.9, 44.0, 32.1, 31.7, 31.2, 28.9;

IR (film) 2949, 2883, 2860, 1600, 1474, 1410, 1131, 1034, 875, 813 cm⁻¹;

EIMS (70 eV) *m/z*: M⁺ 252, 190, 164, 151, 129, 115, 73, 45.

1-Hexyl-6-methylindane (Table 1, entry 1). This compound was prepared according to the General Procedure using (±)-6-methyl-1-bromindane (211 mg, 1.00 mmol) and the appropriate organozinc reagent (~1.6 M in DMA; 1.0 mL, 1.6 mmol). After purification by column chromatography (pentane), the title compound was isolated as a clear, colorless oil.

1st run: 188 mg (87%; 96% ee); 2nd run: 196 mg (91%; 95% ee).

The ee's were determined by chiral GC: CP Chirasil-Dex CB, *t_r* (major) 33.8 min, *t_r* (minor) 33.3 min; heating program: 125 °C; flow: 1.0 mL/min.

$[\alpha]_D^{23} = -23$ (*c* = 0.99, CHCl₃);

¹H NMR (CDCl₃, 500 MHz): δ 7.13-7.11 (m, 1H), 7.04 (s, 1H), 6.99-6.97 (m, 1H), 3.10-3.05 (m, 1H), 2.93-2.75 (m, 2H), 2.36 (s, 3H), 2.33-2.25 (m, 1H), 1.89-1.82 (m, 1H), 1.72-1.63 (m, 1H), 1.48-1.29 (m, 9H), 0.93 (t, *J* = 6.8 Hz, 3H);

¹³C NMR (CDCl₃, 125 MHz): δ 148.1, 141.0, 135.5, 126.9, 124.3, 124.1, 44.8, 35.1, 32.5, 31.9, 31.0, 29.6, 27.7, 22.7, 21.4, 14.2;

IR (film) 3005, 2954, 2924, 2854, 1614, 1491, 1377, 1457, 877, 808, 723 cm⁻¹;

EIMS (70 eV) *m/z*: M⁺ 216, 131, 91.

4-Indan-1-yl-butyronitrile (Table 1, entry 3). This compound was prepared according to the General Procedure using (±)-1-bromindane (197 mg, 1.00 mmol) and the appropriate organozinc reagent (~1.6 M in DMA; 1.0 mL, 1.6 mmol). After purification by column chromatography (6% EtOAc in hexanes), the title compound was isolated as a clear, colorless oil.

1st run: 122 mg (66%; 92% ee); 2nd run: 116 mg (63%; 91% ee).

The ee's were determined via HPLC on an OD-H column (eluent: 1.0% isopropanol in hexanes; flow rate: 1.0 mL/min) with *t_r* (major): 20.8 min, *t_r* (minor): 26.7 min.

$[\alpha]_D^{23} = -4.2$ (*c* = 0.56, CHCl₃);

¹H NMR (CDCl₃, 500 MHz): δ 7.25-7.23 (m, 1H), 7.21-7.17 (m, 3H), 3.17-3.15 (m, 1H), 2.95-2.93 (m, 1H), 2.90-2.85 (m, 1H), 2.41 (t, *J* = 7.0 Hz, 2H), 2.34-2.29 (m, 1H), 2.02-1.98 (m, 1H), 1.83-1.58 (m, 4H);

¹³C NMR (CDCl₃, 125 MHz): δ 146.3, 143.8, 126.6, 126.2, 124.6, 123.5, 119.7, 44.0, 33.9, 31.8, 31.3, 23.5, 17.4;

IR (film) 3068, 3019, 2942, 2850, 2245, 1477, 1457, 1424, 747 cm⁻¹;

EIMS (70 eV) *m/z*: M⁺ 185, 117, 91.

1-(6-Chlorohexyl)indane (Table 1, entry 4) [361541-11-9]. This compound was prepared according to the General Procedure using ($\pm$)-1-bromoindane (197 mg, 1.00 mmol) and the appropriate organozinc reagent ($\sim$ 1.6 M in DMA; 1.0 mL, 1.6 mmol). After purification by column chromatography (pentane), the title compound was isolated as a clear, colorless oil.

1st run: 163 mg (69%; 95% ee); 2nd run: 166 mg (70%; 94% ee).

The ee's were determined via HPLC on an OD-H column (eluent: 0.5% isopropanol in hexanes; flow rate: 1.0 mL/min) with t_r (major): 5.1 min, t_r (minor): 5.6 min.

$[\alpha]_D^{23} = -11$ ($c = 0.79$, CHCl_3);

^1H NMR (CDCl_3 , 500 MHz): δ 7.24-7.14 (m, 4H), 3.56 (t, $J = 6.7$ Hz, 2H), 3.13-3.07 (m, 1H), 2.93 (ddd, $J = 4.6, 8.5, 15.5$ Hz, 1H), 2.87-2.81 (m, 1H), 2.32-2.26 (m, 1H), 1.88-1.78 (m, 3H), 1.72-1.64 (m, 1H), 1.51-1.34 (m, 7H);

^{13}C NMR (CDCl_3 , 125 MHz): δ 147.6, 144.0, 126.2, 125.9, 124.4, 123.5, 45.2, 44.8, 34.9, 32.6, 32.1, 31.4, 29.1, 27.5, 26.9;

IR (film) 2930, 2852, 2359, 2336, 1698, 1648, 1558, 1474, 1457, 1312, 752 cm^{-1} ;

EIMS (70 eV) m/z : M^+ 236, 128, 117, 91.

1-(6-Chlorohexyl)indan-5-carbonitrile (Table 1, entry 5). This compound was prepared according to the General Procedure using ($\pm$)-1-bromo-5-cyanoindane (222 mg, 1.00 mmol) and the appropriate organozinc reagent ($\sim$ 1.6 M in DMA; 1.0 mL, 1.6 mmol). After purification by column chromatography (5% EtOAc in hexanes), the title compound was isolated as a clear, colorless oil.

1st run: 128 mg (49%; 91% ee); 2nd run: 120 mg (46%; 90% ee).

The ee's were determined via HPLC on an AS-H column (eluent: 0.5% isopropanol in hexanes; flow rate: 1.0 mL/min) with t_r (major): 24.6 min, t_r (minor): 28.1 min.

$[\alpha]_D^{23} = +3.8$ ($c = 0.54$, CHCl_3);

^1H NMR (CDCl_3 , 500 MHz): δ 7.47-7.25 (m, 2H), 7.25 (s, 1H), 3.54 (t, $J = 6.7$ Hz, 2H), 3.15-3.07 (m, 1H), 2.98-2.92 (m, 1H), 2.88-2.82 (m, 1H), 2.35-2.28 (m, 1H), 1.85-1.68 (m, 4H), 1.49-1.34 (m, 7H);

^{13}C NMR (CDCl_3 , 125 MHz): δ 153.4, 145.2, 130.4, 128.0, 124.3, 119.6, 109.8, 45.1, 45.1, 34.4, 32.5, 31.8, 31.1, 29.0, 27.3, 26.8;

IR (film) 3380, 2930, 2856, 2226, 1608, 1483, 1442, 827 cm^{-1} ;

EIMS (70 eV) m/z : M^+ 261, 142, 115.

1-Butylacenaphthene (Table 1, entry 6). This compound was prepared according to the General Procedure using ($\pm$)-1-bromoacenaphthene (233 mg, 1.00 mmol) and the appropriate organozinc reagent ($\sim$ 1.6 M in DMA; 1.0 mL, 1.6 mmol). After purification by column chromatography (pentane), the title compound was isolated as a clear, colorless oil.

1st run: 148 mg (70%; 98% ee); 2nd run: 154 mg (73%; 98% ee).

The ee's were determined via HPLC on an OD-H column (eluent: 0.5% isopropanol in hexanes; flow rate: 0.7 mL/min) with t_r (major): 7.6 min, t_r (minor): 8.9 min.

$[\alpha]_D^{23} = -15$ ($c = 0.80$, CHCl_3);

^1H NMR (CDCl_3 , 500 MHz): δ 7.63-7.61 (m, 2H), 7.50-7.45 (m, 2H), 7.30-7.27 (m, 2H), 3.71-3.66 (m, 1H), 3.59 (dd, $J = 8.0, 17.1$ Hz, 1H), 3.08 (dd, $J = 3.4, 17.1$ Hz, 1H), 1.99-1.92 (m, 1H), 1.69-1.61 (m, 1H), 1.54-1.39 (m, 4H), 0.96 (t, $J = 7.1$ Hz, 3H);

^{13}C NMR (CDCl_3 , 125 MHz): δ 149.8, 144.8, 138.6, 131.4, 127.8, 127.7, 122.5, 122.2, 119.1, 118.7, 43.4, 37.5, 36.3, 29.7, 22.9, 14.1;

IR (film) 3036, 2955, 2926, 2856, 1604, 1496, 1466, 1368, 796, 774 cm^{-1} ;

EIMS (70 eV) m/z : M^+ 210, 167, 153.

1-Cyclohexylmethyl-acenaphthene (Table 1, entry 7). This compound was prepared according to the General Procedure at room temperature using ($\pm$)-1-bromoacenaphthene (233 mg, 1.00 mmol) and the appropriate organozinc reagent ($\sim$ 1.6 M in DMA; 1.0 mL, 1.6 mmol). After purification by column chromatography (pentane), the title compound was isolated as a clear, colorless oil.

1st run: 96 mg (38%; 97% ee); 2nd run: 98 mg (39%; 96% ee).

The ee's were determined via HPLC on an OD-H column (eluent: 1.0% isopropanol in hexanes; flow rate: 0.7 mL/min) with t_r (major): 8.6 min, t_r (minor): 9.7 min.

$[\alpha]_D^{23} = -24$ ($c = 0.60$, CHCl_3);

^1H NMR (CDCl_3 , 500 MHz): δ 7.62-7.61 (m, 2H), 7.49-7.44 (m, 2H), 7.28-7.25 (m, 2H), 3.81-3.76 (m, 1H), 3.59 (dd, $J = 7.9, 17.2$ Hz, 1H), 3.05 (dd, $J = 3.2, 17.5$ Hz, 1H), 1.96-1.93 (m, 1H), 1.82-1.72 (m, 5H), 1.66-1.50 (m, 2H), 1.38-1.19 (m, 3H), 1.10-0.99 (m, 2H);

^{13}C NMR (CDCl_3 , 125 MHz): δ 150.3, 144.8, 138.5, 131.5, 127.8, 127.7, 122.4, 122.2,

119.1, 118.7, 45.0, 40.6, 38.0, 35.9, 34.2, 32.9, 26.7, 26.5, 26.4;
IR (film) 3036, 2920, 2849, 1604, 1448, 1369, 795, 771 cm^{-1} ;
EIMS (70 eV) m/z : M^+ 250, 165, 153, 55.

2-(4-Acenaphthen-1-yl-butyl)isoindole-1,3-dione (Table 1, entry 8). This compound was prepared according to the General Procedure at room temperature using ($\pm$)-1-bromoacenaphthene (233 mg, 1.00 mmol), 0.75 mL (rather than 1.75 mL) of DMA, and the appropriate organozinc reagent (~ 0.80 M in DMA; 2.0 mL, 1.6 mmol). After purification by column chromatography (12% EtOAc in hexanes), the title compound was isolated as a clear, colorless oil, which solidified after a period of time.

1st run: 138 mg (39%; 99% ee); 2nd run: 151 mg (42%; 99% ee).

The ee's were determined via HPLC on an OD-H column (eluent: 10% isopropanol in hexanes; flow rate: 1.0 mL/min) with t_r (major): 18.0 min, t_r (minor): 16.9 min.

$[\alpha]_D^{23} = +6.0$ ($c = 0.45$, CHCl_3);

^1H NMR (CDCl_3 , 500 MHz): δ 7.87-7.85 (m, 2H), 7.74-7.72 (m, 2H), 7.61-7.59 (m, 2H), 7.47-7.43 (m, 2H), 7.27-7.26 (m, 2H), 3.73 (t, $J = 7.3$ Hz, 2H), 3.70-3.65 (m, 1H), 3.58 (dd, $J = 8.0, 17.1$ Hz, 1H), 3.05 (dd, $J = 3.4, 17.1$ Hz, 1H), 2.02-1.95 (m, 1H), 1.83-1.65 (m, 3H), 1.61-1.52 (m, 2H);

^{13}C NMR (CDCl_3 , 125 MHz): δ 168.4, 149.3, 144.6, 138.5, 133.9, 132.1, 131.4, 127.8, 127.7, 123.2, 122.6, 122.2, 119.1, 118.8, 43.3, 37.9, 37.5, 36.1, 28.8, 24.8;

IR (CDCl_3 solution) 3155, 2936, 2254, 1772, 1712, 1469, 1398, 1096 cm^{-1} ;

EIMS (70 eV) m/z : M^+ 355, 207, 165, 153, 133, 104, 71;

mp 80–83 $^\circ\text{C}$.

1-(3-Phenylpropyl)benz[f]indane (Table 1, entry 9). This compound was prepared according to the General Procedure using ($\pm$)-1-bromobenz[f]indane (247 mg, 1.00 mmol) and the appropriate organozinc reagent (~ 1.6 M in DMA; 1.0 mL, 1.6 mmol). After purification by column chromatography (2% EtOAc in hexanes), the title compound was isolated as a white solid.

1st run: 211 mg (74%; 97% ee); 2nd run: 227 mg (79%; 99% ee).

The ee's were determined via HPLC on an OD-H column (eluent: 2% isopropanol in hexanes; flow rate: 0.9 mL/min) with *t_r* (major): 11.8 min, *t_r* (minor): 10.8 min.

$[\alpha]_D^{23} = -47$ (*c* = 0.80, CHCl₃);

¹H NMR (CDCl₃, 500 MHz): δ 7.79-7.75 (m, 2H), 7.65 (s, 1H), 7.59 (s, 1H), 7.41-7.37 (m, 2H), 7.33-7.30 (m, 2H), 7.24-7.20 (m, 3H), 3.28-3.23 (m, 1H), 3.10-3.04 (m, 1H), 3.01-2.94 (m, 1H), 2.78-2.66 (m, 2H), 2.40-2.34 (m, 1H), 2.08-2.01 (m, 1H), 1.86-1.71 (m, 3H), 1.61-1.54 (m, 1H);

¹³C NMR (CDCl₃, 125 MHz): δ 146.8, 143.3, 142.6, 133.0, 132.8, 128.4, 128.3, 127.6, 127.4, 125.7, 124.9, 124.8, 122.1, 121.4, 44.3, 36.2, 34.4, 32.8, 31.1, 29.5;

IR (CDCl₃ solution) 3155, 2984, 2254, 1794, 1647, 1472, 1382, 911 cm⁻¹;

EIMS (70 eV) *m/z*: M⁺ 286, 167, 152, 91;

mp 117–119 °C.

(S)-O-Benzyl-4-(p-tolyl)pentan-1-ol (Table 1, entry 10) [85758-06-1]. This compound was prepared according to the General Procedure using (±)-1-(1-bromoethyl)-4-methylbenzene (199 mg, 1.00 mmol) and the appropriate organozinc reagent (~1.6 M in DMA; 1.0 mL, 1.6 mmol). After careful purification by column chromatography (0% → 2% EtOAc in hexanes), the title compound was isolated as a clear, colorless oil.

1st run: 170 mg (63%; 76% ee); 2nd run: 166 mg (62%; 74% ee).

The ee's were determined via HPLC on an OJ-H column (eluent: 1.0% isopropanol in hexanes; flow rate: 1.0 mL/min) with *t_r* (major): 15.7 min, *t_r* (minor): 22.6 min.

$[\alpha]_D^{23} = +12$ (*c* = 0.59, EtOH) {lit.⁴ $[\alpha]_D^{25} = +12.3$ (*c* = 0.91, EtOH); (S)};

¹H NMR (CDCl₃, 400 MHz): δ 7.39-7.28 (m, 5H), 7.12-7.06 (m, 4H), 4.47 (s, 2H), 3.45-3.42 (m, 2H), 2.70-2.62 (m, 1H), 2.33 (s, 3H), 1.70-1.44 (m, 4H), 1.24 (d, *J* = 6.9 Hz, 3H);

¹³C NMR (CDCl₃, 100 MHz): δ 144.4, 138.6, 135.3, 129.0, 128.3, 127.6, 127.4, 126.8, 72.8, 70.5, 39.3, 34.8, 28.0, 22.4, 21.0;

IR (film) 3025, 2924, 2854, 1510, 1491, 1454, 1362, 1101, 816, 733, 697 cm⁻¹;

EIMS (70 eV) *m/z*: M⁺ 268, 177, 159, 119, 91.

4-Indan-1-yl-butyronitrile (eq 3). This compound was prepared (using (±)-1-chloroindane (153 mg, 1.00 mmol)) and assayed as described for entry 3 of Table 1.

1st run: 99 mg (53%; 91% ee); 2nd run: 108 mg (58%; 91% ee).
[α]_D²³ = -4.4 (*c* = 0.54, CHCl₃).

4-(6-Methoxy-indan-1-yl)butyric acid methyl ester (eq 4) [102062-86-2]. This compound was prepared according to the General Procedure using (±)-1-chloro-6-methoxyindane (182 mg, 1.00 mmol) and the appropriate organozinc reagent (~1.6 M in DMA; 1.0 mL, 1.6 mmol). After purification by column chromatography (4% → 6% EtOAc in hexanes), the title compound was isolated as a clear, colorless oil.

1st run: 152 mg (61%; 94% ee); 2nd run: 153 mg (62%; 92% ee).

The ee's were determined via HPLC on an OD-H column (eluent: 0.75% isopropanol in hexanes; flow rate: 1.0 mL/min) with *t_r* (major): 17.2 min, *t_r* (minor): 20.1 min.

[α]_D²³ = -19 (*c* = 0.60, CHCl₃);

¹H NMR (CDCl₃, 400 MHz): δ 7.13-7.11 (m, 1H), 6.75-6.70 (m, 2H), 3.80 (s, 3H), 3.69 (s, 3H), 3.12-3.05 (m, 1H), 2.89-2.73 (m, 2H), 2.42-2.26 (m, 3H), 1.90-1.65 (m, 4H), 1.48-1.39 (m, 1H);

¹³C NMR (CDCl₃, 100 MHz): δ 174.1, 158.6, 148.7, 135.9, 124.8, 111.9, 109.3, 55.4, 51.5, 44.7, 34.4, 34.2, 32.5, 30.5, 23.0;

IR (film) 2948, 2846, 1738, 1608, 1583, 1490, 1436, 1246, 1169, 1034, 808 cm⁻¹;

EIMS (70 eV) *m/z*: M⁺ 248, 160, 147, 115, 91.

3-Bromo-1-indanone [40774-41-2]. A dry 500-mL round-bottomed flask was charged with 1-indanone (11.0 g, 83.2 mmol), *N*-bromosuccinimide (17.8 g, 100 mmol), and benzoyl peroxide (1.0 g, 4.1 mmol), and then it was equipped with a reflux condenser and placed under a nitrogen atmosphere. Dry benzene (200 mL) was added, and the solution was brought to reflux. After 2 h at reflux, the heterogeneous red mixture was allowed to cool to room temperature, and then it was concentrated under vacuum, and passed through Florisil (eluent: benzene). The resulting orange solution was concentrated and stirred under high vacuum for 2 h. Then, hexanes (100 mL) were added, and the resulting mixture was stirred (~3 rotations per min) at 0 °C, leading to the formation of small, dark-orange, granular crystals. The crystals were separated from the brown, gummy residue and resubmitted to the same crystallization conditions. After the second crystallization, pure product was obtained as yellow,

granular crystals (4.7 g, 27%).

^1H NMR (CDCl_3 , 300 MHz): δ 7.78-7.71 (m, 3H), 7.52-7.47 (m, 1H), 5.62 (dd, J = 2.7, 7.2 Hz, 1H), 3.38 (dd, J = 7.2, 19.8 Hz, 1H), 3.07 (dd, J = 2.7, 19.8 Hz, 1H);

^{13}C NMR (CDCl_3 , 75 MHz): δ 201.5, 154.2, 136.0, 135.6, 129.6, 127.5, 123.4, 48.1, 40.6.

(-)-(R)-3-Ethylindan-1-one (eq 5) [16460-88-1]. This compound was prepared according to the General Procedure using ($\pm$)-3-bromo-1-indanone (211 mg, 1.00 mmol) and EtZnBr (~1.6 M in DMA; 1.0 mL, 1.6 mmol). After purification by column chromatography (1 $\rightarrow$ 2% EtOAc in hexanes), the title compound was isolated as a clear, colorless oil.

1st run: 93 mg (58%; 92% ee); 2nd run: 88 mg (55%; 93% ee).

The ee's were determined by chiral GC: CP Chirasil-Dex CB, t_r (major) 22.0 min, t_r (minor) 21.5 min; heating program: 75 $^\circ\text{C}$ $\rightarrow$ 175 $^\circ\text{C}$ @ 3 $^\circ\text{C}/\text{min}$; flow: 1.0 mL/min.

$[\alpha]_D^{23}$ = -27 (c = 2.8, CHCl_3) {lit.⁵ $[\alpha]_D^{25}$ = -15.5 (c = 0.67, EtOH); (R)};

^1H NMR (CDCl_3 , 300 MHz): δ 7.75-7.73 (m, 1H), 7.64-7.58 (m, 1H), 7.52-7.50 (m, 1H), 7.40-7.35 (m, 1H), 3.37-3.29 (m, 1H), 2.86 (dd, J = 7.5, 19.1 Hz, 1H), 2.37 (dd, J = 3.3, 19.1 Hz, 1H), 2.05-1.92 (m, 1H), 1.63-1.48 (m, 1H), 0.99 (t, J = 7.4 Hz, 3H);

^{13}C NMR (CDCl_3 , 75 MHz): δ 206.5, 158.7, 136.9, 134.6, 127.4, 125.6, 123.5, 42.6, 39.6, 28.7, 11.6;

IR (film) 2969, 2936, 1703, 1600, 1464, 1331, 1292, 1245, 765 cm^{-1} ;

EIMS (70 eV) m/z : M^+ 160, 145, 132, 115, 103, 91, 77.

(-)-(R)-3-Methylindan-1-one (Figure 1) [769-14-2]. In the air (no special precautions are necessary), a dry 50-mL Schlenk tube was charged with $\text{NiBr}_2 \cdot \text{diglyme}$ (705 mg, 2.00 mmol), (*R*)-(*i*-Pr)-Pybox (784 mg, 2.60 mmol), and ($\pm$)-3-bromo-1-indanone (4.22 g, 20.0 mmol). The tube was fitted with a septum cap and purged with argon for 15 min. DMA (25 mL) was then added, and the resulting heterogeneous orange mixture was allowed to equilibrate for 15 min under argon in an isopropanol bath cooled to 0 $^\circ\text{C}$. The MeZnI solution (1.6 M in DMA; 20 mL, 32 mmol) was then added in a single portion to the heterogeneous mixture, which rapidly became a clear, brown solution. The reaction mixture was allowed to stir for 48 h at -15 $^\circ\text{C}$. Then, the excess organozinc reagent was quenched by the addition of ethanol (6.0 mL), and the resulting brown-orange liquid was passed through a large pad of silica gel (5% EtOAc in hexanes) to

remove the inorganic salts and much of the DMA. The clear, bright-yellow washings were concentrated and further purified by flash chromatography (1% → 2% EtOAc in hexanes), which afforded the title compound as a clear, colorless oil.

1st run: 1.87 g (64%; 90% ee); 2nd run: 2.04 g (70%; 90% ee).

The ee's were determined by chiral GC: CP Chirasil-Dex CB, *t_r* (major) 18.7 min, *t_r* (minor) 18.0 min; heating program: 75 °C → 175 °C @ 3 °C/min; flow: 1.0 mL/min.

The second run was conducted with (S)-(*i*-Pr)-Pybox.

$[\alpha]_{\text{D}}^{23} = -7.6$ (*c* = 0.63, EtOH) {lit.⁶ $[\alpha]_{\text{D}} = -6.67$ (*c* = 1.05, EtOH); (*R*)};

¹H NMR (CDCl₃, 300 MHz): δ 7.75-7.73 (m, 1H), 7.65-7.59 (m, 1H), 7.53-7.51 (m, 1H), 7.41-7.36 (m, 1H), 3.51-3.40 (m, 1H), 2.95 (dd, *J* = 7.5, 19.1 Hz, 1H), 2.29 (dd, *J* = 3.5, 19.1 Hz, 1H), 1.42 (d, *J* = 7.1 Hz, 3H);

¹³C NMR (CDCl₃, 75 MHz): δ 206.5, 159.9, 136.4, 134.7, 127.4, 125.3, 123.4, 45.3, 32.8, 21.3;

IR (film) 2961, 1713, 1606, 1464, 1464, 1326, 1281, 1239, 1041, 760 cm⁻¹;

EIMS (70 eV) *m/z*: M⁺ 146, 131, 117, 103, 91, 77, 63, 51.

(3R)-cis,trans-3-Methyl-1-indanol [1006-18-4]. This compound was reduced according to the Reduction Procedure using (*R*)-3-methyl-1-indanone (1.85 g, 12.7 mmol) and NaBH₄ (960 mg, 25.4 mmol). The product was obtained as a white solid (1.82 g, 97%) as a mixture of *cis* and *trans* isomers (~20:1).

A second run was performed with (*S*)-3-methyl-1-indanone, and the product was obtained as a white solid (1.80 g, 96%) as a mixture of *cis* and *trans* isomers (~20:1).

¹H NMR (CDCl₃, 500 MHz; *cis* isomer): δ 7.42-7.39 (m, 1H), 7.33-7.22 (m, 3H), 5.22-5.14 (m, 1H), 3.10-3.03 (m, 1H), 2.81-2.73 (m, 1H), 1.91 (d, *J* = 13.0 Hz, 1H), 1.53-1.43 (m, 1H), 1.37 (d, *J* = 11 Hz, 3H);

¹³C NMR (CDCl₃, 125 MHz; *cis* isomer): δ 147.3, 145.0, 128.1, 126.8, 123.6, 123.3, 75.1, 45.7, 36.2, 20.1;

IR (film) 3340, 2958, 2924, 2870, 1645, 1442, 1342, 1211, 1089, 1056, 759 cm⁻¹;

EIMS (70 eV) *m/z*: M⁺ 148, 129, 105, 77, 51;

mp 74–76 °C.

(3R)-cis,trans-1-Bromo-3-methylindane. This compound was prepared according to the Halogenation Procedure using (3R)-cis,trans-3-methyl-1-indanol (20:1 mixture; 1.76

g, 11.9 mmol), PBr₃ (1.61 g, 5.95 mmol), and dry pyridine (0.1 mL). The product was obtained as a clear, colorless oil (2.34 g, 93%) as a mixture of trans and cis isomers (~2:1).

A second run was conducted, and the product was obtained as a clear, colorless oil (2.41 g, 96%) as a mixture of trans and cis isomers (~2:1).

¹H NMR (CDCl₃, 400 MHz; trans isomer): δ 7.41-7.39 (m, 1H), 7.32-7.20 (m, 3H), 5.56 (d, *J* = 6.1 Hz, 1H), 3.54-3.45 (m, 1H), 2.69 (dd, *J* = 6.4, 14.2 Hz, 1H), 2.10 (ddd, *J* = 6.2, 8.9, 14.2 Hz, 1H), 1.35 (d, *J* = 6.9 Hz, 3H);

cis isomer: δ 7.41-7.39 (m, 1H), 7.32-7.20 (m, 3H), 5.46 (dd, *J* = 6.2, 6.7 Hz, 1H), 3.33-3.24 (m, 1H), 3.04-2.97 (m, 1H), 2.24-2.18 (m, 1H), 1.42 (d, *J* = 7.0 Hz, 3H);

¹³C NMR (CDCl₃, 100 MHz; trans isomer): δ 148.4, 143.6, 129.1, 127.1, 124.9, 123.4, 53.8, 46.8, 36.9, 17.8;

cis isomer: δ 147.3, 143.0, 128.7, 127.2, 125.6, 123.5, 51.2, 45.5, 38.5, 20.4;

IR (film) 2958, 2919, 2868, 2364, 2331, 1650, 1634, 1454, 753 cm⁻¹;

EIMS (70 eV) *m/z*: M⁺ 210, 208, 193, 129, 115, 103, 77.

(*R,R*)-(+)-trans-1,3-Dimethylindane [4175-53-5]. In the air (no special precautions are necessary), a dry 50-mL Schlenk tube was charged with NiBr₂·diglyme (368 mg, 1.04 mmol), (*R*)-(*i*-Pr)-Pybox (409 mg, 1.36 mmol), and (3*R*)-1-bromo-3-methylindane (2.20 g, 10.4 mmol). The tube was fitted with a septum cap and purged with argon for 15 min. DMA (15 mL) was then added, and the resulting heterogeneous orange mixture was allowed to equilibrate for 15 min under argon in an isopropanol bath cooled to 0 °C. The MeZnI solution (1.6 M in DMA; 10.4 mL, 16.6 mmol) was then added in a single portion, and the resulting homogeneous brown solution was allowed to stir for 48 h at -15 °C. Then, the excess organozinc reagent was quenched by the addition of ethanol (6.0 mL), and the resulting brown-orange liquid was passed through a large pad of silica gel (pentanes) to remove the inorganic salts and much of the DMA. The solution was then concentrated and further purified by flash chromatography (pentanes), which afforded the title compound as a clear, colorless oil.

1st run: 1.27 g (83%; 93% ee); 2nd run: 1.32 g (87%; 94% ee).

The ee's were determined by chiral GC: CP Chirasil-Dex CB, *t*_r (major) 45.7 min, *t*_r (minor) 45.1 min; heating program: 70 °C ; flow: 0.9 mL/min.

The second run was conducted with (*S*)-(*i*-Pr)-Pybox.

[α]_D²³ = +2.75 (*c* = 1.92, CHCl₃);

¹H NMR (CDCl₃, 500 MHz): δ 7.20 (s, 4H), 3.31-3.27 (m, 2H), 1.93-1.90 (m, 2H), 1.26 (d, *J* = 7.0 Hz, 6H);

¹³C NMR (CDCl₃, 125 MHz): δ 148.3, 126.3, 123.4, 42.9, 37.6, 20.5;

IR (film) 3068, 3019, 2956, 2925, 2866, 1476, 1453, 1373, 1330, 1019, 750 cm⁻¹;

EIMS (70 eV) *m/z*: M⁺ 146, 131, 115, 91.

References

- (1) Arnold, D. R.; Du, X.; Chen, J. *Can. J. Chem.* **1995**, *73*, 307-318.
- (2) (a) Becker, C. L.; McLaughlin, M. L. *Synlett* **1991**, 642. (b) Jones, D. W.; Marmon, R. *J. J. Chem. Soc., Perkin Trans. 1* **1990**, 3271-3275.
- (3) This is based on the work of Huo: Huo, S. *Org. Lett.* **2003**, *5*, 423-425.
- (4) Takano, S.; Sugihara, T.; Samizu, K.; Akiyama, M.; Ogasawara, K. *Chem. Lett.* **1989**, 1781-1784.
- (5) Mani, N. S.; Wu, M. *Tetrahedron: Asymmetry* **2000**, *11*, 4687-4691.
- (6) Yun, J.; Buchwald, S. L. *J. Org. Chem.* **2000**, *65*, 767-774.

1-EthDiox-p-Cl-Indane

Pulse Sequence: szpul

Solvent: CDCl₃

Applic. temperature

INNOVA-500 "butylmkrle"

PULSE SEQUENCE

Relax. delay 5.000 sec

Pulse 65.8 degrees

Acq. time 3.277 sec

Width 3998.8 Hz

16 repetitions

OBSERVE H1 489.753716 MHz

DATA PROCESSING

FT size 65536

Total time 2 min, 12 sec

Current Data Parameters
NAME 6-Me-1-hexInd
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050602
Time_ 12.43
INSTRUM spect
PROBHD 5 mm QNP 1H
PULPROG zg30
TD 65536
SOLVENT CDCl3
NS 16
DS 4
SWH 8278.146 Hz
FIDRES 0.126314 Hz
AQ 3.9584243 sec
RG 128
DW 60.400 usec
DE 6.00 usec
TE 300.0 K
D1 1.00000000 sec

Indanyl-PrCN

Pulse Sequence: zgpg30

Solvent: CDCl3

Acquire Temperature

File: 31857-01

INOVA-500 "z1py"

PULSE SEQUENCE

Relax. delay 5.000 sec

Pulse 85.0 degrees

Acq. time 3.277 sec

Width 9998.8 Hz

16 repetitions

OBSERVE H1, 499.753734 MHz

DATA PROCESSING

File size 65536

Total time 2 min, 12 sec

Pulse Sequence: s2pu1

Solvent: CDCl₃
Ambient temperature
E17a: 3203 cm⁻¹

F116: 3202rp-h1
INOVA-500 "zippy"

PULSE SEQUENCE

Pulse 85.8 degrees
Acq. time 3.277 sec
Width 9998 Hz

16 repetitions
OBSERVE H1 40

OBSERVE H1, 499.7446543 MHZ
DATA PROCESSING
ET 0170 SEC02

Total time 0 min, 52 sec

5-Cyanoindane-HexCl
 Solvent: CDCl3
 Ambient temperature
 INOVA-500 "rocky"
 PULSE SEQUENCE
 Relax. delay 5.000 sec
 Pulse 90.0 degrees
 Acq. time 3.200 sec
 Width 10000.0 Hz
 16 repetitions
 OBSERVE H1, 500.2312734 MHz
 DATA PROCESSING
 FT size 131072
 Total time 2 minutes

1-Bu-Acenaphthene

Pulse Sequence: szpu1

Solvent: CDCl3

Ambient temperature

INOVA-500 "butwinkle"

PULSE SEQUENCE

Relax. delay 5.000 sec

Pulse 65.8 degrees

Acq. time 3.277 sec

Width 8988.8 Hz

18 repetitions

OBSERVE F1, 499.7537718 MHz

DATA PROCESSING

FT size 65556

Total time 2 min, 12 sec

1-Methylcyclohexyl Acenaphthene

Pulse Sequence: szpu1
Solvent: CDCl3
Acq. temperature: INOVA-500 "builtwinkie"
PULSE SEQUENCE
Relax. delay 5.000 sec
Pulse 85.8 degrees
Acq. time 3.277 sec
Width 3998.8 Hz
16 repetitions
OBSERVE H1 499.753722 MHz
DATA PROCESSING
FT size 65536
Total time 2 min, 12 sec

Acenaphthene-8 α -Phthalimide

Pulse Sequence: szpu1

Solvent: CDCl₃

Acquire temperature

File: 20841-p-h1

INOVA-300 "bottle" 101312"

PULSE SEQUENCE

Relax. delay 4.000 sec

Pulse 85.8 degrees

Acq. time 3.277 sec

Width 999.8 Hz

16 repetitions

OBSERVE H1, 499.746546 MHz

DATA PROCESSING

FT size 65536

Total time 1 min, 56 sec

Benzindane-PrPh

Pulse Sequence: szpu1

Solvent: CDCl3

Ambient temperature

File: 318sr-h1

INOVA-500 "zippy"

PULSE SEQUENCE

Relay delay 4.000 sec

Pulse 85.8 degrees

Acq. time 3.277 sec

Width 9998.8 Hz

16 repetitions

OBSERVE H1 499.746534 MHz

DATA PROCESSING

FI size 65536

Total time 1 min, 56 sec

Current Data Parameters
NAME 3174rp
EXPNO 3
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050520

Time 19.49
INSTRUM spect
PROBHD 5mm BBO BB-1
PULPROG zg30
TD 65536
SOLVENT CDCl3
NS 16
DS 2
SWH 8278.146 Hz
FIDRES 0.126314 Hz
AQ 3.9584243 sec
RG 362
DW 60.400 usec
DE 6.00 usec
TE 300.0 K
D1 1.00000000 sec

Current Data Parameters
NAME 3172rp
EXPNO 1
PROCNO 1

F2 - Acquisition Parameters
Date_ 20050520
Time_ 19.32
INSTRUM spect
PROBHD 5mm BBO BB-1
PULPROG zg30
TD 65536
SOLVENT CDCl3
NS 16
DS 2
SWH 8278.146 Hz
FIDRES 0.126314 Hz
AQ 3.9584243 sec
RG 256
DW 60.400 usec
DE 6.00 usec
TE 300.0 K
D1 1.00000000 sec

3-Ethyl-1-Indanone, RP
Pulse Sequence: szpul
Solvent: CDCl3
Ambient temperature
Mercury-300 "mrhah"
Relax. delay 1.000 sec
Pulse 34.1 degrees
Acq. time 1.95 sec
Wdth 4506.5 Hz
15.000 MHz
OBSERVE H1 300.0386328 MHz
DATA PROCESSING
F1 size 32768
Total time 0 min, 0 sec

3-Methyl-1-Indanone, (R)

Pulse Sequence: szpu1

Solvent: CDCl3

Ambient temperature

Mercury-300 "mrhpl"

Relax. delay 1.000 sec

Pulse 34.1 degrees

Acq. time 1.995 sec

Width 4506.5 Hz

16 repetitions

OBSERVE H1: 300.0986331 MHz

DATA PROCESSING

FT size 32768

Total time 0 min, 58 sec

R,R-trans-1,3-Dimethylindene

Pulse Sequence: e2pul

Solvent: CDCl3

Ambient temperature

F1a: 3205P

INOVA-500 "zippy"

PULSE SEQUENCE

Relax delay 5.000 sec

Pulse 85.8 degrees

Acq time 3.277 sec

Width 9398.8 Hz

16 repetitions

OBSERVE H1

DATA PROCESSING

FT size 65536

Total time 2 m 10, 12 sec