Supporting Information

Catalytic Intermolecular Reductive Coupling of Enones and Alkynes

*Ananda Herath, Benjamin Thompson, and John Montgomery**

Department of Chemistry, University of Michigan, Ann Arbor, MI, 48109 – 1055

jmontg@umich.edu

Unless otherwise noted, reagents were commercially available and were used without purification. Tetrahydrofuran (THF) was treated under nitrogen using a solvent purification system (Innovative Technology, Inc., Model # SPS-400-3). Freshly distilled PBU_3 (tributylphosphine) was used in all reactions. $\text{Ni}(\text{COD})_2$ (Strem Chemicals, Inc., used as received) was stored and weighed in an inert atmosphere glovebox. All reactions were conducted in flame-dried glassware under a nitrogen atmosphere. ^1H and ^{13}C spectra were obtained in CDCl_3 at rt, unless otherwise noted, on a Varian Mercury 400 or Varian Unity 500 MHz instrument. Chemical shifts of ^1H NMR spectra were recorded in parts per million (ppm) on the δ scale from an internal standard of residual chloroform (7.27 ppm). Chemical shifts of ^{13}C NMR spectra were recorded in ppm from the central peak of CDCl_3 (77.0 ppm) on the δ scale. High resolution mass spectra (HRMS) were obtained on a VG-70-250-s spectrometer manufactured by Micromass Corp. (Manchester UK) at the University of Michigan Mass Spectrometry Laboratory.

General Procedure for the $\text{Ni}(\text{COD})_2$ / PBU_3 Promoted Reductive Coupling of Enones and Alkynes

To a solution of $\text{Ni}(\text{COD})_2$ (0.1 equiv) in THF (0.6 mL) was added dropwise tributylphosphine (PBU_3) (0.2 equiv) at room temperature. After stirring for 5-10 min at rt, the reaction mixture became bright yellow. A solution of enone (1.0 equiv) and alkyne (1.5 equiv) at rt in MeOH (4.4 mL) was added, and then Et_3B (3.0 equiv) was added. The reaction mixture was stirred at 50 °C until TLC analysis indicated disappearance of the enone. The reaction mixture was quenched with saturated aqueous solution of NH_4Cl and extracted three times with diethyl ether. The combined organic layers were washed with brine, dried over magnesium sulfate, filtered, and concentrated and residue was purified by column chromatography on silica gel. Regioisomeric ratios were determined on crude reaction mixtures using ^1H NMR. The alkene stereochemistry was determined by NOE for the following examples: Table 1, entries 2, 4, 6, 7, 8, 9, 11, and 12 and Table 2, entries 1 and 4. Analysis of coupling constants was satisfactory for the 1,2-disubstituted alkene of Table 1, entry 10.

(E)-5-Methyl-6-phenylhex-5-en-2-one (Table 1, entry 1)

Following the general procedure, 3-buten-2-one (25 μ L, 0.30 mmol), 1-phenyl-1-propyne (52 mg, 0.45 mmol), Ni(COD)₂ (8 mg, 0.03 mmol), PBu₃ (16 μ L, 0.06 mmol), and Et₃B (130 μ L, 0.90 mmol) were stirred for 1 h at 50 °C. The product (48 mg, 85 %, >95:5) was obtained as a colorless oil after SiO₂ chromatography (20 % Et₂O in hexanes). ¹H NMR (400 MHz, CDCl₃) δ 7.33 (m, 2H), 7.22 (m, 3H), 6.29 (s, 1H), 2.67 (t, J = 7.2 Hz, 2H), 2.46 (t, J = 7.6 Hz, 2H), 2.20 (s, 3H), 1.87 (d, J = 0.8 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 208.3, 138.1, 137.3, 128.7, 128.0, 126.0, 125.3, 42.2, 34.4, 29.9, 17.8; IR (film, cm⁻¹) 3055, 2917, 1714, 1599, 1442; GCMS calcd for C₁₃H₁₆O [M⁺] 188, found 188. This compound was previously reported and no spectral data was reported.¹

(E)-2-Methyl-1-phenyldec-1-en-5-one (Table 1, entry 2)

Following the general procedure, 1-octen-3-one (45 μ L, 0.30 mmol), 1-phenyl-1-propyne (52 mg, 0.45 mmol), Ni(COD)₂ (8 mg, 0.03 mmol), PBu₃ (16 μ L, 0.06 mmol), and Et₃B (130 μ L, 0.90 mmol) were stirred for 1 h at 50 °C. The product (57 mg, 78 %, >95:5) was obtained as a colorless oil after SiO₂ chromatography (20 % Et₂O in hexanes). ¹H NMR (400 MHz, CDCl₃) δ 7.32 (m, 2H), 7.22 (m, 3H), 6.29 (s, 1H), 2.64 (m, 2H), 2.46 (m, 4H), 1.87 (d, J = 1.2 Hz, 3H), 1.62 (quint, J = 7.2 Hz, 2H), 1.31 (m, 4H), 0.91 (t, J = 7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 210.7, 138.1, 137.5, 128.7, 128.0, 126.0, 125.3, 42.9, 41.2, 34.4, 31.4, 23.5, 22.4, 17.8, 13.9; IR (film, cm⁻¹) 2931, 1715, 1444; HRMS (EI) m/z calcd for C₁₇H₂₄O [M+Na]⁺ 267.1725, found 267.1722.

(E)-1-Phenyl-3-(1-phenylprop-1-en-2-yl)nonan-1-one (Table 1, entry 3)

Following the general procedure, (E)-1-phenylnon-2-en-1-one (55 mg, 0.25 mmol), 1-phenyl-1-propyne (44 mg, 0.38 mmol), Ni(COD)₂ (7 mg, 0.025 mmol), PBu₃ (14 μ L, 0.05 mmol), and Et₃B (108 μ L, 0.75 mmol) were stirred for 10 h at 50 °C. The product (54 mg, 65 %, >95:5) was obtained as a colorless oil after SiO₂ chromatography (20 % Et₂O in hexanes). ¹H NMR (400 MHz, CDCl₃) δ 7.97 (m, 2H), 7.58 (m, 1H), 7.49 (m,

2H), 7.32 (m, 2H), 7.20 (m, 3H), 6.30 (s, 1H), 3.10 (m, 2H), 2.91 (quint, $J = 7.2$ Hz, 1H), 1.84 (d, $J = 1.2$ Hz, 3H), 1.54 (m, 2H), 1.30 (m, 8H), 0.91 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 199.8, 139.7, 138.2, 137.4, 132.8, 128.9, 128.5, 128.1, 127.9, 126.6, 125.9, 45.6, 43.4, 33.3, 31.8, 29.3, 27.4, 22.6, 14.2, 14.1; IR (film, cm^{-1}) 3058, 2926, 1684, 1598, 1448; HRMS (ESI) m/z calcd for $\text{C}_{24}\text{H}_{30}\text{ONa}$ $[\text{M}+\text{Na}]^+$ 357.2194, found 357.2193.

(E)-4-Methyl-1,3,5-triphenylpent-4-en-1-one (Table 1, entry 4)

Following the general procedure, (E)-chalcone (62 mg, 0.30 mmol), 1-phenyl-1-propyne (52 mg, 0.45 mmol), $\text{Ni}(\text{COD})_2$ (8 mg, 0.03 mmol), PBu_3 (16 μL , 0.06 mmol), and Et_3B (130 μL , 0.90 mmol) were stirred for 12 h at 50 $^\circ\text{C}$. The product (88 mg, 90 %, >95:5) was obtained as a colorless oil after SiO_2 chromatography (10-15 % Et_2O in hexanes). ^1H NMR (400 MHz, CDCl_3) δ 7.98 (m, 2H), 7.58 (m, 1H), 7.48 (m, 2H), 7.38-7.19 (m, 10H), 6.48 (s, 1H), 4.27 (t, $J = 7.2$ Hz, 1H), 3.65 (dd, $J = 7.6, 16.4$ Hz, 1H), 3.54 (dd, $J = 7.2, 16.4$ Hz, 1H), 1.80 (d, $J = 1.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 198.5, 142.6, 140.0, 138.0, 137.2, 133.0, 128.9, 128.6, 128.4, 128.0, 127.96, 127.8, 126.5, 126.1, 125.3, 49.6, 42.4, 17.0; IR (film, cm^{-1}) 3058, 2912, 1687, 1448; HRMS (ESI) m/z calcd for $\text{C}_{24}\text{H}_{22}\text{ONa}$ $[\text{M}+\text{Na}]^+$ 349.1568, found 349.1562.

(E)-3,5-Dimethyl-6-phenylhex-5-en-2-one (Table 1, entry 5)

Following the general procedure, 3-methyl-3-buten-2-one (29 μL , 0.30 mmol), 1-phenyl-1-propyne (52 mg, 0.45 mmol), $\text{Ni}(\text{COD})_2$ (8 mg, 0.03 mmol), PBu_3 (16 μL , 0.06 mmol), and Et_3B (130 μL , 0.90 mmol) were stirred for 1 h at 50 $^\circ\text{C}$. The product (49 mg, 80 %, >95:5) was obtained as a colorless oil after SiO_2 chromatography (20 % Et_2O in hexanes). ^1H NMR (400 MHz, CDCl_3) δ 7.33 (m, 2H), 7.22 (m, 3H), 6.30 (s, 1H), 2.81 (sext, $J = 6.8$ Hz, 1H), 2.55 (dd, $J = 6.4, 13.6$ Hz, 1H), 2.19 (s, 3H), 2.16 (m, 1H), 1.87 (d, $J = 1.2$ Hz, 3H), 1.13 (d, $J = 6.8$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 212.2, 138.0, 135.8, 128.8, 128.0, 127.3, 126.1, 45.3, 43.9, 28.3, 17.6, 15.9; IR (film, cm^{-1}) 3055, 2970, 1712, 1599, 1444; HRMS (EI) m/z calcd for $\text{C}_{14}\text{H}_{18}\text{O}$ $[\text{M}]^+$ 202.1358, found 214.1349.

(*E*)-6-(*tert*-Butyldimethylsilyloxy)-2-methyl-1-phenylundec-1-en-5-one (Table 1, entry 6)

Following the general procedure, 4-(*tert*-butyldimethylsilyloxy)non-1-en-3-one (40 mg, 0.15 mmol), 1-phenyl-1-propyne (26 mg, 0.22 mmol), Ni(COD)₂ (4 mg, 0.015 mmol), PBu₃ (8 μL, 0.03 mmol), and Et₃B (65 μL, 0.45 mmol) were stirred for 1 h at 50 °C. The product (42 mg, 72 %, >95:5) was obtained as a colorless oil after SiO₂ chromatography (5 % Et₂O in hexanes). ¹H NMR (400 MHz, C₆D₆) δ 7.34-7.18 (m, 5H), 6.30 (s, 1H), 4.06 (dd, *J* = 6.8, 5.2 Hz, 1H), 2.77 (m, 2H), 2.43 (t, *J* = 7.6 Hz, 2H), 1.87 (d, *J* = 0.8 Hz, 3H), 1.62 (m, 2H), 1.29 (m, 6H), 0.94 (s, 9H), 0.88 (t, *J* = 6.8 Hz, 3H), 0.08 (s, 3H), 0.07 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 213.5, 138.3, 137.8, 128.8, 128.0, 126.0, 125.2, 78.9, 35.9, 34.9, 33.7, 31.7, 25.8, 24.5, 22.4, 18.1, 17.9, 14.0, -4.9; IR (film, cm⁻¹) 3023, 1719, 1462, 1256; HRMS (ESI) *m/z* calcd for C₂₄H₄₀O₂SiNa [M+Na]⁺ 411.2695, found 411.2689.

(*E*)-3-(2-(*tert*-Butyldimethylsilyloxy)ethyl)-4-methyl-1,5-diphenylpent-4-en-1-one (Table 1, entry 7)

Following the general procedure, (*E*)-5-hydroxy-1-phenylpent-2-en-1-one (53 mg, 0.30 mmol), 1-phenyl-1-propyne (52 mg, 0.45 mmol), Ni(COD)₂ (8 mg, 0.03 mmol), PBu₃ (16 μL, 0.06 mmol), and Et₃B (130 μL, 0.9 mmol) were stirred for 10 hrs at 50 °C. The product (74 mg, 84 %, >95:5) was obtained as a colorless oil after SiO₂ chromatography (50 % Et₂O in hexanes). The alcohol was protected as its TBS ether for characterization. ¹H NMR (400 MHz, CDCl₃) δ 7.96 (m, 2H), 7.57 (m, 1H), 7.47 (m, 2H), 7.30 (m, 2H), 7.18 (m, 3H), 6.28 (s, 1H), 3.65 (m, 2H), 3.11 (m, 3H), 1.84 (d, *J* = 1.2 Hz, 3H), 1.79 (q, *J* = 6.8 Hz, 2H), 0.90 (s, 9H), 0.059 (s, 3H), 0.055 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 199.6, 139.1, 138.0, 137.2, 132.8, 128.9, 128.5, 128.1, 127.9, 126.8, 126.0, 61.1, 43.3, 42.4, 36.2, 25.9, 18.2, 14.3, -5.3; IR (film, cm⁻¹) 3057, 2953, 1686, 1598, 1448; HRMS (ESI) *m/z* calcd for C₂₆H₃₆O₂SiNa [M+Na]⁺ 431.2382, found 431.2378.

(E)-3-(1-Phenylprop-1-en-2-yl)cyclohexanone (Table 1, entry 8)

Following the general procedure, 2-cyclohexen-1-one (29 μ L, 0.30 mmol), 1-phenyl-1-propyne (52 mg, 0.45 mmol), Ni(COD)₂ (8 mg, 0.03 mmol), PBu₃ (16 μ L, 0.06 mmol), and Et₃B (130 μ L, 0.90 mmol) were stirred for 6 h at 50 °C. The product (52 mg, 80 %, >95:5) was obtained as a colorless oil after SiO₂ chromatography (20 % Et₂O in hexanes). ¹H NMR (400 MHz, CDCl₃) δ 7.32 (m, 2H), 7.23 (m, 3H), 6.33 (s, 1H), 2.57-2.41 (m, 4H), 2.34 (m, 1H), 2.15 (m, 1H), 2.01 (m, 1H), 1.87 (d, J = 1.2 Hz, 3H), 1.75 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 211.4, 140.3, 137.9, 128.9, 128.0, 126.2, 124.8, 48.3, 46.7, 41.2, 30.1, 25.2, 15.7; IR (film, cm⁻¹) 3054, 2937, 1710, 1599; HRMS (EI) m/z calcd for C₁₅H₁₈O [M^+] 214.1358, found 214.1358.

(Z)-5,6-Diphenylhex-5-en-2-one (Table 1, entry 9)

Following the general procedure, 3-buten-2-one (25 μ L, 0.30 mmol), diphenylacetylene (80 mg, 0.45 mmol), Ni(COD)₂ (8 mg, 0.03 mmol), PBu₃ (16 μ L, 0.06 mmol), and Et₃B (130 μ L, 0.90 mmol) were stirred for 1 h at 50 °C. The product (47 mg, 63 %) was obtained as a colorless oil after SiO₂ chromatography (20 % Et₂O in hexanes). ¹H NMR (400 MHz, CDCl₃) δ 7.34-6.92 (m, 10H), 6.50 (s, 1H), 2.81 (t, J = 7.2 Hz, 2H), 2.54 (t, J = 7.6 Hz, 2H), 2.11 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 208.1, 141.4, 140.3, 137.0, 129.0, 128.62, 128.56, 127.8, 127.1, 126.9, 126.3, 42.0, 34.5, 30.0; IR (film, cm⁻¹) 3019, 2945, 1597, 1442; GCMS calcd for C₁₈H₁₈O [M^+] 250, found 250. Spectral data for this compound was previously reported and matched with the current data.²

(E)-1-Phenyldec-1-en-5-one (Table 1, entry 10)

Following the general procedure, 1-octen-3-one (45 μ L, 0.30 mmol), phenylacetylene (50 μ L, 0.45 mmol), Ni(COD)₂ (8 mg, 0.03 mmol), PBu₃ (16 μ L, 0.06 mmol), and Et₃B (130 μ L, 0.90 mmol) were stirred for 2 h at 50 °C. The product (51 mg, 74 %, >95:5) was obtained as a yellow oil after SiO₂ chromatography (20 % Et₂O in hexanes). ¹H NMR (400 MHz, CDCl₃) δ 7.31-7.14 (m, 5H), 6.37 (d, J = 15.6 Hz, 1H), 6.16 (dt, J = 16.0, 6.8 Hz, 1H), 2.54 (m, 2H), 2.46 (m, 2H), 2.38 (t, J = 7.6 Hz, 2H), 1.56 (quint, J = 7.2 Hz, 2H), 1.24 (m, 4H), 0.85 (t, J = 6.8 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 210.4, 137.4,

130.6, 129.0, 128.4, 127.0, 125.9, 42.9, 42.2, 31.4, 27.1, 23.5, 22.4, 13.9; IR (film, cm^{-1}) 3026, 2930, 1713, 1448; HRMS (EI) m/z calcd for $\text{C}_{16}\text{H}_{22}\text{O}$ $[\text{M}+\text{Na}]^+$ 253.1568, found 253.1569.

(*E*)-8-(*tert*-Butyldimethylsilyloxy)-4-ethyltridec-3-en-7-one (Table 1, entry 11)

Following the general procedure, 4-(*tert*-butyldimethylsilyloxy)non-1-en-3-one (40 mg, 0.15 mmol), 3-hexyne (26 μL , 0.22 mmol), $\text{Ni}(\text{COD})_2$ (4 mg, 0.015 mmol), PBU_3 (8 μL , 0.03 mmol), and Et_3B (65 μL , 0.45 mmol) were stirred for 1 h at 50 $^\circ\text{C}$. The product (38 mg, 72 %) was obtained as a colorless oil after SiO_2 chromatography (5 % Et_2O in hexanes). ^1H NMR (400 MHz, CDCl_3) δ 5.08 (t, $J = 7.2$ Hz, 1H), 4.02 (dd, $J = 5.2, 7.0$ Hz, 1H), 2.64 (m, 2H), 2.22 (t, $J = 8.0$ Hz, 2H), 2.01 (sext, $J = 7.2$ Hz, 4H), 1.58 (m, 2H), 1.28 (m, 6H), 0.99-0.86 (m, 9H), 0.93 (s, 9H), 0.06 (s, 3H), 0.05 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 213.9, 139.2, 126.2, 78.9, 36.2, 34.9, 31.7, 29.6, 25.7, 24.5, 23.3, 22.5, 20.8, 18.1, 14.6, 14.0, 13.2, -4.92, -4.94; IR (film, cm^{-1}) 2959, 1717, 1463, 1254; HRMS (ESI) m/z calcd for $\text{C}_{21}\text{H}_{42}\text{O}_2\text{SiNa}$ $[\text{M}+\text{Na}]^+$ 377.2852, found 377.2842.

(*E*)-8-Hydroxy-5-methyloct-5-en-2-one (Table 1, entry 12)

Following the general procedure, 3-buten-2-one (25 μL , 0.30 mmol), 3-pentyn-1-ol (42 μL , 0.45 mmol), $\text{Ni}(\text{COD})_2$ (8 mg, 0.03 mmol), PBU_3 (16 μL , 0.06 mmol), and Et_3B (130 μL , 0.90 mmol) were stirred for 1 h at rt. The product (23 mg, 50 %, 3:1) was obtained as a colorless oil after SiO_2 chromatography (50 % Et_2O in hexanes). ^1H NMR (400 MHz, CDCl_3) δ 5.35 (q, $J = 6.8$ Hz, 1H_{min}), 5.14 (t, $J = 7.6$ Hz, 1H_{maj}), 3.67 (t, $J = 6.4$ Hz, 2H_{min}), 3.61 (t, $J = 6.4$ Hz, 2H_{maj}), 2.55 (m, $2\text{H}_{\text{maj}} + 2\text{H}_{\text{min}}$), 2.28 (m, $4\text{H}_{\text{maj}} + 4\text{H}_{\text{min}}$), 2.14 (s, $3\text{H}_{\text{maj}} + 3\text{H}_{\text{min}}$), 1.70 (m, $1\text{H}_{\text{maj}} + 1\text{H}_{\text{min}}$), 1.64 (s, 3H_{maj}), 1.62 (d, $J = 6.4$ Hz, 3H_{min}); ^{13}C NMR (100 MHz, CDCl_3) (major + minor) 208.8, 137.0, 134.7, 121.8, 120.7, 62.3, 60.8, 42.1, 33.6, 33.4, 31.4, 30.5, 30.3, 30.0, 29.9, 16.2, 13.4; IR (film, cm^{-1}) 3411, 2933, 1713, 1440; HRMS (CI- NH_3) m/z calcd for $\text{C}_9\text{H}_{16}\text{O}_2$ $[\text{M}+\text{H}]^+$ 157.1229, found 157.1228.

(E)-3-(5-Hydroxypent-2-en-3-yl)cyclohexanone (Table 1, entry 13)

Following the general procedure, 2-cyclohexen-1-one (29 μL , 0.30 mmol), 3-pentyn-1-ol (42 μL , 0.45 mmol), $\text{Ni}(\text{COD})_2$ (8 mg, 0.03 mmol), PBU_3 (16 μL , 0.06 mmol), and Et_3B (130 μL , 0.90 mmol) were stirred for 5 h at 50 $^\circ\text{C}$. The product (34 mg, 62 %, 53:47) was obtained as a colorless oil after SiO_2 chromatography (20 % Et_2O in hexanes). ^1H NMR (400 MHz, CDCl_3) δ 5.42 (q, $J = 6.8$ Hz, 1H_{maj}), 5.20 (t, $J = 6.8$ Hz, 1H_{min}), 3.62 (m, $2\text{H}_{\text{maj}} + 2\text{H}_{\text{min}}$), 2.33 (m, $7\text{H}_{\text{maj}} + 7\text{H}_{\text{min}}$), 2.06 (m, $1\text{H}_{\text{maj}} + 1\text{H}_{\text{min}}$), 1.88 (m, $1\text{H}_{\text{maj}} + 1\text{H}_{\text{min}}$), 1.65 (m, $6\text{H}_{\text{maj}} + 6\text{H}_{\text{min}}$); ^{13}C NMR (100 MHz, CDCl_3) 211.81, 211.8, 140.2, 138.4, 121.5, 120.1, 62.3, 61.2, 47.6, 47.4, 46.8, 45.3, 41.22, 41.2, 32.6, 31.3, 30.8, 30.1, 25.2, 25.1, 14.2, 13.5; IR (film, cm^{-1}) 3410, 2938, 1710, 1448; HRMS (ESI) m/z calcd for $\text{C}_{11}\text{H}_{18}\text{O}_2\text{Na}$ $[\text{M}+\text{Na}]^+$ 205.1204, found 205.1197.

(E)-Methyl 3-(3-oxobutyl)non-2-enoate (Table 2, entry 1)

Following the general procedure, 3-buten-2-one (25 μL , 0.30 mmol), methyl-2-nonynoate (76 μL , 0.45 mmol), $\text{Ni}(\text{COD})_2$ (8 mg, 0.03 mmol), PBU_3 (16 μL , 0.06 mmol), and Et_3B (130 μL , 0.90 mmol) were stirred for 1 h at rt. The product (43 mg, 60 %, >95:5) was obtained as a colorless oil after SiO_2 chromatography (20 % Et_2O in hexanes). ^1H NMR (400 MHz, CDCl_3) δ 5.58 (s, 1H), 3.67 (s, 3H), 2.58 (m, 4H), 2.42 (m, 2H), 2.17 (s, 3H), 1.42 (m, 2H), 1.29 (m, 6H), 0.87 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 207.1, 166.6, 163.2, 114.9, 50.8, 41.3, 32.4, 31.7, 31.6, 30.0, 29.5, 28.6, 22.6, 14.0; IR (film, cm^{-1}) 2929, 1722, 1643, 1434, 1148; GCMS calcd for $\text{C}_{14}\text{H}_{24}\text{O}_3$ $[\text{M}^+]$ 240, found 240. Spectral data for this compound was previously reported and matched with the current data.²

(E)-Methyl 3-hexyl-6-oxoundec-2-enoate (Table 2, entry 2)

Following the general procedure, 1-octen-3-one (45 μ L, 0.30 mmol), methyl-2-nonynoate (76 μ L, 0.45 mmol), Ni(COD)₂ (8 mg, 0.03 mmol), PBu₃ (16 μ L, 0.06 mmol), and Et₃B (130 μ L, 0.90 mmol) were stirred for 2 h at rt. The product (67 mg, 76 %, 96:4) was obtained as a colorless oil after SiO₂ chromatography (20 % Et₂O in hexanes). ¹H NMR (400 MHz, CDCl₃) δ 5.58 (s, 1H), 3.67 (s, 3H), 2.57 (m, 4H), 2.41 (m, 4H), 1.58 (quint, J = 7.6 Hz, 2H), 1.43 (m, 2H), 1.29 (m, 10H), 0.89 (t, J = 6.8 Hz, 3H), 0.88 (t, J = 6.8 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 209.6, 166.6, 163.4, 114.8, 50.8, 42.9, 40.4, 32.4, 31.8, 31.6, 31.4, 29.5, 28.6, 23.5, 22.6, 22.4, 14.1, 13.9; IR (film, cm⁻¹) 2930, 1720, 1644, 1434, 1149; HRMS (ESI) m/z calcd for C₁₈H₃₂O₃Na [M+Na]⁺ 319.2249, found 319.2241.

(E)-Methyl 3-methyl-6-oxohept-2-enoate (Table 2, entry 3)

Following the general procedure, 3-buten-2-one (25 μ L, 0.30 mmol), methyl 2-butyrate (45 μ L, 0.45 mmol), Ni(COD)₂ (8 mg, 0.03 mmol), PBu₃ (16 μ L, 0.06 mmol), and Et₃B (130 μ L, 0.90 mmol) were stirred for 1 h at rt. The product (38 mg, 74 %, >95:5) was obtained as a colorless oil after SiO₂ chromatography (20 % Et₂O in hexanes). ¹H NMR (400 MHz, CDCl₃) δ 5.66 (q, J = 1.2 Hz, 1H), 3.69 (s, 3H), 2.62 (t, J = 7.2 Hz, 2H), 2.42 (t, J = 7.6 Hz, 2H), 2.18 (s, 3H), 2.17 (d, J = 1.6 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 207.0, 166.9, 158.3, 115.5, 50.8, 41.1, 34.2, 30.0, 18.8; IR (film, cm⁻¹) 2958, 1714, 1688, 1462, 1148; HRMS (EI) calcd for C₉H₁₄O₃ [M]⁺ 170.0943, found 170.0945.

(E)-Methyl 3-methyl-6-oxoundec-2-enoate (Table 2, entry 4)

Following the general procedure, 1-octen-3-one (45 μ L, 0.30 mmol), methyl 2-butyrate (45 μ L, 0.45 mmol), Ni(COD)₂ (8 mg, 0.03 mmol), PBu₃ (16 μ L, 0.06 mmol), and Et₃B (130 μ L, 0.90 mmol) were stirred for 1 h at rt. The product (49 mg, 72 %, >95:5) was obtained as a colorless oil after SiO₂ chromatography (20 % Et₂O in hexanes). ¹H NMR (400 MHz, CDCl₃) δ 5.64 (q, J = 1.2 Hz, 1H), 3.68 (s, 3H), 2.58 (t, J = 7.2 Hz, 2H), 2.41 (t, J = 7.6 Hz, 4H), 2.16 (d, J = 1.2 Hz, 3H), 1.56 (quint, J = 7.2 Hz, 2H), 1.27 (m, 4H), 0.88 (t, J = 6.8 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 209.5, 167.0, 158.6, 115.4, 50.8,

42.9, 40.1, 34.3, 31.3, 23.5, 22.4, 18.5, 13.9; IR (film, cm⁻¹) 2954, 1716, 1650, 1435, 1150; HRMS (EI) *m/z* calcd for C₁₃H₂₂O₃ [M⁺] 226.1569, found 226.1577.

1. Muraoka, T.; Matsuda, I.; Itoh, K. *Tetrahedron Lett.* **2000**, *41*, 8807.
2. Trost, B. M.; Martinez, J. A.; Kulawiec, R. J.; Indolese, A. F. *J. Am. Chem. Soc.* **1993**, *115*, 10402.

Table 1, entry 1

Table 1, entry 2

Table 1, entry 6

Table 1, entry 7

Table 1, entry 8

Table 1, entry 9

Table 1, entry 10

Table 1, entry 11

Table 1, entry 12

Table 1, entry 13

Table 2, entry 1

Table 2, entry 2

Table 2, entry 3