Catalytic Oxidation of Imines Based on Methyltrioxorhenium/Urea Hydrogen Peroxide: A Mild and Easy Chemo- and Regioselective Entry to Nitrones

Gianluca Soldaini, Francesca Cardona and Andrea Goti*

Dipartimento di Chimica Organica “Ugo Schiff”, Laboratorio di Progettazione, Sintesi e Studio di Eterocicli Biologicamente Attivi (HeteroBioLab) and Consorzio INSTM, Università di Firenze, Polo Scientifico, via della Lastruccia 13, I-50019 Sesto Fiorentino (Firenze), Italy

Experimental

Commercial reagents were used as received. All reactions were magnetically stirred and monitored by TLC on 0.25 mm silica gel plates (Merck F₂₅₄) and column chromatography was carried out on Silica Gel 60 (32-63 μ m). Yields refer to spectroscopically and analytically pure compounds unless otherwise stated. ¹H and ¹³C NMR spectra were recorded on Varian Gemini-200 or Varian Mercury-400. Proton and carbon chemical shifts are reported relative to the residue peaks of the solvent CDCl₃ (7.26 ppm for ¹H and 77.0 ppm for ¹³C). Infrared spectra were recorded with a Perkin-Elmer Spectrum BX FT-IR System spectrophotometer. Mass spectra were recorded on a QMD 1000 Carlo Erba instrument by direct inlet; relative percentages are shown in brackets. Elemental analyses were performed with a Perkin-Elmer 2400 analyzer. Optical rotation measurements were performed on a JASCO DIP-370 polarimeter.

Imines **2a-r** were prepared by condensation of the corresponding aldehydes with amines in CH₂Cl₂. Imines **2s-v** were prepared by HCl elimination from the corresponding chloroamines.¹

General Procedure for the Oxidation of Imines to Nitrones

To a stirred solution of the imine **2a-v** (1 mmol) in MeOH (2 mL), UHP (282 mg, 3 mmol) and MTO (5 mg, 0.02 mmol) were added sequentially. The resulting yellow solution was stirred at room temperature until disappearance of the starting material (TLC control) (for reaction time, see Table 1). After removal of the solvent under reduced pressure the reaction mixture was added with CH₂Cl₂ and the undissolved urea filtered off. Removal of the solvent afforded the crude product which was purified by flash column chromatography on silica gel.

***N*-Benzyldiene-benzylamine *N*-oxide (1a)²**

White solid. ¹H NMR (200 MHz, CDCl₃): δ 8.20-8.17 (m, 2 H), 7.49-7.38 (m, 9 H), 5.09 (s, 2 H).

***N*-Benzyldiene-*n*-butylamine *N*-oxide (1b)³**

Pale yellow solid. ¹H NMR (200 MHz, CDCl₃): δ 8.24 (m, 2 H), 7.44-7.39 (m, 4 H), 3.97 (t, 2 H, *J* = 7.0 Hz), 1.98 (quintet, 2 H, *J* = 7.0 Hz), 1.43 (sextet, 2 H, *J* = 7.0 Hz), 0.97 (t, 3 H, *J* = 7.0 Hz).

***N*-Benzyldiene-phenylamine *N*-oxide (1c)⁴**

White solid. ¹H NMR (200 MHz, CDCl₃): δ 8.42-8.37 (m, 2 H), 7.92 (s, 1 H), 7.80-7.75 (m, 2 H), 7.54-7.46 (m, 6 H).

4-Nitrobenzyldiene-benzylamine *N*-oxide (1d)

*R*_f=0.30 (petroleum ether-AcOEt 2:1). Yellow solid; m.p. 115-116 °C. ¹H-NMR (200 MHz, CDCl₃): δ (ppm) 8.38-8.32 (m, 2 H), 8.23-8.19 (m, 2 H), 7.54 (s, 1 H), 7.50-7.40 (m, 5 H), 5.11 (s, 2 H); ¹³C-NMR (50 MHz, CDCl₃): δ (ppm) 147.4 (s), 135.8 (s), 132.3 (s), 132.0 (d), 129.2 (d, 3 C), 128.9 (d, 2 C), 128.6 (d, 2 C), 123.6 (d, 2 C), 72.0 (t); IR (KBr): 3113, 3071, 1596, 1562, 1514, 1341, 1152, 862, 714 cm⁻¹; MS (70 eV): *m/z* (%): 256 (13), 91 (100), 77 (8), 65 (36).

Anal. Calcd for C₁₄H₁₂N₂O₃: C, 65.62; H, 4.72; N, 10.93. Found: C, 65.26; H, 4.78; N, 10.98.

4-Nitrobenzyldiene-butylamine *N*-oxide (1e)

*R*_f=0.31 (petroleum ether-AcOEt 3:2). Pale yellow solid; m.p. 78-79 °C. ¹H-NMR (200 MHz, CDCl₃): δ (ppm) 8.40-8.35 (m, 2 H), 8.24-8.20 (m, 2 H), 7.53 (s, 1 H), 4.00 (t, 2 H, *J* = 7.0 Hz), 1.99 (quintet, 2 H, *J* = 7.0 Hz), 1.44 (sextet, 2 H, *J* = 7.0 Hz), 0.98 (t, 3 H, *J* = 7.0 Hz); ¹³C-NMR (50 MHz, CDCl₃): δ (ppm) 147.4 (s), 135.9 (s), 132.0 (d), 128.5 (d, 2 C), 123.6 (d, 2 C), 67.9 (t), 29.9 (t), 19.8 (t), 13.7 (q); IR (CHCl₃): 2964, 2876, 1599, 1567, 1521, 1165 cm⁻¹; MS (70 eV): *m/z* (%): 222 (30), 205 (20), 163 (82), 151 (34), 117 (40), 104 (22), 77 (64), 63 (56), 51 (100).

Anal. Calcd for C₁₁H₁₄N₂O₃: C, 59.45; H, 6.35; N, 12.60. Found: C, 59.44; H, 6.22; N, 12.69.

2-Naphthylidene-benzylamine *N*-oxide (1f)

*R*_f=0.43 (petroleum ether-AcOEt 2:1). Yellow solid; m.p. 107-108 °C. ¹H-NMR (200 MHz, CDCl₃): δ (ppm) 9.25 (s, 1 H), 7.92-7.77 (m, 4 H), 7.55-7.39 (m, 8 H), 5.09 (s, 2 H); ¹³C-NMR (50 MHz, CDCl₃): δ (ppm) 134.0 (d), 133.8 (s), 133.0 (s), 132.8 (s), 129.0 (d, 2 C), 128.9 (d), 128.7 (d, 3 C), 128.3 (d), 127.6 (d), 127.4 (s), 127.2 (d), 127.1 (d), 126.2 (d), 125.6 (d), 71.1 (t); IR (KBr):

3056, 1595, 1572, 1497, 1451, 1354, 1161, 946, 731 cm^{-1} ; MS (70 eV): m/z (%): 261 (45), 244 (4), 231 (8), 215 (2), 139 (23), 115 (23), 91 (100), 65 (44).

Anal. Calcd for $\text{C}_{18}\text{H}_{15}\text{NO}$: C, 82.73; H, 5.79; N, 5.36. Found: C, 82.69; H, 5.46; N, 5.44.

2-Naphthylidene-butylamine *N*-oxide (1g)

R_f =0.29 (petroleum ether-AcOEt 3:2). White solid; m.p. 42-43 °C. ^1H -NMR (200 MHz, CDCl_3): δ (ppm) 9.26 (s, 1 H), 7.94-7.78 (m, 4 H), 7.55-7.45 (m, 3 H), 4.00 (t, 2 H, $J = 7.0$ Hz), 2.03 (quintet, 2 H, $J = 7.0$ Hz), 1.45 (sextet, 2 H, $J = 7.0$ Hz), 0.99 (t, 3 H, $J = 7.0$ Hz); ^{13}C -NMR (50 MHz, CDCl_3): δ (ppm) 134.0 (d), 133.8 (s), 132.9 (s), 128.9 (d), 128.2 (d), 127.7 (d), 127.5 (s), 127.2 (d), 127.0 (d), 126.2 (d), 125.6 (d), 67.0 (t), 29.8 (t), 19.8 (t), 13.7 (q); IR (CHCl_3): 3059, 2965, 2876, 1579, 1466, 1424, 1155 cm^{-1} ; MS (70 eV): m/z (%): 227 (85), 210 (41), 168 (100), 141 (99), 128 (98), 115 (87), 84 (29).

Anal. Calcd for $\text{C}_{15}\text{H}_{17}\text{NO}$: C, 79.26; H, 7.54; N, 6.16. Found: C, 79.24; H, 7.91; N, 6.16.

3-Methoxybenzylidene-benzylamine *N*-oxide (1h)

R_f =0.33 (petroleum ether-AcOEt 3:2). Pale yellow solid; m.p. 40-42 °C. ^1H -NMR (200 MHz, CDCl_3): δ (ppm) 8.19 (br s, 1 H), 7.52-7.30 (m, 8 H), 7.00-6.95 (m, 1 H), 5.08 (s, 2 H), 3.83 (s, 3 H); ^{13}C -NMR (50 MHz, CDCl_3): δ (ppm) 159.1 (s), 134.0 (d), 133.0 (s), 131.3 (s), 128.9 (d, 3 C), 128.6 (d, 3 C), 121.4 (d), 117.2 (d), 112.1 (d), 71.1 (t), 55.2 (q); IR (CHCl_3): 3052, 2943, 1703, 1595, 1466, 1269, 1174, 1040 cm^{-1} ; MS (70 eV): m/z (%): 241 (41), 224 (7), 211 (8), 135 (4), 91 (100), 65 (53), 51 (26).

Anal. Calcd for $\text{C}_{15}\text{H}_{15}\text{NO}_2$: C, 74.67; H, 6.27; N, 5.81. Found: C, 74.62; H, 6.63; N, 5.89.

3-Methoxybenzylidene-4-methoxybenzylamine *N*-oxide (1i)

R_f =0.28 (petroleum ether-AcOEt 3:1). Colorless oil; ^1H -NMR (200 MHz, CDCl_3): δ (ppm) 8.19 (m, 1 H), 7.44-7.22 (m, 5 H), 6.97-6.87 (m, 3 H), 4.96 (s, 2 H), 3.80 (s, 6 H); ^{13}C -NMR (50 MHz, CDCl_3): δ (ppm) 159.8 (s), 159.1 (s), 133.6 (d), 131.3 (s), 130.6 (d, 2C), 128.9 (d), 124.9 (s), 121.4 (d), 117.1 (d), 114.1 (d, 2 C), 112.0 (d), 70.5 (t), 55.2 (q, 2 C); IR (CHCl_3): 3083, 3004, 2966, 1590, 1613, 1515, 1466, 1253, 1176, 1037 cm^{-1} ; MS (70 eV): m/z (%): 271 (20), 255 (5), 121 (100), 91 (20), 77 (37).

Anal. Calcd for $\text{C}_{16}\text{H}_{17}\text{NO}_3$: C, 70.83; H, 6.32; N, 5.16. Found: C, 70.75; H, 6.69; N, 4.76.

3-Methoxybenzylidene-butylamine *N*-oxide (1j)

$R_f=0.36$ (AcOEt-petroleum ether 2:1). Pale yellow oil. $^1\text{H-NMR}$ (200 MHz, CDCl_3): δ (ppm) 8.22 (m, 1 H), 7.51-7.47 (m, 1 H), 7.37 (s, 1 H), 7.31 (m, 1 H), 7.01-6.95 (m, 1 H), 3.94 (t, 2 H, $J = 7.0$ Hz), 3.85 (s, 3 H), 1.99 (quintet, 2 H, $J = 7.0$ Hz), 1.43 (sextet, 2 H, $J = 7.0$ Hz), 0.98 (t, 3 H, $J = 7.0$ Hz); $^{13}\text{C-NMR}$ (50 MHz, CDCl_3): δ (ppm) 159.2 (s), 134.0 (d), 131.5 (s), 129.0 (d), 121.3 (d), 117.0 (d), 112.1 (d), 67.0 (t), 55.2 (q), 29.7 (t), 19.7 (t), 13.6 (q); IR (CHCl_3): 2964, 2876, 1703, 1591, 1466, 1431, 1270, 1176, 1151, 1040 cm^{-1} ; MS (70 eV): m/z (%): 207 (68), 190 (39), 175 (5), 162 (9), 148 (100), 134 (49), 121 (41), 108 (51), 91 (29), 77 (47), 65 (29), 51 (49).

Anal. Calcd for $\text{C}_{12}\text{H}_{17}\text{NO}_2$: C, 69.54; H, 8.27; N, 6.76. Found: C, 69.18; H, 8.13; N, 7.12.

4-Methoxybenzylidene-benzylamine *N*-oxide (1k)

$R_f=0.22$ (AcOEt-petroleum ether 3:2). White solid; m.p. 96-97 °C. $^1\text{H-NMR}$ (200 MHz, CDCl_3): δ (ppm) 8.20 (m, 2 H), 7.49-7.38 (m, 5 H), 7.33 (s, 1 H), 6.91 (m, 2 H), 5.05 (s, 2 H), 3.84 (s, 3 H); $^{13}\text{C-NMR}$ (50 MHz, CDCl_3): δ (ppm) 160.8 (s), 133.7 (d), 133.2 (s), 130.3 (d, 2 C), 128.9 (d, 2 C), 128.6 (d, 2 C), 128.5 (d), 123.2 (s), 113.5 (d, 2 C), 70.6 (t), 55.2 (q); IR (CHCl_3): 3052, 2938, 2841, 1604, 1508, 1258, 1172, 1031 cm^{-1} ; MS (70 eV): m/z (%): 241 (41), 224 (7), 224 (5), 211 (5), 135 (8), 91 (100), 77 (20), 65 (40), 51 (30).

Anal. Calcd for $\text{C}_{15}\text{H}_{15}\text{NO}_2$: C, 74.67; H, 6.27; N, 5.81. Found: C, 74.49; H, 6.21; N, 5.76.

4-Methoxybenzylidene-4-methoxybenzylamine *N*-oxide (1l)⁵

Pale yellow solid; $^1\text{H NMR}$ (200 MHz, CDCl_3): δ 8.19 (m, 2 H), 7.40 (m, 2 H), 7.26 (s, 1 H), 6.91 (m, 4 H), 4.96 (s, 2 H), 3.83 (s, 3 H), 3.82 (s, 3 H).

4-Methoxybenzylidene-butylamine *N*-oxide (1m)

$R_f=0.25$ (AcOEt-petroleum ether 3:1). Pale yellow oil. $^1\text{H-NMR}$ (200 MHz, CDCl_3): δ (ppm) 8.23 (m, 2 H), 7.31 (s, 1 H), 6.93 (m, 2 H), 3.90 (t, 2 H, $J = 7.0$ Hz), 3.84 (s, 3 H), 1.98 (quintet, 2 H, $J = 7.0$ Hz), 1.41 (sextet, 2 H, $J = 7.0$ Hz), 0.97 (t, 3 H, $J = 7.0$ Hz); $^{13}\text{C-NMR}$ (50 MHz, CDCl_3): δ (ppm) 160.6 (s), 133.7 (d), 130.2 (d, 2 C), 123.3 (s), 113.6 (d, 2 C), 66.4 (t), 55.2 (q), 29.7 (t), 19.8 (t), 13.6 (q); IR (CHCl_3): 2964, 2936, 2875, 1604, 1508, 1464, 1426, 1269, 1172, 1156, 1031 cm^{-1} ; MS (70 eV): m/z (%): 207 (27), 190 (9), 162 (4), 148 (75), 134 (24), 121 (84), 108 (100), 91 (21), 77 (32).

Anal. Calcd for $\text{C}_{12}\text{H}_{17}\text{NO}_2$: C, 69.54; H, 8.27; N, 6.76. Found: C, 69.24; H, 8.17; N, 6.73.

(E)-3-Phenylpropenylidene-benzylamine N-oxide (1n)

$R_f=0.27$ (AcOEt-petroleum ether 3:2). White solid; m.p. 126-127 °C. $^1\text{H-NMR}$ (200 MHz, CDCl_3): δ (ppm) 7.52-7.19 (m, 12 H), 6.94 (d, 1 H, $J = 16.1$ Hz), 4.96 (s, 2 H); $^{13}\text{C-NMR}$ (50 MHz, CDCl_3): δ (ppm) 138.1 (d), 136.2 (d), 135.7 (s), 132.7 (s), 128.9 (d, 2 C), 128.8 (d), 128.7 (d, 2 C), 128.5 (d, 3 C), 127.0 (d, 2 C), 118.1 (d), 69.1 (t); IR (KBr): 3086, 3051, 2998, 1546, 1457, 1424, 1200, 1122, 961, 747, 701 cm^{-1} ; MS (70 eV): m/z (%): 237 (40), 220 (6), 160 (13), 146(6), 130 (3), 115 (21), 105 (7), 91 (100), 77 (14), 65 (41).

Anal. Calcd for $\text{C}_{16}\text{H}_{15}\text{NO}$: C, 80.98; H, 6.37; N, 5.90. Found: C, 80.98; H, 6.65; N, 5.55.

N-Benzylidene-(R)- α -methylbenzylamine N-oxide (1o)⁶

Colorless oil. $^1\text{H NMR}$ (200 MHz, CDCl_3): δ 8.24-8.19 (m, 2 H), 7.53-7.38 (m, 9 H), 5.18 (q, 1 H, $J = 6.9$ Hz), 1.87 (d, 3 H, $J = 6.9$ Hz).

$[\alpha]_D^{21} = -48.8$ (c 1.0, CHCl_3).

N-n-Butylidene-benzylamine N-oxide (1p)⁷

Colorless oil. $^1\text{H NMR}$ (200 MHz, CDCl_3): δ 7.35 (m, 5 H), 6.63 (t, 1 H, $J = 5.8$ Hz), 4.85 (s, 2 H), 2.43 (q, 2 H, $J = 6.9$ Hz), 1.48 (sextet, 2 H, $J = 7.3$ Hz), 0.90 (t, 3 H, $J = 7.3$ Hz).

N-2,2-Dimethylpropylidene-benzylamine N-oxide (1q)

$R_f=0.25$ (petroleum ether-AcOEt 1:1). White solid; m.p. 65-67 °C. $^1\text{H NMR}$ (200 MHz, CDCl_3): δ 7.41 (m, 5 H), 6.55 (s, 1 H), 4.87 (s, 2 H), 1.28 (s, 9 H); $^{13}\text{C NMR}$ (50 MHz, CDCl_3): δ 145.1 (d), 133.4 (s), 128.8 (d, 2 C), 128.7 (d, 2 C), 128.6 (d), 70.5 (t), 32.9 (s), 26.0 (q, 3 C); IR (CHCl_3): 2991, 2960, 2865, 1592, 1497, 1455, 1361, 1142 cm^{-1} ; MS (70 eV): m/z (%): 191 (1), 174 (6), 160 (6), 132(3), 91 (100), 57 (19).

Anal. Calcd for $\text{C}_{12}\text{H}_{17}\text{NO}$: C, 75.35; H, 8.96; N, 7.32. Found: C, 75.64; H, 8.98; N, 7.63.

4-Methoxybenzylidene-1,3,4,6-tetra-O-acetyl-2-glucopyranosylamine N-oxide (1r)

$R_f=0.35$ (AcOEt-petroleum ether 2:1). White solid; 198-199 °C (dec.). $^1\text{H NMR}$ (200 MHz, CDCl_3): δ 8.21 (m, 2 H), 7.26 (s, 1 H), 6.93 (m, 2 H), 6.21 (d, 1 H, $J = 8.2$ Hz), 5.94 (t, 1 H, $J = 9.7$ Hz), 5.18 (t, 1 H, $J = 9.7$ Hz), 4.38 (dd, 1 H, $J = 12.6, 4.4$ Hz), 4.16-4.02 (m, 2 H), 3.97 (ddd, 1 H, $J = 10.0, 4.4, 2.1$ Hz), 3.84 (s, 3 H), 2.08 (s, 3 H), 2.04 (s, 6 H), 1.91 (s, 3 H); $^{13}\text{C NMR}$ (50 MHz, CDCl_3): δ 170.4 (s), 169.8 (s), 169.0 (s), 167.9 (s), 161.5 (s), 136.1 (d), 130.9 (d, 2 C), 122.4 (s), 113.9 (d, 2 C), 90.7 (d), 75.7 (d), 72.4 (d), 70.4 (d), 68.1 (d), 61.4 (t), 55.4 (q), 20.8 (q), 20.7 (q), 20.6 (q), 20.5 (q); IR (CHCl_3): 3018, 2961, 2841, 1756, 1604, 1508, 1368, 1224, 1210, 1074 cm^{-1} ;

MS (70 eV): m/z (%): 481 (11), 422 (6), 362 (2), 229 (12), 211 (17), 194 (18), 169 (28), 152 (100), 134 (31), 109 (23).

$[\alpha]_D^{22} = +91.3$ (c 0.27, CHCl_3).

Anal. Calcd for $\text{C}_{22}\text{H}_{27}\text{NO}_{11}$: C, 54.88; H, 5.65; N, 2.91. Found: C, 54.84; H, 6.05; N, 2.68.

2-Methyl-1-pyrroline N-oxide (1s)⁸

Colorless oil. ^1H NMR (200 MHz, CDCl_3): δ 3.96 (m, 2 H), 2.70 (m, 2 H), 2.15-2.00 (m, 5 H).

(3S,4S)-3,4-O-tert-Butyldimethylsilyl-1-pyrroline N-oxide (1t)⁹

Colorless oil. ^1H NMR (200 MHz, CDCl_3): δ 6.85 (q, 1 H, $J = 1.7$ Hz), 4.67 (br s, 1 H), 4.72-4.60 (m, 1 H), 4.31-4.20 (m, 2 H), 0.87 (s, 18 H), 0.10 (s, 3 H), 0.09 (s, 3 H), 0.08 (s, 3 H), 0.06 (s, 3 H).

2-Methyl-tert-butyldimethylsilyloxy-1-pyrroline N-oxide (1u)

$R_f=0.25$ (CH_2Cl_2 -MeOH 25:1). Colorless oil. ^1H NMR (400 MHz, CDCl_3): δ 4.60 (m, 2 H), 4.02 (m, 2 H), 2.84 (m, 2 H), 2.18-2.10 (m, 2 H), 0.89 (s, 9 H), 0.08 (s, 6 H); ^{13}C NMR (50 MHz, CDCl_3): δ 148.4 (s), 62.5 (t), 58.5 (t), 29.2 (t), 25.5 (q, 3 C), 17.9 (s), 16.9 (t), -5.7 (q, 2 C); IR (CHCl_3): 2956, 2930, 2885, 2858, 1690, 1619, 1472, 1260, 1210, 1122, 1006 cm^{-1} ; MS (70 eV): m/z (%): 229 (1), 214 (17), 172 (100), 156 (42), 129 (87), 115 (10), 101 (41), 75 (90), 59 (60).

Anal. Calcd for $\text{C}_{11}\text{H}_{23}\text{NO}_2\text{Si}$: C, 57.59; H, 10.11; N, 6.11. Found: C, 57.53; H, 10.30; N, 6.23.

(5S)-5-Methyl-tert-butyldimethylsilyloxy-1-pyrroline N-oxide (1v)

$R_f=0.25$ (CH_2Cl_2 -MeOH 25:1). Colorless oil. ^1H NMR (400 MHz, CDCl_3): δ 6.95 (br s, 1 H), 4.31 (dd, 1 H, $J = 10.9, 2.8$ Hz), 4.01 (m, 1 H), 3.07 (ddd, 1 H, $J = 10.9, 2.6, 0.5$ Hz), 2.73-2.56 (m, 2 H), 2.38-2.24 (m, 2 H), 0.80 (s, 9 H), 0.08 (s, 3 H), 0.07 (s, 3 H); ^{13}C NMR (50 MHz, CDCl_3): δ 135.1 (d), 73.9 (d), 61.5 (t), 27.5 (t), 25.8 (q, 3 C), 21.3 (t), 18.2 (s), -5.4 (q), -5.5 (q); IR (CHCl_3): 2956, 2930, 2884, 2857, 1595, 1472, 1462, 1254, 1125 cm^{-1} ; MS (70 eV): m/z (%): 229 (1), 214 (16), 198 (4), 172 (100), 156 (38), 115 (22), 75 (93).

$[\alpha]_D^{22} = -7.2$ (c 0.50, CHCl_3).

Anal. Calcd for $\text{C}_{11}\text{H}_{23}\text{NO}_2\text{Si}$: C, 57.59; H, 10.11; N, 6.11. Found: C, 57.26; H, 10.46; N, 5.93.

References

1. Davis, B. G.; Maughan, M. A. T.; Chapman, T. M.; Villard, R.; Courtney, S. *Org. Lett.* **2002**, *4*, 103.
2. Goti, A.; Cardona, F.; Soldaini, G. *Org. Synth.* **2005**, *81*, 204.
3. Bartoli, G.; Marcantoni, E.; Petrini, M. *J. Org. Chem.* **1992**, *57*, 5834.
4. Murray, R. W.; Iyanar, K.; Chen, J.; Wearing, J. T. *J. Org. Chem.* **1996**, *61*, 8099.
5. Yamada, Y. M. A.; Tabata, H.; Ichinohe, M.; Takahashi, H.; Ikegami, S. *Tetrahedron* **2004**, *60*, 4087.
6. Jost, S.; Gimbert, Y.; Greene, A. E.; Fotiadu, F. *J. Org. Chem.* **1997**, *62*, 6672.
7. Dondoni, A.; Franco, S.; Junquera, F.; Merchan, F.; Merino, P.; Tejero, T. *Synth. Commun.* **1994**, *24*, 2537.
8. Saladino, R.; Neri, V.; Cardona, F.; Goti, A. *Adv. Synth. Catal.* **2004**, *346*, 639.
9. Goti, A.; Cardona, F.; Brandi, A.; Picasso, S.; Vogel, P. *Tetrahedron: Asymmetry* **1996**, *7*, 1659.

1q