A Catalytic Route to the Synthesis of Optically Active β,β -Difluoroglutamic Acid and β,β -Difluoroproline Derivatives

*Atsushi Suzuki, Masayuki Mae, Hideki Amii, and Kenji Uneyama**

Department of Applied Chemistry, Faculty of Engineering, Okayama University

3-1-1 Tsushimanaka, Okayama 700-8530, Japan

uneyamak@cc.okayama-u.ac.jp

^1H and ^{19}F NMR spectra were recorded at 200 and 188 MHz (or 300 and 282 MHz) respectively. The chemical shifts are reported in δ (ppm) values relative to TMS (δ 0 ppm for ^1H NMR), C_6F_6 (δ 0 ppm for ^{19}F NMR). Coupling constants are reported in hertz (Hz). In each case, enantiomeric excess (ee) was determined by HPLC attached chiral stationary phase column, Chiralcel OD-H, OJ, or Chiralpak AD (Daicel Chemical Industries, Ltd.). All air and/or water sensitive reactions were carried out under argon atmosphere with dry, freshly distilled solvents using standard syringe-cannula/septa techniques. All other reagents and solvents were employed without further purification. E. Merk silica gel (Kieselgel 60, 230-400 mesh) was employed for the chromatography. For preparation of **5** and **6**, see Ref. 15a) and 18b).

Ethyl 3-Bromo-2-[*N*-(*p*-methoxyphenyl)]imino-3,3-difluoropropanoate (1**).**

To a solution of **6** (4.47 g, 13.6 mmol) in CH₂Cl₂ (13.6 mL) was added NBS (2.90 g, 16.3 mmol). After stirring at room temperature for 30 min, the solvent was removed under reduced pressure. The crude product was purified by silica gel column chromatography (hexane/AcOEt = 9/1) to afford **1** as an orange oil (4.38 g, 13.1 mmol, 97%); IR (neat) 1740, 1582 cm⁻¹; ¹⁹F NMR (188 MHz, CDCl₃) δ 108.4 (s, 2F); ¹H NMR (200 MHz, CDCl₃) δ 1.17 (t, *J* = 7.0 Hz, 3H), 3.81 (s, 3H), 4.24 (q, *J* = 7.0 Hz, 2H), 6.88 (d, *J* = 9.0 Hz, 2H), 7.01 (d, *J* = 9.0 Hz, 2H); EI-MS (rel. int.) 337 M⁺ (32), 335 M⁺ (29), 162 (46), 134 (100), 29 (48); Anal. Calcd for C₁₂H₁₇BrF₂NO₃: C, 42.88; H, 3.60; N, 4.17. Found: C, 42.89; H, 3.89; N, 4.25.

(*R*)-Ethyl 2-[*N*-(*p*-Methoxyphenylamino)]-3,3-difluoro-5-hexenoate ((*R*)-7**).**

A mixture of Pd(OCOCF₃)₂ (106 mg, 0.32 mmol) and (*R*)-BINAP (299 mg, 0.48 mmol) in acetone (8.0 mL) was stirring at room temperature for 1 h under an argon atmosphere. After concentration, the residue was dissolved in CF₃CH₂OH (8.0 mL) then added **1** (1.34 g, 4.0 mmol). The solution was stirred under hydrogen pressure (100 atm) at room temperature for 48 h. After concentration, the residue was roughly purified by silica-gel column chromatography (hexane/AcOEt = 15/1) to remove the catalyst and byproducts. [(*R*)-**2**: ¹⁹F NMR (282 MHz, CDCl₃) δ 108.5 (dd, *J* = 160.1, 9.3 Hz, 1F), 109.3 (dd, *J* = 160.1, 6.8 Hz, 1F); ¹H NMR (300 MHz, CDCl₃) δ 1.33 (t, *J* = 7.2 Hz, 3H), 3.76 (s, 3H), 4.24-4.37 (m, 2H), 4.37-4.52 (m, 2H), 6.73 (d, *J* = 9.2 Hz, 2H), 6.80 (d, *J* = 9.2 Hz, 2H); EI-MS (rel. int.) 339 M⁺ (24), 337 M⁺ (21), 266 (18), 264 (18), 208 (57), 185 (100), 170 (40), 134 (94). The enantiomeric purity of **2** (87%) was determined by HPLC using a Chiralpak AD column, λ = 230 nm, hexane:ⁱPrOH = 20:1 as eluent, a flow rate of 0.5 mL/min. Under these conditions, base line separation of the two enantiomers was observed with a reference sample containing both enantiomers **2**. The retention times for (*R*)- and (*S*)-enantiomer of **2** are 15.6 and 20.0 min, respectively.] Then, a mixture of (*R*)-**2** and inseparable impurities (total 1.1 g) and allytributyltin (3.0 mL, 9.8 mmol) in toluene (2.2 mL) was added catalytic amount of AIBN several times at 90 °C under an argon atmosphere. After 2 h, the reaction mixture was

concentrated, then added saturated KF aq., AcOEt (50 mL), and stirred at room temperature for 1 h. The organic layer of the reaction mixture was filtered, and concentrated under reduced pressure. The crude product was purified by silica gel column chromatography (hexane/AcOEt = 20/1) and distilled under reduced pressure to afford (*R*)-**7** as a yellow oil (616 mg, 2.1 mmol, 56% from **1**); ee = 87% was determined by HPLC analysis: Chiralpak AD column, hexane:*i*PrOH = 20:1; $[\alpha]_D^{25} = -5.4$ (*c* 0.83, CHCl₃); IR (neat) 3388, 2992, 2916, 1744, 1516 cm⁻¹; ¹⁹F NMR (282 MHz, CDCl₃) δ 54.9 (ddt, *J* = 249, 11.9, 16.1 Hz, 1F), 57.7 (ddt, *J* = 249, 4.5, 18.6 Hz, 1F); ¹H NMR (200 MHz, CDCl₃) δ 1.26 (t, *J* = 7.2 Hz, 3H), 2.74-2.96 (m, 2H), 3.74 (s, 3H), 4.15-4.31 (m, 4H), 5.27 (d, *J* = 17.1 Hz, 1H), 5.30 (d, *J* = 10.3 Hz, 1H), 5.87 (ddt, *J* = 17.1, 10.3, 7.0 Hz, 1H), 6.67 (d, *J* = 9.0 Hz, 2H), 6.79 (d, *J* = 9.0 Hz, 2H); EI-MS (rel. int.) 299 M⁺ (37), 226 (56), 208 (42), 134 (100); Anal. Calcd for C₁₅H₁₉F₂NO₃: C, 60.19; H, 6.40; N, 4.68. Found: C, 60.22; H, 6.71; N, 4.66.

(*R*)-Benzyl 2-[*N*-(*p*-Methoxyphenylamino)]-3,3-difluoro-5-hexenoate ((*R*)-8**).**

A mixture of (*R*)-**7** (1.07 g, 3.6 mmol), CHTD (382 mg, 0.72 mmol) and benzyl alcohol (7.40 mL, 71.6 mmol) in toluene (36 mL) was stirred at refluxing temperature. After 12 h, the reaction mixture was concentrated. The crude product was distilled under reduced pressure (0.6 Torr, 150 °C), then purified by silica-gel column chromatography (hexane/AcOEt = 15/1) to afford (*R*)-**8** as a yellow oil (1.15 g, 3.2 mmol, 89%); ee = 87% was determined by HPLC analysis: Chiralpak AD column, hexane:*i*PrOH = 20:1; $[\alpha]_D^{25} = 10.4$ (*c* 0.91, CHCl₃); IR (neat) 3388, 1750, 1516 cm⁻¹; ¹⁹F NMR (282 MHz, CDCl₃) δ 55.1 (ddt, *J* = 247, 11.9, 16.7 Hz, 1F), 57.9 (ddt, *J* = 247, 4.8, 18.7 Hz, 1F); ¹H NMR (200 MHz, CDCl₃) δ 2.77-2.94 (m, 2H), 3.75 (s, 3H), 4.18-4.40 (m, 2H), 5.11-5.22 (m, 2H), 5.22-5.34 (m, 2H), 5.84 (ddt, *J* = 17.3, 9.9, 7.0 Hz, 1H), 6.65 (d, *J* = 9.0 Hz, 2H), 6.77 (d, *J* = 9.0 Hz, 2H), 7.26-7.35 (m, 5H); EI-MS (rel. int.) 361 M⁺ (22), 226 (37), 91(100); Anal. Calcd for C₂₀H₂₁F₂NO₃: C, 66.47; H, 5.86; N, 3.88. Found: C, 66.24; H, 6.19; N, 3.79.

(*R*)-Benzyl 2-[*N*-(*tert*-Butoxycarbonyl)amino]-3,3-difluoro-5-hexenoate ((*R*)-9a)

To a solution of (*R*)-8 (579 mg, 1.6 mmol) in acetonitrile (16 mL), aqueous solution (16 mL) of CAN (2.64 g, 4.8 mmol) was dropped at 0 °C. The reaction mixture was stirred at 0 °C for 30 min., then neutralized with saturated NaHCO₃ aq.. The mixture was added Na₂S₂O₃ (507 mg, 3.2 mmol), NaHCO₃ (674 mg, 8.0 mmol) and Boc₂O (1.75 g, 8.0 mmol), then stirred at 0 °C to 30 °C for 8 h. The aqueous solution was extracted with EtOAc (50 mL x 4) and the combined organic layer was dried over with MgSO₄. Filtered and removal of the solvent, then the residue was purified by silica gel column chromatography (hexane/EtOAc = 15/1) to afford (*R*)-9a as a colorless oil (538 mg, 1.5 mmol, 94%); ee = 86% was determined by HPLC analysis: Chiralpak AD column, hexane:*i*PrOH = 20:1; [α]_D²⁵ = 9.3 (*c* = 0.80, CHCl₃); IR (neat) 3460, 1762, 1744, 1516 cm⁻¹; ¹⁹F NMR (282 MHz, CDCl₃) δ 55.9 (ddt, *J* = 247, 16.8, 13.0 Hz, 1F), 57.2 (ddt, *J* = 247, 16.8, 10.5 Hz, 1F); ¹H NMR (200 MHz, CDCl₃) δ 1.44 (s, 9H), 2.70 (ddd, *J* = 16.8, 16.8, 7.1 Hz, 2H), 4.68-4.86 (m, 1H), 5.19 (d, *J* = 12.3 Hz, 1H), 5.20-5.40 (m, 3H), 5.26 (d, *J* = 12.3 Hz, 1H), 5.80 (ddt, *J* = 17.0, 10.4, 7.1 Hz, 1H), 7.30-7.40 (s, 5H); Anal. Calcd for C₁₈H₂₃F₂NO₄: C, 60.84; H, 6.52; N, 3.94. Found: C, 61.24; H, 6.64; N, 4.07.

(*R*)-Benzyl 2-[*N*-(Benzyloxycarbonyl)amino]-3,3-difluoro-5-hexenoate ((*R*)-9b)

A solution of (*R*)-8 (1.18 g, 3.3 mmol) in acetonitrile (33 mL) was stirred at 0 °C, and aqueous solution (33 mL) of CAN (5.39 g, 9.8 mmol) was dropped. The reaction mixture was stirred for 30 min. at 0 °C, then neutralized using saturated NaHCO₃ aq.. The mixture was added Na₂S₂O₃ (1.04 g, 6.6 mmol), NaHCO₃ (1.38 g, 16.4 mmol) and Cbz-Cl (1.8 mL, 16.4 mmol), then stirred at 0 °C for 2 h. The aqueous solution was extracted with EtOAc (50 mL x 4) and the combined organic layer was dried over with MgSO₄. Filtered and removal of the solvent, then residue was purified by silica gel column chromatography (hexane/EtOAc = 15/1) to afford (*R*)-9b as a white solid (1.20 g, 3.1 mmol, 94% yield, 87% ee). Recrystallization from hexane/Et₂O gave (*R*)-9b as a colorless crystal (471 mg, 1.2 mmol, 39% yield); ee > 99% was determined by HPLC analysis: Chiralpak AD column, hexane:*i*PrOH = 20:1; [α]_D²⁵

= 17.6 (*c* 0.82, CHCl₃); mp 49.0-49.8 °C; IR (KBr) 3336, 1742, 1696, 1538 cm⁻¹; ¹⁹F NMR (282 MHz, CDCl₃) δ 56.1 (ddt, *J* = 247, 12.9, 16.8 Hz, 1F), 57.2 (ddt, *J* = 247, 10.7, 16.8 Hz, 1F); ¹H NMR (200 MHz, CDCl₃) δ 2.70 (ddd, *J* = 16.8, 16.8, 7.1 Hz, 2H), 4.76-4.93 (m, 1H), 5.10-5.20 (m, 2H), 5.20-5.34 (m, 4H), 5.59 (d, *J* = 9.6 Hz, 1H), 5.79 (ddt, *J* = 17.1, 10.1, 7.1 Hz, 1H), 7.30-7.45 (m, 10H); EI-MS (rel. int.) 254 (1), 180 (3), 107 (48), 91(100); Anal. Calcd for C₂₁H₂₁F₂NO₄: C, 64.77; H, 5.44; N, 3.60. Found: C, 65.01; H, 5.53; N, 3.91.

(*R*)-Ethyl 2-[*N*-(*tert*-Butoxycarbonyl)amino]-3,3-difluoro-5-hexenoate ((*R*)-11)

A solution of (*R*)-7 (500 mg, 1.7 mmol) in acetonitrile (16.7 mL) was stirred at 0 °C, and aqueous solution (16.7 mL) of CAN (2.75 g, 5.0 mmol) was dropped. The reaction mixture was stirred at 0 °C for 30 min., then neutralized with saturated NaHCO₃ aq.. The mixture was added Na₂S₂O₃ (529 mg, 3.3 mmol), NaHCO₃ (702 mg, 8.4 mmol) and Boc₂O (1.82 g, 8.4 mmol), then stirred at 0 °C to 30 °C for 8 h. The aqueous solution was extracted with EtOAc (50 mL x 4) and the combined organic layer was dried over with MgSO₄. Filtration and removal of the solvent, then the residue was purified by silica gel column chromatography (hexane/EtOAc = 15/1) to afford (*R*)-11 as a colorless oil (432 mg, 1.5 mmol, 88%); [α]_D²⁵ = 21.8 (*c* 0.91, CHCl₃).^{10g}

(*R*)-1-Ethyl 2-[*N*-(*tert*-Butoxycarbonyl)amino]-3,3-difluoroglutamate ((*R*)-12)

To a solution of (*R*)-11 (174 mg, 0.59 mmol) and RuO₂·xH₂O (14 mg, 0.070 mmol) in AcOEt (5.9 mL) was added 10% NaIO₄ aq. at 0 °C. The reaction mixture was stirred at 0 °C for 15 min., then stirred at room temperature for 6 h during which several additional aliquots of 10% NaIO₄ aq. were added from time to time to maintain the yellow color of RuO₄. After cooling again at 0 °C, isopropyl alcohol (2 mL) was added to the mixture to quench active RuO₄. The solvent of the reaction mixture was removed under reduced pressure, then the residue was filtered. The filtrate was concentrated under reduced pressure and the residue was purified by silica gel column chromatography (hexane/EtOAc = 1/1 to AcOEt only) to

afford (*R*)-**12** as a colorless oil (153 mg, 0.49 mmol, 83%); $[\alpha]_{\text{D}}^{25} = -2.3$ (*c* 2.16, CHCl₃).

(*R*)-1-Ethyl-5-benzyl 2-[*N*-(*tert*-Butoxycarbonyl)amino]-3,3-difluoroglutamate ((*R*)-13**)**

A mixture of (*R*)-**12** (127 mg, 0.41 mmol) and CsF (87 mg, 0.58 mmol) in DMF (1.5 mL) was stirred at room temperature, then was added benzyl bromide (0.07 mL, 0.58 mmol). After stirring for 24 h at room temperature, the solution was removed under reduced pressure. After filtration, the crude product was purified by silica gel column chromatography (hexane/EtOAc = 9/1) to afford (*R*)-**13** as a colorless oil (143 mg, 0.36 mmol, 88%); ee = 87% was determined by HPLC analysis: Chiralcel OD-H column, hexane:PrOH = 50:1; $[\alpha]_{\text{D}}^{25} = 6.6$ (*c* 1.17, CHCl₃); IR (neat) 3392, 1758, 1754, 1506 cm⁻¹; ¹⁹F NMR (282 MHz, CDCl₃) δ 58.9-59.1 (m, 2F); ¹H NMR (200 MHz, CDCl₃) δ 1.29 (t, *J* = 7.0 Hz, 3H), 1.45 (s, 9H), 3.13 (t, *J* = 15.2 Hz, 2H), 4.25 (q, *J* = 7.0 Hz, 1H), 4.26 (q, *J* = 7.0 Hz, 1H), 5.00-5.20 (m, 1H), 5.20 (s, 2H), 5.35 (br s, 1H), 7.30-7.40 (m, 5H); Anal. Calcd for C₁₉H₂₅F₂NO₆: C, 56.85; H, 6.28; N, 3.49. Found: C, 57.11; H, 6.58; N, 3.54.

(*R*)-Dibenzyl 2-[*N*-(*tert*-Butoxycarbonyl)amino]-3,3-difluoroglutamate ((*R*)-3a**)**

A mixture of (*R*)-**13** (200 mg, 0.50 mmol) and CHTD (293 mg, 0.55 mmol) in benzyl alcohol (5.1 mL) was stirred at 90 °C. After 8 h, benzyl alcohol of the reaction mixture was removed under reduced pressure, then the residue was purified by silica gel column chromatography (hexane/AcOEt = 9/1) for removal of CHTD to afford mixture (*R*)-**13** and (*R*)-**3a**. Subsequently, the resultant mixture of (*R*)-**13** and (*R*)-**3a** was treated with CHTD (293 mg, 0.55 mmol) in benzyl alcohol (5.1 mL) at 90 °C. After 6 h, benzyl alcohol was evaporated under reduced pressure, then residue was purified by silica gel column chromatography (hexane/AcOEt = 9/1) to afford (*R*)-**3a** as a colorless oil (0.36 mmol, 164 mg, 71%); ee = 87% was determined by HPLC analysis: Chiralcel OD-H column, hexane:PrOH = 50:1; $[\alpha]_{\text{D}}^{25} = -10.4$ (*c* 0.64, CH₃OH); IR (neat) 3400, 1756, 1722, 1504, 1458 cm⁻¹; ¹⁹F NMR (282 MHz, acetone-*d*₆) δ 62.2 (dddd, *J* = 251, 15.8, 15.8, 13.8 Hz, 1F), 63.47 (dddd, *J* = 251, 15.8, 15.8, 13.8 Hz, 1F); ¹H NMR

(200 MHz, acetone- d_6) δ 1.40 (s, 9H), 3.30 (dd, J = 15.8, 15.8 Hz, 2H), 5.08-5.32 (m, 5H), 6.73 (br s, 1H), 7.34-7.41 (m, 10H); Anal. Calcd for $C_{24}H_{27}F_2NO_6$: C, 62.20; H, 5.87; N, 3.02. Found: C, 62.03; H, 6.03; N, 3.18.

(*R*)-Dibenzyl 2-[*N*-(Benzyloxycarbonyl)amino]-3,3-difluoroglutamate ((*R*)-3b)

To a solution of (*R*)-3a (70 mg, 0.15 mmol) in CH_2Cl_2 (3.0 mL) was dropped TFA (0.20 mL, 3.0 mmol) at 0 °C. After stirring at 0 °C to room temperature for 4 h, the solution was concentrated under reduced pressure to afford a white solid. The white solid was dissolved in 1,4-dioxane (1.5 mL) and H_2O (1.5 mL), then added $NaHCO_3$ (63 mg, 0.75 mmol) and Cbz-Cl (0.05 mL, 0.45 mmol) at 0 °C. After 2 h, the aqueous solution was extracted with Et_2O (20 mL x 3) and the combined organic layer was dried over with $MgSO_4$, filtered and concentrated under reduced pressure, then the residue was purified by silica gel column chromatography (hexane/ $EtOAc$ = 5/1) to afford (*R*)-3b as a white solid (68 mg, 0.14 mmol, 91%); ee = 87% was determined by HPLC analysis: Chiralcel OD-H column, hexane: i PrOH = 7:1; $[\alpha]_D^{25}$ = -10.2 (c 0.94, CH_3OH); mp 63.9-65.0 °C; IR (neat) 3448, 3392, 1754, 1742, 1522, 1504 cm^{-1} ; ^{19}F NMR (282 MHz, acetone- d_6) δ 62.3 (dddd, J = 254, 15.9, 15.9, 13.5 Hz, 1F), 63.6 (dddd, J = 254, 15.9, 15.9, 13.5 Hz, 1F); 1H NMR (600 MHz, acetone- d_6) δ 3.33 (dd, J = 15.9, 15.9 Hz, 2H), 5.11 (d, J = 12.3 Hz, 1H), 5.13 (d, J = 12.3 Hz, 1H), 5.14 (d, J = 12.3 Hz, 1H), 5.18 (d, J = 12.3 Hz, 1H), 5.24 (d, J = 12.0 Hz, 1H), 5.27 (d, J = 12.0 Hz, 1H), 5.31 (ddd, J = 13.5, 13.5, 9.6 Hz, 1H), 7.23 (d, J = 9.6 Hz, 1H), 7.10-7.40 (m, 15H); Anal. Calcd for $C_{27}H_{25}F_2NO_6$: C, 65.19; H, 5.07; N, 2.82. Found: C, 65.18; H, 5.23; N, 2.50.

(*R*)-Benzyl 1-*N*-(*tert*-Butoxycarbonyl)-3,3-difluoroprolinate ((*R*)-4a)

A solution of 9a (400 mg, 1.1 mmol) in CH_2Cl_2 (4.5 mL) was treated with ozone at -78 °C until the blue color persisted. The solution was purged with N_2 , then portioned Me_2S (1.7 mL, 22.5 mmol). The reaction mixture was stirred at -78 °C to room temperature. After 12 h, the solvent was removed and the

crude mixture was purified by silica gel column chromatography (hexane/EtOAc = 9/1) to afford diastereo mixture of hydroxypyrrolidine **14a** as a colorless oil (362 mg, 1.0 mmol, 90%); ^{19}F NMR (282 MHz, CDCl_3) δ 53.7-56.5 (m), 60.0-62.0 (m), 69.0-70.8 (m), 72.5-74.0 (m).

To a solution of $\text{Ph}_3\text{P}\cdot\text{Cl}_2$ (0.82 mmol, 266 mg) in DMF (1.3 mL), was dropped the solution of **14a** (0.27 mmol, 98 mg) in DMF (1.5 mL) under a nitrogen atmosphere at 0 °C. After 1.5 h at 0 °C, the mixture was stirred for 1 h at room temperature. The solvent was removed under reduced pressure, then the crude product (**15a**) was roughly purified by silica gel column chromatography (hexane/AcOEt = 9/1) to remove triphenylphosphine oxide. Under a hydrogen atmosphere, the crude **15a** was treated with $\text{RhCl}(\text{PPh}_3)_3$ (0.011 mmol, 10 mg) in benzene (2.2 mL) at room temperature. After stirring for 24 h, the solvent was removed and the crude product was purified by silica gel column chromatography (hexane/EtOAc = 9/1) to afford (*R*)-**4a** as a colorless oil (78 mg, 0.23 mmol, 83%); ee = 86% was determined by HPLC analysis: Chiralcel OJ column, hexane:*i*PrOH = 50:1; $[\alpha]_{\text{D}}^{25} = 21.4$ (c 1.08, C_6H_6); IR (neat) 1756, 1714 cm^{-1} ; ^{19}F NMR (282 MHz, acetone- d_6 , a diastereomer ratio of 3:2 was determined by comparing the intensities of the indicated ^{19}F NMR peaks (* =major isomer; ** = minor isomer)) δ **56.4-57.4 (m, 1F), *57.5-58.6 (m, 1F), 67.7-69.0 (m, 1F); ^1H NMR (200 MHz, acetone- d_6 , * =major isomer; ** = minor isomer) δ *1.31 (s, 9H), **1.40 (s, 9H), 2.34-2.59 (m, 2H), 3.47-3.73 (m, 2H), 4.53 (dd, $J = 15.6, 4.0$ Hz, 1H), 5.16-5.34 (m, 2H), 7.36-7.42 (m, 5H); ^{13}C NMR (50 MHz, CDCl_3 , * =major isomer; ** = minor isomer) δ *28.1, **28.3, *32.7 (t, $J = 22.8$ Hz), **33.4 (t, $J = 23.2$ Hz), *42.8 (d, $J = 4.6$ Hz), **43.4 (d, $J = 4.6$ Hz), **64.5 (t, $J = 29.6$ Hz), *33.4 (t, $J = 30.1$ Hz), 67.6, **81.0, *81.1, **125.9 (t, $J = 252$ Hz), *126.5 (t, $J = 253$ Hz), 128.0, 128.2, 128.4, 128.5, *134.9, **135.0, *153.0, **153.5, **167.0, *167.2; Anal. Calcd for $\text{C}_{17}\text{H}_{21}\text{F}_2\text{NO}_4$: C, 59.82; H, 6.20; N, 4.10. Found: C, 59.60; H, 6.54; N, 4.27.

(*R*)-Benzyl 1-*N*-(Benzyloxycarbonyl)-3,3-difluoroprolinate ((*R*)-4b)

Ozonolysis of **9b** (200 mg, 0.51 mmol) was conducted in a same manner as described for **9a** to afford

diastereo mixture of hydroxypyrrolidine **14b** as a colorless oil (180 mg, 0.46 mmol, 89%); ^{19}F NMR (282 MHz, CDCl_3) δ 53.5-56.5 (m), 60.0-61.8 (m), 69.0-71.0 (m), 72.5-74.0 (m).

Dehydration and subsequent hydrogenation of **14b** (0.26 mmol, 100 mg) were conducted in a same manner as described for **14a** to afford (*R*)-**4b** as a colorless oil (83 mg, 0.22 mmol, 87%); ee > 99% was determined by HPLC analysis: Chiralpak AS column, hexane:*i*PrOH = 20:1; $[\alpha]_{\text{D}}^{25} = 19.7$ (*c* 0.86, C_6H_6); IR (neat) 1758, 1724 cm^{-1} ; ^{19}F NMR (282 MHz, acetone- d_6) δ 56.5-59.6 (m, 1F), 67.4-69.0 (m, 1F); ^1H NMR (200 MHz, acetone- d_6) δ 2.41-2.63 (m, 2H), 3.59-3.81 (m, 2H), 4.57-4.73 (m, 1H), 5.05-5.25 (m, 4H), 7.32-7.39 (m, 10H); EI-MS (rel. int.) 180 (3), 107 (18), 91 (100); Anal. Calcd for $\text{C}_{20}\text{H}_{19}\text{F}_2\text{NO}_4$: C, 63.99; H, 5.10; N, 3.73. Found: C, 64.07; H, 5.17; N, 3.83.