Supporting Information

Cationic Rh(I)-catalyst in Fluorinated Alcohol: Mild Intramolecular Cycloaddition Reactions of Ester-tethered Unsaturated Compounds

Akio Saito*, Takamitsu Ono, and Yuji Hanzawa*

*Laboratory of Organic Reaction Chemistry, Showa Pharmaceutical University,
3-3165 Higashi-tamagawagakuen, Machida, Tokyo, 194-8543, Japan
hanzaway@ac.shoyaku.ac.jp (Y. Hanzawa)*

Table of Contents

General Information	S2
General procedure for the preparation of propynyl sorbate derivatives 1	S2
2-Butynyl sorbate (1a), 2-Propynyl sorbate (1b), 3-Phenyl-2-propynyl sorbate (1c), 3- <i>tert</i> -Butyldimethylsilyl-2-propynyl sorbate (1d)	
General procedure for the preparation of 2,4-pentadienyl propiolate derivatives 6	S2-3
Sorbyl propiolate (6a), Sorbyl 2-butynoate (6b), Sorbyl 3-phenylpropiolate (6c)	
Preparation of 2-butynyl (<i>2E</i>)-3-cyclohexenylacrylate (1e)	S3
Preparation of (<i>2E</i>)-3-cyclohexenylallyl but-2-ynoate (5d)	
General procedure for the preparation of propynyl 3-cyclopropylacrylate derivatives 7	S3
(<i>E</i>)-2-butynyl 3-cyclopropylacrylate (7a), (<i>E</i>)-2-propynyl 3-cyclopropylacrylate (7b), (<i>E</i>)-4-methoxy-2-butynyl 3-cyclopropylacrylate (7c), (<i>E</i>)-2-heptynyl 3-cyclopropylacrylate (7d)	
General procedure for the preparation of 3-cyclopropyl-2-propenyl propiolate derivatives 10	S3-S4
(<i>E</i>)-3-cyclopropyl-2-propenyl propiolate (10a), (<i>E</i>)-3-cyclopropyl-2-propenyl 2-butynoate (10b) (<i>E</i>)-3-cyclopropyl-2-propenyl 3-phenylpropiolate (10c), (<i>E</i>)-3-cyclopropyl-2-propenyl 4-methyl-2-pentynoate (10d)	
Preparation of 2-butynyl (<i>2E</i>)-3-cyclohexenylacrylate (12)	S4-5
Preparation of 2-butynyl (<i>2E</i>)-3-cyclohexenylacrylate (15)	S5
Stereochemistry determination of 6a , 6c and 6d	S5
¹ H and ¹³ C NMR spectrum of 1 , 5 , 7 , 10 , 12 and 15	S6-45
¹ H and ¹³ C NMR spectrum of 3 , 4 , 6 , 9 , 11 , 13 , 14 and 16	S46-87
References	S88

General Information. Bis(1,5-cyclooctadiene)- μ,μ' -dichloro dimmer, silver hexafluoroantimonate, 2,2,2-trifluoroethanol and hexafluoroisopropanol are available commercially. All reactions were conducted under an argon atmosphere. For the TLC analysis, Merck precoated TLC plates (silica gel 60 F₂₅₄) were used. Column chromatography was performed on Silica gel 60N (63-200 μm , Kanto Kagaku Co., Ltd.). Medium-pressure liquid chromatography (MPLC) was performed on a 30 cm x 2.2 cm i.d. preppacked column (silica gel, 50 μm) with a UV or RI detector. Melting points were taken on a Yanaco SP-M1 melting point apparatus (Yanagimoto Co.) and were uncorrected. Infrared spectra were taken on a HORIBA FT-710 FT-IR spectrometer. ^1H and ^{13}C NMR spectra were measured at 300.4 and 75.5 MHz in CDCl_3 , and the chemical shifts are given in ppm using Me_4Si (0 ppm) in CDCl_3 for ^1H NMR and CDCl_3 (77.0 ppm) for ^{13}C NMR as internal standard, respectively. Splitting patterns of an apparent multiplet associated with an averaged coupling constant were designed as s (singlet), d (doublet), t (triplet), q (quartet), m (multiplet), and br (broadened). Mass spectra and HRMS were recorded by FAB method on a JMS-HX110 Mass spectrometer. Elemental analysis were measured on a Perkin-Elmer 240B or Elemental Vavio EL.

General procedure for the preparation of propynyl sorbate derivatives 1

2-Butynyl sorbate (1a). To a solution of 2-butyne-1-ol (0.7 g, 10 mmol), *N,N'*-dicyclohexylcarbodiimide (2.3 g, 11 mmol) and 4-(dimethylamino)pyridine (122 mg, 1 mmol) in CH_2Cl_2 (10 mL) was added a solution of sorbic acid (1.1 g, 10 mmol) in CH_2Cl_2 (10 mL) at 0 $^\circ\text{C}$. After being stirred for 4 h at the same temperature, the reaction mixture was diluted with ether and filtered through a celite pad. After concentration of the filtrate to dryness, subsequent purification by silica gel column chromatography (hexane/ Et_2O = 50 : 1) to give **1a** (1.43 g, 87% yield) as a colorless oil. IR (neat) ν cm^{-1} : 1758. ^1H NMR (300 MHz, CDCl_3) δ : 1.80 (s, 3H), 1.80 (d, 3H, J = 4.6 Hz), 4.71 (s, 2H), 5.78 (d, 1H, J = 15.4 Hz), 6.09-6.25 (m, 2H), 7.23 (dd, 1H, J = 10.9, 15.4 Hz). ^{13}C NMR (75 MHz, CDCl_3) δ : 3.5, 18.5, 52.4, 73.2, 82.8, 118.0, 129.6, 139.8, 145.7, 166.4. FAB-LM m/z : 165 ($\text{M}^+\text{+H}$). FAB-HM Calcd for $\text{C}_{10}\text{H}_{13}\text{O}_2$: 165.0897, Found: 165.0915. Anal. Calcd for $\text{C}_{10}\text{H}_{12}\text{O}_2$: C, 73.15; H, 7.37. Found: C, 72.96; H, 7.48.

2-Propynyl sorbate (1b). The ^1H NMR spectra of **1b** was identical to those reported in literature.¹

3-Phenyl-2-propynyl sorbate (1c). Colorless oil. IR (neat) ν cm^{-1} : 1771. ^1H NMR (300 MHz, CDCl_3) δ : 1.77 (d, 3H, J = 5.0 Hz), 4.92 (s, 2H), 5.75 (d, 1H, J = 15.4 Hz), 6.07-6.18 (m, 2H), 7.18-7.29 (m, 4H) 7.37-7.39 (m, 2H). ^{13}C NMR (75 MHz, CDCl_3) δ : 18.6, 52.5, 83.1, 86.2, 117.9, 122.1, 128.2, 128.6, 129.6, 131.8, 140.0, 146.0, 166.4. FAB-LM m/z : 227 ($\text{M}^+\text{+H}$). FAB-HM Calcd for $\text{C}_{15}\text{H}_{15}\text{O}_2$: 227.1057, Found: 227.1072. Anal. Calcd for $\text{C}_{15}\text{H}_{14}\text{O}_2$: C, 79.62; H, 6.24. Found: C, 79.62; H, 6.34.

3-tert-Butyldimethylsilyl-2-propynyl sorbate (1d). White solid. Mp 38-41 $^\circ\text{C}$. IR (neat) ν cm^{-1} : 1718. ^1H NMR (300 MHz, CDCl_3) δ : 0.00 (s, 6H), 0.82 (s, 9H), 1.73 (d, 3H, J = 5.3 Hz), 4.64 (s, 2H), 5.67 (d, 1H, J = 15.3 Hz), 6.03-6.13 (m, 2H), 7.13-7.22 (m, 1H). ^{13}C NMR (75 MHz, CDCl_3) δ : -4.87, 16.4, 18.6, 25.7, 52.4, 96.0, 99.9, 118.0, 129.7, 139.9, 145.9, 166.2. FAB-LM m/z : 265 ($\text{M}^+\text{+H}$). FAB-HM Calcd for $\text{C}_{15}\text{H}_{25}\text{O}_2\text{Si}$: 265.1625, Found: 265.1623. Anal. Calcd for $\text{C}_{15}\text{H}_{24}\text{O}_2\text{Si}$: C, 68.13; H, 9.15. Found: C, 67.81; H, 8.78.

General procedure for the preparation of 2,4-pentadienyl propiolate derivatives 5

Sorbyl propiolate (5a). To a solution of propiolic acid (0.7 mL, 11 mmol) and sorbyl alcohol (1.0 g, 10 mmol) in CH_2Cl_2 (10 mL) was added a solution of *N,N'*-dicyclohexylcarbodiimide (2.3 g, 11 mmol) and 4-(dimethylamino)pyridine (122 mg, 1 mmol) in CH_2Cl_2 (10 mL) at -20 $^\circ\text{C}$. After being stirred at the same temperature for 3 h, the reaction mixture was diluted with ether and filtered through celite pad. The filtrate was concentrated to dryness, and purified by column chromatography on silica gel (hexane/ Et_2O = 50 : 1) to give product **5a** (1.35 g, 90% yield). The ^1H NMR spectra of **5a** was identical to those reported in literature.²

Sorbyl 2-butyrate (5b): The ^1H NMR spectra of **5b** was identical to those reported in literature.³

Sorbyl 3-phenylpropiolate (5c). Colorless oil. IR (neat) ν cm^{-1} : 1708. ^1H NMR (300 MHz, CDCl_3) δ : 1.67 (d, 3H, J = 6.6 Hz), 4.63 (d, 2H, J = 6.6 Hz), 5.53-5.74 (m, 2H), 5.97 (dd, 1H, J = 14.7, 14.7 Hz), 6.22 (dd, 1H, J = 10.4, 14.7 Hz), 7.18-7.46 (m, 5H). ^{13}C NMR (75 MHz, CDCl_3) δ : 18.0, 66.3, 80.4, 86.1, 119.4, 122.4, 128.4, 130.2, 130.5, 131.8, 132.8, 135.9, 153.6. FAB-LM m/z : 227 ($\text{M}^+\text{+H}$). FAB-HM Calcd for $\text{C}_{15}\text{H}_{15}\text{O}_2$: 227.1060, Found: 227.1072. Anal. Calcd for $\text{C}_{15}\text{H}_{14}\text{O}_2$: C, 79.62; H, 6.24. Found: C, 79.54; H, 6.34.

Preparation of 2-butynyl (2E)-3-cyclohexenylacrylate (1e). In similar manner to **1a**, the reaction of (2E)-3-cyclohexenylacrylic acid^{4,5} (341 mg, 2.2 mmol) and 2-butyn-1-ol (125 mg, 2 mmol) gave **1e** (346 mg, 80%) as a colorless oil. IR (neat) ν cm^{-1} ; 1718. ¹H NMR (300 MHz, CDCl_3) δ ; 1.47-1.63 (m, 4H), 1.76 (t, 3H, J = 2.4 Hz), 2.01-2.11 (m, total 4H), 4.60 (d, 1H, J = 2.4 Hz), 4.61 (d, 1H, J = 2.4 Hz), 5.07 (d, 1H, J = 15.5 Hz), 6.07-6.10 (m, 1H), 7.25 (d, 1H, J = 15.5 Hz). ¹³C NMR (75 MHz, CDCl_3) δ ; 3.3, 21.7, 21.8, 23.8, 26.2, 52.2, 73.0, 82.6, 113.5, 134.6, 139.1, 148.6, 166.5. FAB-LM m/z : 205 (M^+ +H). FAB-HM Calcd for $\text{C}_{13}\text{H}_{17}\text{O}_2$: 205.1233, Found: 205.1229. Anal. Calcd for $\text{C}_{13}\text{H}_{16}\text{O}_2$: C, 76.44; H, 7.96. Found: C, 76.71; H, 7.78.

Preparation of (2E)-3-cyclohexenyl-2-propenyl 2-butyrate (5d). In similar manner to **5a**, the reaction of (2E)-3-cyclohexenyl-2-propen-1-ol⁶ (686 mg, 5 mmol) and 2-butyric acid (462 mg, 5.5 mmol) gave **5d** (1.1 g, 98%) as a colorless oil. IR (neat) ν cm^{-1} ; 1708. ¹H NMR (300 MHz, CDCl_3) δ ; 1.55-1.71 (m, 4H), 1.98 (s, 3H), 2.11-2.13 (m, 4H), 4.68 (d, 2H, J = 6.8 Hz), 5.63 (dt, 1H, J = 15.6, 6.8 Hz), 5.77 (br, s, 1H), 6.27 (d, 1H, J = 15.6 Hz). ¹³C NMR (75 MHz, CDCl_3) δ ; 3.3, 21.7, 23.8, 26.2, 51.2, 73.0, 82.6, 113.5, 134.6, 139.1, 148.6, 166.5. FAB-LM m/z : 205 (M^+ +H). FAB-HM Calcd for $\text{C}_{13}\text{H}_{17}\text{O}_2$: 205.1220, Found: 227.1229. Anal. Calcd for $\text{C}_{13}\text{H}_{16}\text{O}_2$: C, 76.44; H, 7.90. Found: C, 76.22; H, 7.76.

General procedure for the preparation of propynyl 3-cyclopropylacrylate derivatives 7

(E)-2-butynyl 3-cyclopropylacrylate (7a). To a solution of 2-butyn-1-ol (0.7 g, 10 mmol), *N,N'*-dicyclohexylcarbodiimide (2.3 g, 11 mmol) and 4-(dimethylamino)pyridine (122 mg, 1 mmol) in CH_2Cl_2 (30 mL) was added a solution of (E)-3-cyclopropylacrylic acid^{7,8} (1.3 g, 12 mmol) in CH_2Cl_2 (10 mL) at 0 °C. After being stirred for 10 h at the same temperature, the reaction mixture was diluted with ether and filtered through a celite pad. After concentration of the filtrate to dryness, subsequent purification by silica gel column chromatography (hexane/AcOEt = 20 : 1) to give **7a** (1.52 g, 80% yield) as a colorless oil. IR (neat) ν cm^{-1} ; 1751. ¹H NMR (300 MHz, CDCl_3) δ ; 0.63-0.68 (m, 2H), 0.93-1.01 (m, 2H), 1.53-1.64 (m, 1H), 4.66 (s, 3H), 4.66 (d, 1H, J = 15.4 Hz), 6.47 (dd, 1H, J = 15.4, 10.4 Hz). ¹³C NMR (75 MHz, CDCl_3) δ ; 3.5, 8.6, 14.4, 52.3, 73.3, 82.7, 117.2, 155.1, 165.8. FAB-LM m/z : 165 (M^+ +H). FAB-HM Calcd for $\text{C}_{10}\text{H}_{13}\text{O}_2$: 165.0912, Found: 165.0916. Anal. Calcd for $\text{C}_{10}\text{H}_{12}\text{O}_2$: C, 73.15; H, 7.37. Found: C, 73.04; H, 7.35.

(E)-2-propynyl 3-cyclopropylacrylate (7b). Colorless oil. IR (neat) ν cm^{-1} ; 1718. ¹H NMR (300 MHz, CDCl_3) δ ; 0.59-0.64 (m, 2H), 0.89-0.98 (m, 2H), 1.48-1.78 (m, 1H), 2.42 (d, 1H, J = 2.4 Hz), 4.67 (d, 1H, J = 2.4 Hz), 5.87 (d, 1H, J = 15.4 Hz), 6.44 (dd, 1H, J = 15.4, 9.9). ¹³C NMR (75 MHz, CDCl_3) δ ; 8.8, 14.5, 51.5, 74.6, 77.9, 116.9, 155.7, 165.6. FAB-LM m/z : 151 (M^+ +H). FAB-HM Calcd for $\text{C}_9\text{H}_{11}\text{O}_2$: 151.0774, Found: 151.0741. Anal. Calcd for $\text{C}_9\text{H}_{10}\text{O}_2$: C, 71.98; H, 6.71. Found: C, 71.76; H, 6.84.

(E)-4-methoxy-2-butynyl 3-cyclopropylacrylate (7c). Colorless oil. IR (neat) ν cm^{-1} ; 1720. ¹H NMR (300 MHz, CDCl_3) δ ; 0.63-0.68 (m, 2H), 0.93-1.00 (m, 2H), 1.53-1.60 (m, 1H), 3.38 (s, 3H), 4.13 (t, 2H, J = 1.8 Hz), 4.77 (t, 1H, J = 1.8 Hz), 5.91 (d, 1H, J = 15.4 Hz), 6.47 (dd, 1H, J = 15.4, 5.1 Hz). ¹³C NMR (75 MHz, CDCl_3) δ ; 8.8, 14.6, 51.9, 57.7, 59.8, 80.8, 82.4, 117.1, 157.6, 165.8. FAB-LM m/z : 195 (M^+ +H). FAB-HM Calcd for $\text{C}_{11}\text{H}_{15}\text{O}_3$: 195.1030, Found: 195.0997. Anal. Calcd for $\text{C}_{11}\text{H}_{14}\text{O}_2$: C, 68.02; H, 7.27. Found: C, 67.84; H, 7.23.

(E)-2-heptynyl 3-cyclopropylacrylate (7d). Colorless oil. IR (neat) ν cm^{-1} ; 1720. ¹H NMR (300 MHz, CDCl_3) δ ; 0.62-0.67 (m, 2H), 0.90 (2H, t, J = 7.1 Hz), 0.92-1.01 (m, 2H), 1.01-2.24 (m, 5H), 2.22 (t, 2H, J = 7.1, 2.2 Hz), 4.71 (d, 1H, J = 15.4 Hz), 6.46 (dd, 1H, J = 15.4, 5.3 Hz). ¹³C NMR (75 MHz, CDCl_3) δ ; 8.8, 13.6, 14.5, 18.5, 21.9, 30.5, 52.5, 74.1, 87.5, 117.3, 155.2, 166.0. FAB-LM m/z : 207 (M^+ +H). FAB-HM Calcd for $\text{C}_{13}\text{H}_{19}\text{O}_2$: 207.1377.0774, Found: 207.1406. Anal. Calcd for $\text{C}_{13}\text{H}_{18}\text{O}_2$: C, 75.69; H, 8.80. Found: C, 75.83; H, 8.67.

General procedure for the preparation of 3-cyclopropyl-2-propenyl propiolate derivatives 10

(E)-3-cyclopropyl-2-propenyl propiolate (10a). To a solution of (E)-3-cyclopropyl-2-propen-1-ol⁹ (390 mg, 4 mmol), *N,N'*-dicyclohexylcarbodiimide (1.0 g, 5 mmol) and 4-(dimethylamino)pyridine (60 mg, 0.5 mmol) in CH_2Cl_2 (10 mL) was added a solution of propiolic acid (310 mg, 4.4 mmol) in CH_2Cl_2 (10 mL) at 0 °C. After being stirred at the same temperature for 10 h, the reaction mixture was diluted with ether and filtered through celite pad. The filtrate was concentrated to dryness, and purified by column chromatography on silica gel (hexane/AcOEt = 20: 1) to give product **10a** (400 mg, 67%) as a colorless oil. IR (neat) ν cm^{-1} ; 1718. ¹H NMR (300 MHz, CDCl_3) δ ; 0.33-0.38 (m, 2H), 0.66-0.75 (m, 2H), 1.40-1.52 (m, 1H), 4.54 (d, 2H, J = 6.6 Hz), 5.27 (dd, 1H, J = 15.4, 8.8 Hz), 5.58 (dt, 1H, J = 15.4, 6.6 Hz). ¹³C NMR (75 MHz, CDCl_3) δ ; 6.8, 13.4, 66.8, 74.6, 75.7, 119.8, 142.2, 152.4. FAB-LM m/z : 151 (M^+ +H). FAB-HM Calcd for $\text{C}_9\text{H}_{11}\text{O}_2$: 151.0745, Found: 151.0735. Anal. Calcd for $\text{C}_9\text{H}_{10}\text{O}_2$: C, 71.98; H, 6.71. Found: C, 71.96; H, 6.71.

(E)-3-cyclopropyl-2-propenyl 2-butynoate (10b). Colorless oil. IR (neat) ν cm^{-1} ; 1706. ^1H NMR (300 MHz, CDCl_3) δ ; 0.33-0.38 (m, 2H), 0.66-0.72 (m, 2H), 1.31-1.40 (m, 1H), 1.92 (s, 3H), 4.51 (d, 2H, $J = 6.6$ Hz), 5.27 (dd, 1H, $J = 15.6, 6.6$ Hz), 5.59 (dt, 1H, $J = 15.6, 6.6$ Hz). ^{13}C NMR (75 MHz, CDCl_3) δ ; 3.7, 6.8, 13.4, 66.3, 72.3, 85.4, 120.2, 141.7, 153.4. FAB-LM m/z : 165 ($\text{M}^+\text{+H}$). FAB-HM Calcd for $\text{C}_{10}\text{H}_{13}\text{O}_2$: 165.0913, Found: 165.0923. Anal. Calcd for $\text{C}_{10}\text{H}_{12}\text{O}_2$: C, 73.15; H, 7.37. Found: C, 73.28; H, 6.52.

(E)-3-cyclopropyl-2-propenyl 3-phenylpropiolate (10c). Colorless oil. IR (neat) ν cm^{-1} ; 1706. ^1H NMR (300 MHz, CDCl_3) δ ; 0.37-0.46 (m, 2H), 0.73-0.80 (m, 2H), 1.37-1.50 (m, 1H), 4.65 (d, 2H, $J = 6.9$ Hz), 5.37 (dd, 1H, $J = 15.4, 8.9$ Hz), 5.71 (dt, 1H, $J = 15.4, 6.9$ Hz). ^{13}C NMR (75 MHz, CDCl_3) δ ; 7.0, 13.6, 66.7, 80.6, 86.2, 119.6, 120.2, 128.5, 130.6, 133.0, 142.1, 153.9. FAB-LM m/z : 226 ($\text{M}^+\text{+H}$). FAB-HM Calcd for $\text{C}_{15}\text{H}_{15}\text{O}_2$: 227.1087, Found: 227.1023. Anal. Calcd for $\text{C}_{15}\text{H}_{14}\text{O}_2$: C, 79.62; H, 6.24. Found: C, 79.43; H, 6.33.

(E)-3-cyclopropyl-2-propenyl 4-methyl-2-pentynoate (10d). In similar manner to **10a**, the reaction of (E)-3-cyclopropyl-2-propen-1-ol (980 mg, 10 mmol) and 4-methyl-2-pentynoic acid¹⁰ (1.2 g, 11 mmol) gave **10d** (823 mg, 43%) as a colorless oil. IR (neat) ν cm^{-1} ; 1712. ^1H NMR (300 MHz, CDCl_3) δ ; 0.32-0.38 (m, 2H), 0.65-0.72 (m, 2H), 1.17 (d, 6H, $J = 7.0$ Hz), 1.30-1.45 (m, 1H), 2.63 (septet, 1H, $J = 7.0$ Hz), 4.54 (d, 2H, $J = 6.9$ Hz), 5.21-5.30 (m, 2H), 5.55-5.65 (m, 2H). ^{13}C NMR (75 MHz, CDCl_3) δ ; 6.7, 13.4, 20.3, 21.6, 66.2, 72.2, 93.8, 120.2, 141.6, 153.6. FAB-LM m/z : 193 ($\text{M}^+\text{+H}$). FAB-HM Calcd for $\text{C}_{12}\text{H}_{17}\text{O}_2$: 193.1212, Found: 193.1232. Anal. Calcd for $\text{C}_{12}\text{H}_{16}\text{O}_2$: C, 74.97; H, 8.39. Found: C, 74.80; H, 8.52.

Preparation of (2E)-3-{1-[(methoxymethyl)oxy]cyclopropyl}-2-propenyl 2-butynoate (12)

Methyl 1-[(methoxymethyl)oxy]cyclopropanecarboxylate (12A). To a solution of methyl 1-hydroxycyclopropanecarboxylate (500 mg, 4.3 mmol) and *N,N*-diisopropylethylamine (1.1 mL, 6.5 mmol) in CH_2Cl_2 (10 mL) was added chloromethyl methyl ether (0.5 mL, 6.5 mmol) at 0°C . After being stirred for 12 h at room temperature, the reaction mixture was quenched by H_2O and extracted with ether. The organic layer was washed with brine, dried over MgSO_4 , and purified by silica gel column chromatography (hexane/ $\text{Et}_2\text{O} = 50 : 1$) to give **12A** (410 mg, 60%) as a colorless oil. IR (neat) ν cm^{-1} ; 1737. ^1H NMR (300 MHz, CDCl_3) δ ; 1.23-1.36 (m, 4H), 3.39 (s, 3H), 3.73 (s, 3H), 4.79 (s, 2H). ^{13}C NMR (75 MHz, CDCl_3) δ ; 15.9, 52.2, 56.3, 59.0, 97.1, 173.6. FAB-LM m/z : 161 ($\text{M}^+\text{+H}$). FAB-HM Calcd for $\text{C}_7\text{H}_{12}\text{O}_4$: 161.0807, Found: 161.1015.

{1-[(methoxymethyl)oxy]cyclopropyl}methanol (12B). Under an argon atmosphere, to a solution of **12A** (400 mg, 2.5 mmol) in CH_2Cl_2 (10 mL) was added diisobutylaluminum hydride (1.0 M in hexane solution, 7.5 mL, 7.5 mmol) at -78°C . After being stirred for 12 h at the same temperature, the reaction mixture was quenched by the addition of MeOH, and filtered through celite pad. The filtrate was concentrated to dryness, and purified by silica gel column chromatography (hexane/ $\text{AcOEt} = 3 : 1$) to give **12B** (301 mg, 91%) as a colorless oil. IR (neat) ν cm^{-1} ; 3392. ^1H NMR (300 MHz, CDCl_3) δ ; 1.28-1.30 (m, 4H), 3.39 (s, 3H), 3.73 (s, 3H), 4.79 (s, 2H). ^{13}C NMR (75 MHz, CDCl_3) δ ; 16.0, 52.2, 55.9, 56.3, 97.1. FAB-LM m/z : 133 ($\text{M}^+\text{+H}$). FAB-HM Calcd for $\text{C}_6\text{H}_{13}\text{O}_3$: 133.0865, Found: 122.0865.

Ethyl (2E)- 3-{1-[(methoxymethyl)oxy]cyclopropyl}acrylate (12C). A solution of **12B** (300 mg, 2.3 mmol), ethyl (triphenylphosphoranyldiene)acetate (1 g, 2.87 mmol), and manganese dioxide (1 g, 11.5 mmol) in CH_2Cl_2 (10 mL) was stirred for 24 h at 50°C . The reaction mixture was diluted with hexane, and filtered through celite pad. The filtrate was

concentrated to dryness, and purified by silica gel column chromatography (hexane/AcOEt = 10 : 1) to give **12C** (281 mg, 62%) as a colorless oil. IR (neat) ν cm^{-1} ; 1716. ^1H NMR (300 MHz, CDCl_3) δ ; 0.95 (dd, 2H, J = 7.9, 5.3 Hz), 1.29 (t, 3H, J = 7.1 Hz), 1.43 (dd, 2H, J = 7.9, 5.3 Hz), 3.37 (s, 3H), 4.19 (q, 2H, J = 7.1 Hz), 4.74 (s, 2H), 5.95 (d, 1H, J = 15.6 Hz), 6.63 (d, 1H, J = 15.6 Hz). ^{13}C NMR (75 MHz, CDCl_3) δ ; 14.3, 16.1, 56.0, 60.3, 61.2, 96.4, 117.3, 151.4, 166.5. FAB-LM m/z : 201 ($\text{M}^+ + \text{H}$). FAB-HM Calcd for $\text{C}_{10}\text{H}_{17}\text{O}_4$: 201.1130, Found: 201.1122. Anal. Calcd for $\text{C}_{10}\text{H}_{16}\text{O}_2$: C, 59.98; H, 8.05. Found: C, 60.04; H, 7.96.

(2E)-3-{1-[(methoxymethyl)oxy]cyclopropyl}-2-propen-1-ol (12D). In a similar manner to **12B**, the treatment of **12C** (250mg, 1.2 mmol) with diisobutylaluminum hydride (1.0 M in hexane solution, 3.6 mL, 3.6 mmol) gave **12D** (173 mg, 92%) as a colorless oil. IR (neat) ν cm^{-1} ; 3300. ^1H NMR (300 MHz, CDCl_3) δ ; 0.72 (d, 2H, J = 5.1 Hz), 0.75 (d, 2H, J = 5.1 Hz), 1.16 (d, 1H, J = 5.1 Hz), 1.56 (d, 1H, J = 5.1 Hz), 3.36 (s, 3H), 4.16 (d, 2H, J = 5.5 Hz), 4.74 (s, 2H), 5.61 (d, 1H, J = 15.6 Hz), 5.77 (dt, 1H, J = 15.6, 5.5). ^{13}C NMR (75 MHz, CDCl_3) δ ; 14.1, 55.8, 60.7, 63.1, 95.9, 76.5, 133.8. FAB-LM m/z : 159 ($\text{M}^+ + \text{H}$). FAB-HM Calcd for $\text{C}_8\text{H}_{15}\text{O}_3$: 159.1004. Found: 159.1021.

(2E)-3-{1-[(methoxymethyl)oxy]cyclopropyl}-2-propenyl 2-butynoate (12)

To a solution of **12D** (158 mg, 1 mmol), *N,N'*-dicyclohexylcarbodiimide (230 mg, 1.1 mmol) and 4-(dimethylamino)pyridine (12.2 mg, 0.1 mmol) in CH_2Cl_2 (3 mL) was added 2-butynoic acid (90mg, 1.1 mmol) at 0 $^\circ\text{C}$. After being stirred at the same temperature for 10 h, the reaction mixture was diluted with ether and filtered through celite pad. The filtrate was concentrated to dryness, and purified by column chromatography on silica gel (hexane/AcOEt = 20: 1) to give product **12** (168mg, 75%) as a colorless oil. IR (neat) ν cm^{-1} ; 1714. ^1H NMR (300 MHz, CDCl_3) δ ; 0.75 (dd, 2H, J = 7.7, 5.4 Hz), 0.75 (dd, 2H, J = 7.7, 5.4 Hz), 1.99 (m, 3H), 3.35 (s, 3H), 1.92 (s, 3H), 4.64 (d, 2H, J = 4.9 Hz), 4.72 (s, 2H), 5.68-5.69 (m, 2H). ^{13}C NMR (75 MHz, CDCl_3) δ ; 3.8, 14.3, 55.9, 60.6, 65.6, 72.3, 85.8, 96.0, 120.3, 138.3, 153.4. FAB-LM m/z : 225 ($\text{M}^+ + \text{H}$). FAB-HM Calcd for $\text{C}_{10}\text{H}_{17}\text{O}_4$: 225.1131, Found: 225.1126. Anal. Calcd for $\text{C}_{12}\text{H}_{17}\text{O}_4$: C, 64.27; H, 7.19. Found: C, 64.19; H, 7.21.

(E)-3-cyclobutyl-2-propenyl 2-butynoate (15)

To a solution of (*E*)-3-cyclobutyl-2-propen-1-ol¹¹ (1.12 g, 10 mmol), *N,N'*-dicyclohexylcarbodiimide (2.26g, 11 mmol) and 4-(dimethylamino)pyridine (122mg, 1mmol) in CH_2Cl_2 (30 mL) was added a solution of 2-butynoic acid (924mg, 11 mol) in CH_2Cl_2 (10 mL) at 0 $^\circ\text{C}$. After being stirred at the same temperature for 10 h, the reaction mixture was diluted with ether and filtered through celite pad. The filtrate was concentrated to dryness, and purified by column chromatography on silica gel (hexane/AcOEt = 20: 1) to give product **15** (1.3 g, 68%) as a colorless oil. IR (neat) ν cm^{-1} ; 1716. ^1H NMR (300 MHz, CDCl_3) δ ; 1.75-2.00 (m, 4H), 1.93 (s, 3H), 2.04-2.18 (m, 2H), 4.58-4.63 (m, 2H), 4.60 (t, 2H, J = 5.4 Hz), 5.49 (dt, 1H, J = 15.4, 5.4 Hz), 5.90 (dd, 1H, J = 6.6, 15.4 Hz). ^{13}C NMR (75 MHz, CDCl_3) δ ; 3.8, 18.5, 28.2, 37.8, 66.4, 72.4, 85.5, 120.7, 141.6, 153.5. FAB-LM m/z : 179 ($\text{M}^+ + \text{H}$). FAB-HM Calcd for $\text{C}_{11}\text{H}_{15}\text{O}_2$: 179.1087, Found: 179.1072. Anal. Calcd for $\text{C}_{11}\text{H}_{14}\text{O}_2$: C, 74.13; H, 7.92. Found: C, 74.03; H, 7.95.

Stereochemistry determination of 6a, 6c and 6d

Relative configurations of **6a**, **6c** and **6d** could be determined by means of NOE experiment, the results of which are shown in Figure 1.

Figure 1. NOE experiments

References

- ¹ Nilsson, Y. I. M.; Gatti, R. G. P.; Andersson, P. G.; Backvall, J. E. *Tetrahedron* **1996**, 52, 7511.
- ² Birtwhistle, D. H.; Brown, J. M.; Foxton, M. W. *Tetrahedron* **1998**, 44, 7309.
- ³ White, J. D.; Sheldon, B. G.. *J. Org. Chem.* **1981**, 46, 2273.
- ⁴ Piva, O.; Comesse S. *Eur. J. Org. Chem.* **2000**, 13, 2417.
- ⁵ Aoki, M.; Kaneko, M. Izumi, S.; Ukai, K.; Iwasawa, N. *Chem. Commun.* **2004**, 2568.
- ⁶ Norsikian, S.; Baudry M.; Jean, F. *Tetrahedron Lett.* **2000**, 41, 6575.
- ⁷ Blackburn, L.; Wei X.; Taylor, R. *Chem. Commun.* **1999**, 1337.
- ⁸ Ma, Z.; Wang, S.; Cooper, C. S.; Fung, A. K.; Lynch, L.; Plagge, F.; Chu, D. *Tetrahedron: Asymmetry* **1997**, 883.
- ⁹ Chatani, N.; Inoue, H.; Morimoto, T.; Muto, T.; Murai, S. *J. Org. Chem.* **2001**, 66, 12, 4433.
- ¹⁰ Marshall, J. A.; Wang, X. *J. Org. Chem.* **1992**, 57, 2747.
- ¹¹ Wang, X.; Stankovich, S. Z.; Widenhoefer, R. A. *Organometallics* **2002**, 21, 901.