This item was submitted to [Loughborough's Research Repository](#) by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

Changes in invertebrate assemblage composition in benthic and hyporheic zones during a severe supraseasonal drought

PLEASE CITE THE PUBLISHED VERSION

<http://dx.doi.org/10.1086/679467>

PUBLISHER

University of Chicago Press / © Society for Freshwater Science

VERSION

AM (Accepted Manuscript)

PUBLISHER STATEMENT

This work is made available according to the conditions of the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) licence. Full details of this licence are available at:
<https://creativecommons.org/licenses/by-nc-nd/4.0/>

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Stubbington, Rachel, A.J. Boulton, Sally Little, and Paul J. Wood. 2019. "Changes in Invertebrate Assemblage Composition in Benthic and Hyporheic Zones During a Severe Supraseasonal Drought". figshare.
<https://hdl.handle.net/2134/17352>.

Changes in invertebrate assemblage composition in benthic and hyporheic zones during a severe supraseasonal drought

Rachel Stubbington^{1,5}, Andrew J. Boulton^{2,6}, Sally Little^{3,7}, Paul J. Wood^{4,8}

¹School of Science and Technology, Nottingham Trent University, Clifton Campus, Nottingham
NG11 8NS UK

²Ecosystem Management, University of New England, Armidale, New South Wales, Australia

³Institute of Estuarine & Coastal Studies, University of Hull, Cottingham Road, Hull
HU6 7RX UK

⁴Department of Geography, Loughborough University, Loughborough, Leicestershire LE11 3TU
UK

E-mail addresses: ⁵rachel.stubbington@ntu.ac.uk; ⁶aboulton@une.edu.au;

⁷sally.little@hull.ac.uk; ⁸To whom correspondence should be addressed, p.j.wood@lboro.ac.uk

Abstract: Droughts are unpredictable disturbances characterized in streams by declining flow, reduced habitat availability, and deteriorating abiotic conditions. Such events typically reduce benthic invertebrate taxon richness and modify assemblage composition, but little is known about how hyporheic invertebrate assemblages respond to drought or how these responses relate to changes in benthic assemblages. We hypothesized that taxon richness (α diversity) and variability (as within-site β diversity) in benthic assemblage composition would decline as drought proceeded, whereas concurrent changes in hyporheic assemblages would be lower in this more stable environment. We predicted that benthic assemblage composition between sites would converge as epigeal taxa were selectively eliminated, whereas between-site hyporheic β diversity would change little. We sampled benthic and hyporheic invertebrates concurrently from 4 sites along a groundwater-fed stream during the final stages of a severe suprasedasonal drought punctuated by a record heat wave. Abiotic conditions in benthic habitats deteriorated as flow declined but changes were less pronounced in the hyporheic zone. Benthic α diversity declined during drought, whereas hyporheic α diversity changed little. However, benthic within-site β diversity increased as the drought progressed because of localized variation in the abundance of common taxa. Temporal trends in hyporheic β diversity were less consistent. Benthic assemblages at individual sites became more similar, especially during the heat wave, reflecting low α diversity and abundance. Hyporheic assemblages changed markedly because of temporary increases in abundances of epigeal and hypogean amphipods. These contrasting responses of benthic and hyporheic assemblages to drought should be recognized when developing management strategies for drought-impacted streams.

Key words: low flows, benthos, hyporheos, macroinvertebrate, alpha-diversity, beta-diversity, *Gammarus*, *Niphargus*

In all lotic ecosystems, supraseasonal droughts are severe and unpredictable disturbances that gradually intensify over space and time (Lake 2003). In-stream changes during drought are complex and variable but typically include reduction in wetted habitat availability and heterogeneity, slowing and often cessation of flow, and an increase in the influence of groundwater and solar radiation on surface-water temperatures (Boulton 2003, Dewson et al. 2007). However, few investigators have examined the effects of these in-stream changes on aquatic biota during severe supraseasonal droughts (Lake 2011) largely because of the challenges of capturing the early stages of these unpredictable, slow-onset events (Fleig et al. 2010).

The few studies conducted were focused on benthic invertebrates in surface sediments (reviews in Dewson et al. 2007, Lake 2011). However, the hyporheic zone (HZ), between the surface sediments and phreatic groundwater, often supports a diverse invertebrate assemblage (Boulton et al. 1998), comprising groundwater and hyporheic specialists and taxa that typically inhabit the benthic zone (BZ) (Williams et al. 2010). Predominantly benthic taxa are proposed to migrate into the HZ during adverse conditions in the surface stream (Williams and Hynes 1974, Dole-Olivier 2011), including drought. These migrations may be triggered by intensified biotic interactions within contracting areas of wetted benthic habitat (Stubbington et al. 2011a) or by thermal stress during periods of elevated water temperature (Wood et al. 2010).

In addition to disturbance-related drivers, seasonal variability and multiple environmental variables influence the distribution and movements of invertebrates between the benthic and hyporheic zones, collectively resulting in a spatiotemporally variable community (Brunke and Gonser 1999). Disturbances, such as drought or high-flow events, may have different effects on BZ and HZ habitats, eliciting different responses by their respective invertebrate faunas. However, few investigators have concurrently sampled invertebrate assemblages in the BZ and

65 HZ (e.g., James et al. 2008, Young et al. 2011), and we are aware of only 4 studies in which
66 paired benthic–hyporheic invertebrate samples were collected repeatedly during severe low
67 flows or drying: Belaidi et al. (2004), Stubbington et al. (2011a), Datry (2012), and our study in
68 the Little Stour, a groundwater-fed stream in southeastern England.

69 Severe low flows and drying of surface water may have greater effects on invertebrates in
70 the BZ than in the HZ, reducing taxon richness (α diversity) and selectively eliminating surface
71 species that require reliable flow and good water quality (Boulton 2003). As drought progresses,
72 variation in benthic community composition among sites and over time (both spatial and
73 temporal β diversity; sensu Anderson et al. 2011) also is likely to decline as conditions become
74 progressively less favorable to most stream invertebrates. In other aquatic habitats, such as
75 ponds, drought can reduce β diversity by acting as a strong ‘environmental filter’ (Chase 2007).
76 Similar changes might be predicted in streams as a suprasedational drought proceeds (Lake 2011).

77 We concurrently sampled benthic and hyporheic invertebrate assemblages in riffle
78 habitats at 2 perennial and 2 historically semiperennial sites along the Little Stour River, UK, at
79 monthly intervals during the latter stages of a severe suprasedational drought to test 3 hypotheses:
80 1) benthic taxon richness (α diversity) would decline steadily as the drought proceeded, whereas
81 hyporheic α diversities would change little because subsurface environmental conditions would
82 be buffered in this groundwater-fed stream; 2) within-site variation in benthic community
83 composition, as fine-scale patchiness among replicates (β diversity), would decline as the
84 drought proceeded, especially at the 2 semiperennial sites (Wood and Petts 1999, Wood and
85 Armitage 2004), whereas no significant temporal change in hyporheic β diversity would occur
86 within sites because subsurface conditions would be buffered from drought; 3) benthic
87 community composition would converge among sites over time as a result of the environmental

filter of drought, whereas less convergence would occur in hyporheic community composition over time, but some divergence would be possible because of site-specific movements of benthos into the HZ.

METHODS

Study site and drought characteristics

The Little Stour River (lat 51.275°N, long 1.168°E) is a 11.5 km long lowland groundwater-dominated stream that drains a predominantly agricultural catchment of ~213 km². The geology in the headwaters is chalk overlain by alluvial gravels in the river's middle and lower reaches. The sedimentary calcareous rocks result in relatively high conductivities (~580 µm/cm). Further details regarding the river and its catchment were provided by Wood and Petts (1999).

Mean annual precipitation in the catchment is ~650 mm, which is usually sufficient to maintain perennial flow below the spring head (lat 51.265°N, long 1.153°E). However, a 1-km section within the study area dried during 3 previous supraseasonal droughts in 1949, 1991–1992, and 1996–1997 (Wood and Armitage 2004). Our study was conducted during the latter stages of a severe (return period exceeding 1-in-100 y; Marsh 2007) supraseasonal drought that affected southern and eastern England between 2004 and 2006. In the Little Stour, groundwater and surface-water levels fell during the extended period of rainfall deficit between November 2004 and October 2006 (Marsh 2007). Mean daily discharge between 2003 and the end of 2006 was 0.181 m³/s but throughout 2006, discharge never reached this average (Fig. 1A). In July 2006, the rainfall deficit was accompanied by a heat wave that included the highest monthly mean air temperatures recorded in the 348-y Central England Temperature series (Prior and

Beswick 2007; Fig. 1B). We sampled from April to October 2006, capturing the final months of this supraseasonal drought and the 2 mo after the meteorological drought broke. Mean daily discharge then increased gradually until mid-December. The first winter storms in January 2007 caused a rapid rise in discharge to above the 2003–2006 average.

Each month, the benthic and hyporheic zones of 2 perennial sites (sites 1 and 4) and 2 semiperennial sites (sites 2 and 3) in the upper 5 km of the Little Stour were sampled. The semiperennial sites were in a 1-km reach that dried during 4 of the previous 16 y (Wood et al. 2000), but neither site dried completely during our study. Riffle crests were gradually exposed at both sites as flow declined, whereas sampling points remained inundated. All sites were predominantly riffle habitat, 5 to 10 m wide, and with a pebble–gravel bed (4–32 mm).

Invertebrate sampling and community characterization

At each site, 5 replicate sampling points were selected in riffles. The same points were used on each sampling occasion. Based on observations made during previous supraseasonal droughts (Wood and Petts 1999, Wood and Armitage 2004), sampling points were in areas that would dewater last. Riffle crests and stream margins were avoided. Sampling points were ≥ 3 m apart to avoid overlap between the sediments sampled at 1 point with those at a neighboring point, particularly during hyporheic sampling. At each point, invertebrates were sampled from the BZ with a Surber sampler (0.1-m² frame, 250- μ m-mesh net) by manually disturbing the sediments to a depth of 5 cm for 30 s and inspecting larger clasts for attached invertebrates.

Hyporheic invertebrates were then sampled 1.5 to 2 m upstream of each benthic Surber sample. On the 1st sampling date, polyvinyl chloride (PVC) wells (25-mm internal diameter) were inserted 20 cm into the HZ at each of the 5 sampling points with a stainless steel T-bar.

Each month, 6-L samples were extracted from the base of the well with a manual bilge pump following the procedure of Boulton and Stanley (1995). Six liters provides a standardized volumetric sample that previous investigators (Boulton et al. 2004, Stubbington et al. 2011b, Datry 2012) showed was adequate to characterize differences in hyporheic community composition among sites and sampling points. Samples were passed through a sieve (90- μ m mesh) and preserved in 4% formaldehyde. For comparison with data from the benthic samples, only the invertebrates retained by a 250- μ m sieve are considered in our paper. The PVC wells were left in situ and sampled monthly, ensuring that repeated temporal sampling was not confounded by spatial variability.

Conductivity (μ S/cm), pH, dissolved O₂ (DO; mg/L) and water temperature ($^{\circ}$ C) were recorded in surface water and hyporheic water using standard meters (Hanna Instruments, Leighton Buzzard, UK). Water depth and mean flow velocity (at 0.6 $\times$ depth) were recorded at each sampling point with an ADS SENSE-RC2 flow meter (ADS Environmental Services, Huntsville, Alabama). In the laboratory, most taxa were identified to species. However, larvae and adults of *Helophorus* (Coleoptera:Helophoridae) and *Hydraena* (Coleoptera:Hydraenidae) and larvae of *Riolus* (Coleoptera:Elmidae) were identified to genus; Planariidae (Tricladida), Sphaeriidae (Bivalvia), larval Dytiscidae and Haliplidae (Coleoptera), Baetidae (Ephemeroptera), and all Hemiptera and Diptera were identified to family; and Oligochaeta and Hydracarina were recorded at higher levels.

Data analysis

Two-way repeated-measures analysis of variance (RM ANOVA) with month as the within-subjects factor and site as the between-subjects factor was used to compare hydrological

and water-chemistry variables spatially and temporally for benthic and hyporheic habitats. For water-chemistry variables, RM ANOVA tests using habitat (benthic vs hyporheic) as the between-subjects factor were conducted on a combined benthic–hyporheic data set. Mauchly’s tests were used to verify the assumption of sphericity, and where this assumption was violated, p values calculated by Greenhouse–Geisser tests were used to assess significance. Where RM ANOVAs identified significant differences between groups, paired t -tests with Bonferroni adjustments were used to examine differences between individual months or sites. Where significant interactions of sites $\times$ months were identified by RM ANOVAs, temporal changes at individual sites were examined using 1-way ANOVAs with Tukey’s post hoc tests. All HZ and BZ assemblage analyses were undertaken separately because different sampling methods were used.

Studies in the Little Stour during the 2004–2006 drought showed significant changes in total taxon richness, total invertebrate abundance, and abundances of particular taxa in benthic and hyporheic zones (Stubbington et al. 2009, Wood et al. 2010). However, no site-level analysis of α diversity and β diversity has been conducted, nor have hypotheses about temporal changes in assemblage composition within sites and habitats been tested. Habitat-specific comparisons within and between sites over time can be explored in light of the differences in environmental conditions between the 2 zones as the drought proceeded.

To test our 1st hypothesis, site-level taxon richness (α diversity) was calculated for each sampling occasion as the mean of the 5 replicate samples and compared over time among sites with RM ANOVAs. Where significant site $\times$ month interactions were found, 1-way ANOVAs were conducted for individual sites, as described for environmental variables.

Many authors have argued that spatial and temporal *variation* in assemblage composition,

expressed as β diversity (in both its spatial and temporal senses; Anderson et al. 2011), is a more informative measure of the effects of disturbances than changes in α diversity (review in Séguin et al. 2014). Therefore, to test our 2nd hypothesis, benthic and hyporheic β diversity were assessed at each site with the Index of Multivariate Dispersion (IMD) proposed by Anderson et al. (2006). The IMD expresses variation in assemblage composition as the average dissimilarity between individual samples and their group centroid in multivariate ordination space (see below). One-way ANOVAs were then used to detect temporal changes in β diversity (as IMD).

Our 3rd hypothesis was tested by plotting sequential sample centroids in 2-dimensional ordination space to reveal temporal trajectories of changes in assemblage composition at each site. Data from each zone (i.e., 5 replicates $\times$ 7 sampling times $\times$ 4 sites = 140 samples/zone) were ordinated with nonmetric multidimensional scaling (NMDS) of a Bray–Curtis similarity matrix with 100 restarts. We used analysis of similarities (ANOSIM) of site-level data followed by pairwise comparisons (10,000 permutations) to test whether assemblage composition differed between consecutive sampling dates (i.e., divergence) or not (i.e., convergence). The main taxa contributing to differences between sites and over time within each habitat were ascertained with similarity percentages (SIMPER). Pearson correlation coefficients were calculated to examine relationships between NMDS axis scores and environmental variables.

All univariate analyses were done with IBM SPSS Statistics (version 19; IBM Corporation, Armonk, New York). All multivariate analyses (NMDS, IMD, ANOSIM, SIMPER) were done using $\sqrt[4]{(x)}$ -transformed data in PRIMER (version 6; PRIMER-E, Ivybridge, UK).

RESULTS

Environmental conditions during the drought

River discharge was below the long-term average ($0.181 \text{ m}^3/\text{s}$) in April and May, decreased rapidly during June and July, and was lowest in August (Fig. 1A). Local management of weirs and in-stream macrophytes between August and October meant that changes to mean water depth were limited during this period (Table 1) but depths varied among sites while flow velocity remained locally variable. The greatest changes were declines of $>5 \text{ cm}$ at the semiperennial sites, where riffle crests were exposed between July and September, although depth was never $<2 \text{ cm}$ at any sampling point (Table 1). After a lagged response to increased rainfall during August, discharge began to recover during September and remained relatively stable until the end of the study (Fig. 1A).

Water temperatures were higher in the surface stream ($15.7 \pm 0.3^\circ\text{C}$) than in the HZ ($14.9 \pm 0.3^\circ\text{C}$; RM ANOVA, $p = 0.001$) and peaked in July in both habitats (Table 1). In July, site-specific surface-water means varied from 20.0 to 21.9°C , and peak values reached 22.6°C , whereas in hyporheic water, site means were 19.3 to 21.3°C , and temperatures peaked at 21.7°C . Temperatures declined in August but experienced a 2nd, lower peak in September (Table 1, Fig. 1B). DO concentrations were lower in hyporheic ($7.3 \pm 0.2 \text{ mg/L}$) than in surface water ($9.4 \pm 0.1 \text{ mg/L}$, $p < 0.001$). Surface-water DO concentrations were higher in April ($12.2 \pm 0.1 \text{ mg/L}$) than in subsequent months ($p \leq 0.001$), whereas hyporheic DO concentrations remained high in May before declining in June (RM ANOVA, $p < 0.001$). This decline continued until August (Table 1). Surface and hyporheic water had similar pH (7.4 – 8.6 , RM ANOVA, $p = 0.952$), and monthly means ranged from 7.7 to 8.2 in both zones (Table 1). Conductivity was higher in hyporheic ($576 \text{ }\mu\text{S/cm}$) than in surface water ($568 \text{ }\mu\text{S/cm}$, RM ANOVA, $p < 0.001$) and was significantly higher in April than other months in both zones (RM ANOVAs, $p < 0.001$; Table 1).

Benthic assemblage composition

In total, 53,918 individuals from 89 taxa were recorded from 140 Surber samples. *Gammarus pulex* (Amphipoda:Gammaridae) dominated the assemblage (35% of total abundance), followed by Chironomidae larvae (Diptera, 14% of total abundance). Oligochaeta, *Agapetus fuscipes* (Trichoptera:Glossosomatidae), larval and adult *Elmis aenea* (Coleoptera:Elmidae), *Serratella ignita* (Ephemeroptera:Ephemerellidae); and Baetidae (Ephemeroptera) each accounted for >3% of the assemblage (data not shown). Two hypogean amphipods (<0.02% of all invertebrates) occurred in the BZ: *Crangonyx subterraneus* (Amphipoda:Crangonyctidae) (data not shown) and *Niphargus aquilex* (Amphipoda:Niphargidae) (Table 2).

Hyporheic assemblage composition

A total of 3140 invertebrates from 47 taxa were present in 140 hyporheic samples, with 1 to 13 taxa present per sample. *Gammarus pulex* was the most abundant taxon (46% of total abundance; Table 2). Oligochaeta, Chironomidae, and *A. fuscipes* also were abundant (11.8–12.6% each). *Elmis aenea* larvae, Baetidae, and *N. aquilex* were common (2–4% each). Four other hypogean taxa (*C. subterraneus*, *Niphargus fontanus*, *N. kochianus kochianus*, and *Proasellus cavaticus* [Isopoda:Asellidae]) were uncommon (collectively 1.0% of total hyporheic abundance) (data not shown).

Temporal trends in site-level benthic and hyporheic taxon richness (α diversity)

Benthic taxon richness (α diversity) varied significantly over time and among sites (RM ANOVA, $p < 0.001$), and was highest at perennial site 4 in all months except September (Fig.

2A). At all sites, α diversity declined gradually until July, when taxon richness was lowest at sites 2–4 (Fig. 2A). This decline was across multiple groups, including Diptera, Mollusca, and Trichoptera (Table 2). Taxon richness increased in August, mainly because of increases in Diptera, Coleoptera, Mollusca, and Trichoptera. A 2nd, moderate dip occurred in September at sites 1, 3, and 4 (Fig. 2A), reflecting minor reductions in the richness of several groups (Table 2). Temporal trends in α diversity varied among sites (RM ANOVA interaction, $p < 0.001$). Semiperennial site 2 deviated from the overall pattern, largely because richness increased from July to a September peak (Tukey's test, $p < 0.001$) at this site.

Hyporheic α diversity did not differ among sites from April to July ($p = 1$ for all pairwise comparisons; Fig. 2B, Table 2). Taxon richness decreased in August, with a significant decline at site 2 (1-way ANOVA, $p < 0.001$) caused largely by the absence of 4 uncommon crustaceans. Richness then rose sharply at all sites in September (RM ANOVA $p < 0.001$, Fig. 2B), especially in the Trichoptera and Diptera, and also in the Coleoptera and Crustacea (Table 2). In October, richness at all sites had returned to levels similar to those recorded in April (Fig. 2B). Taxa present (at low abundance in all cases) in September but absent in August and October included *Glossiphonia complanata* (Hirudinea:Glossiphoniidae), *Asellus aquaticus*, *A. merianus*, and *Proasellus cavaticus* (Crustacea:Asellidae), *N. fontanus*, the trichopterans *Hydropsyche angustipennis* (Hydropsychidae), *Athripsodes cinereus* (Leptoceridae) and *Polycentropus flavomaculatus* (Polycentropodidae), and the dipteran family Empididae. The α diversity of semiperennial sites did not differ from that of perennial sites, nor was the time $\times$ site interaction significant (RM ANOVA, $p > 0.05$).

Temporal trends in within-site variation in benthic and hyporheic assemblage composition

(β diversity)

β diversity, as spatial variation of assemblage composition among replicate samples collected monthly at each site, was expressed by the IMD. Considering all sites, and despite relatively stable values at site 1, the IMD varied significantly over time in the BZ (1-way ANOVA, $p = 0.023$) with higher values occurring in September than in June (Tukey's test, $p = 0.038$; Fig. 3A). Some patterns were consistent across sites, particularly at sites 2, 3 and 4. Benthic IMD was generally low between April and June, increased considerably in July, and remained fairly high after July (Fig. 3A). Hyporheic β diversity also varied among months (1-way ANOVA, $p = 0.032$), but temporal changes were not consistent among sites until July (Fig. 3B). At Sites 1–3, hyporheic IMD increased between July and August, decreased significantly in September (Tukey test, $p = 0.011$), and rose in October. A different pattern was observed at site 4, where a steady decline began in June and continued until September (Fig. 3B).

Temporal trends in between-site variation in benthic assemblage composition (β diversity)

NMDS ordination of benthic samples revealed marked differences in assemblage composition between sites 1, 4, and the 2 semiperennial sites (2 and 3), which overlapped slightly (Fig. 4A). Differences between all site pairs were significant (ANOSIM, global $R = 0.563$, $p < 0.001$) and reflected variation in the densities of *S. ignita*, Baetidae, *A. fuscipes*, and *E. aenea* larvae (SIMPER). Samples from site 4 had high axis-1 scores (Fig. 4A), which were significantly correlated ($p \leq 0.002$) with increasing depth and flow and declining pH. Semiperennial sites typically had higher axis-2 scores, which were significantly correlated ($p \leq 0.029$) with temperature and DO.

Temporal trajectories of change in benthic assemblage composition were broadly similar

across sites, with NMDS axis-2 scores declining until month 4 (July), when centroids of sites 1–3 plotted in close proximity. This assemblage was characterized by low richness of insect and noninsect taxa and low abundance of common taxa, such as *Oligochaeta* (Table 2). After July, trajectories indicated a partial return to the assemblage composition seen in April (Fig. 4A). The pattern at perennial site 1 was the most variable mainly because of divergence in September. This divergence also was evident at semiperennial site 3. Nonetheless, β diversity over time (evident as the spread of the 7 centroids representing each site) was comparable at perennial sites 1 and 4 and was lower than at the semiperennial sites (Fig. 4A).

Temporal change in assemblage composition was significant at site 1 (ANOSIM, global $R = 0.596$, $p < 0.001$), and the only significant overlap between consecutive months was for April and May ($p = 0.103$). Temporal change in assemblage composition also was significant at the other perennial site (4) ($R = 0.533$, $p < 0.001$), and the only significant overlap was between September and October ($p = 0.103$). At semiperennial site 2, assemblage composition changed over time ($R = 0.679$, $p < 0.001$), with no overlap between consecutive sampling dates. At semiperennial site 3, temporal changes were significant ($R = 0.518$, $p < 0.001$) with marked overlap between September and October. Despite the similar assemblage composition at sites 2 and 3 on the first 3 sampling dates (Fig. 4A), composition diverged at these sites after June.

Temporal trends in between-site variation in hyporheic assemblage composition (β diversity)

Hyporheic assemblage composition overlapped considerably among all sites (Fig. 4B), as indicated by nonsignificant ANOSIM scores between sites 1, 2, and 4 (pairwise global $R < 0.075$, Bonferroni-corrected $p > 0.05$). However, the temporal trajectories of change at individual sites

were similar. Axis 1 scores decreased at all sites between April and May. In June, axis 1 scores increased slightly at sites 1, 3, and 4. When the heat wave occurred in July, all site centroids plotted in a single cluster with low axis 1 and high axis 2 scores (Fig. 4B) associated with relatively high densities of *G. pulex* and hypogean crustaceans, especially *N. aquilex*. Between July and August, axis 2 scores declined considerably because many taxa, including *G. pulex* and *N. aquilex* became less abundant at both perennial and semiperennial sites. Scores then increased on both axes for all sites, and by September, samples had formed a distinct cluster with high axis 1 scores, associated with a 2nd peak in the abundance of *G. pulex* but not *N. aquilex*. Axis 2 scores then declined between September and October (Fig. 4B).

ANOSIM indicated significant temporal change in hyporheic assemblage composition at site 1 (global $R = 0.364$, $p < 0.001$), mainly between July and August and August and September ($p < 0.01$). These differences were associated with *N. aquilex*, *G. pulex*, *A. fuscipes*, and Chironomidae densities, all of which were considerably higher in July than August. The 2 amphipods and *A. fuscipes* were also more abundant in the HZ in September than October, although chironomid abundance increased during this period. Temporal changes also were significant in the HZ of perennial site 4 (global $R = 0.309$, $p < 0.001$) but only between the final 4 mo. Fluctuations in densities of oligochaetes, *A. fuscipes*, and larval *E. aenea* (particularly as September peaks) were primarily responsible for these significant differences (SIMPER).

Hyporheic assemblage composition also changed over time at the 2 semiperennial sites (global $R = 0.536$ for site 2, 0.355 for site 3, $p < 0.001$ for both) but assemblage composition did not differ among the first 3 sampling occasions within sites ($p \geq 0.151$). At site 3, a nonsignificant difference ($p = 0.357$) occurred between hyporheic assemblage composition in April and October, indicating potential convergence.

DISCUSSION

Benthic assemblage responses to supraseasonal drought

Our results supported our 1st hypothesis that benthic taxon richness (α diversity) would decrease steadily as the drought proceeded, whereas hyporheic α diversities would change little. A general decrease in benthic α diversity during the final stages of the drought (April–July) was apparent despite site-specific fluctuations in richness. The loss of some taxa can be attributed to seasonal aquatic insect emergence (Wood et al. 2010), but other losses pertain to drought-related environmental changes, including declining discharge and elevated water temperatures (Graeber et al. 2013). Gradual decline in α diversity is a common benthic invertebrate assemblage response to flow recession during drought and is caused by gradual reduction in habitat availability and heterogeneity (Wood and Armitage 2004, Dewson et al. 2007). Richness reductions reflect the loss of taxa susceptible to low flows and dewatering (Datry 2012).

Benthic α diversity was lowest in July because of low taxon richness of noninsect groups, such as Crustacea, Hirudinea, Mollusca (mainly gastropods), and many Insecta. This low richness coincided with the record heat wave conditions, indicating that water temperatures exceeded the thermal tolerance of many lotic taxa (Woodward et al. 2010), including common, eurytopic taxa, such as *G. pulex* (Maazouzi et al. 2011). Mouthon and Daufresne (2006) recorded similar significant reductions in gastropod and bivalve taxon richness in a large lowland river during a heat wave and suggested that these declines were caused by high water temperatures.

Benthic α diversity increased at all sites in August, coinciding with the lowest discharge and after the heat wave ended. Groups contributing to this increase included multiple insect orders, Hirudinea and in particular, Mollusca. This rapid partial recovery after the heat wave is in

contrast to the results of Mouthon and Daufresne (2006), who noted poor recovery of the mollusk population after a more prolonged and severe heat wave, and implies persistence in refugia (Robertson 2000). Very few snails or leeches occurred in the HZ in the Little Stour, but our riffle-focused sampling design excluded cooler, deeper habitats, such as pools that may have served as sources of recolonists, as shown by Robson et al. (2011).

Our 2nd hypothesis, that within-site variation in β diversity of the benthic assemblage would decline as the drought proceeded, especially at semiperennial sites, was rejected. At all sites except perennial site 1, within-site β diversity (expressed as IMD) typically was higher from July to October than before the heat wave. These increases in benthic IMD values reflected variable within-site abundances of common taxa, such as *G. pulex* and *A. fuscipes*. These taxa reached high densities at some sampling points, possibly because of patches of preferred flow velocities or abundant food resources (Dewson et al. 2007, Stubbington et al. 2011b).

At the larger spatial scale of between-site β diversity, NMDS ordination revealed marked differences in benthic assemblage composition between each perennial site and the 2 semiperennial sites. Despite these differences among sites, temporal trajectories of assemblage change were broadly consistent across sites during the drought. In particular, during the July heat wave, assemblage composition of 3 sites converged in the same quadrant of the ordination because of shared declines in taxon richness and abundance. However, this convergence was not statistically significant, and support for our 3rd hypothesis is equivocal. Postdrought, temporal trajectories generally began to return to an assemblage composition resembling that of the 1st sample, reflecting the described recovery in α diversity and increases in abundance.

Hyporheic assemblage responses to supraseasonal drought

Our hypothesis that hyporheic taxon richness (α diversity) would be stable over time was largely supported because no overall significant change in richness occurred between April and August (the last month before the drought broke). We expected little change in hyporheic α diversity because drought-induced environmental extremes would be dampened in the HZ of the groundwater-fed Little Stour. Hyporheic water temperatures were cooler than in the surface stream and hyporheic DO concentrations remained within the tolerance of most hypogean taxa until August (Malard and Hervant 1999, Issartel et al. 2005). In September, a marked but temporary peak in richness was observed because of the occurrence of multiple taxa at low abundance, including epigean and hypogean insect and noninsect taxa. The presence of epigean insects may reflect transient use of the HZ as a nursery for early instars (Malard et al. 2003), but the influx of the epigean isopods *A. aquaticus* and *A. meridianus* is unexplained. Hypogean crustaceans (*P. cavaticus*, *N. fontanus*) probably were carried into the upper HZ by an increase in the strength of upwelling groundwater concurrent with the increase in river discharge, as described by Plénet et al. (1995).

Hyporheic β diversity (as IMD) varied significantly over time at all sites but, as hypothesized, did not show any consistent response as the drought proceeded. As for α diversity, the environmental conditions in the HZ of this groundwater-fed stream apparently ameliorated the stressful extremes of water temperature (Robertson and Wood 2010, Stubbington and Wood 2013). In predominantly downwelling surface-fed streams, effects on the hyporheic invertebrate assemblage would be expected to be greater than observed in our study because of the higher temperature of downwelling water (Malard et al. 2002) and the greater influxes of benthic refugees (Davy-Bowker et al. 2006, Stubbington et al. 2011a). These predictions acknowledge the importance of the predominant direction of vertical hydrological exchange in determining

likely interactions between the surface stream and the underlying HZ (Brunke and Gosner 1997).

Our 3rd hypothesis was that little or no convergence would occur in hyporheic assemblage composition over time. This hypothesis was rejected. The NMDS plot of the temporal trajectories of monthly centroids representing site-level assemblage composition showed that despite the significant overlap in assemblage composition at sites 1, 2, and 4, some convergence was evident during the heat wave in July, when centroids from all 4 sites shared low axis 1 and high axis 2 scores. This convergence was associated with high hyporheic densities of the epigeal amphipod *G. pulex* and the hypogean amphipod *N. aquilex* in July (Wood et al. 2010). The influx of predominantly benthic amphipods indicated that the HZ served as a thermal refuge when water temperatures were $\geq 3.0^{\circ}\text{C}$ cooler than in the BZ, a result supporting the concept of the HZ as a refuge for benthos during multiple disturbance types (Dole-Olivier 2011, Stubbington 2012). The influx of the hypogean *N. aquilex* into the shallow HZ may reflect declining O_2 availability in deeper sediments and groundwater. Hyporheic sediments are typically more highly oxygenated than underlying aquifers, and reduced flows that increased groundwater residence times may have further depleted O_2 concentrations during drought (Malard and Hervant 1999, McGuire et al. 2002). As a result, vertical migrants escaping either thermally stressful surface water or poorly oxygenated groundwater colonized the intervening HZ. This result elegantly illustrates the complexity of faunal responses to drought disturbances in benthic, hyporheic, and phreatic habitats, and the importance of interhabitat connectivity in allowing migrations between adjacent ecosystem components (Dole-Oliver 2011, Stubbington 2012). Experiments in which benthic and hyporheic invertebrates are tagged with nontoxic biochemical markers and tracked along migration pathways within the sediments would permit direct observation of movements and refuge use during disturbances, such as drought.

In August, discharge was at its lowest, and temporal trajectories of hyporheic assemblage composition diverged widely from the July cluster as epigeal and hypogean refugees returned to their respective principal habitats when the heat wave ended. After the drought broke, the centroids of all sites converged and had high axis 1 scores, associated with a 2nd peak in *G. pulex* abundance, which, unlike the first peak, cannot be attributed to thermal stress or any other recorded environmental variable. Ordination scores then declined along axis 2 between the last 2 months of sampling. In October, the centroids for both semiperennial sites plotted near their April equivalents, indicating resilience of these assemblages to drought disturbance. In contrast, trajectories of assemblage composition continued to diverge from the initial composition at both perennial sites. However, the suprasedonal drought we studied in 2006 had its origins in a winter rainfall deficit in 2004–2005 (Marsh 2007) and changes to invertebrate assemblages had almost certainly taken place prior to the onset of our sampling.

Implications for river ecosystem management

Management of lotic ecosystems in the face of increasing climatic extremes typically has focused on flooding because of its dramatic impacts on human societies (Jones 2013). However, effective management also must preserve ecological resistance and resilience to the ‘creeping disaster’ of drought, especially in parts of the world where temperatures are rising and precipitation is declining (Lake 2005, Robson et al. 2011). Such management strategies should encompass protection of all instream components, including the fauna and biogeochemical processes occurring in hyporheic habitats. This protection is especially important because the HZ can act as a source of propagules for recolonization after floods and droughts (Dole-Olivier 2011, Stubbington and Datry 2013) and is a major site for the microbially mediated transformation of

nutrients and other materials in lotic ecosystems.

Our demonstration of the contrasting responses of benthic and hyporheic invertebrate assemblages to extreme drought, even in a groundwater-fed stream, highlights the need to consider the entire lotic assemblage—benthic and hyporheic—across a range of sites with differing flow histories when planning adaptation strategies for mitigating the effects of climate change or restoration programs following anthropogenic disturbances. Researchers and managers should not assume that benthic assemblage responses will resemble those of the hyporheos or other stream biota. Instead, these assemblages are likely to be connected and to share taxa that migrate between zones and seek different refugia at different times depending on the type and magnitude of a given disturbance. Therefore, we urge river managers to adopt strategies that explicitly recognize the vertical linkages connecting benthic biota and associated ecological processes to those in the underlying HZ and adjacent alluvial aquifers. For example, plans for the restoration of impaired river reaches should include specific predictions about recovering damaged hyporheic linkages and processes (Boulton 2007), especially where the effects of anthropogenic changes to streams are superimposed on large natural disturbances, such as suprasedasonal drought.

ACKNOWLEDGEMENTS

This research was supported by a Natural Environment Research Council Urgency Grant (NE/E001769/1) entitled ‘The response of aquatic invertebrate fauna to supra-seasonal drought and drying in a largely perennial chalk stream’. The authors acknowledge the support of the Environment Agency of England and Wales and particularly Ian Humpheryes, Shelagh Wilson, and Kevin Grimmett. Meteorological data for Manston (Kent) were provided by the British Atmospheric Data Centre (BADC). We are very grateful to Kathleen Rugel, Pam Silver and 3 referees for constructive advice on earlier drafts of this manuscript, Cath Leigh for assistance with data analyses, Sam Lake for discussion about the likely effects of drought on β diversity in streams, and Mark Szegner for the production of Fig. 2.

LITERATURE CITED

- Anderson, M. J., K. E. Ellingsen, and B. H. McArdle. 2006. Multivariate dispersion as a measure of beta diversity. *Ecology Letters* 9:683–693.
- Anderson, M. J., T. O. Crist, J. M. Chase, M. Vellend, B. D. Inouye, A. L. Freestone, N. J. Sanders, H. V. Cornell, L. S. Comita, K. F. Davies, S. P. Harrison, N. J. B. Kraft, J. C. Stegen, and N. G. Swenson. 2011. Navigating the multiple meanings of β diversity: a roadmap for the practicing ecologist. *Ecology Letters* 14:19–28.
- Belaidi, N., A. Taleb, and J. Gagneur. 2004. Composition and dynamics of hyporheic and surface fauna in a semi-arid stream in relation to the management of a polluted reservoir. *Annales de Limnologie – International Journal of Limnology* 40:237–248.
- Boulton, A. J. 2003. Parallels and contrasts in the effects of drought on stream macroinvertebrate assemblages. *Freshwater Biology* 48:1173–1185.
- Boulton, A. J. 2007. Hyporheic rehabilitation in rivers: restoring vertical connectivity. *Freshwater Biology* 52:632–650.
- Boulton, A. J., M.-J. Dole-Olivier, and P. Marmonier. 2004. Effects of sample volume and taxonomic resolution on assessment of hyporheic assemblage composition sampled using a Bou-Rouch pump. *Archiv für Hydrobiologie* 159:327–355.
- Boulton, A. J., S. Findlay, P. Marmonier, E. H. Stanley, and H. M. Valett. 1998. The functional significance of the hyporheic zone in streams and rivers. *Annual Review of Ecology and Systematics* 29:59–81.
- Boulton, A. J., and E. H. Stanley. 1995. Hyporheic processes during flooding and drying in a Sonoran Desert stream. I. Hydrologic and chemical dynamics. *Archiv für Hydrobiologie* 134:1–26.

509 Brunke, M., and T. Gonser. 1997. The ecological significance of exchange processes between
 510 rivers and groundwater. *Freshwater Biology* 37:1–33.

511 Brunke, M., and T. Gonser. 1999. Hyporheic invertebrates: the clinal nature of interstitial
 512 communities structured by hydrological exchange and environmental gradients. *Journal*
 513 *of the North American Benthological Society* 18:344–362.

514 Chase, J. M. 2007. Drought mediates the importance of stochastic community assembly.
 515 *Proceedings of the National Academy of Sciences of the United States of America*
 516 104:17430–17434.

517 Davy-Bowker, J., W. Sweeting, N. Wright, R. T. Clarke, and S. Arnott. 2006. The distribution of
 518 benthic and hyporheic macroinvertebrates from the heads and tails of riffles.
 519 *Hydrobiologia* 563:109–123.

520 Datry, T. 2012. Benthic and hyporheic invertebrate assemblages along a flow intermittence
 521 gradient: effects of duration of dry events. *Freshwater Biology* 57:563–574.

522 Dewson, Z. S., A. B. W. James, and R. G. Death. 2007. A review of the consequences of
 523 decreased flow for instream habitat and macroinvertebrates. *Journal of the North*
 524 *American Benthological Society* 26:401–415.

525 Dole-Olivier, M.-J. 2011. The hyporheic refuge hypothesis reconsidered: a review of
 526 hydrological aspects. *Marine and Freshwater Research* 62:1281–1302.

527 Fleig, A. K., L. M. Tallaksen, H. Hisdal, and D. M. Hannah. 2010. Regional hydrological
 528 drought in north-western Europe: linking a new Regional Drought Area Index with
 529 weather types. *Hydrological Processes* 25:1163–1179.

530 Graeber, D., M. T. Pusch, S. Lorenzo, and M. Brauns. 2013. Cascading effects of flow reduction
 531 on the benthic invertebrate community in a lowland river. *Hydrobiologia* 717:147–159.

532 Issartel, J., F. Hervant, Y. Voituron, D. Renault, and P. Vernon. 2005. Behavioural, ventilator and
533 respiratory responses of epigean and hypogean crustaceans to different temperatures.
534 *Comparative Biochemistry and Physiology: Part A* 141:1–7.

535 James, A. B. W., Z. S. Dewson, and R. G. Death. 2008. Do stream macroinvertebrates use
536 instream refugia in response to severe short-term flow reduction in New Zealand
537 streams? *Freshwater Biology* 53:1316–1334.

538 Jones, I. (editor). 2013. The impact of extreme events on freshwater ecosystems. *Ecological*
539 *Issues* no. 12. British Ecological Society, London, UK.

540 Lake, P. S. 2003. Ecological effects of perturbation by drought in flowing waters. *Freshwater*
541 *Biology* 48:1161–1172.

542 Lake, P. S. 2005. Perturbation, restoration and seeking ecological sustainability in Australian
543 flowing waters. *Hydrobiologia* 552:109–120.

544 Lake, P. S. 2011. Drought and aquatic ecosystems: effects and responses. Wiley-Blackwell,
545 Oxford, UK.

546 Maazouzi, C., C. Piscart, F. Legier, and F. Hervant. 2011. Ecophysiological responses to
547 temperature of the “killer shrimp” *Dikerogammarus villosus*: Is the invader really
548 stronger than the native *Gammarus pulex*? *Comparative Biochemistry and Physiology*
549 *Part A: Molecular and Integrative Physiology* 159:268–274.

550 Malard, F., D. Galassi, M. Lafont, S. Dolédec, and J. V. Ward. 2003. Longitudinal patterns of
551 invertebrates in the hyporheic zone of a glacial river. *Freshwater Biology* 48:1709–1725.

552 Malard, F., and F. Hervant. 1999. Oxygen supply and the adaptations of animals in groundwater.
553 *Freshwater Biology* 41:1–30.

554 Malard F., K. Tockner, M.-J. Dole-Oliver, and J. V. Ward. 2002. A landscape perspective of

555 surface-subsurface hydrological exchanges in river corridors. *Freshwater Biology*
556 47:621–640.

557 Marsh, T. 2007. The 2004–2006 drought in southern Britain. *Weather* 62:191–196.

558 McGuire, K. J., D. R. DeWalle, and W. J. Gburek. 2002. Evaluation of mean residence time in
559 subsurface waters using oxygen-18 fluctuations during drought conditions in the mid-
560 Appalachians. *Journal of Hydrology* 261:132–149.

561 Mouthon, J., and M. Daufresne. 2006. Effects of the 2003 heatwave and climatic warming on
562 mollusc communities of the Saône: a large lowland river and of its two main tributaries
563 (France). *Global Change Biology* 12:441–449.

564 Plénet, S., J. Gibert, and P. Marmonier. 1995. Biotic and abiotic interactions between surface and
565 interstitial systems in rivers. *Ecography* 18:296–309.

566 Prior, J., and M. Beswick. 2007. The record-breaking heat and sunshine of July 2006. *Weather*
567 62:174–182.

568 Robertson, A. L. 2000. Lotic meiofaunal community dynamics: colonisation, resilience and
569 persistence in a spatially and temporally heterogeneous environment. *Freshwater Biology*
570 44:135–147.

571 Robertson, A. L., and P. J. Wood. 2010. Ecology of the hyporheic zone: origins, current
572 knowledge and future directions. *Fundamental and Applied Limnology* 176:279–289.

573 Robson, B. J., E. T. Chester, and C. M. Austin. 2011. Why life history information matters:
574 drought refuges and macroinvertebrate persistence in non-perennial streams subject to a
575 drier climate. *Marine and Freshwater Research* 62:801–810.

576 Séguin, A., D. Gravel, and P. Archambault. 2014. Effect of disturbance regime on alpha and beta
577 diversity of rock pools. *Diversity* 6:1–17.

578 Stubbington, R. 2012. The hyporheic zone as an invertebrate refuge: a review of variability in
579 space, time, taxa and behaviour. *Marine and Freshwater Research* 63:293–311.

580 Stubbington, R., and T. Datry. 2013. The macroinvertebrate seedbank promotes community
581 persistence in temporary rivers across climate zones. *Freshwater Biology* 58:1202–1220.

582 Stubbington, R., and P. J. Wood. 2013. Benthic and interstitial habitats of a lentic spring as
583 invertebrate refuges during supra-seasonal drought. *Fundamental and Applied Limnology*
584 182:61–73.

585 Stubbington, R., P. J. Wood, and A. J. Boulton. 2009. Low flow controls on benthic and
586 hyporheic macroinvertebrate assemblages during supra-seasonal drought. *Hydrological*
587 *Processes* 23:2252–2263.

588 Stubbington, R., P. J. Wood, and I. Reid. 2011a. Spatial variability in the hyporheic zone
589 refugium of temporary streams. *Aquatic Sciences* 73:499–511.

590 Stubbington, R., P. J. Wood, I. Reid, and J. Gunn. 2011b. Benthic and hyporheic invertebrate
591 community responses to seasonal flow recession in a groundwater-dominated stream.
592 *Ecohydrology* 4:500–511.

593 Williams, D. D., C. M. Febria, and J. C. Y. Wong. 2010. Ecotonal and other properties of the
594 hyporheic zone. *Fundamental and Applied Limnology* 176:349–364.

595 Williams, D. D., and H. B. N. Hynes. 1974. The occurrence of benthos deep in the substratum of
596 a stream. *Freshwater Biology* 4:233–256.

597 Wood, P. J., M. D. Agnew, and G. E. Petts. 2000. Flow variations and macroinvertebrate
598 community responses in a small groundwater-dominated stream in south-east England.
599 *Hydrological Processes* 14:3133–3147.

600 Wood, P. J., and P. D. Armitage. 2004. The response of the macroinvertebrate community to low-

601 flow variability and supra-seasonal drought within a groundwater dominated stream.
602 Archiv für Hydrobiologie 161:1–20.

603 Wood, P. J., A. J. Boulton, S. Little, and R. Stubbington. 2010. Is the hyporheic zone a refugium
604 for aquatic macroinvertebrates during severe low flow conditions? Fundamental and
605 Applied Limnology 176:377–390.

606 Wood, P. J., and G. E. Petts. 1999. The influence of drought on chalk stream macroinvertebrates.
607 Hydrological Processes 13:387–399.

608 Woodward, G., D. M. Perkins, and L. E. Brown. 2010. Climate change and freshwater
609 ecosystems: impacts across multiple levels of organization. Philosophical Transactions of
610 the Royal Society of London Series B: Biological Sciences 365:2093–2106.

611 Young, B. A., R. H. Norris, and F. Sheldon. 2011. Is the hyporheic zone a refuge for
612 macroinvertebrates in drying perennial streams? Marine and Freshwater Research
613 62:1373–1382.

614

FIGURE CAPTIONS

Fig. 1. Environmental conditions in the Little Stour River in 2006. A.—Daily discharge and precipitation at Littlebourne (lat 51.277°N, long 1.167°E). B.—Twelve-hourly maximum temperatures at Manston (lat 51.344°N, long 1.369°E).

Fig. 2. Mean (± 1 SE) taxon richness (α diversity) of 5 replicate samples from 2 perennial sites (1 and 4) and 2 semiperennial sites (2 and 3) in the benthic zone (A) and hyporheic zone (B) in the Little Stour River. Note the difference in vertical scales between panels.

Fig. 3. Temporal changes in Index of Multivariate Dispersion (IMD; β diversity) of 5 replicate samples from 2 perennial sites (1 and 4) and 2 semiperennial sites (2 and 3) in the benthic zone (A) and hyporheic zone (B) in the Little Stour River.

Fig. 4. Two-dimensional plot of centroids derived from nonmetric multidimensional scaling ordination of benthic (A) and hyporheic (B) invertebrate assemblage composition from 2 perennial sites (1 and 4) and 2 semiperennial sites (2 and 3). Site-level centroids of 5 replicates for each month (April–October: 1–7) are linked by lines representing temporal trajectories of change during the final stages of drought (months 1–5) and after the drought broke (months 6 and 7).

633 Table 1. Monthly mean (± 1 SE) values for hydrological and water chemistry variables ($n = 5$ per
634 site each month) at 4 sites along the Little Stour River. SW = surface water, HW = hyporheic
635 water.

Variable	SW/HW	April	May	June	July	August	September	October
Depth (cm)	SW	9.1 ± 0.7	8.9 ± 0.6	8.0 ± 0.7	8.1 ± 0.6	9.5 ± 1.0	6.5 ± 0.7	7.1 ± 0.5
Velocity (cm/s)	SW	36 ± 3	36 ± 5	23 ± 2	24 ± 3	29 ± 4	27 ± 4	18 ± 2
Temperature ($^{\circ}\text{C}$)	SW	9.9 ± 0.1	14.1 ± 0.2	19.0 ± 0.5	21.2 ± 0.2	15.8 ± 0.1	16.9 ± 0.1	13.0 ± 0.1
	HW	9.3 ± 0.1	13.3 ± 0.2	18.0 ± 0.5	20.0 ± 0.2	14.9 ± 0.2	15.8 ± 0.2	12.8 ± 0.1
Dissolved O_2 (mg/L)	SW	12.2 ± 0.1	9.9 ± 0.2	8.8 ± 0.2	9.1 ± 0.2	7.8 ± 0.3	8.7 ± 0.1	9.5 ± 0.2
	HW	9.5 ± 0.4	9.3 ± 0.3	8.1 ± 0.2	7.0 ± 0.2	5.3 ± 0.2	5.7 ± 0.2	6.3 ± 0.2
pH	SW	7.7 ± 0.01	8.2 ± 0.06	8.0 ± 0.05	7.9 ± 0.05	7.8 ± 0.05	7.8 ± 0.05	8.0 ± 0.05
	HW	7.7 ± 0.03	8.2 ± 0.04	8.0 ± 0.05	7.9 ± 0.05	7.8 ± 0.04	7.8 ± 0.05	8.1 ± 0.05
Conductivity ($\mu\text{S}/\text{cm}$)	SW	649 ± 7	526 ± 2	556 ± 3	560 ± 1	562 ± 3	563 ± 3	563 ± 2
	HW	662 ± 5	539 ± 1	563 ± 3	565 ± 1	567 ± 3	572 ± 2	565 ± 2

636

637

638 Table 2. Monthly mean (± 1 SE) total and taxon-specific invertebrate abundances and higher-
639 level taxon richness on the Little Stour River. BZ = benthic zone, HZ = hyporheic zone. Values
640 are per sample: either /0.1 m² (BZ) or /6 L (HZ).

	Zone	April	May	June	July	August	September	October
Abundance								
All invertebrates	BZ	363 $\pm$ 42	504 $\pm$ 55	432 $\pm$ 38	198 $\pm$ 31	375 $\pm$ 43	438 $\pm$ 47	386 $\pm$ 47
	HZ	14 $\pm$ 2	13 $\pm$ 2	19 $\pm$ 3	38 $\pm$ 5	11 $\pm$ 1	51 $\pm$ 5	13 $\pm$ 2
<i>Erpobdella octoculata</i>	BZ	3.7 $\pm$ 0.7	6.3 $\pm$ 1.5	4.4 $\pm$ 0.9	1.7 $\pm$ 0.4	5.7 $\pm$ 1.4	3.3 $\pm$ 0.9	4.0 $\pm$ 0.9
<i>Gammarus pulex</i>	BZ	80 $\pm$ 8	154 $\pm$ 30	204 $\pm$ 26	117 $\pm$ 21	122 $\pm$ 22	129 $\pm$ 30	125 $\pm$ 13
	HZ	3.4 $\pm$ 0.4	5.1 $\pm$ 1.7	8.5 $\pm$ 2.1	22.7 $\pm$ 2.4	3.7 $\pm$ 0.6	26.3 $\pm$ 2.5	3.5 $\pm$ 0.6
Gastropoda	BZ	1.9 $\pm$ 0.5	2.5 $\pm$ 0.9	1.5 $\pm$ 0.4	0.3 $\pm$ 0.1	2.5 $\pm$ 0.5	1.8 $\pm$ 0.5	0.8 $\pm$ 0.4
<i>Niphargus aquilex</i>	HZ	0 $\pm$ 0	0.3 $\pm$ 0.1	0.2 $\pm$ 0.1	3.0 $\pm$ 0.8	0.05 $\pm$ 0.05	0.1 $\pm$ 0.1	0 $\pm$ 0
Oligochaeta	BZ	43 $\pm$ 10	51 $\pm$ 14	51 $\pm$ 14	8 $\pm$ 2	35 $\pm$ 7	66 $\pm$ 20	43 $\pm$ 12
	HZ	3.4 $\pm$ 0.6	2.8 $\pm$ 0.7	3.9 $\pm$ 0.7	1.1 $\pm$ 0.4	1.4 $\pm$ 0.4	5.0 $\pm$ 0.7	1.1 $\pm$ 0.5
Planariidae	BZ	5.5 $\pm$ 1.5	3.2 $\pm$ 0.8	3.3 $\pm$ 0.9	1.0 $\pm$ 0.3	4.7 $\pm$ 0.8	3.0 $\pm$ 1.1	7.4 $\pm$ 3.1
Higher-level taxon richness								
Coleoptera	BZ	3.4 $\pm$ 0.3	3.3 $\pm$ 0.3	3.1 $\pm$ 0.4	2.5 $\pm$ 0.3	3.3 $\pm$ 0.3	3.0 $\pm$ 0.4	2.9 $\pm$ 0.3
	HZ	0.5 $\pm$ 0.2	0.2 $\pm$ 0.1	0.4 $\pm$ 0.2	0.2 $\pm$ 0.1	0.5 $\pm$ 0.1	1.5 $\pm$ 0.1	0.5 $\pm$ 0.1
Crustacea	BZ	1.7 $\pm$ 0.2	2.1 $\pm$ 0.2	1.9 $\pm$ 0.2	1.5 $\pm$ 0.2	1.2 $\pm$ 0.1	1.4 $\pm$ 0.1	2.3 $\pm$ 0.3
	HZ	1.1 $\pm$ 0.1	1.4 $\pm$ 0.2	1.3 $\pm$ 0.1	2.2 $\pm$ 0.2	1.1 $\pm$ 0.1	1.4 $\pm$ 0.2	1.2 $\pm$ 0.1
Diptera	BZ	2.3 $\pm$ 0.2	2.5 $\pm$ 0.2	2.6 $\pm$ 0.2	1.8 $\pm$ 0.2	2.5 $\pm$ 0.2	1.9 $\pm$ 0.2	1.9 $\pm$ 0.2
	HZ	1.0 $\pm$ 0.2	1.1 $\pm$ 0.1	1.1 $\pm$ 0.2	0.9 $\pm$ 0.2	0.8 $\pm$ 0.1	1.3 $\pm$ 0.1	1.0 $\pm$ 0.1
Ephemeroptera	BZ	3.0 $\pm$ 0.2	2.7 $\pm$ 0.2	2.1 $\pm$ 0.1	1.6 $\pm$ 0.2	1.7 $\pm$ 0.2	1.5 $\pm$ 0.2	2.1 $\pm$ 0.2
Hirudinea	BZ	1.8 $\pm$ 0.2	1.8 $\pm$ 0.2	1.9 $\pm$ 0.2	1.4 $\pm$ 0.2	1.9 $\pm$ 0.2	1.6 $\pm$ 0.2	1.7 $\pm$ 0.3
Mollusca	BZ	1.9 $\pm$ 0.2	1.0 $\pm$ 0.2	1.4 $\pm$ 0.3	0.3 $\pm$ 0.1	2.1 $\pm$ 0.4	1.3 $\pm$ 0.4	0.6 $\pm$ 0.2
Trichoptera	BZ	6.2 $\pm$ 0.4	5.1 $\pm$ 0.4	3.5 $\pm$ 0.3	2.3 $\pm$ 0.3	3.6 $\pm$ 0.3	3.8 $\pm$ 0.5	3.6 $\pm$ 0.4
	HZ	0.9 $\pm$ 0.2	0.3 $\pm$ 0.1	0.4 $\pm$ 0.1	0.7 $\pm$ 0.2	0.6 $\pm$ 0.1	1.4 $\pm$ 0.2	0.7 $\pm$ 0.2

Fig. 1A

Fig. 1B

Fig. 3

Fig. 4