Supporting Information

A Chelation-Assisted Transformation : Synthesis of 1,4-Dicarboxylate Esters by the Rh-Catalyzed Carbonylation of Internal Alkynes with 2-Pyridinylmethanol

Satoshi Inoue, Kazuhiko Yokota, Hiroto Tatamidani, Yoshiya Fukumoto, and Naoto Chatani*

*Department of Applied Chemistry, Faculty of Engineering, Osaka University,
Suita, Osaka 565-0871, Japan*

General Information ^1H NMR and ^{13}C NMR spectra were recorded on a JEOL JMN-270 spectrometer in CDCl_3 with tetramethylsilane as an internal standard. Data are reported as follows: chemical shift in ppm (δ), multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, and c = complex), coupling constant (Hz), integration, and interpretation. Infrared spectra (IR) were obtained on a Horiba FT-700 spectrometer. Mass spectra were obtained on a Shimadzu GCMS-QP 5000 instrument with ionization voltages of 70 eV. Elemental analyses were performed by the Elemental Analysis Section of Osaka University. High resolution mass spectra (HRMS) were obtained on a JEOL JMS-DX303. Analytical GC was carried out on a Shimadzu GC-14B gas chromatography, equipped with a flame ionization detector. Melting points were determined on a Yamato melting point apparatus and are uncorrected. Flash column chromatography was performed with SiO_2 (Merck SilicaGel 60 (230-400 mesh)).

Materials Unless otherwise noted, all reagents were obtained commercial suppliers and used as received. Toluene was distilled over CaH_2 . Pyridin-2-yl-methanol, $\text{HRh}(\text{CO})(\text{PPh}_3)_3$ and 3-Hexyne were purchased from Ardrich Chemical Co. $\text{Rh}_4(\text{CO})_{12}$, $\text{Rh}_6(\text{CO})_{16}$, $\text{RhCl}(\text{PPh}_3)_3$ and $\text{Co}_2(\text{CO})_8$ was purchased from Strem Chemicals Inc. 4-Octyne, 3-Octyne, 2-Octyne, 1-Octyne, 1-(trimethylsilyl)-1-propyne and PdCl_2 were purchased from Wako Pure Chemical Industries, Ltd. 1-Phenyl-1-hexyne, Diphenylacetylene and Diethyl acetylenedicarboxylate were purchased from Tokyo Kasei Kogyo Co., Ltd. Benzyl alcohol and Pyridine were purchased from Nacalai Tesque, Inc. 2,2-Dimethyl-3-pentyne was purchased from Lancaster Synthesis, Inc. 1-Hex-1-ynyl-4-methoxybenzene, 1-Hex-1-ynyl-4-trifluoromethylbenzene and 3,3-dimethyl-1-but-1-ynylbenzene were prepared by a sonogashira reaction. Cyclododecyne was prepared by the method of Brandsma with minor modification. (L. Brandsma; H. D. Verkruijsse, *Synthesis of Acetylenes, Allenes and Cumulenes*; Elsevier, Amsterdam 1981, Vol 8, pp119.)

Typical Procedure for Hydroesterification of Alkynes. $\text{Rh}_4(\text{CO})_{12}$ (30.0 mg, 0.04 mmol), pyridin-2-yl-methanol (218 mg, 2.00 mmol), 4-octyne (220 mg, 2.00 mmol) and toluene (1 mL) were placed in a 50-mL stainless steel autoclave under N_2 . The system was flushed with 10 atm of carbon monoxide three times. Finally, it was pressurized to 3 atm. The autoclave was heated in an oil bath at 100 °C for 20 h, followed by cooling for rt. Then CO was released. The contents were transferred to a round-bottom flask with EtOAc or CH_2Cl_2 , and the volatiles were removed in vacuo. The residue was subjected to flash column chromatography on silica-gel (eluent; Hexane/EtOAc = 5/1→0/1) to give *meso*-2,3-dipropyl-succinic acid dipyridin-2-yl-methyl ester (**3a**) (119 mg, 31 % yield based on

pyridin-2-yl-methanol) as a pale solid and *dl*-2,3-dipropyl-succinic acid dipyridin-2-yl-methyl ester (**3a'**) (237 mg, 62 % yield based on pyridin-2-yl-methanol) as a pale solid. The stereochemistry was assigned by ^{13}C NMR. (R. Quermann, R. Maletz, and H. J. Schäfer, *Liebigs Ann. Chem.* **1993**, 1219-1223.)

Procedure for equation 4 $\text{Rh}_4(\text{CO})_{12}$ (15.0 mg, 0.02 mmol), pyridin-2-yl-methanol (109 mg, 1.00 mmol), 2-Propyl-hex-2-enoic acid pyridin-2-yl-methylester (**4a**) (247 mg, 1.00 mmol) and toluene (1 mL) were placed in a 50-mL stainless steel autoclave under N_2 . The system was flushed with 10 atm of carbon monoxide three times. Finally, it was pressurized to 3 atm. The autoclave was heated in an oil bath at 100 °C for 20 h, followed by cooling for rt. Then CO was released. The contents were transferred to a round-bottom flask with EtOAc and the volatiles were removed in vacuo. The residue was subjected to flash column chromatography on silica-gel (eluent; Hexane/EtOAc = 8/1→2/1) to give 2-Propyl-hex-2-enoic acid pyridin-2-yl-methylester (**4a**) (234 mg, 95.5 % yield) as a colorless oil.

Procedure for equation 5 $\text{Rh}_4(\text{CO})_{12}$ (30.0 mg, 0.04 mmol), pyridin-2-yl-methanol (109 mg, 1.00 mmol), 2-Propyl-hex-2-enoic acid pyridin-2-yl-methylester (**4a**) (247 mg, 1.00 mmol), 3-hexyne (82 mg, 1.00 mmol) and toluene (1 mL) were placed in a 50-mL stainless steel autoclave under N_2 . The system was flushed with 10 atm of carbon monoxide three times. Finally, it was pressurized to 3 atm. The autoclave was heated in an oil bath at 100 °C for 20 h, followed by cooling for rt. Then CO was released. The contents were transferred to a round-bottom flask with EtOAc and the volatiles were removed in vacuo. The residue was subjected to flash column chromatography on silica-gel (eluent; Hexane/EtOAc = 6/1→1/5) to give 2-Propyl-hex-2-enoic acid pyridin-2-yl-methylester (**4a**) (205 mg, 83% yield) as a colorless oil and 2,3-diethyl-succinic acid dipyridin-2-yl-methyl ester (**3b**) (100 mg, 56% yield based on pyridin-2-yl-methanol) as a colorless oil.

Procedure for equation 6 $\text{Rh}_4(\text{CO})_{12}$ (15.0 mg, 0.02 mmol), pyridin-2-yl-methanol (109 mg, 1.00 mmol), 3,4-dipropyl-5-(pyridin-2-yl-methoxy)-5*H*-furanone (**5a**) (128 mg, 0.46 mmol) and toluene (0.5 mL) were placed in a 50-mL stainless steel autoclave under N_2 . The system was flushed with 10 atm of carbon monoxide three times. Finally, it was pressurized to 3 atm. The autoclave was heated in an oil bath at 100 °C for 20 h, followed by cooling for rt. Then CO was released. The contents were transferred to a round-bottom flask with EtOAc and the volatiles were removed in vacuo. The residue was subjected to flash column chromatography on silica-gel (eluent; Hexane/EtOAc = 4/1→3/2) to give 3,4-dipropyl-5-(pyridin-2-yl-methoxy)-5*H*-furanone (**5a**) (121 mg, 94% yield) as a pale yellow oil.

Procedure for equation 7 $\text{Rh}_4(\text{CO})_{12}$ (30.0 mg, 0.04 mmol), pyridin-2-yl-methanol (109 mg, 1.00 mmol), 3,4-dipropyl-5-(pyridin-2-yl-methoxy)-5*H*-furanone (**5a**) (124 mg, 0.45 mmol), 3-hexyne (82 mg, 1.00 mmol) and toluene (1 mL) were placed in a 50-mL stainless steel autoclave under N_2 . The system was flushed with 10 atm of carbon monoxide three times. Finally, it was pressurized to 3 atm. The autoclave was heated in an oil bath at 100 °C for 20 h, followed by cooling for rt. Then CO was released. The contents were transferred to a round-bottom flask with EtOAc and the volatiles were removed in vacuo. The residue was subjected to flash column chromatography on silica-gel (eluent; Hexane/EtOAc = 6/1→0/1) to give 3,4-dipropyl-5-(pyridin-2-yl-methoxy)-5*H*-furanone (**5a**) (107 mg, 86% yield) as a pale yellow oil and 2,3-diethyl-succinic acid dipyridin-2-yl-methyl ester (**3b**) (124 mg, 69% yield based on pyridine-2-yl-methanol) as a colorless oil.

***dl*-2,3-dipropyl-succinic acid dipyridin-2-yl-methyl ester (3a)**

orange solid. mp 67 - 68 °C. *R_f* 0.14 (hexane/EtOAc = 1/1). ¹H NMR (CDCl₃, TMS) δ: 8.55 (d, *J*=4.9 Hz, 2H), 7.67 (td, *J*=7.6, 2.7 Hz, 2H), 7.35 (d, *J*=8.1 Hz, 2H), 7.21 (m, 2H), 5.25 (d, *J*=13.5 Hz, 1H), 5.19 (d, *J*=13.5 Hz, 1H), 2.93 – 1.84 (m, 2H), 1.67 – 1.60 (m, 4H), 1.38 – 1.22 (m, 4H), 0.90 (m, 6H). ¹³C NMR (CDCl₃) δ: 174.38, 157.75, 148.99, 136.99, 122.85, 121.81, 66.66, 46.31, 31.27, 20.71,

14.05. IR (KBr): 2958 m, 2875 m, 1729 s, 1594 s, 1571 w, 1438 m, 1344 s, 1311 s, 1261 s, 1213 w, 1166 s, 1099 w, 1049 w, 1020 w, 985 s, 943 w, 757 s. MS, *m/z* (relative intensity, %): 384 (M⁺, 1.5), 276 (111), 192 (12), 164 (14), 110 (17), 109 (11), 108 (13), 93 (19), 92 (100), 69 (15), 65 (26), 55 (21). Anal. Calcd for C₂₂H₂₈N₂O₄ : C,68.73; H,7.34; N,7.29. Found: C,68.70; H,7.30; N,7.25.

***meso*-2,3-dipropyl-succinic acid dipyridin-2-yl-methyl ester (3a)**

pale yellow solid. mp 77- 78 °C. *R_f* 0.22 (hexane/EtOAc = 1/1). ¹H NMR (CDCl₃, TMS) δ: 8.59 (d, *J*=5.1 Hz, 2H), 7.74 (td, *J*=7.7, 1.9 Hz, 2H), 7.39 (d, *J*=7.7 Hz, 2H) 7.26 (m, 2H), 5.27 (s, 4H), 2.82 – 2.79 (m, 2H), 1.68 – 1.64 (m, 4H), 1.42 – 1.20 (m, 4H), 0.88 – 0.84 (m, 6H). ¹³C NMR (CDCl₃) δ: 173.87, 155.41, 148.92, 137.23, 123.07, 122.14, 66.50, 48.12, 32.89, 20.59, 13.82. IR (KBr): 2958 m,

2861 m, 1729 s, 1594 s, 1571 w, 1436 m, 1344 s, 1311 s, 1261 s, 1213 w, 1166 s, 1099 w, 1049 w, 1020 w, 985 s, 943 w, 881 w, 759 s, 655 w, 626 w, 582 s. MS, *m/z* (relative intensity, %): 384 (M⁺, 1.5), 276 (11), 192 (12), 164 (14), 110 (17), 109 (11), 108 (13), 93 (19), 92 (100), 69 (15), 65 (26), 55 (21). Anal. Calcd for C₂₂H₂₈N₂O₄ : C,68.73; H,7.34; N,7.29. Found: C,68.62; H,7.25; N,7.27.

2-Propyl-hex-2-enoic acid pyridine-2-yl-methylester (4a)

colorless oil. *R_f* 0.48 (hexane/EtOAc = 1/1). ¹H NMR (CDCl₃) δ: 8.72 (d, *J*=4.3 Hz, 1H), 7.70 (td, *J*=7.6, 1.9 Hz, 1H), 7.35 (d, *J*=7.8 Hz, 1H), 7.22 (t, *J*=5.1 Hz, 1H), 5.87 (t, *J*=7.6 Hz, 1H), 5.30 (s, 2H), 2.32 (t, *J*=7.3 Hz, 2H), 2.18 (q, *J*=7.3 Hz, 2H), 1.52 – 1.40 (m, 4H), 0.97 – 0.89 (c, 6H). ¹³C NMR (CDCl₃) δ: 167.42, 156.23, 149.12, 143.56, 136.65, 131.86, 122.56, 121.36, 66.67, 30.75, 28.84, 22.62, 22.15, 14.09, 14.06. IR (neat): 2960 s, 2933 w, 2871 m, 1714 s, 1643 w,

1592 m, 1459 m, 1438 m, 1378 m, 1280 s, 1214 s, 1143 s, 1108 m, 757 s. MS, *m/z* (relative intensity, %): 247 (M⁺, 5.3), 139 (11), 138 (41), 123 (53), 110 (15), 109 (28), 108 (100), 95 (18), 93 (41), 92 (56), 82 (22), 81 (31), 79 (13), 69 (47), 68 (17), 67 (32), 66 (11), 65 (49), 55 (45), 53 (24). Anal. Calcd for C₁₅H₂₁NO₂ : C,72.84; H,8.56; N,5.66. Found: C,72.56; H,8.32; N,5.66.

3,4-dipropyl-5-(pyridin-2-yl-methoxy)-5H-furanone (5a)

pale yellow oil. *R_f* 0.34 (hexane/EtOAc = 1/1). ¹H NMR (CDCl₃) δ: 8.57 (d, *J*=4.9 Hz, 1H), 7.71 (td, *J*=7.8, 1.9 Hz, 1H), 7.41 (d, *J*=7.8 Hz, 1H), 7.23 (m, 1H), 5.86 (s, 1H), 4.95 (d, *J*=12.5 Hz, 1H), 4.83 (d, *J*=12.5 Hz, 1H), 2.39 (t, *J*=7.3 Hz, 2H), 2.25 (t, *J*=7.3 Hz, 2H), 1.61 – 1.47 (m, 4H), 0.94 (m, 6H). ¹³C

NMR (CDCl₃) δ : 172.01, 158.00, 156.68, 149.26, 136.75, 130.85, 122.84, 121.86, 101.50, 71.97, 28.09, 25.52, 21.39, 20.92, 14.17, 13.95. IR (neat): 2962 s, 2935 m, 2873 w, 1768 s, 1679 m, 1592 m, 1463 m, 1366 s, 1344 s, 1257 w, 1232 w, 1180 s, 1105 s, 1049 m, 971 s, 985 s, 960 s, 759 s. MS, m/z (relative intensity, %): 275 (M⁺, 2.4), 93 (100), 92 (17), 65 (13). Exact Mass. Calcd for C₁₆H₂₁NO₃ : 275.1721. Found: 275.1519.

2-Propyl-hex-2-enoic acid benzyl ester (6a)

Rh₄(CO)₁₂ (30.0 mg, 0.04 mmol), Benzyl alcohol (216 mg, 2.00 mmol), 4-octyne (220 mg, 2.00 mmol), Pyridine (158 mg, 2 mmol) and toluene (1 mL) were placed in a 50-mL stainless steel autoclave under N₂. The system was flushed with 10 atm of carbon monoxide three times. Finally, it was pressurized to 3 atm. The autoclave was heated in an oil bath at 100 °C for 20 h, followed by cooling for rt. Then CO was released. The contents were transferred to a round-bottom flask with EtOAc and the volatiles were removed in vacuo. The residue was subjected to flash column chromatography on silica-gel (eluent; Hexane/EtOAc = 8/1→6/1) to give 2-Propyl-hex-2-enoic acid benzyl ester (6a) (379 mg, 77 % yield based on benzyl alcohol) as a colorless oil.

colorless oil. R_f 0.51 (hexane/EtOAc = 5/1). ¹H NMR (CDCl₃, TMS) δ : 7.36 – 7.31(m, 5H), 6.81(t, J =7.6 Hz, 1H), 5.18 (s, 2H), 2.24 (t, J =7.6 Hz, 2H), 2.16 (q, J =7.6 Hz, 2H), 1.14 (m, J =7.6 Hz, 4H), 0.92 (c, 6H). ¹³C NMR (CDCl₃) δ : 167.92, 143.26, 136.49, 132.20, 127.49, 127.96, 127.91, 66.08, 30.65, 28.74, 22.55, 22.09, 14.01, 13.98. IR (neat): 2960 s, 2871 s, 1706 s, 1643 s, 1587 w, 1498 w, 1457 s, 1378 s, 1278 s, 1214 s, 1141 s, 1106 s, 1064 m, 1022 m, 916 w,

883 w, 848 w, 813 w, 752 s, 698 s, 601 w. MS, m/z (relative intensity, %): 246 (M⁺, 1.2), 139 (25), 92 (10), 91 (100), 69 (35), 67 (10), 65 (18), 55 (25). Anal. Calcd for C₁₈H₂₂O₂ : C, 78.01; H, 9.00. Found: C, 77.92; H, 8.85.

dl-2,3-diethyl-succinic acid dipyridin-2-yl-methyl ester (3b)

colorless oil. R_f 0.10 (hexane/EtOAc = 1/1). ¹H NMR (CDCl₃) δ : 8.54 (d, J =5.1 Hz, 2H), 7.66 (td, J =7.6, 1.6 Hz, 2H), 7.35 (d, J =7.8 Hz, 2H), 7.23 (dd, J =6.8, 4.9 Hz, 2H), 5.24 (d, J =13.5 Hz, 2H), 5.17 (d, J =13.5 Hz, 2H), 2.92 – 2.83 (c, 2H), 1.73 – 1.49 (m, 4H), 0.85 (t, J =7.3 Hz, 6H). ¹³C NMR (CDCl₃) δ : 174.27, 155.80, 149.12,

136.83, 122.76, 121.68, 66.74, 46.88, 22.01, 10.95. IR (neat): 2967 m, 2938 w, 2879 w, 1735 s, 1594 m, 1573 w, 1438 w, 1386 w, 1348 w, 1268 m, 1234 w, 1160 s, 995 m, 761 s. MS, m/z (relative intensity, %): 356 (M⁺, 0), 248 (11), 110 (14), 109 (12), 108 (15), 93 (17), 92 (100), 65 (24), 55 (19). Exact Mass. Calcd for C₂₀H₂₄N₂O₄ : 356.1736. Found: 356.1725.

meso-2,3-diethyl-succinic acid dipyridin-2-yl-methyl ester (3b)

colorless oil. R_f 0.17 (hexane/EtOAc = 1/1). ¹H NMR (CDCl₃) δ : 8.58 (d, J =4.9 Hz, 2H), 7.73 (td, J =7.8, 1.9 Hz, 2H), 7.35 (d, J =8.1 Hz, 2H), 7.23 (dd, J =7.3, 5.7 Hz, 2H), 5.24 (s, 4H), 2.75 (c, 2H), 1.73 – 1.49 (m, 4H), 0.87 (t, 7.6 Hz, 6H). ¹³C NMR (CDCl₃) δ : 173.74, 155.60, 149.37, 136.79, 122.90, 121.85, 66.84, 49.59,

24.07, 11.73. IR (neat): 2965 m, 2937 w, 2877 w, 1733 s, 1594 m, 1573 w, 1438 w, 1382 w, 1340 w,

1274 m, 1230 w, 1151 s, 761 s, 728 m, 601 m. MS, m/z (relative intensity, %): 356 (M^+ , 0), 248 (11), 110 (14), 109 (12), 108 (15), 93 (17), 92 (100), 65 (24), 55 (19). Exact Mass. Calcd for $C_{22}H_{28}N_2O_4$: C,68.73; H,7.34; N,7.29. Found: C,68.70; H,7.30; N,7.25.

***threo*-2-butyl-3-ethyl-succinic acid dipyridin-2-ylmethyl ester (3c)**

colorless oil. R_f 0.11 (hexane/EtOAc = 1/1). 1H NMR ($CDCl_3$) δ : 8.55 (d, $J=4.6$ Hz, 2H), 7.65 (td, $J=7.6$, 1.4 Hz, 2H), 7.34 (d, $J=7.8$ Hz, 2H), 7.21 (dd, $J=6.8$, 5.1 Hz, 2H), 5.24 (d, $J=15.7$ Hz, 2H), 5.17 (d, $J=13.7$ Hz, 2H), 2.91 – 2.81 (c, 2H), 1.80 – 1.65 (c, 4H), 1.28 – 1.26 (c, 4H), 0.93 – 0.83 (c, 6H).

^{13}C NMR ($CDCl_3$) δ : 174.42, 174.21, 155.84, 149.23, 136.69, 136.67, 122.72, 121.66, 66.81, 66.80, 47.44, 45.75, 28.76, 28.73, 22.62, 22.05, 13.86, 11.03. IR (neat): 2983 m, 2942 m, 1671 s, 1592 m, 1455 m, 1440 m, 1378 w, 1288 s, 1268 s, 1224 m, 1195 m, 1174 m, 1097 s, 1033 s, 985 w, 900 w, 856 w, 763 s, 701 m, 605 w. MS, m/z (relative intensity, %): 384 (M^+ , 1), 276 (10), 206 (10), 178 (11), 164 (16), 110 (23), 109 (21), 108 (17), 93 (21), 92 (100), 69 (21), 65 (28), 55 (24). Anal. Calcd for $C_{22}H_{28}N_2O_4$: C,68.73; H,7.34; N,7.29. Found: C,68.70; H,7.30; N,7.25.

***erythro*-2-butyl-3-ethyl-succinic acid dipyridin-2-ylmethyl ester (3c)**

colorless oil. R_f 0.26 (hexane/EtOAc = 1/1). 1H NMR ($CDCl_3$) δ : 8.58 (d, $J=4.3$ Hz, 2H), 7.69 (t, $J=6.8$ Hz, 2H), 7.35 (d, $J=8.1$ Hz, 2H), 7.23 (dd, $J=7.3$, 5.4 Hz, 2H), 5.25 (d, $J=1.6$ Hz, 4H), 2.82 – 2.79 (m, 2H), 1.74 – 1.46 (c, 4H), 1.23 – 1.20 (c, 4H), 0.91 – 0.78 (c, 6H). ^{13}C NMR ($CDCl_3$) δ : 173.74, 173.57, 155.53, 149.30, 136.59, 122.80, 121.82, 121.78, 68.89, 49.92, 48.15, 30.73, 29.51, 24.18, 22.51, 13.92, 11.85. IR (neat): 2960 m, 2933 m, 2358 w, 1735 s,

1594 m, 1473 m, 1438 m, 1340 w, 1272 m, 1151 s, 1099 w, 987 w, 935 w, 900 w, 761 s, 636 m, 611 m. MS, m/z (relative intensity, %): 384 (M^+ , 1), 276 (10), 206 (10), 178 (11), 164 (16), 110 (23), 109 (21), 108 (17), 93 (21), 92 (100), 69 (21), 65 (28), 55 (24). Anal. Calcd for $C_{22}H_{28}N_2O_4$: C,68.73; H,7.34; N,7.29. Found: C,68.70; H,7.30; N,7.25.

2-Methyl-3-pentyl-succinic acid dipyridin-2-ylmethyl ester (3d)

Supectral data was obtain from a mixture of *threo* and *erythro*: colorless oil. R_f 0.11 (hexane/EtOAc = 1/1). 1H NMR ($CDCl_3$) δ : 8.55 (d, $J=5.1$ Hz, 2H), 7.71 – 7.62 (c, 2H), 7.33 (c, 2H), 7.20 (dd, $J=11.8$, 6.5 Hz, 2H), 5.26 – 5.15 (c, 4H), 3.02 – 2.75 (m, 2), 1.71 – 1.40 (c, 2H), 1.30 – 1.15 (c, 9H), 0.90 – 0.75 (c, 3H). ^{13}C NMR ($CDCl_3$) δ : 174.71, 174.17, 173.52, 155.71, 155.66, 155.53, 149.29, 149.16, 149.14, 136.58, 136.55, 122.77, 122.65, 121.78, 121.64, 121.59, 121.46, 66.95, 66.90, 66.83, 48.54, 47.33, 42.10, 40.76, 31.78, 31.57, 30.70, 28.68, 27.13, 26.47, 22.50, 22.47, 15.23, 14.41, 14.07. IR (KBr): 2954 m, 2931 m, 2861 w, 1737 s, 1594 m, 1573 w, 1475 m, 1438 m, 1382 w, 1338 s, 1267 m, 1160 s, 1105 w, 1051 w, 997 w, 829 w, 761 m, 601 w. MS, m/z (relative intensity, %): 384 (M^+ , 1), 276 (10), 206 (10), 178 (11), 164 (16), 110 (23), 109 (21), 108 (17), 93 (21), 92 (100), 69 (21), 65 (28), 55 (24). Exact Mass. Calcd for $C_{22}H_{28}N_2O_4$: 384.2049. Found: 384.2044.

Cyclododecane-1,2-dicarboxylic acid dipyridine-2-ylmethyl ester (3e)

Spectral data were obtained from a mixture of *cis* and *trans* isomers: colorless oil. *R_f* 0.085 (hexane/EtOAc = 1/1). ¹H NMR (CDCl₃) δ: 8.53 (d, *J*=4.3 Hz, 2H), 7.70 – 7.55 (c, 2H), 7.35 – 7.28 (c, 2H), 7.20 – 7.15 (c, 2H), 5.27 – 5.13 (c, 4H), 3.10 – 2.90 (c, 2H), 1.90 – 1.60 (c, 4H), 1.60 – 1.25 (c, 16H). ¹³C NMR (CDCl₃) δ: 175.02, 173.75, 155.95, 155.78, 149.17, 149.07, 136.74, 122.73, 122.66, 121.74, 121.58, 67.00, 66.74, 42.98, 25.76, 24.21, 23.97, 23.92, 22.49, 21.96. IR (neat): 2935 s, 2861 m, 1737 s, 1592 m, 1573 w, 1473 m, 1440 m, 1384 w, 1340 m, 1263 m, 1236 m, 1174 s, 1155 s, 1097 w, 1049 w, 995 w, 759 m, 725 w, 628 w, 601 w. MS, *m/z* (relative intensity, %): 438 (M⁺, 2.0), 330 (11), 110 (19), 109 (12), 108 (12), 93 (23), 92 (100), 67 (10), 65 (21), 55 (23). Anal. Calcd for C₂₂H₂₈N₂O₄: C, 71.21; H, 7.81; N, 6.39. Found: C, 70.92; H, 7.58; N, 6.14.

***threo*-2-Butyl-3-phenyl-succinic acid dipyridine-2-ylmethyl ester (3f)**

colorless oil *R_f* 0.07 (hexane/EtOAc = 1/1). ¹H NMR (CDCl₃) δ: 8.53 (dt, *J*=4.9, 0.8 Hz, 1H), 8.43 (dt, *J*=4.9, 0.8 Hz, 1H), 7.63 (td, *J*=7.6, 1.4 Hz, 1H), 7.47 (td, *J*=7.6, 1.4 Hz, 1H), 7.36 (d, *J*=7.8 Hz, 1H), 7.34 – 7.27 (m, 5H), 7.18 – 7.14 (c, 1H), 7.07 (c, 1H), 6.93 (d, *J*=7.8 Hz, 1H), 5.29 (d, *J*=13.5 Hz, 1H), 5.21 (d, *J*=13.5 Hz, 1H), 5.16 (d, *J*=13.2 Hz, 1H), 5.11 (d, *J*=13.2 Hz, 1H), 4.03 (d, *J*=11.9 Hz, 1H), 3.34 (m, 1H), 1.41 – 1.04 (c, 6H), 0.75 – 0.71 (t, *J*=6.8 Hz, 3H). ¹³C NMR (CDCl₃) δ: 174.54, 172.63, 155.76, 155.73, 149.01, 148.88, 136.60, 136.48, 128.85, 128.56, 128.42, 127.82, 122.64, 122.37, 121.55, 120.60, 66.80, 66.74, 52.73, 47.29, 29.03, 28.03, 22.24, 13.61. IR (neat): 2956 m, 1735 s, 1594 m, 1573 w, 1477 m, 1454 w, 1438 m, 1274 w, 1232 w, 1199 w, 1157 s, 995 w, 759 m, 701 m, 605 m. MS, *m/z* (relative intensity, %): 432 (M⁺, 2.5), 324 (11), 117 (18), 110 (10), 93 (43), 92 (100), 91 (20), 65 (42). Exact Mass. Calcd for C₂₆H₂₈N₂O₄: 432.1736. Found: 432.2051.

***erythro*-2-Butyl-3-phenyl-succinic acid dipyridine-2-ylmethyl ester (3f)**

colorless oil. *R_f* 0.14 (hexane/EtOAc = 1/1). ¹H NMR (CDCl₃) δ: 8.53 (d, *J*=4.3 Hz, 1H), 8.48 (d, *J*=4.6 Hz, 1H), 7.75 (t, *J*=7.8 Hz, 1H), 7.49 (t, *J*=7.8 Hz, 1H), 7.38 – 7.11 (c, 7H), 7.06 (d, *J*=8.1 Hz, 1H), 6.65 (d, *J*=7.6 Hz, 1H), 5.29 – 5.18 (c, 2H), 5.03 (d, *J*=13.5 Hz, 1H), 4.92 (d, *J*=13.5 Hz, 1H), 3.97 (d, *J*=11.3 Hz, 1H), 3.39 (td, *J*=11.1, 3.8 Hz, 1H), 1.83 – 1.63 (c, 2H), 1.40 – 1.15 (c, 4H), 0.84 (t, *J*=6.8 Hz, 3H). ¹³C NMR (CDCl₃) δ: 172.92, 171.55, 155.29, 155.27, 149.07, 148.83, 136.43, 136.37, 136.00, 128.46, 128.37, 127.67, 122.59, 122.38, 121.11, 121.08, 67.02, 66.49, 54.23, 49.50, 31.56, 29.38, 22.39, 13.82. IR (neat): 2956 m, 1735 s, 1592 m, 1573 w, 1475 w, 1455 m, 1376 w, 1278 s, 1228 w, 1149 s, 1049 w, 1008 m, 757 m, 700 m, 601 m. MS, *m/z* (relative intensity, %): 432 (M⁺, 2.5), 324 (11), 117 (18), 110 (10), 93 (43), 92 (100), 91 (20), 65 (42). Exact Mass. Calcd for C₂₆H₂₈N₂O₄: 432.1736. Found: 432.2042.

***threo*-2-Butyl-3-(4-methoxyphenyl)-succinic acid dipyridine-2-ylmethyl ester (3g)**

orange oil. *R_f* 0.07 (hexane/EtOAc = 1/1). ¹H NMR (CDCl₃) δ: 8.56 (d, *J*=4.9 Hz, 1H), 8.48 (d, *J*=4.9 Hz, 1H), 7.66 (td, *J*=7.6, 1.6 Hz, 1H), 7.53 (td, *J*=7.6, 1.6 Hz, 1H), 7.39 (d, *J*=8.1 Hz, 1H), 7.27 – 7.17 (c, 3H), 7.12 (dd, *J*=7.0, 4.9 Hz, 1H), 7.01 (d, *J*=7.8 Hz, 1H), 6.88 (d, *J*=8.6 Hz, 1H), 5.31 (d, *J*=13.5 Hz, 1H), 5.23 (d, *J*=13.5 Hz, 1H), 5.20 (d, *J*=14.0 Hz, 1H), 5.12 (d, *J*=14.0 Hz, 1H), 3.99 (d, *J*=11.6 Hz, 1H), 3.81 (s, 3H), 3.33 – 3.24 (m, 1H), 1.45 – 1.15 (c, 6H), 0.75 (t, *J*=6.2 Hz, 3H). ¹³C NMR (CDCl₃) δ: 174.70, 172.99, 159.19, 155.92, 155.86, 149.18, 148.98, 136.69, 136.58, 129.52, 127.94, 122.71, 122.45, 121.63, 120.74, 114.27, 66.85, 66.79, 55.23, 51.95, 47.48, 29.12, 28.15, 22.37, 13.72. IR (neat): 2956 m, 2933 m, 2865 w, 1733 s, 1610 m, 1592 m, 1575 w, 1511 s, 1473 m, 1438 m, 1376 w, 1330 m, 1251 s, 1178 s, 1149 s, 1103 w, 1033 m, 998 w, 831 m, 794 w, 759 m, 601 w. MS, *m/z* (relative intensity, %): 462 (*M*⁺, 4.4), 261 (26), 217 (22), 216 (30), 187 (15), 148 (10), 147 (24), 121 (34), 108 (10), 93 (94), 92 (100), 91 (13), 65 (40), 55 (10). Exact Mass. Calcd for C₂₇H₃₀N₂O₅ : 462.2155. Found: 462.2159.

erythro-2-Butyl-3-(4-methoxyphenyl)-succinic acid dipyridin-2-ylmethyl ester (3g)

orange oil. *R_f* 0.14 (hexane/EtOAc = 1/1). ¹H NMR (CDCl₃) δ: 8.54 (d, *J*=4.6 Hz, 1H), 8.48 (d, *J*=4.6 Hz, 1H), 7.60 (td, *J*=7.6, 1.4 Hz, 1H), 7.49 (td, *J*=7.6, 1.4 Hz, 1H), 7.28 (d, *J*=8.6 Hz, 2H), 7.21 – 7.12 (m, 2H), 7.09 (d, *J*=8.1 Hz, 1H), 6.78 (d, *J*=8.6 Hz, 2H), 6.66 (d, *J*=8.1 Hz, 1H), 5.26 (d, *J*=13.5 Hz, 1H), 5.19 (d, *J*=13.5 Hz, 1H), 5.08 (d, *J*=13.2 Hz, 1H), 4.91 (d, *J*=13.2 Hz, 1H), 3.90 (d, *J*=11.6 Hz, 1H), 3.77 (s, 3H), 3.33 (td, *J*=10.8, 4.1 Hz, 1H), 1.80 – 1.64 (c, 2H), 1.32 – 1.27 (c, 4H), 0.85 (t, *J*=6.9 Hz, 3H). ¹³C NMR (CDCl₃) δ: 173.14, 171.91, 159.01, 155.44, 149.19, 148.93, 136.54, 136.35, 129.54, 128.15, 122.68, 122.47, 121.27, 121.20, 113.91, 67.08, 66.54, 55.20, 53.51, 49.79, 31.64, 29.52, 22.49, 13.91. IR (KBr): 2956 m, 2935 m, 2865 w, 1735 s, 1610 m, 1592 m, 1575 w, 1511 s, 1473 m, 1438 m, 1380 w, 1299 m, 1255 s, 1176 s, 1157 s, 1106 w, 1033 m, 995 w, 831 w, 796 w, 761 m, 601 w. MS, *m/z* (relative intensity, %): 462 (*M*⁺, 4.4), 261 (26), 217 (22), 216 (30), 187 (15), 148 (10), 147 (24), 121 (34), 108 (10), 93 (94), 92 (100), 91 (13), 65 (40), 55 (10). Exact Mass. Calcd for C₂₇H₃₀N₂O₅ : 462.2155. Found: 462.2151.

threo-2-Butyl-3-(4-trifluoromethylphenyl)-succinic acid dipyridin-2-ylmethyl ester (3h)

orange oil. *R_f* 0.07 (hexane/EtOAc = 1/1). ¹H NMR (CDCl₃) δ: 8.57 (d, *J*=4.6 Hz, 1H), 8.47 (d, *J*=4.6 Hz, 1H), 7.70 – 7.56 (c, 3H), 7.53 (td, *J*=7.6, 1.6 Hz, 1H), 7.46 – 7.37 (c, 3H), 7.24 – 7.19 (c, 1H), 7.16 – 7.15 (c, 1H), 6.99 (d, *J*=7.8 Hz, 1H), 5.32 (d, *J*=13.5 Hz, 1H), 5.24 (d, *J*=13.5 Hz, 1H), 5.19 (d, *J*=13.5 Hz, 1H), 5.13 (d, *J*=13.5 Hz, 1H), 4.13 (d, *J*=11.6 Hz, 1H), 3.38 – 3.30 (m, 1H), 1.45 – 1.10 (c, 6H), 0.76 – 0.74 (c, 3H). ¹³C NMR (CDCl₃) δ: 173.96, 171.83, 155.49, 155.28, 149.12, 148.98, 139.89, 136.44, 130.09 (q, *J*=32.3 Hz), 128.84, 125.78 (q, *J*=3.3 Hz), 123.80 (q, *J*=269.7 Hz), 122.70, 122.56, 121.61, 120.83, 67.18, 67.06, 52.65, 47.32, 29.15, 28.15, 22.40, 13.71. IR (neat): 2958 w, 2937 w, 1737 s, 1617 w, 1594 w, 1573 w, 1440 w, 1376 w, 1326 s, 1234 m, 1164 s, 1128 s, 1068 m, 1018 w, 997 w, 894 w, 846 w, 759 m, 727 w, 603 w. MS, *m/z* (relative intensity, %): 500 (*M*⁺, 2), 206 (12), 110 (19), 109 (12), 108 (12), 93 (25), 92 (100), 65 (29). Exact Mass. Calcd for C₂₇H₂₇F₃N₂O₄ : 500.1923. Found: 500.1919.

erythro-2-Butyl-3-(4-trifluoromethylphenyl)-succinic acid dipyridin-2-ylmethyl ester (3h)

pale yellow solid. mp 47 °C. R_f 0.19 (hexane/EtOAc = 1/1). ^1H NMR (CDCl_3) δ : 8.53 (d, $J=4.6$ Hz, 1H), 8.48 (d, $J=4.6$ Hz, 1H), 7.59 (td, $J=7.6$, 1.6 Hz, 1H), 7.51 - 7.47 (m, 5H), 7.22 - 7.13 (c, 2H), 7.03 (d, $J=7.8$ Hz, 1H), 6.71 (d, $J=7.8$ Hz, 1H), 5.25 (d, $J=13.5$ Hz, 1H), 5.19 (d, $J=13.5$ Hz, 1H), 5.05 (d, $J=13.2$ Hz, 1H), 4.92 (d, $J=13.2$ Hz, 1H), 4.01 (d, $J=11.6$ Hz, 1H), 3.36 (td, $J=11.3$, 4.1 Hz, 1H), 1.77 - 1.67 (c, 2H), 1.33 - 1.28 (m, 4H), 0.85 (t, $J=6.8$ Hz, 3H). ^{13}C NMR (CDCl_3) δ : 172.63, 170.95, 154.92, 149.25, 149.06, 140.01, 136.49, 136.31, 129.84 (q, $J=32.2$ Hz), 128.83, 125.27 (q, $J=3.6$ Hz), 123.78 (q, $J=269.9$ Hz), 122.81, 122.69, 121.52, 121.35, 67.37, 66.77, 53.96, 49.31, 31.53, 29.28, 22.39, 13.82. IR (KBr): 2958 w, 2937 w, 1737 s, 1617 w, 1594 w, 1573 w, 1440 w, 1376 w, 1326 s, 1234 m, 1164 s, 1128 s, 1068 m, 1018 w, 997 w, 894 w, 846 w, 759 m, 727 w, 603 w. MS, m/z (relative intensity, %): 500 (M^+ , 2), 206 (12), 110 (19), 109 (12), 108 (12), 93 (25), 92 (100), 65 (29). Exact Mass. Calcd for $\text{C}_{27}\text{H}_{27}\text{F}_3\text{N}_2\text{O}_4$: 500.1923. Found: 500.1929.

dl-2,3-diphenyl-succinic acid dipyridin-2-ylmethyl ester (3i)

orange oil. R_f 0.085 (hexane/EtOAc = 1/1). ^1H NMR (CDCl_3 , TMS) δ : 8.49 (d, $J=4.1$ Hz, 2H), 7.54 (t, $J=7.6$ Hz, 2H), 7.14 - 7.04 (c, 14H), 5.30 (d, $J=14.2$ Hz, 2H), 5.21 (d, $J=14.2$ Hz, 2H), 4.45 (s, 2H). ^{13}C NMR (CDCl_3) δ : 172.50, 155.62, 148.84, 136.57, 135.04, 128.45, 128.34, 127.51, 122.47, 120.75, 67.05, 54.61. IR (neat): 1731 s, 1592 w, 1571 w, 1494 w, 1475 w, 1455 w, 1436 m, 1371 w, 1294 s, 1226 w, 1141 s, 1047 w, 991 m, 775 w, 757 w, 701 m, 609 w. MS, m/z (relative intensity, %): 452 (M^+ , 5.4), 344 (10), 226 (11), 180 (12), 179 (18), 178 (11), 108 (13), 93 (57), 92 (100), 91 (15), 65 (38). Exact Mass. Calcd for $\text{C}_{28}\text{H}_{24}\text{N}_2\text{O}_4$: 452.1736. Found: 452.1731.

meso-2,3-diphenyl-succinic acid dipyridin-2-ylmethyl ester (3i)

white solid. mp 147 °C. R_f 0.13 (hexane/EtOAc = 1/1). ^1H NMR (CDCl_3 , TMS) δ : 8.44 (d, $J=4.6$ Hz, 2H), 7.55 (t, $J=4.1$ Hz, 4H), 7.43 - 7.32 (c, 8H), 7.10 (t, $J=5.4$ Hz, 2H), 6.46 (d, $J=7.6$ Hz, 2H), 5.09 (d, $J=14.0$ Hz, 2H), 4.92 (d, $J=14.0$ Hz, 2H), 4.58 (s, 2H). ^{13}C NMR (CDCl_3) δ : 170.81, 155.31, 148.91, 136.44, 135.81, 128.74, 128.59, 128.02, 122.38, 120.60, 66.97, 55.04. IR (KBr): 1731 s, 1592 w, 1571 w, 1494 w, 1475 w, 1455 w, 1436 m, 1371 w, 1294 s, 1226 w, 1141 s, 1047 w, 991 m, 775 w, 757 w, 701 m, 609 w. MS, m/z (relative intensity, %): 452 (M^+ , 5.4), 344 (10), 226 (11), 180 (12), 179 (18), 178 (11), 108 (13), 93 (57), 92 (100), 91 (15), 65 (38). Exact Mass. Calcd for $\text{C}_{28}\text{H}_{24}\text{N}_2\text{O}_4$: 452.1736. Found: 452.1732.

2,3-dipropyl-succinic acid benzyl ester pyridin-2-yl-methyl ester (7a)

$\text{Rh}_4(\text{CO})_{12}$ (30.0 mg, 0.04 mmol), pyridin-2-yl-methanol (109 mg, 1.00 mmol), Benzyl alcohol (108 mg, 1.00 mmol), 4-octyne (220 mg, 2.00 mmol) and toluene (1 mL) were placed in a 50-mL stainless steel autoclave under N_2 . The system was flushed with 10 atm of carbon monoxide three times. Finally, it was pressurized to 3 atm. The autoclave was heated in an oil bath at 100 °C for 20 h, followed by cooling for rt. Then CO was released. The contents were transferred to a round-bottom flask with EtOAc and the volatiles were removed in vacuo. The residue was subjected to flash column

chromatography on silica-gel (eluent; hexane/EtOAc = 8/1→0/1) to give 2,3-dipropyl-succinic acid benzyl ester pyridin-2-yl-methyl ester (**7a**) (201 mg, 52 % yield based on **2**) as a colorless oil and 2,3-dipropyl-succinic acid dipyridin-2-yl-methyl ester (**3a**) (29 mg, 8% yield based on pyridin-2-yl-methanol) as a pale yellow oil.

colorless oil. *R_f* 0.17 (hexane/EtOAc = 1/1). ¹H NMR (CDCl₃, TMS)

δ: 8.56 (d, *J*=4.6 Hz, 1H), 7.65 (t, *J*=8.6 Hz, 1H), 7.34 – 7.31 (c, 6H), 7.22 – 7.18 (c, 1H), 5.23 – 5.07 (c, 4H), 2.85 – 2.76 (c, 2H), 1.62 – 1.59 (c, 4H), 1.33 – 1.24 (c, 4H), 0.90 – 0.81 (c, 6H). ¹³C NMR (CDCl₃) δ: 174.28, 174.05, 155.78, 149.30, 149.12, 136.54, 135.75, 128.41, 128.36, 128.25, 128.14, 128.04, 122.78, 122.62, 121.60,

66.82, 66.32, 48.32, 46.46, 46.43, 31.30, 20.63, 20.21, 20.19, 14.12, 13.91. IR (neat): 2960 s, 2873 m, 1735 s, 1592 w, 1573 w, 1498 w, 1459 m, 1382 w, 1346 w, 1265 m 1238 w, 1213 m, 1160 s, 1105 w, 1052 w, 754 m, 700 m, 601 w. MS, *m/z* (relative intensity, %): 383 (*M*⁺, 1.6), 206 (19), 164 (17), 110 (23), 109 (76), 108 (35), 93 (29), 92 (56), 91 (100), 69 (17), 65 (31), 55 (29). Anal. Calcd for C₂₂H₂₈N₂O₄ : C,68.73; H,7.34; N,7.29. Found: C,71.85; H,7.57; N,3.64..

MENUF	NON
OBNUC	1H
OFIR	270.05 MHz
OBSET	112.00 KHz
OBFIN	5800.0 Hz
PW1	5.3 us
DEADT	108.5 us
PREDL	0.2000 ms
WT	1.0 sec
POINT	16384
SPO	16384
TIMES	16
DUMMY	1
FREOU	5405.4 Hz
FLT	2700 Hz
DELAY	74.1 us
ACQTM	3.031 sec
PD	3.967 sec
ADBIT	16.00
RGAIN	19
BF	0.12 Hz
T1	0.00
T2	0.00
T3	90.00
T4	100.00
EXMOD	NON
EXPCM	NON:Single coupled PW1_ACQTM.
IRNUC	1H
IFR	270.05 MHz
IRSET	112.00 KHz
IRFIN	5800.0 Hz
IRRPW	50 us
IRATN	511
DFILE	THSAT
SF	64.60 KHz
LKSET	59.0 Hz
LKFIN	200
LKLEV	23
LGAIN	117
LKPHS	594
LKSIG	17 Hz
CSPED	
FILDC

MENU	NON
OBNUC	1H
OPR	270.05 MHz
OBSET	112.00 KHz
OBFIN	5800.0 Hz
PW1	5.3 us
DEADT	108.5 us
PREDL	0.2000 ms
WT	1.0 sec
POINT	16384
SPO	16384
TIMES	8
DUMMY	1
FREQ	5402.4 Hz
FLT	2700 Hz
DELAY	74.1 us
ACQTM	3.033 sec
PD	3.967 sec
ADBIT	16.00
RGAIN	19
BF	0.12 Hz
T1	0.00
T2	0.00
T3	90.00
T4	100.00
EXMOD	NON
EXPCM	NON;Single,coupled,PW1,ACQTM
IRNUC	1H
IFR	270.05 MHz
IRSET	112.00 KHz
IRFIN	5800.0 Hz
IRAPW	50 us
IRATN	511
DFILE	C:\WINNMR98\COMMON_DEFAU
SF	THSAT
LKSET	64.60 KHz
LKFN	59.0 Hz
LKLEV	200
LGAIN	20
LKPHS	117
LKSIG	557
CSPED	17 Hz
FILDC	
FILDF

MENU	NON
OBNUC	1H
OFFR	270.05 MHz
OBSET	112.00 KHz
OBFIN	5800.0 Hz
PW1	5.3 us
DEADT	108.5 us
PREDL	0.2000 ms
IWT	1.0 sec
POINT	16384
SPO	16384
TIMES	8
DUMMY	1
FREOU	5405.4 Hz
FLT	2700 Hz
DELAY	74.1 us
ACQTM	3.031 sec
PD	3.967 sec
ADBIT	16.00
RGAIN	19
BF	0.12 Hz
T1	0.00
T2	0.00
T3	90.00
T4	100.00
EXMOD	NON
EXPCM	NON:Single coupled:PW1_ACQTM
IRNUC	1H
IFR	270.05 MHz
IRSET	112.00 KHz
IRFIN	5800.0 Hz
IRRPW	50 us
IRATN	511
DFILE	C:\My Documents\S. Inoue\Spectra\THSAT
SF	64.60 KHz
LKSET	59.0 Hz
LKFIN	200
LKLEV	23
LGAIN	117
LKPHS	583
LKSIG	16 Hz
CSPED	
FILDC	
FILDF

MENUF	NON
OBNUC	¹ H
OFR	270.05 MHz
OBSET	112.00 KHz
OBFIN	5800.0 Hz
PW1	5.3 us
DEADT	108.5 us
PREDL	0.2000 ms
INT	1.0 sec
POINT	16384
SPO	16384
TIMES	8
DUMMY	1
FREOU	5402.4 Hz
FLT	2700 Hz
DELAY	74.1 us
ACQTM	3.033 sec
PD	3.967 sec
ADBIT	16.00
RGAIN	18
BF	0.12 Hz
T1	0.00
T2	0.00
T3	90.00
T4	100.00
EXMOD	NON
EXPCM	NON:Single-coupled PW1_ACQTM
IRNUC	¹ H
IFR	270.05 MHz
IRSET	112.00 KHz
IFEN	5800.0 Hz
IRRPW	50 us
IRATN	511
DFILE	THSAT
SF	64.60 KHz
LKSET	59.0 Hz
LKFIN	200
LKLEV	23
LGAIN	117
LKPHS	636
LKSIG	14 Hz
CPRED	
FILDC	
FILDF

MENUF 1H NON
OBNUC 270.05 MHz
OFR 112.00 KHz
OBSET 5800.0 Hz
OBFIN 5.3 us
PW1 108.5 us
DEADT 0.2000 ms
PREDL 1.0 sec
WT 16394
POINT 16394
SPO 16394
TIMES 16
DUMMY 1
FREQU 5405.4 Hz
FLT 2700 Hz
DELAY 74.1 us
ACQTM 3.031 sec
PD 3.967 sec
ADBIT 16.00
RGAIN 12
BF 0.12 Hz
T1 0.00
T2 0.00
T3 90.00
T4 100.00
EXMOD NON
EXPCM NON:Single coupled PW1_ACQTM
IRNUC 1H
IFR 270.05 MHz
IRSET 112.00 KHz
IRFIN 5800.0 Hz
IRRPW 50 us
IRATN 511
DFILE C:\My Documents\15 Inoue\15 Spectra
SF TH5AT
LKSET 64.60 KHz
LKFIN 59.0 Hz
LKLEV 200
LGAIN 23
LKPHS 117
LKSIG 505
CSPED 15 Hz
FLDC
FLDF

MENUF NON
OBNUC 1H
OFR 270.05 MHz
OBSET 112.00 KHz
OBFIN 5800.0 Hz
PW1 5.3 us
DEADT 108.5 us
PREDL 0.2000 ms
WIT 1.0 sec
POINT 16384
SPO 16394
TIMES 8
DUMMY 1
FREOU 5402.4 Hz
FLT 2700 Hz
DELAY 74.1 us
ACQTM 3.033 sec
PD 3.967 sec
ADBIT 16.00
RGAIN 12
BF 1.20 Hz
T1 0.00
T2 0.00
T3 90.00
T4 100.00
EXMOD NON
EXPCM NON, Single coupled PW1, ACQTM, 1H
IRNUC 270.05 MHz
IFR 112.00 KHz
IRSET 5800.0 Hz
IRFIN 50 us
IRAPW 511
IRATN 511
DFILE
SF THSAT
LKSET 64.60 KHz
LKFN 59.0 Hz
LKLEV 200
LGAIN 23
LKPHS 117
LKSIG 483
CSPED 16 Hz
FLDC
FLDF

MENUF	NON
OBNUC	1H
OFR	270.05 MHz
OBSET	112.00 KHz
OBFIN	5800.0 Hz
PW1	5.3 us
DEADT	108.5 us
PREDL	0.2000 ms
WT	1.0 sec
POINT	16394
SPO	16394
TIMES	12
DUMMY	1
FREQU	5405.4 Hz
FLT	2700 Hz
DELAY	74.1 us
ACQTM	3.031 sec
PD	3.967 sec
ADBIT	16.00
RGAIN	17
BF	0.12 Hz
T1	0.00
T2	0.00
T3	90.00
T4	100.00
EXMOD	NON
EXPCM	NON:Single,coupled,PW1_ACQTM
IRNUC	1H
IFR	270.05 MHz
IRSET	112.00 KHz
IRPIN	5800.0 Hz
IRRPW	50 us
IRATN	511
DFILE	C:\My Documents\S:\Inoue\Spectra\
SF	THSAT
LKSET	64.60 KHz
LKFEN	59.0 Hz
LKLEV	180
LGAIN	20
LKPHS	117
LKSIG	596
CSPED	17 Hz
FLDC	
FILDF

MENUF	NON
OBNUC	1H
OFR	270.05 MHz
OBSET	112.00 KHz
OBFIN	5800.0 Hz
PW1	5.3 us
DEADT	108.5 us
PREDL	0.2000 ms
IWT	1.0 sec
POINT	16384
SPO	16384
TIMES	8
DUMMY	1
FREQU	5402.4 Hz
FLT	2700 Hz
DELAY	74.1 us
ACQTM	3.033 sec
PD	3.967 sec
ADBIT	16.00
RGAIN	18
BF	1.20 Hz
T1	0.00
T2	0.00
T3	90.00
T4	100.00
EXMOD	NON
EXPCM	NON:Single coupled-PW1_ACQTM
IRNUC	1H
IFR	270.05 MHz
IRSET	112.00 KHz
IRFIN	5800.0 Hz
IRRPW	50 us
IRALT	511
DFILE	C:\My Documents\1\Inoue\Spectra\TH5AT
SF	64.60 KHz
LKSET	59.0 Hz
LKFIN	200
LKLEV	20
LGAIN	117
LKPHS	639
LKSIG	17 Hz
CSPED	
FILDC	
FILDF

MENUP	NON
OBNUC	1H
OFR	270.05 MHz
OBSET	112.00 KHz
OBFIN	5800.0 Hz
PW1	5.3 us
DEADT	108.5 us
PREDL	0.2000 ms
WT	1.0 sec
POINT	16384
SPO	16384
TIMES	8
DUMMY	1
FREQU	5402.4 Hz
FLT	2700 Hz
DELAY	74.1 us
ACQTM	3.033 sec
PD	3.967 sec
ADBIT	16.00
RGAIN	14
BF	1.20 Hz
T1	0.00
T2	0.00
T3	90.00
T4	100.00
EXMOD	NON
EXPCM	NON
IRNUC	NON,Single,coupled,PW1,ACQTM
IFR	270.05 MHz
IRSET	112.00 KHz
IRPIN	5800.0 Hz
IRRPW	50 us
IRATN	511
DFILE	C:\My Documents\S:\Inoue\Spectra\
SF	THSAT
LKSET	64.60 KHz
LKFIN	59.0 Hz
LKLEV	200
LGAIN	23
LKPHS	117
LKSIG	476
CSPED	14 Hz
FILDC	
FILDF

MENUF	NON
OBNUC	1H
OFR	270.05 MHz
OBSET	112.00 KHz
OBFIN	5800.0 Hz
PW1	5.3 us
DEADT	108.5 us
PREDL	0.2000 ms
INT	1.0 sec
POINT	16394
SPO	16394
TIMES	8
DUMMY	1
FREOU	5402.4 Hz
FLT	2700 Hz
DELAY	74.1 us
ACQTM	3.033 sec
PD	3.967 sec
ADBIT	16.00
RGAIN	15
BF	1.20 Hz
T1	0.00
T2	0.00
T3	90.00
T4	100.00
EXMOD	NON
EXPCM	NON
IRNUC	NON:Single,coupled,PW1_ACQTM.
IFR	1H
IRSET	270.05 MHz
IRFIN	112.00 KHz
IRRPW	5800.0 Hz
IRATN	50 us
DFILE	511
SF	C:\My Documents\15 Inoue\15 Inoue Spectra\data\0901094-a\094-a-H.als
LKSET	64.60 KHz
LKFIN	59.0 Hz
LKLEV	200
LGAIN	22
LKPHS	117
LKSIG	600
CSPED	17 Hz
FILDC	
FILDF

MENUF	NON
OBNUC	1H
OFPR	270.05 MHz
OBSET	112.00 KHz
OBFIN	5800.0 Hz
PW1	5.3 us
DEADT	108.5 us
PREDL	0.2000 ms
WWT	1.0 sec
POINT	16394
SPO	16394
TIMES	8
DUMMY	1
FREOU	5402.4 Hz
FLT	2700 Hz
DELAY	74.1 us
ACQTM	3.033 sec
PD	3.967 sec
ADBIT	16.00
RGAIN	17
BF	1.20 Hz
T1	0.00
T2	0.00
T3	90.00
T4	100.00
EXMOD	NON
EXPCM	NON:Single,coupled,PW1_ACQTM
IRNUC	1H
IFR	270.05 MHz
IRSET	112.00 KHz
IRFIN	5800.0 Hz
IRRPW	50 us
IRATN	511
DFILE	
SF	THSAT
UKSET	64.60 KHz
LKFIN	59.0 Hz
LKLEV	200
LGAIN	23
LKPHS	117
LKSIG	482
CSPED	20 Hz
FILDC	
FILDF

MENUF	NON
OBNUC	1H
OFI	270.05 MHz
OBSET	112.00 KHz
OBFIN	5800.0 Hz
PW1	5.3 us
DEADT	108.5 us
PREDL	0.2000 ms
WT	1.0 sec
POINT	16384
SPO	16384
TIMES	8
DUMMY	1
FREOU	5402.4 Hz
FLT	2700 Hz
DELAY	74.1 us
ACQTM	3.033 sec
PD	3.967 sec
ADBIT	16.00
RGAIN	20
BF	1.20 Hz
T1	0.00
T2	0.00
T3	90.00
T4	100.00
EXMOD	NON
EXPCM	NON, Single coupled PW1, ACQTM
IRNUC	1H
IFR	270.05 MHz
IRSET	112.00 KHz
IRFIN	5800.0 Hz
IRRPW	50 us
IRATN	511
DFILE	C:\WINNMR98\COMMON_DEFAULTS
SF	THSAT
LKSET	64.60 KHz
LKEN	59.0 Hz
LKLEV	200
LGAIN	23
LKPHS	117
LKSIG	520
CSPED	16 Hz
FILDC	
FILDF	