Supporting Information

Chemoenzymatic Dynamic Kinetic Resolution of Primary Amines

Jens Paetzold and Jan. E. Bäckvall*

Department of Organic Chemistry, Arrhenius Laboratory
Stockholm University, SE-10691 Stockholm, Sweden

jeb@organ.su.se

General.

Unless stated otherwise all manipulations were performed under argon. Flash chromatography was carried out on 60 Å (35-70 µm) silica gel. ¹H and ¹³C NMR spectra were recorded at 400 or 300 MHz and at 100 or 75 MHz, respectively. Chemical shifts (δ) are reported in ppm, using the residual solvent peak in CDCl₃ (δ_H 7.26 and δ_C 77.00) as internal standard, and coupling constants (*J*) are given in Hz.

The racemic amines **1a**, **1c**, **1d**, **1f**, **1j** and **1k** were obtained from commercial sources, distilled and stored over molecular sieves before use. Amines **1b**, **1e**, **1g-i** were synthesized according to the literature¹ and stored over molecular sieves before use. *Candida antarctica* lipase B (CALB) was used as immobilized and thermostable Novozym 435. Isopropyl acetate was distilled over K₂CO₃ and stored over molecular sieves.

Complex **4** was purchased from Strem and used as received. Complex **3** was synthesized according to the literature (ref 15b of the paper). The complexes **5a-c** were obtained from the corresponding cyclopentadieneone and Ru₃CO₁₂ by refluxing in methanol.

The enantiomeric excess was determined by a Varian 3800 analytical gas chromatograph employing a CP-Chirasil-DEX CB, 25 X mm column using racemic compounds as references. The absolute stereochemical assignment was established by comparison of the sign of optical rotation with values found in literature to be the (*R*)-configured product.²⁻⁴

General procedure for racemization :

A flame dried 20 mL reaction vessel is charged with the ruthenium precursor (0.02 mmol Ru); the vessel is closed, evacuated and backfilled with argon three times. Toluene (2 mL), 1-(*S*)-Phenylethyl amine ((*S*)-**1a**, 0.50 mmol, 65 µl) and 30 µl pentadecane are added subsequently via syringe. The mixture is stirred at 100 °C for 24h. After the reaction, samples of 200 µl are taken, transferred to a GC-vial and diluted with toluene. The amine **1a** was transformed to the corresponding amide **2a** by treatment with two drops of acetic anhydride and two drops of triethylamine to the GC vial.

Chiral GC-analysis:

Injector 250 °C

Program: 125 °C / 4 min / 140 °C / 3 °C min⁻¹ / 200 °C / 40 °C min⁻¹, 5 min.

t_S= 10.39 min, t_R= 10.62 min

General Procedure for Dynamic Kinetic Resolution

Synthesis of N-((R)-1-phenylethyl)acetamide (2a): A flame dried 20 ml reaction vessel is charged with complex **5c** (26.5 mg, 0.02 mmol), Novozym 435 (20 mg) and Na₂CO₃ (0.20 mmol, 20 mg). The vessel is closed, evacuated and backfilled with argon three times. Toluene (8 mL), 1-phenyl ethylamine (**1a**, 0.50 mmol, 65 µl), isopropyl acetate (3.50 mmol, 400 µl) and 30 µl pentadecane are added subsequently via syringe under argon. The reaction mixture is stirred at 90 °C. After three days the reaction is cooled to room temperature, filtered and washed dichloromethane (3x 20 mL). The solvents are removed *in vacuo* and the crude product is purified by column chromatography (SiO₂, pentane/ethyl acetate, 1:4) yielding 75 mg of **2a** (90 %), as a white solid in 98% *ee*. The product was identified by means of ¹H, ¹³C NMR and optical rotation to be *N-((R)-1-phenylethyl)acetamide*. The spectroscopic data of the product are identical to those reported in the literature.²

Chiral GC-analysis:

Injector 250 °C

Program: 125 °C / 4 min / 140 °C / 3 °C min⁻¹ / 200 °C / 40 °C min⁻¹, 5 min.

$t_s = 10.39$ min, $t_R = 10.62$ min
[α_D] = +127.1 ($c = 1.0$, CH_2Cl_2).

Synthesis of N-((R)-1-m-tolyethyl)acetamide (2b):

Compound **2b** was synthesized from 1-*m*-tolylethyl amine (**1b**, 0.50 mmol, 73 μl) following the general procedure above using complex **5c** (26.5 mg, 0.02 mmol), CALB (30 mg) and Na_2CO_3 (0.30 mmol, 30 mg). After purification by column chromatography (SiO_2 , pentane / ethyl acetate, 1:4) **2b** was obtained as a white solid (61 mg, 0.35 mmol, 69 %, 97.5 *ee*). The spectroscopic data of the product were identical with those reported in the literature.²

Chiral GC-analysis:

Injector 250 °C

Program: 100 °C / 5 min / 155 °C / 3 °C min⁻¹, 5 min/ 200 °C / 20 °C min⁻¹, 5 min.

$t_s = 23.33$ min, $t_R = 23.86$ min

Synthesis of N-((R)-1-(4-fluorophenyl)ethyl)acetamide (2c):

Compound **2c** was synthesized from 1-(4-fluorophenyl)ethyl amine (**1c**, 0.50 mmol, 72 μl) following the general procedure above. After purification by column chromatography (SiO_2 , pentane / ethyl acetate, 1:4) **2c** was obtained as a white solid (75 mg, 0.42 mmol, 83 %, 98.5 *ee*). The spectroscopic data of the product were identical to those reported in the literature.²

Chiral GC-analysis:

Injector 250 °C

Program: 100 °C / 5 min / 155 °C / 3 °C min⁻¹, 5 min/ 200 °C / 20 °C min⁻¹, 5 min.

$t_s = 22.01$ min, $t_R = 22.57$ min

Synthesis of N-((R)-1-(4-bromophenyl)ethyl)acetamide (2d):

Compound **2d** was synthesized from 1-(4-bromophenyl)ethyl amine (**1d**, 0.50 mmol, 73 μl) following the general procedure above. After purification by column chromatography (SiO_2 , pentane / ethyl acetate, 1:4) **2d** was obtained as a white solid (95 mg, 0.39 mmol, 78 %, 99 *ee*). The spectroscopic data of the product were identical to those reported in the literature.²

Chiral GC-analysis:

Injector 250 °C

Program: 100 °C / 5 min / 155 °C / 3 °C min⁻¹, 5 min/ 200 °C / 20 °C min⁻¹, 5 min.

$t_s = 31.31$ min, $t_R = 31.48$ min

Synthesis of N-((R)-1-(4-methoxyphenyl)ethyl)acetamide (2e):

Compound **2e** was synthesized from 1-(4-methoxyphenyl)ethyl amine (**1e**, 0.50 mmol, 76 μl) following the general procedure above using complex **5c** (26.5 mg, 0.02 mmol), CALB (30 mg) and Na_2CO_3 (0.30 mmol, 30 mg). After purification by column chromatography (SiO_2 , pentane / ethyl acetate, 1:4) **2e** was obtained as a white solid (95 mg, 0.47 mmol, 95 %, 99 *ee*). The spectroscopic data of the product were identical to those reported in the literature.²

Chiral GC-analysis:

Injector 250 °C

Program: 100 °C / 5 min / 155 °C / 3 °C min⁻¹, 5 min/ 200 °C / 20 °C min⁻¹, 5 min.

$t_s = 30.31$ min, $t_R = 30.48$ min

Synthesis of N-((R)-1-(naphthalen-2-yl)ethyl)acetamide (2f):

Compound **2f** was synthesized from 1-(4-methoxyphenyl)ethyl amine (**1f**, 0.50 mmol, 82 μl) following the general procedure above using complex **5c** (26.5 mg, 0.02 mmol), CALB (30 mg) and Na_2CO_3 (0.30 mmol, 30 mg). The reaction is stirred for 5 days at 90 °C and after purification by column chromatography (SiO_2 , pentane / ethyl acetate, 1:4) **2f** was obtained as

a white solid (85 mg, 0.42 mmol, 80 %, >99.5 *ee*). The spectroscopic data of the product were identical to those reported in the literature.²

Chiral GC-analysis:

Injector 250 °C

Program: 150 °C / 5 min / 155 °C / 1 °C min⁻¹, 5 min/ 200 °C / 20 °C min⁻¹, 10 min.

*t*_S = 31.20 min, *t*_R = 31.43 min

[α _D] = +115.6 (c = 1.0, CH₂Cl₂).

Synthesis of N-((R)-1-phenylpropyl)acetamide (2g):

Compound **2g** was synthesized from 1-phenylpropyl amine (**1g**, 0.50 mmol, 76 μ l) following the procedure above using complex **5c** (26.5 mg, 0.02 mmol), CALB (30 mg) and Na₂CO₃ (0.30 mmol, 30 mg). The reaction is stirred for 5 days at 90 °C and after purification by column chromatography (SiO₂, pentane / ethyl acetate, 1:4) **2g** was obtained as a white solid (80 mg, 0.45 mmol, 90 %, 99 *ee*). The spectroscopic data of the product were identical to those reported in the literature.²

Chiral GC-analysis:

Injector 250 °C

Program: 100 °C / 5 min / 155 °C / 3 °C min⁻¹, 5 min/ 200 °C / 20 °C min⁻¹, 5 min.

*t*_S = 22.81 min, *t*_R = 23.22 min

[α _D] = +130.4 (c = 1.0, CH₂Cl₂).

Synthesis of N-((R)-2,3-dihydro-1H-inden-3-yl)acetamide (2h):

Compound **2h** was synthesized from 2,3-dihydro-1H-inden-1-amine (**1h**, 0.50 mmol, 68 μ l) following the procedure above using complex **5c** (26.5 mg, 0.02 mmol), CALB (30 mg) and Na₂CO₃ (0.30 mmol, 30 mg). After purification by column chromatography (SiO₂, pentane / ethyl acetate, 1:4) **2h** was obtained as a white solid (85 mg, 0.46 mmol, 92 %, 95 % *ee*). The spectroscopic data of the product were identical to those reported in the literature.³

Chiral GC-analysis:

Injector 250 °C

Program: 100 °C / 5 min / 155 °C / 3 °C min⁻¹, 5 min/ 200 °C / 20 °C min⁻¹, 5 min.

*t*_S = 29.17 min, *t*_R = 29.56 min

[α _D] = +74.0 (c = 1.0, CH₂Cl₂).

Synthesis of N-((R)-1-(4-(trifluoromethyl)phenyl)ethyl)acetamide (2i):

Compound **2i** was synthesized from 4-trifluoromethylphenyl ethyl amine (**1i**, 0.50 mmol, 95 μ l) following the procedure above using complex **5c** (26.5 mg, 0.02 mmol), CALB (30 mg) and Na₂CO₃ (0.30 mmol, 30 mg). The reaction is stirred for 5 days at 90 °C and after purification by column chromatography (SiO₂, pentane / ethyl acetate, 1:4) **2i** was obtained as a white solid (115 mg, 0.45 mmol, 91 %, 99 % *ee*). The spectroscopic data of the product were identical to those reported in the literature.²

Chiral GC-analysis:

Injector 250 °C

Program: 100 °C / 5 min / 155 °C / 3 °C min⁻¹, 5 min/ 200 °C / 20 °C min⁻¹, 5 min.

*t*_S = 29.17 min, *t*_R = 29.56 min

Synthesis of N-((R)-octan-2-yl)acetamide (2j):

Compound **2j** was synthesized from 2-octylamine (**1j**, 0.50 mmol, 85 μ l) following the procedure above using complex **5c** (26.5 mg, 0.02 mmol), CALB (20 mg) and Na₂CO₃ (0.20 mmol, 20 mg). After purification by column chromatography (SiO₂, pentane / ethyl acetate, 1:4) **2j** was obtained as a white solid (72 mg, 0.42 mmol, 85 %, 93 % *ee*).

¹H-NMR (400.1 MHz, CDCl₃): δ = 5.71 (brs, 1H), 3.93-3.83 (m, 1H), 1.89 (s, 3H), 1.39-1.30 (m, 2H), 1.29-1.15 (m, 8H) 1.05 (t, ³J = 7 Hz, 3H), 0.81 (t, 3H, ³J = 7 Hz) ppm.

¹³C-NMR (100 MHz, CDCl₃): δ = 169.3, 45.2, 36.8, 31.6, 29.0, 25.9, 23.3, 22.5, 20.8 ppm.

One carbon atom missing due to overlapping.

Chiral GC-analysis:

Injector 250 °C

Program: 100 °C / 5 min / 155 °C / 3 °C min⁻¹, 5 min/ 200 °C / 20 °C min⁻¹, 5 min.

t_S = 17.04 min, t_R = 17.71 min

[α_D] = +4.4 (c = 1.0, CH₂Cl₂).

Synthesis of N-((R)-4-phenylbutan-2-yl)acetamide (2k):

Compound **2k** was synthesized from 4-phenylbutan-2-amine (**1k**, 0.50 mmol, 82 μl) following the procedure above using complex **5c** (26.5 mg, 0.02 mmol), CALB (15 mg) and Na₂CO₃ (0.15 mmol, 15 mg). After purification by column chromatography (SiO₂, pentane / ethyl acetate, 1:4) **2k** was obtained as a white solid (75 mg, 0.38 mmol, 78 %, 95 % ee). The spectroscopic data of the product were identical to those reported in the literature.⁴

Chiral GC-analysis:

Injector 250 °C

Program: 100 °C / 5 min / 155 °C / 3 °C min⁻¹, 5 min/ 200 °C / 20 °C min⁻¹, 5 min.

t_S = 28.41 min, t_R = 28.79 min

[α_D] = +45.5 (c = 1.0, CH₂Cl₂).

Spectroscopic data for **5c**

¹H-NMR (400.1 MHz, CDCl₃): δ = 5.71 (brs, 1H), 3.93-3.83 (m, 1H), 1.89 (s, 3H), 1.39-1.30 (m, 2H), 1.29-1.15 (m, 8H) 1.05 (t, ³J = 7 Hz, 3H), 0.81 (t, 3H, ³J = 7 Hz) ppm.

¹³C-NMR (100 MHz, CDCl₃): δ = 169.3, 45.2, 36.8, 31.6, 29.0, 25.9, 23.3, 22.5, 20.8 ppm.

References

1. Kinbara, K.; Harada, Y.; Saigo, K. *J. Chem. Soc., Perkin Trans. 2* **2000**, 1339-1347.
2. Burk, M. J.; Wang, Y. M.; Lee, J. R. *J. Am. Chem. Soc.*, **118**, **1996**, 5142-5143.
3. Zhang, Z.; Zhu, G.; Jiang, Q.; Xiao, D.; Zhang, X.; *J. Org. Chem.* **1999**; **64**; 1774-1775.
4. Gonzalez-Sabin, J.; Gotor, V.; Rebollo, F. *Tetrahedron: Asymmetry* **2002**, **13**, 1315-1320.