Supporting Information

Chemoselective Hydrogenation of Imides Catalyzed by Cp*Ru(PN) Complexes and Its Application to the Asymmetric Synthesis of Paroxetine

Masato Ito, Ayaka Sakaguchi, Chika Kobayashi, and Takao Ikariya*

Department of Applied Chemistry, Graduate School of Science and Engineering, Tokyo Institute of Technology, 2-12-1, O-okayama, Meguro-ku, Tokyo 152-8552, Japan

CONTENTS

1) General methods.	S2
2) Preparation of imides (3).	S3
3) General procedure for the hydrogenation of imides (3) catalyzed by Cp*Ru(PN) complexes and KOt-Bu.	S4
4) Conversion of (R)-(-)-4s to (R)-(+)-5s.	S7
5) CAN-mediated dearylation of (R)-(+)-5s.	S8
6) X-ray diffraction study of 3s and (R)-(+)-5s.	S8
7) References.	S9

1) General methods. All manipulation of oxygen and moisture-sensitive materials were conducted under purified argon atmosphere (BASF–Catalyst R3-11) by the use of standard Schlenk techniques. All reactions were performed in commercially available anhydrous solvents unless otherwise noted. Column chromatography was performed using Fuji Silysia silica gel FL100D (neutral). ^1H (300 MHz) and ^{13}C (75 MHz) spectra were recorded on a JEOL JNM–LA300 spectrometer and chemical shifts were referenced to tetramethylsilane (0.0 ppm). Infrared spectra were recorded on a JASCO FT/IR-610 spectrometer. Optical rotation was measured on JASCO DIP-370. HPLC was performed on JASCO Gulliver system with UV detector. HRMS were measured on a JEOL JMS-T100CS (ESI-TOF), calibrating with reserpine as an internal standard. Elemental analyses were performed by the Analytical Facility at the Research Laboratory of Resources Utilization, Tokyo Institute of Technology or on a Perkin-Elmer 2400II CHN analyzer.

Chemicals. Unless otherwise noted, all reagents and solvents were purchased and used as delivered. Dimethyl glutaconate was purchased from Fluka as an 83:17 trans-cis mixture. $\text{Cp}^*\text{RuCl}(\text{cod})$,^{S1} and Ru complexes **2b**^{S2} and **2f**^{S3} were prepared as reported. Ligands **1c**,^{S2} **1d**,^{S4} **1e**,^{S5} and $\text{Ph}_2\text{P}(\text{CH}_2)_2\text{NMe}_2$ ^{S2} were prepared by the literature methods. 3-Phenylglutaric anhydride^{S6} and *N*-pivaloyl- δ -valerolactam **3m**^{S7} were prepared according to the literature procedures.

Preparation of 3-(4-fluorophenyl)glutaric anhydride. This compound was newly prepared from dimethyl glutaconate in three steps by Rh-catalyzed addition of the 4-fluorophenylboronic acid and subsequent saponification and dehydrative cyclization. The method of Miyaura and co-workers^{S8} was slightly modified for the first step. To a solution of 4-fluorophenylboronic acid (9.69 g, 69.2 mmol) and $[\text{RhCl}(\text{cod})]_2$ (0.68 g, 1.4 mmol) in degassed aqueous 1,4-dioxane (6/1 v/v, 140 mL) was added triethylamine (6.4 mL, 46.0 mmol) and dimethyl glutaconate (6.5 mL, 46.2 mmol). The orange solution was heated at 50 °C for 16 h, during which white precipitates were gradually formed. The resulting mixture was quenched with 1 M citric acid aq. and extracted with diethyl ether. The organic layer was washed with brine, dried over MgSO_4 , filtered through a silica plug, and evaporated. The crude oil was purified by bulb-to-bulb distillation to give the dimethyl 3-(4-fluorophenyl)glutarate as a colorless liquid, which crystallized upon standing (9.5 g, 81% yield). ^1H NMR (CDCl_3) δ 2.58 (dd, J = 15.0, 8.0 Hz, 2H), 2.69 (dd, J = 15.0, 7.0 Hz, 2H), 3.56 (s, 6H), 3.59–3.72 (m, 1H), 6.92–7.04 (m, 2H), 7.14–7.33 (m, 2H).^{S9a} To a solution of the diester (13.0 g, 51 mmol) in 95% ethanol (300 mL) was added a solution of KOH (6.0 g, 107 mmol) in 95% ethanol (150 mL). The mixture was heated under reflux overnight and then evaporated. The residue was dissolved in water, acidified with 1 N HCl aq., and extracted with ethyl acetate. The combined organic layer was washed with brine, dried over MgSO_4 , and filtered through a pad of Celite. Concentration of the filtrate followed by recrystallization from *n*-hexane/ethyl acetate to give 3-(4-fluorophenyl)glutaric acid (9.75 g, 84% yield). ^1H NMR (CDCl_3) δ 2.65 (dd, J = 15.7, 8.2 Hz, 2H), 2.72 (dd, J = 15.7, 7.0 Hz, 2H), 3.58–3.72 (m, 1H), 6.98–7.00 (m, 2H), 7.18–7.23 (m, 2H).^{S9b} Thus obtained acid (9.60 g, 42 mmol) was dissolved in acetyl chloride (15 mL) and heated at 65 °C overnight. The mixture was concentrated *in vacuo* to give the red solid, which was then dissolved in ethyl acetate and filtered through a plug of silica gel. Concentration of the filtrate followed by recrystallization from *n*-hexane/ethyl acetate gave the title compound as a yellow solid (8.22 g, 93% yield). ^1H NMR (CDCl_3) δ 2.82 (dd, J = 17.2, 11.1 Hz, 2H), 3.11 (dd, J = 17.2, 4.2 Hz, 2H), 3.42 (tt, J = 11.1, 4.2 Hz, 1H), 7.06–7.20 (m, 4H).^{S9b}

2) Preparation of imides (3).

Preparation of 3c. To a solution of *N*-benzylmaleimide (1.15 g, 6.1 mmol) in 95% ethanol (95%, 70 mL) and CH₂Cl₂ (10 mL) was added 5% Pd/C (55 mg) and the resulting mixture was vigorously stirred under hydrogen atmosphere at 30 °C for 16 h. The mixture was filtered through a plug of Celite and the filtrate was concentrated *in vacuo* to give a crude product, which was purified by recrystallization from *n*-hexane/ethyl acetate to give the title compounds as white needles (1.07 g, 93% yield). ¹H NMR (CDCl₃) δ 2.71 (s, 4H), 4.66 (s, 2H), 7.27–7.41 (m, 5H).^{S10}

Preparation of 3d–h, 3j–l, 3n, and 3p–s. These compounds were prepared by the condensation of the corresponding acid anhydrides with primary amines. The literature method^{S11} was slightly modified: To a solution of the appropriate acid anhydride (10 mmol) and triethylamine (10 mmol) in THF (50 mL) was slowly added a solution of the appropriate primary amine (10 mmol) in THF (50 mL) at 0 °C. The mixture was heated under reflux overnight and cooled. The mixture was washed with 1 N HCl aq. and brine, dried over MgSO₄, filtered through a pad of Celite. Evaporation of the filtrate left a crude amide acid, which was dissolved in acetyl chloride. The resulting solution was heated at 65 °C overnight and cooled. Concentration of the mixture under reduced pressure gave a crude imide, which was purified either by bulb-to-bulb distillation or by recrystallization from *n*-hexane/ethyl acetate. Yields (unoptimized) were 59% (3d), 78% (3e), 54% (3f), 73% (3g), 70% (3h), 20% (3j), 46% (3k), 22% (3l), 97% (3n), 94% (3p), 59% (3q), 94% (3r), and 98% (3s). Spectral data and elemental analyses for each compound are as follows.

- 3d** ¹H NMR (CDCl₃) δ 1.90–1.98 (m, 2H), 2.67 (t, *J* = 6.6 Hz, 4H), 4.96 (s, 2H), 7.21–7.40 (m, 5H).^{S12}
- 3e** ¹H NMR (CDCl₃) δ 1.06 (d, *J* = 6.1 Hz, 3H), 2.19–2.36 (m, 3H), 2.77 (dd, *J* = 16.6, 3.5 Hz, 2H), 4.94 (s, 2H), 7.22–7.36 (s, 5H).^{S11}
- 3f** ¹H NMR (CDCl₃) δ 2.80 (dd, *J* = 16.9, 11.8 Hz, 2H), 2.99 (dd, *J* = 16.9, 4.1 Hz, 2H), 3.35 (tt, *J* = 11.8, 4.1 Hz, 1H), 4.97 (s, 2H), 7.14–7.35 (m, 10H).^{S13}
- 3g** ¹H NMR (CDCl₃) δ 2.75 (dd, *J* = 17.1, 11.5 Hz, 2H), 2.97 (dd, *J* = 17.1, 4.4 Hz, 2H), 3.32 (tt, *J* = 11.5, 4.4 Hz, 1H), 4.95 (s, 2H), 6.96–7.35 (m, 9H).^{S14}
- 3h** ¹H NMR (CDCl₃) δ 1.04 (s, 6H), 2.52 (s, 4H), 4.96 (s, 2H), 7.21–7.30 (m, 3H), 7.34–7.37 (m, 2H).^{S13}
- 3j** ¹H NMR (CDCl₃) δ 2.11 (quint, *J* = 6.6 Hz, 2H), 2.83 (t, *J* = 6.6 Hz, 4H), 7.08–7.11 (m, 2H), 7.38–7.49 (m, 3H).^{S15}
- 3k** ¹H NMR (CDCl₃) δ 2.10 (quint, *J* = 6.8 Hz, 2H), 2.82 (t, *J* = 6.8 Hz, 4H), 3.82 (s, 3H), 6.72–7.02 (m, 4H).^{S16}
- 3l** ¹H NMR (acetone-*d*₆) δ 2.04–2.06 (m, 2H), 2.69–2.75 (m, 4H), 7.78 (d, *J* = 9.0 Hz, 2H), 8.10 (d, *J* = 9.0 Hz, 2H).^{S17}
- 3n** ¹H NMR (CDCl₃) δ 1.21 (s, 6H), 1.75 (t, *J* = 6.8 Hz, 2H), 2.66 (t, *J* = 6.8 Hz, 2H), 4.88 (s, 2H), 7.16–7.27 (m, 5H).^{S18}
- 3p** ¹H NMR (CD₂Cl₂) δ 2.75 (dd, *J* = 17.0, 11.5 Hz, 2H), 2.93 (dd, *J* = 17.0, 4.4 Hz, 2H), 3.33 (tt, *J* = 11.5, 4.4 Hz, 1H), 5.32 (s, 2H), 6.90–7.00 (m, 2H), 7.02–7.07 (m, 1H), 7.08–7.11 (m, 2H), 7.25 (t, *J* = 8.3 Hz, 1H), 7.37–7.45 (m, 2H), 7.64 (d, *J* = 8.3 Hz, 1H), 7.75 (d, *J* = 9.0 Hz, 1H), 7.99 (d, *J* = 8.5 Hz, 1H); ¹³C{¹H} NMR (CD₂Cl₂) δ 33.8, 40.1, 40.5, 115.9 (d, *J* = 21.7 Hz), 123.3, 125.0, 125.2, 125.7, 126.2, 127.9 (d, *J* = 8.0 Hz), 128.0, 128.7, 131.2, 131.8, 133.7, 136.1 (d, *J* = 3.4 Hz), 162.0, (d, *J* = 246.4 Hz), 171.5; IR (KBr) 1678, 1728 cm⁻¹ (ν_{C=O}); Anal. Calcd for C₂₂H₁₈FNO₂: C, 76.06; H, 5.22; N, 4.03. Found: C, 75.93; H, 5.13; N, 3.89.

- 3q** ^1H NMR (CD_2Cl_2) δ 2.95 (dd, $J = 17.1, 11.7$ Hz, 2H), 3.10 (dd, $J = 17.1, 4.4$ Hz, 2H), 3.54 (tt, $J = 11.7, 4.4$ Hz, 1H), 7.09–7.48 (m, 9H); $^{13}\text{C}\{^1\text{H}\}$ NMR (CD_2Cl_2) δ 34.0, 40.1, 116.1 (d, $J = 21.4$ Hz), 120.1, 127.9 (d, $J = 8.0$ Hz), 128.3, 128.7, 129.3, 136.1 (d, $J = 3.1$ Hz), 162.0 (d, $J = 246.8$ Hz), 171.4; IR (KBr) 1681, 1730 cm^{-1} ($\nu_{\text{C}=\text{O}}$); Anal. Calcd for $\text{C}_{17}\text{H}_{14}\text{FNO}_2$: C, 72.07; H, 4.98; N, 4.94. Found: C, 71.88; H, 4.84; N, 4.93.
- 3r** ^1H NMR (CD_2Cl_2) δ 2.95 (dd, $J = 16.8, 11.5$ Hz, 2H), 3.10 (dd, $J = 16.8, 4.4$ Hz, 2H), 3.54 (tt, $J = 11.5, 4.4$ Hz, 1H), 3.84 (s, 3H), 6.96–7.04 (m, 4H), 7.09–7.14 (m, 2H), 7.26–7.31 (m, 2H); $^{13}\text{C}\{^1\text{H}\}$ NMR (CD_2Cl_2) δ 34.1, 40.2, 55.5, 114.4, 115.9 (d, $J = 21.3$ Hz), 127.8, 128.2 (d, $J = 8.0$ Hz), 129.4, 136.7 (d, $J = 3.4$ Hz), 159.6, 162.0 (d, $J = 244.9$ Hz), 171.7; IR (KBr) 1677, 1733 cm^{-1} ($\nu_{\text{C}=\text{O}}$); Anal. Calcd for $\text{C}_{18}\text{H}_{16}\text{FNO}_3$: C, 69.00; H, 5.15; N, 4.47. Found: C, 68.97; H, 4.96; N, 4.40.
- 3s** ^1H NMR (CD_2Cl_2) δ 2.92 (dd, $J = 16.2, 11.3$ Hz, 2H), 3.08 (dd, $J = 16.2, 4.4$ Hz, 2H), 3.54 (tt, $J = 11.3, 4.4$ Hz, 1H), 6.03 (s, 2H), 6.55 (d, $J = 7.8$ Hz, 1H), 6.56 (s, 1H), 6.87 (d, $J = 7.8$ Hz, 1H), 7.08–7.13 (m, 2H), 7.25–7.30 (m, 2H); $^{13}\text{C}\{^1\text{H}\}$ NMR (CD_2Cl_2) δ 34.0, 40.1, 101.8, 108.6, 109.2, 116.1 (d, $J = 21.7$ Hz), 121.7, 127.9 (d, $J = 8.4$ Hz), 128.1, 136.1 (d, $J = 3.4$ Hz), 147.9, 148.3, 162.1 (d, $J = 246.6$ Hz), 171.6; IR (KBr) 1677, 1736 cm^{-1} ($\nu_{\text{C}=\text{O}}$); Anal. Calcd for $\text{C}_{18}\text{H}_{14}\text{FNO}_4$: C, 66.05; H, 4.31; N, 4.28. Found: C, 66.09; H, 4.34; N, 4.23.

Preparation of 3i. To a mixture of 3-ethyl-3-methylglutarimide (5.19 g, 33.4 mmol) and K_2CO_3 (4.65 g, 33.6 mmol) in H_2O (45 mL) and DMF (15 mL) was slowly added benzyl bromide (4.11 mL, 34.5 mmol) and the resulting mixture was heated at 90 °C for 2 days. The mixture was diluted with H_2O and extracted with CHCl_3 . The combined organic layer was washed with 1 N NaOH aq. and brine, dried over MgSO_4 , and filtered through a pad of Celite. The filtrate was evaporated to leave a crude product, which was purified by silica gel chromatography eluting with *n*-hexane/ethyl acetate (9/1) to give **3i** as a colorless liquid (5.91 g, 72% yield). ^1H NMR (CDCl_3) δ 0.88 (t, $J = 7.3$ Hz, 3H), 0.96 (s, 3H), 1.35 (q, $J = 7.3$ Hz, 2H), 2.49 (s, 4H), 4.94 (s, 2H), 7.21–7.67 (m, 5H).^{S19}

Preparation of 3o. To a suspension of L-phenylalanine methyl ester hydrochloride (4.31 g, 20.0 mmol) in THF (50 mL) was added triethylamine (2.5 mL, 20 mmol), and the mixture was stirred vigorously at room temperature for 1 h to give another precipitates, which were removed by filtration through a pad of Celite. To the filtrate was added phthalic anhydride (2.96 g, 20.0 mmol) and subsequently carbonyl-1,1'-carbonylbis-1*H*-imidazole) (3.24 g, 20.0 mmol). After stirring the resulting mixture at room temperature for 14 h, the solvent was removed under reduced pressure and the residue was purified by silica gel chromatography eluting with *n*-hexane/ethyl acetate (9/1) to give **3o** as a viscous oil, which crystallized upon standing (4.07 g, 66%). ^1H NMR (CDCl_3) δ 3.50 (dd, $J = 14.4, 11.0$ Hz, 1H), 3.59 (dd, $J = 14.4, 5.6$ Hz, 1H), 3.76 (s, 3H), 5.13 (dd, $J = 11.0, 5.6$ Hz, 1H), 7.13–7.18 (m, 5H), 7.65–7.77 (m, 4H).^{S20}

3) General procedure for the hydrogenation of imides (3) catalyzed by $\text{Cp}^*\text{Ru}(\text{PN})$ complexes and KO*t*-Bu. A degassed solution of **3** in 2-propanol was transferred to a mixture either of $\text{Cp}^*\text{RuCl}(\text{cod})$, **1b**, and KO*t*-Bu, or of **2b** and KO*t*-Bu, or of **2f** and KO*t*-Bu placed in a 50-mL stainless autoclave. Argon present in the autoclave was replaced by hydrogen. Then hydrogen (1–3 MPa) was introduced into the autoclave. The mixture was stirred at 80 °C for the time indicated in the Table 1 or 2. After

carefully venting hydrogen, the solvent was removed under reduced pressure and the residue was filtered through a plug of silica gel. The filtrate was concentrated under reduced pressure, and the residue was analyzed by ^1H NMR spectroscopy. Yields of **4** indicated in Table 1 or 2 were determined by integrating well-resolved characteristic resonances of **3** and **4**.

- 4a** ^1H NMR (CDCl_3) δ 1.20 (br, 1H, D_2O -exchangeable), 4.64 (s, 2H), 4.66 (d, $J = 5.6$ Hz, 2H), 6.56 (br, 1H, D_2O -exchangeable), 7.26–7.49 (m, 8H), 7.55 (d, $J = 7.6$ Hz, 1H).^{S21}
- 4b** ^1H NMR (CDCl_3) δ 4.07 (br, 1H, D_2O -exchangeable), 4.64 (s, 2H), 7.15 (t, $J = 7.5$ Hz, 1H), 7.33–7.49 (m, 5H), 7.61–7.71 (m, 3H), 8.62 (br, 1H, D_2O -exchangeable).^{S22}
- 4c** ^1H NMR (CDCl_3) δ 1.84 (tt, $J = 6.8, 5.8$ Hz, 2H), 2.33 (t, $J = 6.8$ Hz, 2H), 3.11 (br, 1H, D_2O -exchangeable), 3.63 (t, $J = 5.8$ Hz, 2H), 4.37 (d, $J = 5.9$ Hz, 2H), 6.19 (br, 1H, D_2O -exchangeable), 7.21–7.29 (m, 5H).^{S11,S23}
- 4d** ^1H NMR (CDCl_3) δ 1.18 (d, 1H, D_2O -exchangeable), 1.59–1.76 (m, 4H), 2.25 (t, $J = 7.2$ Hz, 2H), 3.62–3.64 (m, 2H), 4.42 (d, $J = 5.7$ Hz, 2H), 5.76 (br, 1H, D_2O -exchangeable), 7.24–7.34 (m, 5H).^{S11,S24}
- 4e** ^1H NMR (CDCl_3) δ 0.96 (d, $J = 6.5$ Hz, 3H), 1.50–1.53 (m, 2H), 2.09–2.23 (m, 3H), 2.45 (br, 1H, D_2O -exchangeable), 3.63–3.64 (m, 2H), 4.41 (d, $J = 5.8$ Hz, 2H), 5.99 (br, 1H, D_2O -exchangeable), 7.24–7.30 (m, 5H).^{S11}
- 4f** ^1H NMR (CDCl_3) δ 1.73 (br, 1H, D_2O -exchangeable), 1.88–1.96 (m, 2H), 2.47 (dd, $J = 14.4, 7.8$ Hz, 1H), 2.59 (dd, $J = 14.4, 6.8$ Hz, 1H), 3.32–3.37 (m, 1H), 3.52–3.57 (m, 2H), 4.26 (dd, $J = 14.7, 5.4$ Hz, 1H), 4.37 (dd, $J = 14.7, 5.9$ Hz, 1H), 5.51 (br, 1H, D_2O -exchangeable), 6.99–7.30 (m, 10H); ^{13}C NMR (CDCl_3) δ 39.4, 39.5, 43.9, 44.3, 60.9, 127.1, 127.7, 127.8, 128.0, 129.0, 129.1, 138.3, 144.2, 172.0; IR (neat) 3063, 1730 cm^{-1} ; HRMS m/z ($\text{M}+\text{Na}$)⁺ calcd for $\text{C}_{18}\text{H}_{21}\text{NO}_2$, 306.1470; found 306.1472.
- 4g** ^1H NMR (CDCl_3) δ 1.50 (br, 1H, D_2O -exchangeable), 1.81–1.91 (m, 2H), 2.38 (dd, $J = 14.2, 7.3$ Hz, 1H), 2.58 (dd, $J = 14.2, 6.4$ Hz, 1H), 3.30–3.39 (m, 1H), 3.41–3.56 (m, 2H), 4.21 (dd, $J = 14.7, 5.1$ Hz, 1H), 4.37 (dd, $J = 14.7, 6.1$ Hz, 1H), 5.68 (br, 1H, D_2O -exchangeable), 6.91–7.24 (m, 9H); $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3) δ 38.4, 39.0, 43.5, 44.1, 60.4, 115.5 ($J = 21.0$ Hz), 127.5 (d, $J = 8.3$ Hz), 128.6, 128.8, 128.9, 137.9, 139.5 (d, $J = 3.1$ Hz), 161.6 (d, $J = 244.5$ Hz), 171.3; IR (neat) 3300, 1671 cm^{-1} ; HRMS m/z ($\text{M}+\text{Na}$)⁺ calcd for $\text{C}_{18}\text{H}_{20}\text{FNO}_2$, 324.1376; found 324.1352.; (+)-**4g** (64% ee), $[\alpha]_{\text{D}} +3.0$ (c 1.3, CHCl_3); Daicel Chiralcel OD–H column, df = 4.6 mm i.d. x 250 mm; eluent, 90/10 = *n*-hexane/2-propanol; temp. 30 °C, flow rate, 0.5 mL/min; detection (UV), $\nu = 254$ nm-light, $t_{\text{R}} = 36.9$ min for the major isomer and $t_{\text{R}} = 45.8$ min for the minor isomer.
- 4h** ^1H NMR (CDCl_3) δ 1.02 (s, 6H), 1.65 (t, $J = 5.8$ Hz, 2H), 2.33 (s, 2H), 3.75 (t, $J = 5.8$ Hz, 2H), 4.41 (d, $J = 5.6$ Hz, 2H), 6.65 (br, 1H, D_2O -exchangeable), 7.24–7.45 (m, 5H); $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3) δ 29.0, 32.9, 42.5, 43.7, 48.1, 59.5, 127.6, 127.9, 128.7, 138.2, 172.3; IR (neat) 3295, 1645 cm^{-1} ; HRMS m/z ($\text{M}+\text{Na}$)⁺ calcd for $\text{C}_{14}\text{H}_{21}\text{NO}_2$, 258.1470; found 258.1429.
- 4i** ^1H NMR (CDCl_3) δ 0.85 (t, $J = 7.6$ Hz, 3H), 0.95 (s, 3H), 1.38 (q, $J = 7.6$ Hz, 2H), 1.68 (t, $J = 5.7$ Hz, 2H), 2.24 (d, $J = 13.0$ Hz, 1H), 2.32 (d, $J = 13.0$ Hz, 1H), 3.50 (br, 1H, D_2O -exchangeable), 3.77 (t, $J = 5.7$ Hz, 2H), 4.44 (d, $J = 5.6$ Hz, 2H), 6.29 (br, 1H, D_2O -exchangeable), 7.25–7.37 (m, 5H); $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3) δ 8.0, 25.1, 33.3, 35.4, 39.8, 43.6, 46.3, 59.1, 127.5, 127.8, 128.7,

- 138.2, 172.4; IR (neat) 3290, 1645 cm^{-1} ; HRMS m/z ($\text{M}+\text{Na}$)⁺ calcd for $\text{C}_{15}\text{H}_{23}\text{NO}_2$, 272.1626; found 272.1625.
- 4j** ^1H NMR (CDCl_3) δ 1.60–1.68 (m, 2H), 1.78–1.87 (m, 2H), 2.41 (t, J = 7.2 Hz, 2H), 3.68 (t, J = 6.0 Hz, 2H), 7.08 (t, J = 7.3 Hz, 1H), 7.26–7.32 (m, 2H), 7.51–7.71 (m, 2H), 7.85 (br, 1H, D_2O -exchangeable).^{S25}
- 4k** ^1H NMR (CDCl_3) δ 1.56–1.60 (m, 2H), 1.72–1.76 (m, 2H), 2.29–2.34 (m, 2H), 3.57–3.60 (m, 2H), 3.72 (s, 3H), 6.75–6.80 (m, 2H), 7.35–7.39 (m, 2H), 8.00 (br, 1H, D_2O -exchangeable).^{S25a}
- 4l** ^1H NMR (CDCl_3) δ 1.64–1.73 (m, 2H), 1.83–1.92 (m, 2H), 2.00 (t, J = 4.4 Hz, 1H, D_2O -exchangeable), 2.51 (t, J = 7.2 Hz, 2H), 3.75 (td, J = 5.7, 4.4 Hz, 2H), 7.72 (d, J = 9.3 Hz, 2H), 8.16 (br, 1H, D_2O -exchangeable), 8.20 (d, J = 9.3 Hz, 2H); $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3) δ 22.0, 31.2, 37.2, 62.5, 118.9, 125.1, 143.4, 143.9, 171.9; IR (KBr) 3671, 1687 cm^{-1} ; HRMS m/z ($\text{M}+\text{Na}$)⁺ calcd for $\text{C}_{11}\text{H}_{14}\text{N}_2\text{O}_4$, 261.0851; found 261.0846.
- 4m** ^1H NMR (CDCl_3) δ 1.52–2.00 (m, 4H), 2.35 (t, J = 6.4 Hz), 3.25–3.35 (m, 2H), 5.75 (br, 1H, D_2O -exchangeable).
- 4n** ^1H NMR (CDCl_3) δ 1.19 (s, 6H), 1.49–1.55 (m, 2H), 1.58–1.65 (m, 2H), 2.36 (br, 1H, D_2O -exchangeable), 3.56 (t, J = 6.1 Hz, 2H), 4.41 (d, J = 5.6 Hz, 2H), 6.15 (br, 1H, D_2O -exchangeable), 7.23–7.35 (m, 5H); $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3) δ 25.6, 28.0, 36.7, 41.8, 43.6, 62.5, 127.6, 127.7, 128.6, 138.5, 177.7; IR (neat) 2966, 1679 cm^{-1} ; HRMS m/z ($\text{M}+\text{Na}$)⁺ calcd for $\text{C}_{14}\text{H}_{21}\text{NO}_2$, 258.1470; found 258.1475.
- 4p** ^1H NMR (CDCl_3) δ 1.89–1.93 (m, 2H), 2.41 (dd, J = 14.2, 7.8 Hz, 1H), 2.56 (dd, J = 14.2, 6.7 Hz, 1H), 3.32–3.40 (m, 1H), 3.48–3.55 (m, 2H), 4.78 (d, J = 5.1 Hz, 2H), 5.48 (br, 1H, D_2O -exchangeable), 6.90 (t, J = 8.6 Hz, 2H), 7.09–7.51 (m, 6H), 7.78–7.85 (m, 3H); $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3) δ 38.3, 38.9, 41.7, 43.9, 60.4, 115.4 (d, J = 20.9 Hz), 123.3, 126.0, 126.6, 128.7 (d, J = 8.0 Hz), 131.2, 133.1, 133.8, 139.3 (d, J = 3.2 Hz), 161.5 (d, J = 244.2 Hz), 171.0; IR (KBr) 3312, 1627 cm^{-1} ; HRMS m/z ($\text{M}+\text{Na}$)⁺ calcd for $\text{C}_{22}\text{H}_{22}\text{FNO}_2$, 374.1532; found 374.1531.; (–)-**4p** (75% ee), $[\alpha]_{\text{D}} -1.7$ (c 1.1, CHCl_3); Daicel Chiralcel OD column, df = 4.6 mm i.d. x 250 mm; eluent, 90/10 = *n*-hexane/2-propanol; temp. 30 °C, flow rate, 0.5 mL/min; detection (UV), ν = 254 nm-light), t_{R} = 36.6 min for the major isomer and t_{R} = 49.7 min for the minor isomer.
- 4q** ^1H NMR (CDCl_3) δ 1.90–2.00 (m, 2H), 1.95 (br, 1H), 2.60 (dd, J = 14.5, 7.3 Hz, 1H), 2.73 (dd, J = 14.5, 7.1 Hz, 1H), 3.38–3.45 (m, 1H), 3.50–3.63 (m, 2H), 6.97–7.11 (m, 3H), 7.19–7.38 (m, 6H); $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3) δ 38.3, 38.7, 44.9, 60.4, 115.6 (d, J = 21.4 Hz), 120.0, 124.5, 128.9 (d, J = 7.6 Hz), 129.0, 137.5, 139.4 (d, J = 3.4 Hz), 161.6 (d, J = 245.3 Hz), 169.9; IR (neat) 3299, 1662 cm^{-1} ; HRMS m/z ($\text{M}+\text{Na}$)⁺ calcd for $\text{C}_{17}\text{H}_{18}\text{FNO}_2$, 310.1219; found 310.1200.; (–)-**4q** (93% ee), $[\alpha]_{\text{D}} -28.6$ (c 0.5, CHCl_3); Daicel Chiralpak AS column, df = 4.6 mm i.d. x 250 mm; eluent, 88/12 = *n*-hexane/2-propanol; temp. 30 °C, flow rate, 0.5 mL/min; detection (UV), ν = 254 nm-light), t_{R} = 25.1 min for the minor isomer and t_{R} = 30.1 min for the major isomer.
- 4r** ^1H NMR (CDCl_3) δ 1.75–2.00 (m, 2H), 2.47 (dd, J = 14.3, 7.5 Hz, 1H), 2.64 (dd, J = 14.3, 7.1 Hz, 1H), 2.95 (br, 1H, D_2O -exchangeable), 3.30–3.40 (m, 1H), 3.45–3.51 (m, 2H), 3.72 (s, 3H), 6.73–6.76 (m, 2H), 6.90–6.95 (m, 2H), 7.08–7.13 (m, 2H), 7.20–7.23 (m, 2H), 7.95 (br, 1H, D_2O -exchangeable); $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3) δ 38.2, 38.6, 44.2, 55.3, 60.0, 114.0, 115.3 (d, J = 20.9 Hz), 122.2, 128.8 (d, J = 8.0 Hz), 130.6, 139.4 (d, J = 3.5 Hz), 156.4, 161.4 (d,

$J = 244.2$ Hz), 170.4; IR (neat) 3230, 1650 cm^{-1} ; Anal. Calcd for $\text{C}_{18}\text{H}_{20}\text{FNO}_3$: C, 68.12; H, 6.35; N, 4.41. Found: C, 67.93; H, 6.34; N, 4.43.; (–)-**4r** (91% ee), $[\alpha]_{\text{D}} -33.0$ (c 0.5, CHCl_3); Daicel Chiralcel OD column, df = 4.6 mm i.d. x 250 mm; eluent, 90/10 = *n*-hexane/2-propanol; temp. 30 °C, flow rate, 0.5 mL/min; detection (UV), $\nu = 254$ nm-light), $t_{\text{R}} = 61.0$ min for the major isomer and $t_{\text{R}} = 75.3$ min for the minor isomer.

4s ^1H NMR (CDCl_3) δ 1.90–1.97 (m, 2H), 2.56 (dd, $J = 14.4, 7.3$ Hz, 1H), 2.70 (dd, $J = 14.4, 7.1$ Hz, 1H), 3.39–3.44 (m, 1H), 3.56–3.61 (m, 2H), 5.92 (s, 2H), 6.62 (d, $J = 8.3$ Hz, 1H), 6.69 (d, $J = 8.3$ Hz, 1H), 6.98–7.03 (m, 2H), 7.08 (s, 1H), 7.19–7.24 (m, 2H); $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3) δ 38.4, 38.7, 44.8, 60.4, 101.3, 103.1, 108.0, 113.3, 115.6 (d, $J = 21.0$ Hz), 128.9 (d, $J = 8.0$ Hz), 131.7, 139.4 (d, $J = 3.1$ Hz), 144.4, 147.8, 161.6 (d, $J = 244.9$ Hz), 169.7; IR (neat) 3288, 1656 cm^{-1} ; HRMS m/z ($\text{M}+\text{Na}$)⁺ calcd for $\text{C}_{18}\text{H}_{18}\text{FNO}_4$, 354.1118; found 354.1088.; (R)-(–)-**4s** (98% ee), $[\alpha]_{\text{D}} -39.9$ (c 0.9, CHCl_3); Daicel Chiralpak AD column, df = 4.6 mm i.d. x 250 mm; eluent, 90/10 = *n*-hexane/2-propanol; temp. 30 °C, flow rate, 0.5 mL/min; detection (UV), $\nu = 254$ nm-light), $t_{\text{R}} = 60.0$ min for (R)-isomer and $t_{\text{R}} = 72.8$ min for (S)-isomer.

Deprotection of 3o. Following the general procedure, the hydrogenation of **3o** (0.26 mmol, 0.02 M in 2-propanol, $P_{\text{H}_2} = 1$ MPa) was carried out using the ternary catalyst system of $\text{Cp}^*\text{RuCl}(\text{cod})$, **1b**, and KOt-Bu (10 mol%) at 80 °C for 2 h. ^1H NMR analysis of the aliquot showed the complete consumption of **3o** and the formation of **4o**. ^1H NMR (CDCl_3) δ 3.17 (dd, $J = 13.9, 6.3$ Hz, 1H), 3.31 (dd, $J = 13.9, 5.6$ Hz, 1H), 3.79 (s, 3H), 4.05–4.08 (m, 1H, D_2O -exchangeable), 4.52 (d, $J = 6.6$ Hz, 2H), 5.03–5.09 (m, 1H), 6.75 (br, 1H, D_2O -exchangeable), 7.12–7.46 (m, 9H). Thus obtained crude **4o** was diluted with 1,4-dioxane (8.0 mL) and to this was added 4 N HCl solution in 1,4-dioxane (0.3 mL, 1.2 mmol). The resulting mixture was stirred at 30 °C for 1 h and then concentrated under reduced pressure. The residual white solids were washed with diethyl ether to leave the L-phenylalanine methyl ester hydrochloride (57.6 mg, >99% yield), $[\alpha]_{\text{D}} +36.6$ (c 2.0, $\text{CH}_3\text{CH}_2\text{OH}$). Concentration of the diethyl ether washings afforded phthalide (35.2 mg, >99% yield).

4) Conversion of (R)-(–)-4s to (R)-(+)–5s. To a solution of (R)-(–)-**4s** (>99% ee, 358.5 mg, 1.08 mmol) in CH_2Cl_2 (15 mL) was added PPh_3 (354.7 mg, 1.35 mmol), and CBr_4 (448.5 mg, 1.35 mmol). The resulting mixture was stirred at room temperature for 14 h. The solvent was removed under reduced pressure and the residue was purified by silica gel chromatography eluting with *n*-hexane/ethyl acetate (4/1) to give the bromide (278.3 mg, 65% yield, unoptimized) as colorless liquid, which was used without further purification for the next step. ^1H NMR (CDCl_3) δ 2.10–2.38 (m, 2H), 2.50–2.65 (m, 2H), 3.04–3.12 (m, 1H), 3.12–3.40 (m, 1H), 3.42–3.50 (m, 1H), 5.93 (s, 2H), 6.57 (d, $J = 8.3$ Hz, 1H), 6.69 (d, $J = 8.3$ Hz, 1H), 7.00–7.05 (m, 3H), 7.20–7.49 (m, 2H). Thus obtained bromide was dissolved in THF (40 mL) and to this was added NaH (60%, 71.0 mg, 1.77 mmol). The resulting mixture was heated at 85 °C for 24 h. The reaction was quenched by adding methanol (0.76 mL) and the solvent was removed under reduced pressure. The residue was diluted with H_2O and extracted with ethyl acetate. The combined organic layer was washed with brine, dried over Na_2SO_4 , and filtered through a pad of Celite. The filtrate was evaporated to give the crude product, which was purified by recrystallization from *n*-hexane/ethyl acetate to afford (R)-(+)–**5s** as a colorless solid (119.1 mg, 54% yield, unoptimized). ^1H NMR (CDCl_3) δ 2.00–2.10 (m, 1H), 2.10–2.20 (m, 1H), 2.63 (dd, $J = 17.6, 10.7$ Hz, 1H), 2.86 (dd, $J = 17.6, 5.4$ Hz,

1H), 3.21–3.27 (m, 1H), 3.55–3.62 (m, 1H), 3.67–3.76 (m, 1H), 5.98 (s, 2H), 6.69 (d, $J = 8.0$ Hz, 1H), 6.75 (s, 1H), 6.82 (d, $J = 8.0$ Hz, 1H), 7.02–7.08 (m, 2H), 7.19–7.24 (m, 2H); $^{13}\text{C}\{^1\text{H}\}$ NMR (CDCl_3) δ 30.8, 38.1, 39.9, 51.0, 101.5, 107.9, 108.5, 115.6, (d, $J = 20.8$ Hz), 119.5, 127.9 (d, $J = 8.0$ Hz), 136.8, 138.9 (d, $J = 3.1$ Hz), 146.5, 148.1, 161.7 (d, $J = 244.5$ Hz), 169.4; IR (KBr) 1639 cm^{-1} ($\nu_{\text{C}=\text{O}}$); $[\alpha]_{\text{D}} +84.6$ (c 0.9, CHCl_3); Anal. Calcd for $\text{C}_{18}\text{H}_{16}\text{FNO}_3$: C, 69.00; H, 5.15; N, 4.47. Found: C, 68.85; H, 5.16; N, 4.55.

5) CAN-mediated dearylation of (*R*)-(+)-5s. To a solution of (*R*)-(+)-5s (88.2 mg, 0.28 mmol) in acetonitrile (27.0 mL) was slowly added a solution of $(\text{NH}_4)_2\text{Ce}(\text{NO}_3)_6$ (CAN, 612.5 mg, 1.12 mmol) in H_2O (9.0 mL) at room temperature. The resulting mixture was stirred at room temperature overnight and quenched by adding 1 N NaOH aq. The mixture was extracted with CH_2Cl_2 , washed with brine, dried over Na_2SO_4 , and filtered through a pad of Celite. The filtrate was concentrated under reduced pressure to give the crude product, which was purified by recrystallization from *n*-hexane/ethyl acetate to give (*R*)-(+)-6s (19.0 mg, 35% yield, unoptimized). ^1H NMR (CDCl_3) δ 1.80–1.93 (m, 1H), 1.98–2.03 (m, 1H), 2.44 (dd, $J = 17.6, 11.1$ Hz, 1H), 2.67 (dd, $J = 17.6, 4.2$ Hz, 1H), 3.02–3.04 (m, 1H), 3.38–3.44 (m, 2H), 6.17 (br, 1H, D_2O -exchangeable), 6.93–6.99 (m, 2H), 7.08–7.18 (m, 2H).^{S26}

6) X-ray diffraction study of 3s and (*R*)-(+)-5s. Single crystals suitable for X-ray analyses were mounted on glass fibers. Diffraction experiments were performed on a Rigaku Saturn CCD area detector with graphite monochromated Mo- $\text{K}\alpha$ radiation ($\lambda = 0.710\ 70\ \text{\AA}$). Intensity data were corrected for Lorentz-polarization effects and for absorption. Structure solution and refinements were carried out by using the CrystalStructure program package.^{S27} The heavy-atom positions were determined by a direct method SIR92^{S28} and remaining non-hydrogen atoms were found by subsequent Fourier syntheses. All non-hydrogen atoms were refined anisotropically by full-matrix least-squares techniques based on F^2 . All hydrogen atoms were placed at calculated positions. The absolute structures were determined on the basis of the Flack absolute structure parameters.^{S29} The atomic scattering factors were taken from ref. S30, and anomalous dispersion effects were included; the values of Δf and $\Delta f'$ were taken from ref. S31.

Figure S1. Crystal Structure of 3s. Hydrogen atoms are omitted for clarity. Thermal ellipsoids are drawn at 50% probability.

Figure S2. Crystal Structure of (*R*)-(+)-**5s**. Hydrogen atoms are omitted for clarity. Thermal ellipsoids are drawn at 50% probability.

7) References

- (S1) Oshima, N.; Suzuki, H.; Moro-oka, Y. *Chem. Lett.* **1984**, 1161–1164.
- (S2) Ito, M.; Hirakawa, M.; Osaku, A.; Ikariya, T. *Organometallics* **2003**, 22, 4190–4192.
- (S3) Ito, M.; Kitahara, S.; Ikariya, T. *J. Am. Chem. Soc.* **2005**, 127, 6172–6173.
- (S4) (a) Cadogan, J. I. G.; Sears, D. J.; Smith, D. M. *J. Chem. Soc. C* **1969**, 1314–1318
(b) Cooper, M. K.; Downes, J. M. *Inorg. Chem.* **1978**, 17, 880–884.
- (S5) (a) Coote, S. J.; Dawson, G. J.; Frost, C. G.; Williams, M. J. *Synlett* **1993**, 509–510. (b) Brunner, A.; Kühnle, F. N. M.; Seebach, D. *Helv. Chim. Acta.* **1996**, 79, 319–345. (c) Hingst, M.; Tepper, M.; Stelzer, O. *Eur. J. Inorg. Chem.* **1998**, 73–82.
- (S6) (a) Reed, P. E.; Katzenellenbogen, J. A. *J. Med. Chem.* **1991**, 34, 1162–1176.
(b) Theisen, P. D.; Heathcock, C. H. *J. Org. Chem.* **1993**, 58, 142–146.
- (S7) Moody, C. J.; Taylor, R. J. *J. Chem. Soc., Perkin Trans. 1* **1989**, 721–731.
- (S8) Itooka, R.; Iguchi, Y.; Miyaura, N. *J. Org. Chem.* **2003**, 68, 6000–6004.
- (S9) (a) Yu, M. S.; Lantos, I.; Peng, Z.-Q.; Yu, J.; Cacchio, T. *Tetrahedron Lett.* **2000**, 41, 5647–5651. (b) Liu, L. T.; Hong, P.-C.; Huang, H.-L.; Chen, S.-F.; Wang, C.-L. J.; Wen, Y.-S. *Tetrahedron: Asymmetry* **2001**, 12, 419–426.
- (S10) Pal, B.; Pradhan, P. K.; Jaisankar, P.; Giri, V. S. *Synthesis* **2003**, 1549–1552.
- (S11) Watanabe, T.; Hamaguchi, F.; Ohki, S. *YakugakuZasshi* **1973**, 93, 845–853.
- (S12)(a) Bettoni, G.; Franchini, C.; Morlacchi, F.; Tangari, N.; Tortorella, V. *J. Org. Chem.* **1976**, 41, 2780–2782. (b) Reddy, P. Y.; Kondo, S.; Toru, T.; Ueno, Y. *J. Org. Chem.* **1997**, 62, 2652–2654.
- (S13) Paden, J. H.; Adkins, H. *J. Am. Chem. Soc.* **1936**, 58, 2487–2499.
- (S14) Gill, C. D.; Greenhalgh, D. A.; Simpkins, N. S. *Tetrahedron* **2003**, 59, 9213–9230.
- (S15)(a) Devlin, J. P.; Ollis, W. D.; Thorpe, J. E.; Wood, R. J.; Broughton, B. J.; Warren, P. J.; Wooldridge, K. R. H.; Wright, D. E. *J. Chem. Soc., Perkin Trans. 1* **1975**, 830–841. (b) Aubert, T.; Farnier, M.; Guillard, R. *Tetrahedron* **1991**, 47, 53–60.
- (S16) Wanner, M. J.; Koomen, G.-J. *Tetrahedron* **1991**, 47, 8431–8442.
- (S17) Mederski, W. W. K. R.; Baumgarth, M.; Germann, M.; Kux, D.; Weitzel, T. *Tetrahedron Lett.* **2003**, 44, 2133–2136.
- (S18)(a) Cannizzo, L. F.; Grubbs, R. H. *J. Org. Chem.* **1985**, 50, 2316–2323. (b) Hayashi, T.; Sawamura, M.; Ito, Y. *Tetrahedron* **1992**, 48, 1999–2012.
- (S19) Cameron, D. W.; Down, J. G.; Kissane, P. J.; Laycock, G. M.; Shulman, A. *Aust. J. Chem.* **1977**, 30, 1157–1159

- (S20) Greenlee, W. J.; Thorsett, E. D. *J. Org. Chem.* **1981**, *46*, 5351–5353.
- (S21) Uhle, C. *J. Org. Chem.* **1961**, *26*, 2998–3001.
- (S22) Johnson, C. D.; Lane, S.; Edwards, P. N.; Taylor, P. J. *J. Org. Chem.* **1988**, *53*, 5130–5139.
- (S23) (a) Decker, M.; Nguyen, T. T. H.; Lehmann, J. *Tetrahedron* **2004**, *60*, 4567–4578.
 (b) Conejo-Garcia, A.; Schofield, C. J. *Bioorg. Med. Chem. Lett.* **2005**, *15*, 4004–4009.
- (S24) Evans, D. A.; Fu, G. C.; Hoveyoda, A. H. *J. Am. Chem. Soc.* **1992**, *114*, 6671–6679.
- (S25) (a) Knobler, Y.; Bonni, E.; Sheradsky, T. *J. Org. Chem.* **1964**, *29*, 1229–1235.
 (b) Matsumoto, K.; Hashimoto, S.; Uchida, T.; Okamoto, T.; Otani, S. *Bull. Chem. Soc. Jpn.*, **1989**, *62*, 3138–3142.
- (S26) Senda, T.; Ogasawara, M.; Hayashi, T. *J. Org. Chem.* **2001**, *66*, 6852–6856.
- (S27) *CrystalStructure 3.6: Crystal Structure Analysis Package*, Rigaku and Rigaku/MSO, The Woodlands TX 77381 USA, **2000–2004**.
- (S28) Altomare, A.; Cascarano, G.; Giacovazzo, C.; Guagliardi, A.; Burla, M.; Polidori, G.; Camalli, M. *J. Appl. Cryst.* **1994**, *27*, 435.
- (S29) Flack, H. D. *Acta Crystallogr., Sect. A* **1983**, *39*, 876.
- (S30) *International Tables for X-ray Crystallography*; Kynoch Press: Birmingham, England, 1974; Vol. IV.
- (S31) *International Tables for X-ray Crystallography*; Kluwer Academic Publishers: Boston, MA, 1992; Vol. C.