Chiral Bis- π -allylpalladium Complex Catalyzed Asymmetric Allylation of Imines: Enhancement of Enantioselectivity and Chemical Yield in the Presence of Water

Rodney A. Fernandes, Anton Stimac, and Yoshinori Yamamoto*

Department of Chemistry, Graduate School of Science, Tohoku University,
Sendai 980-8578, Japan

Supporting Information

(32 pages)

General Information. Commercially available imines were procured from Aldrich Chemical Co.; while most were prepared by refluxing equimolar amount of corresponding aldehyde and amine in toluene or benzene with Dean-Stark removal of water. They were then purified by either high vacuum distillation or recrystallization. Most commercially supplied chemicals were freshly distilled or recrystallized before use. Solvents were dried by standard procedures. Dry THF refers to commercial dehydrated THF procured from Kanto Chem. Co., Inc, Japan. ^1H and ^{13}C NMR spectra were recorded on a Jeol JNM LA-300 spectrometer. The chemical shifts are reported in δ units relative to internal tetramethylsilane for ^1H NMR and CDCl_3 was used as internal standard ($\delta = 77.0$) for ^{13}C NMR. IR spectra were recorded on a Shimadzu FTIR-8200 A spectrometer. Mass spectra were recorded on Jeol JMS-HX 110 in EI mode. Optical rotations were measured on a Jasco DIP-1000 digital polarimeter. HPLC was performed on Shimadzu LC9A using Daicel CHIRALCEL OD or CHIRALCEL OD-R columns (0.46 cm x 25 cm). The absolute configuration was determined by converting amine **13a** into 1-phenylbutylamine which exhibited optical rotation with + value. The literature value of (*S*)-1-phenylbutylamine is $[\alpha]_{\text{D}} -21.3$ (c 1.3, CHCl_3)¹. Hence the absolute configuration of **13a** is confirmed to be (*R*). Thus, by analogy the homoallylamines obtained are assumed to have (*R*) configuration² (unless otherwise indicated).

Preparation of the catalyst, **2a**

(1*R*)-(+)-nopinone is available in Aldrich Chemical Co. in 90% ee. But its precursor (1*S*)-(-)- β -pinene is available in 97% ee. So we preferred to prepare (1*R*)-(+)-nopinone from (1*S*)-(-)- β -pinene. The ozonolysis of exocyclic double bond of (1*S*)-(-)- β -pinene by known procedure³ afforded (1*R*)-(+)-nopinone in 75% yield, $[\alpha]_{\text{D}}^{22} + 39.4$ (c =1, MeOH). Wittig reaction with ethyltriphenylphosphonium bromide and $t\text{BuOK}$ by known procedure⁴ gave the olefin **7** (*E*:*Z* = 1:1) in 83% yield.

From olefin **7** we followed the literature procedure⁵ to prepare the **2a:2b** mixture. To a solution of Pd(OCOCF₃)₂ (5 g, 15.04 mmol) in dry acetone (125 mL) in a two necked round bottom flask under argon atmosphere was added olefin **7** (2.26 g, 15.04 mmol). The reaction mixture was stirred for 1 h at room temperature (TLC monitored). ⁿBu₄NCl (4.56 g, 16.55 mmol) in acetone (30 mL) was added and the reaction mixture was stirred for 1 h. The clear brown-orange solution was then filtered through a plug of celite to remove suspended Pd-black. The filtrate was concentrated to thick amber colored oil. This was purified by silica gel column chromatography using hexane/EtOAc = 5:1 as eluent to give **2a/2b** mixture as semi solid (6.92 g). ¹H NMR indicated **2a:2b** = 1.3:1. This was dissolved in minimum CH₂Cl₂ and hexane (30 mL) was added. The crystallized solid was filtered through sintered funnel, washed with hexane and dried with suction to afford **2a:2b** = 4.8:1 (3.2 g) as yellow powder (**I**). Further upgradation is as shown in the flow chart below (**II-V**). During propionitrile recrystallization boiling of solvent to dissolve the solid should be avoided to prevent catalyst decomposition. To recover the catalyst the filtrates after each recrystallization **F1-F3** as shown below were combined and concentrated to yellow colored solid foam. This was subjected to benzene column chromatography. Elution with benzene gave in first few fractions the catalyst **2b** (0.45 g) having [α]_D²² +47.5 (c = 0.4, CHCl₃). The later fractions were mixture of **2a** and **2b**. The filtrates **F4** and **F5** were combined and concentrated to yellow powder (1 g). This on successive three recrystallizations in propionitrile gave **2a:2b** = 100:1 (0.452 g), [α]_D²² -17.2 (c = 0.4, CHCl₃). Further three repeated recrystallizations in propionitrile afforded **2a:2b** >400:1 (0.215 g), [α]_D²² -19.8 (c = 0.4, CHCl₃).

Recovery of the catalyst

Filtrates **F1 - F3** → Combined and subjected to benzene column separation
2b ligand was separated (0.45 g), $[\alpha]_D^{22} + 47.5$ (c = 0.4, CHCl_3)

Filtrates **F4 and F5** [1.0 g] → Three repeated recrystallizations in EtCN
 Yellow needles, 0.452 g [**2a:2b** = 100:1]
 $[\alpha]_D^{22} = -17.2$ (c = 0.4, CHCl_3)

↓ Three repeated
 recrystallizations in EtCN

Yellow needles, 0.215 g [2a:2b** = >400:1]**
 $[\alpha]_D^{22} = -19.8$ (c = 0.4, CHCl_3)

^1H NMR of **2a:2b** = 1.3:1

^1H NMR of 2a:2b = 100:1

^1H NMR of 2a:2b = >400:1

2a: (CDCl_3) δ 3.84–3.80 (m, 1H), 3.78–3.68 (m, 1H), 2.7–2.64 (m, 1H), 2.55 (brt, $J = 5.5$ Hz, 1H), 2.44–2.33 (m, 1H), 2.08–2.03 (m, 1H), 1.77–1.74 (m, 2H), 1.34 (s, 3H), 1.13 (d, $J = 6.5$ Hz, 3H), 0.91 (s, 3H).

¹³C NMR 2a: (CDCl₃) δ 136.0, 72.2, 71.0, 40.5, 39.9, 37.8, 33.8, 29.9, 26.1, 21.7, 13.9.

IR (KBr) 2982, 2951, 2912, 2870, 1499, 1466, 1448, 1429, 1367, 1265, 1219, 1101, 1051, 1032, 974, 860, 758 cm^{-1} .

^1H NMR of **2b**: (CDCl_3) δ 4.47–4.41 (m, 1H), 4.33–4.22 (m, 1H), 2.67–2.52 (m, 1H), 2.36–2.3 (m, 1H), 2.2–2.13 (m, 1H), 2.08–2.02 (m, 1H), 1.87–1.74 (m, 2H), 1.29 (s, 3H), 1.03 (d, $J = 6.8$ Hz, 3H), 0.71 (s, 3H).

^{13}C NMR 2b: (CDCl_3) δ 132.5, 72.7, 70.8, 46.5, 39.9, 38.0, 34.4, 29.9, 26.0, 21.7, 16.2.

IR (KBr) 2980, 2918, 2872, 2833, 1468, 1454, 1429, 1369, 1265, 1221, 1099, 1042, 968, 920, 762 cm^{-1} .

General equation for allylation of imines with allyltributylstannane and catalyst 2a

Table. Catalytic asymmetric allylation of imines with allyltributylstannane in the presence of the catalyst**2a^a**

entry	substrate	product	time (h)	% yield	% ee
	12	13			
1	12a R ₁ = R ₂ = R ₃ = H	13a	48	86	85
2	12b R ₁ = R ₃ = H, R ₂ = OMe	13b	98	82	85
3	12c R ₁ = R ₃ = H, R ₂ = Me	13c	67	92	90
4	12d R ₁ = R ₂ = H, R ₃ = OMe	13d	73	78	88
5	12e R ₁ = R ₂ = OMe, R ₃ = H	13e	116	76	70
6	14	15	74	84	50
7	16	17	72	84	69
	18	19			
8	18a X = O	19a	73	90	67
9	18b X = S	19b	90	90	53
	10	11			
10	10a R ₁ = R ₂ = H, R ₃ = OMe	11a	73	89	90
11	10b R ₁ = R ₃ = OMe, R ₂ = H	11b	88	78	85
12	10c R ₁ = Me, R ₂ = H, R ₃ = OMe	11c	72	81	86
13	10d R ₁ = NO ₂ , R ₂ = H, R ₃ = OMe	11d	70	94	42
14	10e R ₁ = R ₃ = H, R ₂ = OMe	11e	78	82	89
	20	21			

15	20a R ₁ = H	21a	96	70	91
16	20b R ₁ = OMe	21b	74	81	82

17	22a R = allyl	23a	78	82	78
18	22b R = SO ₂ Ph	23b	144	<10 ^b	---
19			168	42	55
20			120	75	34

21	8a X = R ₁ = R ₂ = R ₃ = H	9a	84	76	90
22	8b X = R ₁ = R ₂ = H, R ₃ = OMe	9b	72	76	89
23	8c X = R ₁ = R ₃ = H, R ₂ = OMe	9c	79	68	89
24	8d X = Cl, R ₁ = R ₂ = R ₃ = H	9d	89	84	87
25	8e X = Br, R ₁ = R ₂ = R ₃ = H	9e	99	87	88
26	8f X = R ₂ = R ₃ = H, R ₁ = Ph	9f	126	30	83
27			20	38	2
		20	63	18 ^c
28			41	77	7 ^d
		43	73	19 ^{c,d}

^aAll reactions were carried out with 1.25 equivalent of allyltributylstannane and one equivalent of water in the presence of 5 mol% of **2a** and at 0°C for the specified time. In some cases additional 0.5 equivalent of allyltributylstannane was added if it was completely consumed and the reaction was not complete (checked by GC-MS); ^bdetermined by ¹H NMR of the crude reaction mixture with CH₃NO₂ as internal standard. ^cusing catalyst **2b**. ^dee based on known [α]_D values (see reference 6)

N-Benzyl-1-phenyl-3-butenylamine, 13a

Colorless oil; $[\alpha]_D^{23} = +51.2$ (c 1.45, CHCl_3); IR, ^1H NMR, ^{13}C NMR, HRMS and analysis data (see ref. 2). The enantiomeric excess was determined by using CHIRALCEL OD-R column, $\text{CH}_3\text{CN}/1\text{M NaClO}_4$ aq = 40/60, flow rate = 0.8 mL/min, UV detection at 254 nm, $t_R = 16.38$ min (major enantiomer), $t_R = 18.48$ min (minor enantiomer); 85% ee.

N-Benzyl-1-(4-methoxyphenyl)-3-butenylamine, 13b

Colorless oil; $[\alpha]_D^{22} = +56.1$ (c 0.745, CHCl_3); IR, ^1H NMR, ^{13}C NMR, HRMS and analysis data (see ref. 2). The enantiomeric excess was determined by using CHIRALCEL OD-R column, $\text{CH}_3\text{CN}/1\text{M NaClO}_4$ aq = 40/60, flow rate = 0.8 mL/min, UV detection at 254 nm, $t_R = 18.02$ min (major enantiomer), $t_R = 21.5$ min (minor enantiomer); 85% ee.

N-Benzyl-1-(4-methylphenyl)-3-butenylamine, 13c

Colorless oil; $[\alpha]_D^{22} = +50.4$ (c 1.0, CHCl_3); IR (neat) 3325, 3061, 3025, 2977, 2921, 1638, 1603, 1513, 1495, 1454, 1350, 1304, 1176, 1029, 995, 916, 816, 735 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.28–7.13 (m, 7H), 7.09 (d, $J = 8$ Hz, 2H), 5.6 (ddt, $J = 17.1$ Hz, 10.3 Hz, 7.1 Hz, 1H), 5.02 (ddt, $J = 17.1$ Hz, 2 Hz, 2 Hz, 1H), 4.97 (ddt, $J = 10.3$ Hz, 2 Hz, 2 Hz, 1H), 3.61–3.59 (m, 1H), 3.6 (d, $J = 13.4$ Hz, 1H), 3.45 (d, $J = 13.4$ Hz, 1H), 2.34–2.32 (m, 2H), 2.3 (s, 3H), 1.66 (brs, 1H);

^{13}C NMR (CDCl_3) δ 140.7, 136.5, 135.5, 129.0, 128.2, 128.0, 127.2, 126.7, 117.4, 61.2, 51.3, 43.1, 21.0; MS (EI) m/z (relative intensity) 210 [M^+ -allyl] (5.7), 132 (0.7), 118 (1.2), 105 (2.9), 91 (100); Anal. Calcd for $\text{C}_{18}\text{H}_{21}\text{N}$ (251.37): C, 86.00; H, 8.42; N, 5.57. Found: C, 85.82; H, 8.66; N, 5.52. The enantiomeric excess was determined using CHIRALCEL OD-R column, $\text{CH}_3\text{CN}/1\text{M NaClO}_4$ aq = 40/60, flow rate = 0.8 mL/min, UV detection at 254 nm, t_R = 22.82 min (major enantiomer), t_R = 27.20 min (minor enantiomer); 90% ee.

N-Benzyl-1-(2-methoxyphenyl)-3-butenylamine, 13d

Colorless oil; $[\alpha]_D^{22} = +37.2$ (c 1.0, CHCl_3); IR (neat) 3330, 3063, 3027, 3001, 2935, 2835, 1638, 1600, 1586, 1489, 1437, 1363, 1281, 1239, 1173, 1095, 1050, 1029, 996, 914, 754 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.38–7.35 (m, 1H), 7.26–7.16 (m, 6H), 6.95–6.83 (m, 2H), 5.72 (ddt, J = 16.1 Hz, 10.2 Hz, 6.7 Hz, 1H), 5.03–4.93 (m, 2H), 4.07–4.03 (m, 1H), 3.77 (s, 3H), 3.65 (d, J = 13 Hz, 1H), 3.51 (d, J = 13 Hz, 1H), 2.5–2.34 (m, 2H), 1.8 (brs, 1H);

N-Benzyl-1-(3,4-dimethoxyphenyl)-3-butenylamine, 13e

Colorless oil; [α]_D²³ = +38.6 (c 1.75, CHCl₃); IR (neat) 3325, 3062, 2998, 2932, 2833, 1638, 1593, 1514, 1463, 1417, 1358, 1262, 1233, 1139, 1029, 916, 856, 808, 747 cm⁻¹; ¹H NMR (CDCl₃) δ 7.32–7.19 (m, 5H), 6.95–6.81 (m, 3H), 5.72 (ddt, *J* = 16.2 Hz, 10.2 Hz, 7.2 Hz, 1H), 5.06 (ddt, *J* = 16.2 Hz, 2 Hz, 2 Hz, 1H), 5.03 (ddt, *J* = 10.2 Hz, 2 Hz, 2 Hz, 1H), 3.9 (s, 3H), 3.88 (s, 3H), 3.69 (d, *J* = 13.2 Hz, 1H), 3.63–3.59 (m, 1H), 3.54 (d, *J* = 13.2 Hz, 1H), 2.41–2.34 (m, 2H), 1.81 (brs, 1H);

^{13}C NMR (CDCl_3) δ 149.0, 147.9, 140.6, 136.3, 135.5, 128.3, 128.1, 126.8, 119.5, 117.5, 110.8, 109.9, 61.2, 55.8, 51.3, 43.2; MS (EI) m/z (relative intensity) 256 [M^+ -allyl] (14.8), 239 (5), 214 (9.4), 151 (20.8), 91 (100); Anal. Calcd for $\text{C}_{19}\text{H}_{23}\text{NO}_2$ (297.39): C, 76.73; H, 7.79; N, 4.71. Found: C, 76.68; H, 8.06; N, 4.72. The enantiomeric excess was determined by using CHIRALCEL OD-R column, $\text{CH}_3\text{CN}/1\text{M NaClO}_4 \text{ aq} = 30/70$, flow rate = 0.8 mL/min, UV detection at 254 nm, $t_R = 28.57$ min (major enantiomer), $t_R = 32.58$ min (minor enantiomer); 70% ee.

N-Benzyl-1-cyclohexyl-3-butenylamine, 15

Colorless oil; $[\alpha]_D^{21} = -5.4$ (c 0.5, CHCl_3); IR, ^1H NMR, ^{13}C NMR, HRMS and analysis data (see ref. 2). The enantiomeric excess was determined by converting to trifluoroacetylamine form, $[\alpha]_D^{22} = -7.8$ (c 0.6, CHCl_3) and using CHIRALCEL OD column, hexane/*i*-PrOH = 800/1, flow rate = 0.6 mL/min, UV detection at 254 nm, $t_R = 26.04$ min (minor enantiomer), $t_R = 30.41$ min (major enantiomer); 50% ee.

N-Benzyl-1-phenyl-1,5-hexadien-3-ylamine, 17

Colorless oil; $[\alpha]_D^{23} = +77.2$ (c 1.0, CHCl_3); IR, ^1H NMR, ^{13}C NMR, HRMS and analysis data (see ref. 2). The

enantiomeric excess was determined using CHIRALCEL OD-R column, CH₃CN/1M NaClO₄ aq = 60/40, flow rate = 0.6 mL/min, UV detection at 254 nm, *t_R* = 10.03 min (major enantiomer), *t_R* = 12.17 min (minor enantiomer); 69% ee.

N-Benzyl-1-(2-furfuryl)-3-butenylamine, 19a

Colorless oil; $[\alpha]_D^{22} = +55.9$ (c 1.0, CHCl₃); IR (neat) 3330, 3063, 3028, 2978, 2914, 2837, 1640, 1604, 1512, 1495, 1455, 1346, 1248, 1176, 1148, 1108, 1073, 1028, 918, 884, 807, 734 cm⁻¹; ¹H NMR (CDCl₃) δ 7.31–7.13 (m, 6H), 6.25 (dd, *J* = 3.2 Hz, 2 Hz, 1H), 6.11 (d, *J* = 3.2 Hz, 1H), 5.68 (ddt, *J* = 17.1 Hz, 10.2 Hz, 6.6 Hz, 1H), 5.05 (ddt, *J* = 17.1 Hz, 2.2 Hz, 2 Hz, 1H), 5.03 (ddt, *J* = 10.2 Hz, 2.2 Hz, 2 Hz, 1H), 3.72–3.70 (m, 1H), 3.69 (d, *J* = 13.2 Hz, 1H), 3.54 (d, *J* = 13.2 Hz, 1H), 2.46 (dd, *J* = 6.6 Hz, 1.3 Hz, 2H), 1.65 (brs, 1H);

¹³C NMR (CDCl₃) δ 156.0, 141.4, 140.1, 134.8, 128.2, 128.0, 126.7, 117.5, 109.7, 106.5, 54.7, 50.93, 39.2; MS (EI) *m/z* (relative intensity) 186 [*M*⁺–allyl] (33.9), 131 (1.5), 95 (1.8), 91 (100); Anal. Calcd for C₁₅H₁₇NO (227.3): C, 79.26; H, 7.53; N, 6.16. Found: C, 79.11; H, 7.84; N, 6.15. The enantiomeric excess was determined by converting to trifluoroacetylamine form, $[\alpha]_D^{22} = +54.7$ (c 1.0, CHCl₃) and using CHIRALCEL OD column, hexane/*i*-PrOH = 500/1, flow rate = 0.5 mL/min, UV detection at 254 nm, *t_R* = 22.18 min (major enantiomer), *t_R* = 25.2 min (minor enantiomer); 67% ee.

N-Benzyl-1-(2-thiophenyl)-3-butenylamine, 19b

Colorless oil; $[\alpha]_D^{22} = +24.5$ (c 1.0, CHCl_3); IR (neat) 3325, 3064, 3027, 2977, 2910, 2835, 1639, 1603, 1495, 1454, 1434, 1368, 1225, 1168, 1029, 995, 917, 850, 735 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.27–7.14 (m, 6H), 6.9–6.84 (m, 2H), 5.69 (ddt, $J = 16.1$ Hz, 10.0 Hz, 6.9 Hz, 1H), 5.05 (ddt, $J = 16.1$ Hz, 2.1 Hz, 2 Hz, 1H), 5.0 (ddt, $J = 10.0$ Hz, 2.1 Hz, 2 Hz, 1H), 3.92 (t, $J = 6.7$ Hz, 1H), 3.74 (d, $J = 13.2$ Hz, 1H), 3.55 (d, $J = 13.2$ Hz, 1H), 2.44–2.34 (m, 2H), 1.73 (brs, 1H);

^{13}C NMR (CDCl_3) δ 149.2, 140.2, 134.8, 128.3, 128.2, 126.9, 126.3, 124.2, 123.9, 117.9, 57.0; MS (EI) m/z (relative intensity) 202 [M^+ -allyl] (18.9), 111 (2), 106 (1.3), 97 (7.9), 91 (100); Anal. Calcd for $\text{C}_{15}\text{H}_{17}\text{NS}$ (243.37): C, 74.03; H, 7.04; N, 5.75; S, 13.17. Found: C, 73.96; H, 7.32; N, 5.66; S, 13.18. The enantiomeric excess was determined by converting to trifluoroacetylamine form, $[\alpha]_D^{22} = +43.4$ (c 1.0, CHCl_3) and using CHIRALCEL OD column, hexane/*i*-PrOH = 500/1, flow rate = 0.5 mL/min, UV detection at 254 nm, $t_R = 27.55$ min (major enantiomer), $t_R = 30.59$ min (minor enantiomer); 53% ee.

N-(4-Methoxybenzyl)-1-phenyl-3-butenylamine, 11a

Colorless oil; $[\alpha]_D^{23} = +49.1$ (c 1.0, CHCl_3); IR, ^1H NMR, ^{13}C NMR, HRMS and analysis data (see ref. 2). The enantiomeric excess was determined by converting to trifluoroacetamide form, $[\alpha]_D^{22} = +75.0$ (c 1.75, CHCl_3) and using CHIRALCEL OD column, hexane/*i*-PrOH = 100/1, flow rate = 0.6 mL/min, UV detection at 254 nm, $t_R = 17.02$ min (major enantiomer), $t_R = 20.19$ min (minor enantiomer); 90% ee.

N-(4-Methoxybenzyl)-1-(4-methoxyphenyl)-3-butenylamine, 11b

Colorless oil; $[\alpha]_D^{22} = +48.3$ (c 1.0, CHCl_3); IR (neat) 3325, 3072, 2998, 2932, 2907, 2834, 2059, 1638, 1611, 1585, 1512, 1463, 1351, 1301, 1245, 1174, 1106, 1036, 997, 917, 831, 781, 756 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.26 (d, $J = 8.6$ Hz, 2H), 7.15 (d, $J = 8.6$ Hz, 2H), 6.89 (d, $J = 8.6$ Hz, 2H), 6.84 (d, $J = 8.6$ Hz, 2H), 5.7 (ddt, $J = 16.2$ Hz, 10.1 Hz, 6.5 Hz, 1H), 5.07–4.9 (m, 2H), 3.8 (s, 3H), 3.78 (s, 3H), 3.63–3.61 (m, 1H), 3.58 (d, $J = 13$ Hz, 1H), 3.44 (d, $J = 13$ Hz, 1H), 2.38–2.33 (m, 2H), 1.69 (brs, 1H);

^{13}C NMR (CDCl_3) δ 158.5, 135.7, 135.6, 132.7, 129.2, 128.2, 117.3, 113.6, 60.7, 55.2, 55.1, 50.6, 43.1; MS (EI) m/z (relative intensity) 256 [M^+ -allyl] (9.4), 160 (6), 122 (8.8), 121 (100), 91 (6.7), 77 (2.9); Anal. Calcd for $\text{C}_{19}\text{H}_{23}\text{NO}_2$ (297.39): C, 76.73; H, 7.79; N, 4.71. Found: C, 76.6; H, 7.98; N, 4.68. The enantiomeric excess was determined using CHIRALCEL OD-R column, $\text{CH}_3\text{CN}/1\text{M NaClO}_4$ aq = 35/65, flow rate = 0.8 mL/min, UV detection at 254 nm, $t_R = 31.9$ min (major enantiomer), $t_R = 36.22$ min (minor enantiomer); 85% ee.

N-(4-Methoxybenzyl)-1-(4-methylphenyl)-3-butenylamine, 11c

Colorless oil; $[\alpha]_D^{22} = +48.5$ (c 1.0, CHCl_3); IR (neat) 3325, 3002, 2910, 2833, 1638, 1612, 1584, 1512, 1463, 1349, 1301, 1247, 1173, 1105, 1037, 996, 917, 817, 781, 757 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.32 (d, $J = 8.5$ Hz, 2H), 7.24 (d, $J = 8$ Hz, 4H), 6.92 (d, $J = 8.5$ Hz, 2H), 5.78 (ddt, $J = 16$ Hz, 10.2 Hz, 6.8 Hz, 1H), 5.07–5.15 (m, 2H), 3.85 (s, 3H), 3.72–3.7 (m, 1H), 3.69 (d, $J = 13$ Hz, 1H), 3.53 (d, $J = 13$ Hz, 1H), 2.47–2.45 (m, 2H), 2.42 (s, 3H), 1.73 (brs, 1H);

^{13}C NMR (CDCl_3) δ 158.4, 140.7, 136.4, 135.6, 132.7, 129.2, 129.0, 127.1, 117.4, 113.6, 61.0, 55.1, 50.6, 43.1, 21.0; MS (EI) m/z (relative intensity) 240 [M^+ -allyl] (13.3), 160 (4.9), 121 (100), 119 (4.9), 91 (12.4), 77 (5.3); Anal. Calcd for $\text{C}_{19}\text{H}_{23}\text{NO}$ (281.39): C, 81.10; H, 8.24; N, 4.97. Found: C, 81.02; H, 8.47; N, 4.97. The enantiomeric excess was determined using CHIRALCEL OD-R column, $\text{CH}_3\text{CN}/1\text{M NaClO}_4$ aq = 35/65, flow rate = 0.8 mL/min, UV detection at 254 nm, $t_R = 41.41$ min (major enantiomer), $t_R = 46.64$ min (minor enantiomer); 86% ee.

N-(4-Methoxybenzyl)-1-(4-nitrophenyl)-3-butenylamine, 11d

Pale yellow oil; $[\alpha]_D^{22} = +26.9$ (c 1.0, CHCl_3); IR (neat) 3331, 3075, 3001, 2933, 2835, 1639, 1609, 1515, 1463, 1346, 1301, 1247, 1176, 1108, 1035, 1013, 996, 920, 856, 825, 753 cm^{-1} ; ^1H NMR (CDCl_3) δ 8.14 (d, $J = 8.6\text{ Hz}$, 2H), 7.48 (d, $J = 8.6\text{ Hz}$, 2H), 7.07 (d, $J = 8.6\text{ Hz}$, 2H), 6.78 (d, $J = 8.6\text{ Hz}$, 2H), 5.62 (ddt, $J = 16.4\text{ Hz}$, 10.1 Hz, 6.4 Hz, 1H), 5.52–4.96 (m, 2H), 3.74–3.73 (m, 1H), 3.72 (s, 3H), 3.52 (d, $J = 13\text{ Hz}$, 1H), 3.38 (d, $J = 13\text{ Hz}$, 1H), 2.37–2.21 (m, 2H), 1.7 (brs, 1H);

^{13}C NMR (CDCl_3) δ 158.6, 151.8, 147.0, 134.2, 131.9, 129.1, 128.0, 123.6, 118.5, 113.7, 60.9, 55.2, 50.9, 42.8; MS (EI) m/z (relative intensity) 271 [M^+ -allyl] (2), 186 (14.6), 150 (4.3), 121 (44.6), 91 (100), 81 (10.2), 77 (11.8); Anal. Calcd for $\text{C}_{18}\text{H}_{20}\text{N}_2\text{O}_3$ (312.37): C, 69.21; H, 6.45; N, 8.96. Found: C, 68.98; H, 6.73; N, 8.8. The enantiomeric excess was determined by converting to trifluoroacetylamine form, $[\alpha]_D^{22} = +30.6$ (c 1.5, CHCl_3) and using CHIRALCEL OD column, hexane/*i*-PrOH = 50/1, flow rate = 0.8 mL/min, UV detection at 254 nm, $t_R = 27.43\text{ min}$ (minor enantiomer), $t_R = 29.75\text{ min}$ (major enantiomer); 42% ee.

N-(2-Methoxybenzyl)-1-phenyl-3-butenylamine, 11e

Colorless oil; $[\alpha]_D^{22} = +51.6$ (c 1.0, CHCl_3); IR (neat) 3343, 3062, 3026, 3001, 2934, 2835, 1638, 1601, 1588, 1492, 1463, 1358, 1288, 1241, 1175, 1117, 1048, 1029, 955, 916, 836, 754, 702 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.34–7.18 (m, 6H), 7.1–7.05 (m, 1H), 6.89–6.82 (m, 2H), 5.66 (ddt, $J = 16.8$ Hz, 10.1 Hz, 6.8 Hz, 1H), 5.07 (ddt, $J = 16.8$ Hz, 2.2 Hz, 2 Hz, 1H), 5.02 (ddt, $J = 10.1$ Hz, 2.2 Hz, 2 Hz, 1H), 3.8 (s, 3H), 3.72 (d, $J = 13.4$ Hz, 1H), 3.63–3.6 (m, 1H), 3.58 (d, $J = 13.4$ Hz, 1H), 2.39–2.35 (m, 2H), 2.12–1.91 (brs, 1H);

^{13}C NMR (CDCl_3) δ 157.6, 143.8, 135.5, 129.8, 128.2, 128.1, 128.0, 127.3, 126.8, 120.1, 117.3, 110.0, 60.9, 54.9, 47.1, 43.1; MS (EI) m/z (relative intensity) 226 [M^+ -allyl] (11.9), 160 (3.9), 122 (9.2), 121 (100), 105 (2.3), 94 (3.1), 93 (37.6), 91 (7.2), 79 (1.3), 77 (1); Anal. Calcd for $\text{C}_{18}\text{H}_{21}\text{NO}$ (267.37): C, 80.86; H, 7.91; N, 5.23. Found: C, 80.83; H, 8.18; N, 5.2. The enantiomeric excess was determined using CHIRALCEL OD-R column, $\text{CH}_3\text{CN}/1\text{M NaClO}_4$ aq = 50/50, flow rate = 0.8 mL/min, UV detection at 254 nm, $t_R = 10.03$ min (minor enantiomer), $t_R = 12.66$ min (major enantiomer); 89% ee.

N-Benzyl-1-(2-naphthyl)-3-butenylamine, 21a

Colorless oil; $[\alpha]_D^{21} = +64.0$ (c 1.5, CHCl_3); IR, ^1H NMR, ^{13}C NMR, HRMS and analysis data (see ref. 2). The enantiomeric excess was determined by using CHIRALCEL OD-R column, $\text{CH}_3\text{CN}/1\text{M NaClO}_4$ aq = 40/60, flow rate = 0.8 mL/min, UV detection at 254 nm, $t_R = 58.3$ min (major enantiomer), $t_R = 67.45$ min (minor enantiomer); 91% ee.

N-(4-Methoxybenzyl)-1-(2-naphthyl)-3-butenylamine, 21b

Colorless oil; $[\alpha]_D^{22} = +54.3$ (c 1.0, CHCl_3); IR (neat) 3330, 3076, 3026, 2978, 2911, 2833, 1640, 1602, 1492, 1454, 1415, 1357, 1115, 1070, 1027, 994, 759 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.95–7.86 (m, 4H), 7.63–7.51 (m, 3H), 7.26 (d, $J = 8.6$ Hz, 2H), 6.94 (d, $J = 8.6$ Hz, 2H), 5.85 (ddt, $J = 16.4$ Hz, 10.2 Hz, 6.9 Hz, 1H), 5.2–5.1 (m, 2H), 3.95–3.91 (m, 1H), 3.87 (s, 3H), 3.74 (d, $J = 13$ Hz, 1H), 3.5 (d, $J = 13$ Hz, 1H), 2.62–2.52 (m, 2H), 1.87 (brs, 1H);

^{13}C NMR (CDCl_3) δ 158.4, 141.1, 135.3, 133.3, 132.7, 132.5, 129.1, 128.0, 127.6, 126.1, 125.8, 125.3, 125.2, 117.5, 113.6, 61.4, 55.0, 50.6, 42.8; MS (EI) m/z (relative intensity) 276 [M^+ -allyl] (7.7), 160 (6.6), 122 (8.7), 121 (100), 91 (4.8), 77 (2.4); Anal. Calcd for $\text{C}_{22}\text{H}_{23}\text{NO}$ (317.43): C, 83.24; H, 7.3; N, 4.41. Found: C, 83.19; H, 7.49; N, 4.58. The enantiomeric excess was determined using CHIRALCEL OD-R column, $\text{CH}_3\text{CN}/1\text{M NaClO}_4$ aq = 50/50, flow rate = 0.7 mL/min, UV detection at 254 nm, t_R = 38.16 min (major enantiomer), t_R = 42.87 min (minor enantiomer); 82% ee.

N-allyl-1-phenyl-3-butenylamine, 23a

Colorless oil; $[\alpha]_D^{23} = +24.8$ (c 1.0, CHCl_3); IR (neat) 3329, 3077, 3026, 3003, 2977, 2910, 2833, 1640, 1602, 1542, 1492, 1454, 1357, 1307, 1235, 1199, 1069, 1027, 994, 917, 758 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.35–7.23 (m, 5H), 5.86–5.73 (m, 1H), 5.73–5.7 (m, 1H), 5.14–5.03 (m, 4H), 3.72 (dd, $J = 6.2$ Hz, 1.2 Hz, 1H), 3.11–3.02 (m, 2H), 2.44–2.39 (m, 2H), 1.71 (brs, 1H);

^{13}C NMR (CDCl_3) δ 143.4, 136.4, 135.2, 128.5, 126.9, 126.7, 117.3, 115.4, 61.4, 49.7, 42.8; MS (EI) m/z (relative intensity) 188 [M^++1] (0.9), 146 [M^+ -allyl] (16.1), 129 (14.9), 119 (2.6), 104 (4.8), 91 (100), 77 (2.1); Anal. Calcd for $\text{C}_{13}\text{H}_{17}\text{N}$ (187.28): C, 83.37; H, 9.14; N, 7.47. Found: C, 83.07; H, 9.35; N, 7.4. The enantiomeric excess was determined by converting to trifluoroacetylamine form, $[\alpha]_D^{22} = +93.2$ (c 1.0, CHCl_3)

and using CHIRALCEL OD column, hexane/*i*-PrOH = 300/1, flow rate = 0.6 mL/min, UV detection at 254 nm, t_R = 16.17 min (major enantiomer), t_R = 17.85 min (minor enantiomer); 78% ee.

N-Methyl-1-(3-pyridyl)-3-butenylamine, 25

Yellow liquid; $[\alpha]_D^{22} = +43.1$ (c 1.66, CHCl_3); IR (neat) 3324, 3082, 2979, 2936, 1666, 1640, 1578, 1476, 1428, 1098, 1027, 996, 734 cm^{-1} ; ^1H NMR (CDCl_3) δ 8.45–8.41 (m, 2H), 7.62–7.58 (m, 1H), 7.23–7.2 (m, 1H), 5.69–5.59 (m, 1H), 5.08–5.01 (m, 2H), 3.64–3.60 (m, 1H), 3.51 (t, J = 6.8 Hz, 1H), 2.47–2.4 (m, 2H), 2.2 (s, 3H);

^{13}C NMR (CDCl_3) δ 149.2, 148.5, 138.6, 134.7, 134.3, 123.5, 118.3, 62.1, 42.4, 34.7; MS (EI) m/z (relative intensity) 163 [$\text{M}^+ + \text{H}$] (3.5), 161 [$\text{M}^+ - \text{H}$] (6.3), 121 [$\text{M}^+ - \text{allyl}$] (100), 119 (9.6), 94 (92), 80 (18.7), 78 (9.4); Anal. Calcd for $\text{C}_{10}\text{H}_{14}\text{N}_2$ (162.23): C, 74.04; H, 8.69; N, 17.27. Found: C, 74.24; H, 8.85; N, 16.94. The enantiomeric excess was determined by converting to trifluoroacetylamine form, $[\alpha]_D^{22} = +90.5$ (c 0.5, CHCl_3) and using CHIRALCEL OD column, hexane/*i*-PrOH = 10/1, flow rate = 0.7 mL/min, UV detection at 254 nm, t_R = 24.13 min (minor enantiomer), t_R = 32.58 min (major enantiomer); 55% ee.

N-Benzyl-1-(3,4,5-trimethoxyphenyl)-3-butenylamine, 27

Colorless syrupy liquid; $[\alpha]_D^{22} = +15.3$ (c 1.0, CHCl_3); IR (neat) 3325, 3062, 2997, 2936, 2835, 1638, 1590, 1505, 1462, 1419, 1325, 1233, 1183, 1128, 1009, 917, 831, 739 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.34–7.21 (m, 5H), 6.6 (s, 2H), 5.73 (ddt, $J = 16.6$ Hz, 10.0 Hz, 6.9 Hz, 1H), 5.13–5.05 (m, 2H), 3.87 (s, 6H), 3.85 (s, 3H), 3.71 (d, $J = 13.2$ Hz, 1H), 3.63–3.59 (m, 1H), 3.56 (d, $J = 13.2$ Hz, 1H), 2.45–2.31 (m, 2H), 1.81 (brs, 1H);

^{13}C NMR (CDCl_3) δ 153.2, 140.5, 139.6, 136.6, 135.4, 128.3, 128.1, 126.8, 117.7, 103.8, 61.8, 60.8, 56.0, 51.4, 43.3; MS (EI) m/z (relative intensity) 286 [M^+ -allyl] (14.9), 254 (8.4), 238 (23.2), 209 (6.1), 195 (6.9), 181 (13.8), 169 (4), 125 (1.2), 91 (100); Anal. Calcd for $\text{C}_{20}\text{H}_{25}\text{NO}_3$ (327.42): C, 73.36; H, 7.69; N, 4.27. Found: C, 73.12; H, 7.89; N, 4.57. The enantiomeric excess was determined using CHIRALCEL OD-R column, $\text{CH}_3\text{CN}/1\text{M NaClO}_4$ aq = 30/70, flow rate = 0.8 mL/min, UV detection at 254 nm, $t_R = 31.15$ min (major enantiomer), $t_R = 34.64$ min (minor enantiomer); 34% ee.

N-Benzyl-1-piperonyl-3-butenylamine, 9a

Colorless oil; $[\alpha]_D^{22} = +59.1$ (c 1.0, CHCl_3); IR (neat) 3327, 3063, 3026, 2976, 2894, 1638, 1607, 1468, 1439, 1376, 1243, 1183, 1100, 1040, 996, 865, 736 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.32–7.21 (m, 5H), 6.91 (s, 1H), 6.77 (s, 2H), 5.95 (s, 2H), 5.69 (ddt, $J = 17.0$ Hz, 10.2 Hz, 7.2 Hz, 1H), 5.09 (ddt, $J = 17.0$ Hz, 2.2 Hz, 2 Hz, 1H), 5.03 (ddt, $J = 10.2$ Hz, 2.2 Hz, 2 Hz, 1H), 3.66 (d, $J = 13.2$ Hz, 1H), 3.6 (t, $J = 6.8$ Hz, 1H), 3.51 (d, $J = 13.2$ Hz, 1H), 2.38–2.28 (m, 2H), 1.7 (brs, 1H);

^{13}C NMR (CDCl_3) δ 147.8, 146.4, 140.5, 137.8, 135.3, 128.3, 128.0, 126.8, 120.5, 117.5, 107.9, 107.2, 100.8, 61.3, 51.3, 43.2; MS (EI) m/z (relative intensity) 240 [M^+ -allyl] (100), 148 (3.9), 117 (2.5), 91 (81), 65 (5.7); Anal. Calcd for $\text{C}_{18}\text{H}_{19}\text{NO}_2$ (281.35): C, 76.84; H, 6.8; N, 4.97. Found: C, 76.52; H, 6.68; N, 5.21. The enantiomeric excess was determined using CHIRALCEL OD-R column, $\text{CH}_3\text{CN}/1\text{M NaClO}_4$ aq = 40/60, flow rate = 0.8 mL/min, UV detection at 254 nm, $t_R = 18.4$ min (major enantiomer), $t_R = 22.2$ min (minor enantiomer); 90% ee.

N-(4-Methoxybenzyl)-1-piperonyl-3-butenylamine, 9b

Colorless oil; $[\alpha]_D^{22} = +54.9$ (c 1.0, CHCl_3); IR (neat) 3327, 3072, 2902, 2834, 1638, 1611, 1584, 1512, 1376, 1300, 1244, 1177, 1101, 1038, 997, 814, 759 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.26 (d, $J = 8.4$ Hz, 2H), 7.0 (s, 1H), 6.95 (d, $J = 8.4$ Hz, 2H), 6.85 (s, 2H), 6.04 (s, 2H), 5.78 (ddt, $J = 16.1$ Hz, 10.1 Hz, 6.6 Hz, 1H), 5.18 (ddt, $J = 16.1$ Hz, 2.1 Hz, 2 Hz, 1H), 5.12 (ddt, $J = 10.1$ Hz, 2.1 Hz, 2 Hz, 1H), 3.87 (s, 3H), 3.71–3.7 (m, 1H), 3.67 (d, $J = 13$ Hz, 1H), 3.55 (d, $J = 13$ Hz, 1H), 2.46–2.42 (m, 2H), 1.77 (brs, 1H);

^{13}C NMR (CDCl_3) δ 158.4, 147.7, 146.4, 137.8, 135.4, 132.6, 129.2, 120.5, 117.4, 113.6, 107.8, 107.2, 100.8, 61.1, 55.2, 50.6, 43.1; MS (EI) m/z (relative intensity) 270 [M^+ -allyl] (8.4), 160 (5.4), 122 (9.8), 121 (100), 91 (7.4), 77 (2.8); Anal. Calcd for $\text{C}_{19}\text{H}_{21}\text{NO}_3$ (311.38): C, 73.28; H, 6.79; N, 4.49. Found: C, 73.07; H, 6.9; N, 4.78. The enantiomeric excess was determined using CHIRALCEL OD-R column, $\text{CH}_3\text{CN}/1\text{M NaClO}_4$ aq = 35/65, flow rate = 0.8 mL/min, UV detection at 254 nm, $t_R = 34.37$ min (major enantiomer), $t_R = 38.25$ min (minor enantiomer); 89% ee.

N-(2-Methoxybenzyl)-1-piperonyl-3-butenylamine, 9c

Colorless oil; $[\alpha]_D^{22} = +57.8$ (c 1.0, CHCl_3); IR (neat) 3341, 3073, 3001, 2902, 2834, 1638, 1601, 1588, 1490, 1376, 1326, 1289, 1180, 1098, 996, 868, 755 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.18–7.12 (m, 1H), 7.03–7.0 (m, 1H), 6.85–6.76 (m, 3H), 6.69 (s, 2H), 5.87 (s, 2H), 5.6 (ddt, $J = 16.4$ Hz, 10.2 Hz, 6.6 Hz, 1H), 5.0 (ddt, $J = 16.4$ Hz, 2.1 Hz, 2 Hz, 1H), 4.96 (ddt, $J = 10.2$ Hz, 2.1 Hz, 2 Hz, 1H), 3.75 (s, 3H), 3.65 (d, $J = 13.4$ Hz, 1H), 3.48–3.46 (m, 1H), 3.42 (d, $J = 13.4$ Hz, 1H), 2.3–2.17 (m, 2H), 1.95 (brs, 1H);

^{13}C NMR (CDCl_3) δ 157.5, 147.0, 146.2, 137.9, 135.5, 129.8, 128.2, 128.0, 120.5, 120.1, 117.2, 110.0, 107.7, 107.3, 100.6, 60.7, 54.9, 46.9, 43.2; MS (EI) m/z (relative intensity) 270 [M^+ -allyl] (10.5), 160 (5.9), 148 (1.4), 122 (8.9), 121 (100), 93 (23), 91 (3.5); Anal. Calcd for $\text{C}_{19}\text{H}_{21}\text{NO}_3$ (311.38): C, 73.28; H, 6.79; N, 4.49. Found: C, 73.1; H, 7.01; N, 4.4. The enantiomeric excess was determined using CHIRALCEL OD-R column, $\text{CH}_3\text{CN}/1\text{M}$ NaClO_4 aq = 50/50, flow rate = 0.7 mL/min, UV detection at 254 nm, $t_R = 13.3$ min (minor enantiomer), $t_R = 17.11$ min (major enantiomer); 89% ee.

N-Benzyl)-1-(2-chloropiperonyl-3-butenylamine, 9d

Colorless oil; $[\alpha]_D^{22} = +48.0$ (c 1.0, CHCl_3); IR (neat) 3330, 3075, 3027, 2977, 2896, 1639, 1603, 1502, 1475, 1408, 1391, 1375, 1269, 1237, 1159, 1110, 1038, 996, 935, 841, 796, 734 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.92–7.21 (m, 5H), 7.2 (s, 1H), 6.77 (s, 1H), 5.94–5.92 (m, 2H), 5.74 (ddt, $J = 16.8$ Hz, 10.2 Hz, 6.8 Hz, 1H), 5.07–5.0 (m, 2H), 4.15 (dd, $J = 4.8$ Hz, 3.8 Hz, 1H), 3.62 (d, $J = 13.2$ Hz, 1H), 3.5 (d, $J = 13.2$ Hz, 1H), 2.43–2.35 (m, 1H), 2.23–2.13 (m, 1H), 1.66 (brs, 1H);

^{13}C NMR (CDCl_3) δ 147.1, 146.7, 140.3, 134.9, 134.2, 128.2, 128.0, 126.8, 124.9, 117.8, 109.5, 107.4, 101.5, 57.1, 51.4, 41.4; MS (EI) m/z (relative intensity) 274 [M^+ -allyl] (7.5), 169 (2), 115 (2.2), 91 (100); Anal. Calcd for $\text{C}_{18}\text{H}_{18}\text{ClNO}_2$ (315.8): C, 68.46; H, 5.75; N, 4.43; Cl, 11.23. Found: C, 68.4; H, 5.85; N, 4.46; Cl, 11.11. The enantiomeric excess was determined using CHIRALCEL OD-R column, $\text{CH}_3\text{CN}/1\text{M NaClO}_4$ aq = 60/40, flow rate = 0.6 mL/min, UV detection at 254 nm, $t_R = 17.13$ min (major enantiomer), $t_R = 19.3$ min (minor enantiomer); 87% ee.

N-Benzyl-1-(2-bromopiperonyl-3-butenylamine, 9e

Colorless oil; $[\alpha]_D^{22} = +40.2$ (c 1.0, CHCl_3); IR (neat) 3330, 3074, 3027, 2976, 2896, 1638, 1604, 1500, 1473, 1406, 1388, 1373, 1234, 1158, 1038, 996, 981, 837, 735 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.38–7.31 (m, 5H), 7.3 (s, 1H), 7.03 (s, 1H), 6.02–6.01 (m, 2H), 5.81 (ddt, $J = 16.8$ Hz, 10.2 Hz, 6.7 Hz, 1H), 5.16–5.09 (m, 2H), 4.19 (dd, $J = 4.6$ Hz, 3.9 Hz, 1H), 3.70 (d, $J = 13.2$ Hz, 1H), 3.58 (d, $J = 13.2$ Hz, 1H), 2.52–2.43 (m, 1H), 2.29–2.18 (m, 1H), 1.76 (brs, 1H);

^{13}C NMR (CDCl_3) δ 147.7, 147.0, 140.3, 135.7, 134.9, 128.2, 128.0, 126.8, 117.8, 113.9, 112.4, 107.8, 101.5, 59.6, 51.4, 41.5; MS (EI) m/z (relative intensity) 318 [M^+ -allyl] (1.8), 238 (2.9), 214 (2.1), 148 (2), 107 (2.2), 91 (100); Anal. Calcd for $\text{C}_{18}\text{H}_{18}\text{BrNO}_2$ (360.25): C, 60.01; H, 5.04; N, 3.89; Br, 22.18. Found: C, 59.71; H, 5.17; N, 3.98; Br, 22.48. The enantiomeric excess was determined using CHIRALCEL OD-R column, $\text{CH}_3\text{CN}/1\text{M}$ NaClO_4 aq = 60/40, flow rate = 0.6 mL/min, UV detection at 254 nm, $t_R = 21.97$ min (major enantiomer), $t_R = 24.92$ min (minor enantiomer); 88% ee.

N-Diphenylmethylene-1-piperonyl-3-butenylamine, 9f

Colorless oil; $[\alpha]_D^{22} = +53.5$ (c 1.58, CHCl_3); IR (neat) 3325, 3063, 3027, 2978, 2887, 2775, 1639, 1600, 1547, 1504, 1487, 1445, 1377, 1324, 1246, 1184, 1098, 1075, 1042, 941, 912, 864, 762 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.32–7.13 (m, 10H), 6.81 (d, $J = 2\text{ Hz}$, 1H), 6.74 (d, $J = 8\text{ Hz}$, 1H), 6.64 (dd, $J = 8\text{ Hz}$, 2Hz, 1H), 5.92 (s, 2H), 5.65 (ddt, $J = 16.8\text{ Hz}$, 10.1 Hz, 6.8 Hz, 1H), 5.05 (ddt, $J = 16.8\text{ Hz}$, 2 Hz, 2 Hz, 1H), 4.97 (ddt, $J = 10.1\text{ Hz}$, 2 Hz, 2 Hz, 1H), 4.59 (s, 1H), 3.46 (t, 6.5 Hz, 1H), 2.38–2.34 (m, 2H), 2.0–1.94 (brs, 1H);

^{13}C NMR (CDCl_3) δ 147.7, 146.4, 144.5, 143.3, 137.8, 135.4, 128.4, 128.2, 127.7, 127.3, 126.9, 126.6, 120.5, 117.3, 107.9, 107.2, 100.8, 63.0, 58.9, 43.0; MS (EI) m/z (relative intensity) 316 [M^+ -allyl] (8.6), 167 (100), 165 (17.3), 148 (1.8), 122 (3), 115 (4.2), 76 (5.7); Anal. Calcd for $\text{C}_{24}\text{H}_{23}\text{NO}_2$ (357.45): C, 80.64; H, 6.48; N, 3.92. Found: C, 80.52; H, 6.51; N, 3.8. The enantiomeric excess was determined by converting to trifluoroacetylamide form, $[\alpha]_D^{22} = +78.2$ (c 1.0, CHCl_3) and using CHIRALCEL OD column, hexane/*i*-PrOH = 50/1, flow rate = 0.8 mL/min, UV detection at 254 nm, $t_R = 14.44$ min (major enantiomer), $t_R = 18.95$ min (minor enantiomer); 83% ee.

(S)-2-Allylpiperidine

^1H NMR of the hydrochloride ($\text{D}_2\text{O} + \text{CH}_3\text{NO}_2$ internal standard) δ 5.877–5.73 (m, 1H), 5.26–5.2 (m, 2H), 3.39–3.35 (m, 1H), 3.17–3.13 (m, 1H), 3.06–2.87 (m, 1H), 2.52–2.29 (m, 2H), 2.00–1.43 (m, 6H);

^{13}C NMR (CDCl_3) δ 135.5, 117.2, 56.0, 47.0, 41.9, 32.6, 26.2, 24.8. The enantiomeric excess was determined by converting into N-tosylamide, pale yellow oil; $[\alpha]_{\text{D}}^{21} = -0.43$ (c, 0.9, CH_2Cl_2); and using CHIRALCEL OD column, hexane/*i*-PrOH = 50/1, flow rate = 1.0 mL/min, UV detection at 254 nm, $t_{\text{R}} = 9.91$ min (minor enantiomer), $t_{\text{R}} = 11.12$ min (major enantiomer); 2% ee. Configurational assignment was made by comparison with the reported optical rotation.^{6,7}

(R)-2-Allylpiperidine

^1H NMR and ^{13}C NMR same as above. The enantiomeric excess was determined by converting into N-tosylamide, pale yellow oil; $[\alpha]_{\text{D}}^{23} = +4.1$ (c, 0.995, CH_2Cl_2); and using CHIRALCEL OD column, hexane/*i*-PrOH = 50/1, flow rate = 1.0 mL/min, UV detection at 254 nm, $t_{\text{R}} = 9.92$ min (major enantiomer), $t_{\text{R}} = 11.42$ min (minor enantiomer); 18% ee.

(R)-1-Allyl-1,2,3,4-tetrahydroisoquinoline

Yellow oil; $[\alpha]_D^{22} = +6.2$ (c 0.61, THF); 7% ee based on $[\alpha]_D$ reported in literature.⁶ Configurational assignment was made by comparison with the reported optical rotation.⁶ IR (neat) 3072, 3018, 2920, 2833, 2803, 2728, 1638, 1492, 1454, 1430, 1315, 1132, 997, 914, 741 cm^{-1} ; ^1H NMR (CDCl_3) δ 7.17–7.07 (m, 4H), 5.85–5.79 (m, 1H), 5.2–5.12 (m, 2H), 4.05 (dd, $J = 9.0$ Hz, 3.5 Hz, 1H), 3.26–3.2 (m, 1H), 3.01–2.93 (m, 1H), 2.83–2.63 (m, 3H), 2.55–2.48 (m, 1H), 1.81 (brs, 1H);

^{13}C NMR (CDCl_3) δ 138.7, 135.8, 129.1, 126.4, 125.5, 117.8, 55.3, 41.2, 29.8; MS (EI) m/z (relative intensity) 173 [M^+] (0.3), 172 [$\text{M}^+ - \text{H}$] (1.9), 132 [$\text{M}^+ - \text{allyl}$] (100), 117 (6.9), 105 (5.1), 91 (1.2), 77 (3.5).

(S)-1-Allyl-1,2,3,4-tetrahydroisoquinoline

Yellow oil; $[\alpha]_D^{22} = -16.1$ (c 0.61, THF); 19% ee based on $[\alpha]_D$ reported in literature.⁶ IR, ^1H NMR, ^{13}C NMR and MS data same as above.

References and Notes

1. Yamamoto, Y.; Shimota, H.; Oda, J.; Inouye, Y. *Bull. Chem. Soc. Jpn.* **1976**, *49*, 3247.
2. Nakamura, H.; Nakamura, K.; Yamamoto, Y. *J. Am. Chem. Soc.* **1998**, *120*, 4242.
3. Brown, H. C.; Weissman, S. A.; Perumal, P. T.; Dhokte, U. P. *J. Org. Chem.* **1990**, *55*, 1217.
4. Trost, B. M.; Strege, P. E.; Weber, L.; Fullerton, T. J.; Dietsche, T. J. *J. Am. Chem. Soc.* **1978**, *100*, 3407.
5. Trost, B. M.; Metzner, P. J. *J. Am. Chem. Soc.* **1980**, *102*, 3572.
6. (a) Nakamura, M.; Hirai, A.; Nakamura, E. *J. Am. Chem. Soc.* **1996**, *118*, 8489. (b) Meyers, A. I.; Dickman, D. A.; Boes, M. *Tetrahedron* **1987**, *43*, 5095.
7. Aketa, K.-I.; Terashima, S.; Yamada, S.-I. *Chem. Pharm. Bull.* **1976**, *24*, 621.