Supporting information (27 pages)

Cobalt-catalyzed hydrohydrazination reaction of dienes and enynes: convenient access to allylic and propargylic hydrazines

Jérôme Waser, José C. González-Gómez, Hisanori Nambu, Pascal Huber and Erick M. Carreira*

Laboratorium für Organische Chemie, ETH Hönggerberg, CH-8093 Zürich, Switzerland.

carreira@org.chem.ethz.ch

General. All reactions were carried out in oven dried glassware under an atmosphere of argon or nitrogen. For flash chromatography technical grade solvents were used, which were distilled prior to use. Solvents for reactions were analytical grade purchased from Merck or Baker and used without further purifications. All chemicals were purchased from Acros, Aldrich, Fluka, Merck or Lancaster and used as such unless stated otherwise.

Chromatographic purification was performed as flash chromatography using Brunschwig silica 32-63, 60Å, using pentanes/diethylether or hexanes/ethyl acetate as eluent with 0.3-0.5 bar pressure.

TLC was performed on Merck silica gel 60 F₂₅₄ TLC glass plates and visualized with UV light and permanganate stain.

Melting points were measured on a Büchi 510 melting point apparatus using open glass capillaries, the data is uncorrected.

¹H-NMR spectra were recorded on a VARIAN Mercury 300 MHz spectrometer in chloroform-d or DMSO-d₆, all signals are reported in ppm with the internal chloroform signal at 7.26 ppm or the internal DMSO signal at 2.50 ppm as standard. The data is being reported as (s = singlet, d = doublet, t = triplet, m = multiplet or unresolved, br = broad signal, coupling constant(s) in Hz, integration, interpretation).

¹³C-NMR spectra were recorded with ¹H-decoupling on a VARIAN Mercury 75 MHz spectrometer in chloroform-d or DMSO-d₆, all signals are reported in ppm with the internal chloroform signal at 77.0 ppm or the internal DMSO signal at 39.5 ppm as standard.

Many NMR measurements were done at 52 °C to minimize signal broadening due to rotamers mixture (temperature given).

Infrared spectra were recorded on a Perkin Elmer Spectrum RX-I FT-IR spectrophotometer as thin films unless stated otherwise and are reported as cm⁻¹ (w=weak, m=medium, s=strong, sh=shoulder).

Mass spectrometric measurements were performed by the mass spectrometry service of the LOC at the ETHZ on a Finnigan TSQ 7000 ESI spectrometer for low resolution measurements and on an IONSPEC Ultima ESI-FT-ICR spectrometer at 4.7 Tesla for high resolution measurements.

Elemental analysis was performed by the Mikroelementaranalytisches Laboratorium der ETHZ.

Part 1: Catalyst Preparation

N-salicylidene-2-amino-isobutyrate –derivated cobalt (III) complexes

Procedure A:¹

Salicylaldehyde (1.06 mL, 10.0 mmol, 2.00 equiv) and 2-amino-isobutyric acid (1.03 g, 10.0 mmol, 2.00 equiv) were added to a solution of sodium hydroxide (0.21 g, 5.0 mmol, 1.0 equiv) in ethanol (10 mL) at 23 °C under N₂. The resulting thick yellow suspension was cooled to 0 °C and cobalt(II) nitrate hexahydrate (1.45 g, 5.0 mmol, 1.0 equiv) was added as a cold (0 °C) solution in ethanol (10 mL) and the suspension turned to orange. After stirring 15 min at 0 °C, hydrogen peroxide (35%, 7 mL) was added dropwise over 20 min, whereas

(1) Burrows, R. C.; Bailar, J. C. *J. Am. Chem. Soc.* **1966**, 88, 4150.

the reaction mixture immediately turned dark red under gas evolution. The reaction mixture was further stirred at 0 °C for 2 h and then warmed to 23 °C over 3 h. The resulting nearly clear dark red solution was filtered, the solvent was removed under reduced pressure and the residues co- evaporated with methanol (2x) and CH₂Cl₂ (2x). The resulting dark red-brown residues were suspended in Et₂O, triturated for 1 h and filtered, the obtained fine dark red-brown powder was dried in high vacuo at 70 °C for 12 h to afford a dark-red brown solid (2.04 g, 4.14 mmol, 83%).

The identity of the isolated compound was established to be a mixture by NMR analysis. Further repeated attempts to purify it via column chromatography or recrystallization were unsuccessful. The catalyst was used without further purifications and several batches furnished reproducible results. No loss of activity was observed after several months when the compound was kept in dessicator at 23 °C.

Procedure B:

[N-salicylidene-2-amino-isobutyrate]-[2-amino-isobutyrate]-cobalt (III) (1)

Co(OAc)₂·4H₂O (12.5 g, 50.2 mmol, 1.00 equiv) was added to a suspension of salicylaldehyde (5.2 mL, 50 mmol, 1.0 equiv) and 2-amino-isobutyric acid (5.1g, 50 mmol, 1.0 equiv) in water (50 mL) at 23 °C under argon and the reaction mixture was heated to reflux. After refluxing for 2 h, a thick suspension was obtained. The reaction mixture was filtered, the solid washed with water (60 mL) and dried 12 h in high vacuo.

The isolated solid (1.0 g, about 3 mmol) was then suspended in water (12 mL) and ethanol (30 mL) under air and 2-amino-isobutyric acid (0.32 g, 3.1 mmol) was added. The reaction mixture was stirred under air at 23 °C and slowly turned to a dark red solution. After 5 h, the dark red solution was filtered and the solvent was removed under reduced pressure. The residues were co-evaporated with CH₂Cl₂ (2x), dissolved in ethanol, filtered, the solvent was removed under reduced pressure and dried 6 h in high vacuo at 80 °C. Further purification was achieved via column chromatography (CH₂Cl₂/MeOH, solvent gradient 1:0 to 10:1 to 1:1) afforded 920 mg dark red solid. The complex obtained by this procedure contained pure **1** (L=MeOH) as a mixture of isomers (2:1). It shows comparable activity as the complex obtained following procedure A.

¹H NMR (CD₃OD, 300 MHz) δ 8.61 (s, 1H (33%), imine H 1. isomer), 8.30 (s, 1H (66%), imine H 2. isomer), 7.56-7.48 (m, 2H, Ar H), 7.29-7.19 (m, 2H, Ar H), 7.09 (d, *J* = 8.7 Hz, 2H, Ar H), 6.70-6.62 (m, 2H, Ar H), 1.84 (s, 3H (33%), CH₃ 1.isomer), 1.82 (s, 3H (33%), CH₃ 1.isomer), 1.79 (s, 3H (66%), CH₃ 2. isomer), 1.70 (s, 3H (33%), CH₃ 1.isomer), 1.67 (s, 3H (66%), CH₃ 2. isomer), 1.62 (s, 3H (33%), CH₃ 1.isomer), 1.32 (s, 3H (66%), CH₃ 2. isomer), 1.26 (s, 3H (66%), CH₃ 2. isomer);

¹³C NMR (DMSO-*d*₆, 75 MHz) δ 185.5, 184.3, 183.8, 183.2, 166.3, 165.8, 163.8, 163.5, 157.6, 134.7, 134.6, 133.6, 121.9, 121.7, 118.6, 118.2, 114.4, 114.0, 69.6, 69.4, 60.3, 58.2, 29.0, 28.3, 28.2, 27.9, 27.1, 26.6;

IR (KBr pellets) ν 3410 (m), 3242 (m), 3130 (m), 2978 (m), 1661 (s, sh), 1644 (s), 1600 (s), 1546 (s, sh), 1467 (m), 1450 (s), 1387 (s), 1345 (s), 1310 (m), 1205 (m), 1174 (m), 1147 (m), 1128 (w), 1032 (w), 995 (w), 901 (m), 847 (w), 807 (w), 756 (m), 705 (w), 661 (w), 626 (w), 588 (w), 552 (w), 530 (w), 468 (w), 419 (w);

HRMS (MALDI) calcd for $C_{15}H_{20}CoN_2O_5^+$ (M+H): 367.0704, found 367.0695.

Part 2: Preparation of enynes and dienes

2-Methyl-hex-5-en-3-yn-2-ol^{2,3} (**6**)

CuI (0.14 g, 0.72 mmol, 0.014 equiv) and $Pd(PPh_3)_4$ (0.14 g, 0.12 mmol, 0.0023 equiv) were dissolved in diethylamine (25 mL, freshly distilled over KOH) at 10 °C under argon. 2-Methyl-3-in-2-ol (5.0 mL, 52 mmol, 1.0 equiv, freshly distilled) and vinyl bromide (1 M in THF, 65 mL, 65 mmol, 1.3 equiv) were added dropwise and the resulting suspension stirred at 23 °C for 2.5 h. The reaction mixture was then poured into water (100 mL) at 0 °C and extracted with Et_2O (3x125 mL), the combined organic layers were washed with 2 M HCl (2x 50 mL), dried over Na_2SO_4 and concentrated under reduced pressure. The crude mixture was distilled (p = 20 mbar, T = 58°C) to afford **6** (4.84 g, 43.5 mmol, 84%) as a colorless liquid.

1H NMR ($CDCl_3$, 300 MHz) δ 5.81 (dd, 1H, J = 17.4, 10.9 Hz, alkene H), 5.62 (dd, 1H, J = 17.7, 2.5 Hz, alkene H), 5.46 (dd, 1H, J = 10.9, 2.5 Hz), 1.97 (bs, OH), 1.54 (s, 6H, $(CH_3)_2C$);

^{13}C NMR ($CDCl_3$, 75 MHz) δ 126.8, 116.6, 94.4, 80.7, 65.4, 31.3;

IR ν 3350 (s), 3101 (w), 3011 (w), 2982 (s), 2933 (w), 2871 (w), 1844 (w), 1609 (w), 1455 (w), 1411 (m), 1376 (m), 1363 (m), 1240 (s), 1165 (s), 995 (w), 973 (m), 944 (s), 846 (m), 678 (w), 551 (w).

4-Methoxy-benzoic acid 1,1-dimethyl-pent-4-en-2-ynyl ester (**7**)

2-Methyl-hex-5-en-3-yn-2-ol (**6**) (2.24 g, 20.2 mmol, 1.00 equiv) was dissolved in 10 mL pyridine (distilled) at 23 °C under argon. The reaction mixture was cooled to 0 °C and anisoyl chloride (3.5 mL, 26 mmol, 1.3 equiv) was added dropwise. The resulting suspension was heated to 90 °C for 14 h. cooled to 0 °C and quenched with 10 mL water. The mixture was

(2) Kawasaki, T.; Saito, S.; Yamamoto, Y. *J. Org. Chem.* **2002**, 67, 2653.

(3) Sotnichenko, T. V.; Kruglikova, R. I.; Varga, M.; Vasilenko, I. A.; Unkovskii, B. V. *Zh. l Org. Khim.* **1979**, 15, 254.

diluted with water (30 mL) and extracted with Et₂O (3x30 mL). The combined organic layers were washed with 2 M HCl (3x 50 mL), ges. NaHCO₃ (3X30 mL), brine (2x30 mL), dried over Na₂SO₄ and the solvent removed under reduced pressure. The crude mixture was purified by column chromatography (AcOEt/hexane 1:13) to afford **7** (4.40 g, 17.9 mmol, 89%) as a colorless oil. **7** polymerised slowly when kept in the air at 23 °C, but is stable for months when kept at -20 °C under argon.

¹H NMR (CDCl₃, 300 MHz) δ 7.97 (dd, 2H, *J* = 9.0, 1.9 Hz, Ar. H), 6.90 (dd, 2H, *J* = 9.0, 1.9 Hz), 5.82 (dd, 1H, *J* = 17.4, 10.9 Hz, alkene H), 5.64 (dd, 1H, *J* = 17.7, 2.2 Hz, alkene H), 5.46 (dd, 1H, *J* = 10.9, 2.2 Hz), 3.85 (s, 3H, OCH₃), 1.81 (s, 6H, (CH₃)₂COC);

¹³C NMR (CDCl₃, 75 MHz) δ 164.4, 163.0, 131.5, 127.1, 123.3, 116.7, 113.4, 90.9, 82.7, 72.5, 55.4, 29.2;

IR ν 2987 (m), 2938 (m), 2840 (w), 2044 (w), 1922 (w), 1849 (w), 1719 (s), 1606 (s), 1511 (s), 1465 (m), 1420 (w), 1364 (m), 1316 (m), 1283 (s), 1257 (s), 1170 (s), 1141 (s), 1097 (s), 1030 (m), 921 (m), 848 (m), 770 (s), 697 (m);

HRMS(EI) calcd for C₁₅H₁₆O₃⁺ (M) 244.1099, found 244.1090;

Anal. calcd for C₁₅H₁₆O₃: C, 73.75; H, 6.60. Found: C, 73.83; H, 6.79.

But-3-en-1-ynyl-trimethyl-silane⁴ (**8**)

CuI (46 mg, 0.24 mmol, 0.015 equiv) and Pd(PPh₃)₄ (45 mg, 0.039 mmol, 0.0025 equiv) were dissolved in diethylamine (7.5 mL, freshly distilled over KOH) at 10 °C under argon. trimethylsilylacetylene (2.2 mL, 16 mmol, 1.0 equiv, freshly distilled) and vinyl bromide (1 M in THF, 20 mL, 20 mmol, 1.3 equiv) were added dropwise and the resulting suspension stirred at 23 °C for 2.5 h. The reaction mixture was then poured into water (50 mL) at 0 °C and extracted with Et₂O (3x40 mL), the combined organic layers were washed with 2 M HCl (2x 20 mL), dried over Na₂SO₄ and concentrated under reduced pressure. The crude mixture was distilled (p = 104 mbar, T = 54°C) to afford enyne **8** (1.03 g, 8.29 mmol, 53%) as a colorless liquid.

¹H NMR (CDCl₃, 300 MHz) δ 5.82 (dd, 1H, *J* = 17.4, 10.9 Hz, alkene H), 5.69 (dd, 1H, *J* = 17.7, 2.5 Hz, alkene H), 5.50 (dd, 1H, *J* = 10.9, 2.5 Hz), 0.20 (s, 9H, SiCH₃);

¹³C NMR (CDCl₃, 75 MHz) δ 127.8, 117.1, 103.7, 95.0, 0.0;

IR ν 2960 (m), 2900 (w), 2177 (w), 1688 (w), 1412 (w), 1357 (w), 1319 (w), 1251 (s), 1026 (s), 957 (w), 844 (s), 761 (s), 700 (w), 638 (w), 524 (w), 441 (w), 425 (w).

(4) Belyaev, N. N.; Komarov, V. Y.; Komissarova, E. V.; Stadnichuk, M. D. *Zh. Obshch. Khim.* **1982**, 52, 679.

But-3-en-1-ynyl-benzene⁵ (9)

CuI (55 mg, 0.29 mmol, 0.019 equiv) and Pd(PPh₃)₄ (55 mg, 0.048 mmol, 0.0032 equiv) were dissolved in diethylamine (7.5 mL, freshly distilled over KOH) at 10 °C under argon. Phenyl acetylene (1.6 mL, 15 mmol, 1.0 equiv) and vinyl bromide (1 M in THF, 20 mL, 20 mmol, 1.3 equiv) were added dropwise and the resulting suspension stirred at 23 °C for 15 h. The reaction mixture was then poured into water (50 mL) at 0 °C and extracted with Et₂O (3x40 mL), the combined organic layers were washed with 2 M HCl (2x20 mL), dried over Na₂SO₄ and concentrated under reduced pressure. The crude mixture was distilled (p = 10 mbar, T = 73 °C) to afford enyne **9** (1.55 g, 12.1 mmol, 83%) as a colorless liquid.

¹H NMR (CDCl₃, 300 MHz) δ 7.44 (m, 2H, Ar H), 7.32 (m, 3H, Ar H), 6.03 (dd, 1H, *J* = 17.4, 11.2 Hz, alkene H), 5.74 (dd, 1H, *J* = 17.4, 2.2 Hz), 5.55 (dd, 1H, *J* = 11.2, 2.2 Hz);

¹³C NMR (CDCl₃, 75 MHz) δ 131.4, 128.2, 126.8, 123.0, 117.1, 89.9, 88.0;

IR ν 3099 (w), 3080 (w), 3055 (w), 3034 (w), 3009 (w), 2970 (w), 2217 (w), 1950 (w), 1844 (w), 1752 (w), 1606 (m), 1573 (w), 1489 (s), 1442 (m), 1411 (w), 1291 (w), 1262 (w), 1176 (w), 1157 (w), 1070 (w), 1026 (w), 970 (m), 918 (m), 755 (s), 690 (s), 552 (w), 523 (m), 498 (w).

Oct-1-en-3-yne⁶ (10)

CuI (45 mg, 0.24 mmol, 0.015 equiv) and Pd(PPh₃)₄ (45 mg, 0.039 mmol, 0.0025 equiv) were dissolved in diethylamine (7.5 mL, freshly distilled over KOH) at 10 °C under argon. 1-Hexyne (1.7 mL, 15 mmol, 1.0 equiv) and vinyl bromide (1 M in THF, 20 mL, 20 mmol, 1.3 equiv) were added dropwise and the resulting suspension stirred at 23 °C for 4 h. The reaction mixture was then poured into water (50 mL) at 0 °C and extracted with Et₂O (3x40 mL), the combined organic layers were washed with 2 M HCl (2x20 mL), dried over Na₂SO₄ and concentrated under reduced pressure. The crude mixture was distilled (p = 100 mbar, T = 65 °C) to afford enyne **10** (1.11 g, 10.3 mmol, 69%) as a colorless liquid.

¹H NMR (CDCl₃, 300 MHz) δ 5.78 (ddt, 1H, *J* = 17.4, 10.9, 1.9 Hz, alkene H), 5.54 (ddt, 1H, *J* = 17.4, 2.5, 0.6 Hz, alkene H), 5.37 (dd, 1H, *J* = 10.9, 2.5 Hz, alkene H), 2.31 (m, 2H, CH₂CC), 1.57-1.38 (m, 4H, CH₃CH₂CH₂), 0.92 (t, 3H, *J* = 7.3 Hz, CH₃);

¹³C NMR (CDCl₃, 75 MHz) δ 125.3, 117.6, 91.1, 79.2, 30.8, 22.1, 19.1, 13.7;

IR ν 2958 (s), 2934 (s), 2872 (m), 1608 (m), 1465 (w), 1327 (w), 1162 (w), 973 (w), 914 (m).

(5) Miller, K. M.; Luanphaisarnnont, T.; Molinaro, C.; Jamison, T. F. *J. Am. Chem. Soc.* **2004**, 126, 4130.

(6) Jacobson, R. A.; Carothers, W. H. *J. Am. Chem. Soc.* **1933**, 55, 1622.

Trimethyl-pent-3-en-1-ynyl-silane (Z/E 3:1)⁷ (11)

CuI (45 mg, 0.24 mmol, 0.015 equiv) and Pd(PPh₃)₄ (45 mg, 0.039 mmol, 0.0025 equiv) were dissolved in diethylamine (7.5 mL, freshly distilled over KOH) at 10 °C under argon. Trimethylsilylacetylene (2.2 mL, 15 mmol, 1.0 equiv) and 1-bromo-1-propene (Z/E 3:1, 1.7 mL, 20 mmol, 1.3 equiv) were added dropwise and the resulting suspension stirred at 23 °C for 4 h. The reaction mixture was then poured into water (50 mL) at 0 °C and extracted with Et₂O (3x40 mL), the combined organic layers were washed with 2 M HCl (2x20 mL), dried over Na₂SO₄ and concentrated under reduced pressure. The crude mixture was distilled (p = 26 mbar, T = 48 °C) to afford enyne **11** (1.45 g, 10.5 mmol, 68%) as a colorless liquid.

¹H NMR (CDCl₃, 300 MHz) δ 6.22 (dq, 1H, *J* = 15.9, 6.9 Hz, E-alkene H), 6.02 (dq, 1H, *J* = 10.6, 6.9 Hz, Z-alkene H), 5.51 (m, 1H, Z and E-alkene H), 1.89 (dd, 3H, *J* = 6.8, 1.9 Hz, Z-alkene CH₃), 1.78 (dd, 3H, *J* = 6.8, 1.9 Hz, E-alkene CH₃), 0.21 (s, 9H, Z-alkene-SiCH₃), 0.18 (s, 9H, E-alkene-SiCH₃);

¹³C NMR (CDCl₃, 75 MHz) δ 140.9, 139.9, 110.8, 110.1, 104.0, 101.8, 98.9, 92.2, 18.7, 16.1, 0.2, 0.1;

IR ν 3029 (w), 2961 (m), 2913 (w), 2152 (m), 1618 (w), 1441 (w), 1398 (w), 1362 (w), 1250 (s), 1102 (w), 1016 (m), 994 (w), 916 (w), 843 (s), 760 (m), 722 (m), 698 (w), 628 (w), 426 (m).

(Z)-5-Trimethylsilyl-pent-2-en-4-ynoic acid methyl ester^{8,9} (12)

Methylpropiolate (1.8 mL, 20 mmol, 1.0 equiv) and acetic acid (1.3 mL, 22 mmol, 1.1 equiv) were added to a suspension of dry LiBr (2.09 g, 24.1 mmol, 1.20 equiv, dried with heat gun under high vacuo) in dry CH₃CN (20 mL) at 23 °C under argon. The reaction mixture was heated to reflux for 23 h, cooled to 0 °C, trimethylsilylacetylene (3.3 mL, 23 mmol, 1.2 equiv), Pd(PPh₃)₂Cl₂ (0.14 g, 0.20 mmol, 0.010 equiv), CuI (80 mg, 0.42 mmol, 0.021 equiv) and triethylamine (60 mL) were added successively and the resulting suspension was stirred for 4 h at 23 °C protected from light. The resulting dark purple suspension was poured into water (100 mL) at 0 °C, the mixture was extracted with Et₂O (3x120 mL), the combined organic layers were washed with brine, dried over Na₂SO₄ and the solvent was removed under reduced pressure. The crude product was purified by distillation (p = 9 mbar, T = 82 °C) to afford enyne **12** (2.49 g, 13.7 mmol, 68%) as a slightly yellow oil.

¹H NMR (CDCl₃, 300 MHz) δ 6.16 (d, 1H, *J* = 11.6 Hz, alkene H), 6.10 (d, 1H, *J* = 11.6 Hz, alkene H), 3.77 (s, 3H, CO₂CH₃), 0.24 (s, 9H, SiCH₃);

(7) Pornet, J.; Princet, B.; Mevaa, L. M.; Miginiac, L. *Synth. Commun.* **1996**, 26, 2099.

(8) Lu, X. Y.; Huang, X. L.; Ma, S. M. *Tetrahedron Lett.* **1992**, 33, 2535.

(9) Rossi, R.; Bellina, F.; Catanese, A.; Mannina, L.; Valensin, D. *Tetrahedron* **2000**, 56, 479.

¹³C NMR (CDCl₃, 75 MHz) δ 164.9, 129.1, 122.7, 108.2, 100.6, 51.3, -0.5;

IR ν 2958 (m), 2900 (w), 2150 (w), 2068 (w), 1951 (w), 1885 (w), 1732 (s), 1716 (s), 1608 (s), 1437 (m), 1396 (m), 1289 (m), 1251 (s), 1227 (s), 1192 (s), 1173 (s), 1037 (s), 990 (m), 910 (m), 845 (s), 819 (s), 761 (s), 701 (w), 636 (m), 592 (w).

(Z)-5-Trimethylsilyl-pent-2-en-4-yn-1-ol¹⁰ (13)

Ester **12** (375 mg, 2.06 mmol, 1.00 equiv) was diluted in dry Et₂O (5 mL) at 23 °C under argon and the mixture was cooled to -20 °C at which temperature DIBALH (1 M in CH₂Cl₂, 6.5 mL, 6.5 mmol, 3.2 equiv) was added dropwise. The reaction mixture was then allowed to warm to 23 °C over 2 h and stirred 1 h at 23 °C, then quenched with MeOH (0.5 mL) and H₂O (0.7 mL). After gas evolution was finished, potassium/sodium tartrate (1 g) and Et₂O (10 mL) were added and the resulting suspension stirred overnight, whereas the aluminium salts precipitated. The suspension was decanted, the solid extracted with Et₂O (3x10 mL), the combined organic layers dried over Na₂SO₄ and the solvent removed under reduced pressure. The crude product was purified by column chromatography (AcOEt/hexane 1:5) to afford enyne **13** (289 mg, 1.87 mmol, 91%).

¹H NMR (CDCl₃, 300 MHz) δ 6.11 (dt, 1H, *J* = 11.2, 6.5 Hz, alkene H), 5.61 (dt, 1H, *J* = 11.2, 1.6 Hz, alkene H), 4.42 (td, 2H, *J* = 6.2, 1.6 Hz, CH₂OH), 1.67 (bm, 1H, OH), 0.20 (s, 9H, SiCH₃);

¹³C NMR (CDCl₃, 75 MHz) δ 142.6, 110.3, 100.7, 100.5, 60.9, -0.1;

IR ν 3327 (m), 3030 (w), 2959 (m), 2899 (w), 2148 (m), 1612 (w), 1408 (w), 1334 (w), 1250 (s), 1029 (s), 994 (s), 932 (w), 843 (s), 760 (s), 700 (m), 631 (m), 452 (w).

(Z)-5-(tert-Butyl-dimethyl-silanyloxy)-1-trimethylsilyl-pent-3-en-1-yne (14)

Alcohol **13** (308 mg, 2.00 mmol, 1.00 equiv) was added to a solution of imidazole (0.16 g, 2.4 mmol, 1.2 equiv) in DMF (3 mL) at 23 °C under argon. After further addition of *tert*-butyl-dimethyl-silyl chloride (0.36 g, 2.4 mmol, 1.2 equiv) the reaction mixture was stirred 20 h, quenched with water (15 mL) and extracted with AcOEt (3x20 mL). The combined organic layers were washed with water (4x10 mL) and brine (10 mL), dried over Na₂SO₄ and the solvent was removed under reduced pressure. The crude product was purified by column chromatography (pentane/Et₂O 100:1) to afford **14** (460 mg, 1.71 mmol, 86%).

¹H NMR (CDCl₃, 300 MHz) δ 6.05 (dt, 1H, *J* = 11.0, 6.6 Hz, alkene H), 5.51 (dt, 1H, *J* = 11.0, 1.6 Hz, alkene H), 4.43 (dd, 2H, *J* = 6.3, 1.4 Hz, CH₂OTBS), 0.91 (s, 9H, CCH₃), 0.19 (s, 9H, TMS-SiCH₃), 0.1 (s, 6H, TBS-SiCH₃);

(10) Feldman, K. S. *Tetrahedron Lett.* **1982**, 23, 3031.

^{13}C NMR (CDCl₃, 75 MHz) δ 143.8, 109.2, 100.9, 100.2, 61.6, 25.9, 18.3, -0.1, -5.2;

IR ν 3032 (w), 2957 (m), 2929 (m), 2897 (w), 2857 (m), 2149 (w), 1472 (w), 1463 (w), 1400 (w), 1361 (w), 1345 (w), 1251 (s), 1105 (s), 1073 (m), 1001 (w), 969 (w), 939 (w), 842 (s), 776 (m), 760 (m), 721 (w), 698 (w), 662 (w), 628 (w);

HRMS (EI) calcd for C₁₀H₁₉OSi₂⁺ (M-C₄H₉): 211.0969, found 211.0970;

Anal. calcd for C₁₄H₂₈OSi₂: C, 62.62; H, 10.51. Found: C, 62.52; H, 10.55.

Trimethyl-(3-methyl-but-3-en-1-ynyl)-silane¹¹ (15)

n-Butyllithium (1.6 M in hexane, 16.9 mL, 27.0 mmol, 1.00 equiv) was added dropwise over 30 min to a solution of 2-methyl-but-1-en-3-in (2.87 mL, 30.0 mmol, 1.11 equiv) in THF (20 mL) at -78 °C under argon. TMSCl (3.81 mL, 30.0 mmol, 1.11 equiv) was added dropwise at -78 °C and the reaction mixture was let to warm to 23 °C over 1.5 h and stirred at 23 °C for 7 h. The reaction mixture was quenched with water (50 mL), extracted with Et₂O (3x50 mL) and hexane (2x50 mL), the combined organic layers were washed with brine (50 mL), dried over Na₂SO₄ and the solvents were removed under reduced pressure. The crude product was purified by distillation (p = 76 mbar, T = 61 °C) to afford enyne **15** (1.97 g, 14.2 mmol, 53%) as a colorless liquid.

^1H NMR (CDCl₃, 300 MHz) δ 5.34 (m, 1H, alkene H), 5.24 (m, 1H, alkene H), 1.88 (m, 3H, alkene CH₃), 0.18 (s, 9H, SiCH₃);

^{13}C NMR (CDCl₃, 75 MHz) δ 126.8, 122.8, 106.4, 92.9, 23.3, -0.1;

IR ν 2960 (m), 2174 (w), 1450 (w), 1409 (w), 1372 (w), 1251 (s), 1208 (w), 1176 (w), 1131 (w), 1042 (w), 968 (w), 843 (s), 760 (m), 700 (w).

(E)-3-Methyl-5-trimethylsilyl-pent-2-en-4-ynoic acid ethyl ester¹² (16)

Pd(OAc)₂ (67 mg, 0.30 mmol, 0.031 equiv) and tris(2,6-dimethoxyphenyl)phosphine (0.13 g, 0.30 mmol, 0.031 equiv) were dissolved in dry THF (10 mL) at 10 °C under argon. Ethyl-2-butyrate (1.1 g, 9.8 mmol, 1.0 equiv) and trimethylsilylacetylene (1.41 mL, 10.0 mmol, 1.02 equiv) were added dropwise to the red solution which turned dark purple. After stirring 1

(11) Shapiro, E. A.; Lunkova, G. V.; Dolgii, I. E.; Nefedov, O. M. *Bull. Acad. Sci. USSR Div. Chem. Sci.* **1981**, 30, 1047.

(12) Trost, B. M.; Sorum, M. T.; Chan, C.; Harms, A. E.; Ruhter, G. *J. Am. Chem. Soc.* **1997**, 119, 698.

h at 23 °C, the solvent was removed and the crude mixture purified by column chromatography (AcOEt/hexane 1:40) to afford **16** (1.58 g, 7.51 mmol, 77%)

¹H NMR (CDCl₃, 300 MHz) δ 6.09 (q, 1H, *J* = 1.6 Hz, alkene H), 4.16 (q, 2H, *J* = 7.2 Hz, OCH₂CH₃), 2.27 (d, 3H, *J* = 1.6 Hz, alkene CH₃), 1.27 (t, 3H, *J* = 7.2 Hz, OCH₂CH₃), 0.20 (s, 9H, SiCH₃);

¹³C NMR (CDCl₃, 75 MHz) δ 165.7, 137.3, 124.9, 106.4, 99.0, 60.0, 19.7, 14.3, -0.2;

IR ν 2962 (w), 2903 (w), 2147 (w), 1719 (s), 1616 (m), 1446 (w), 1367 (w), 1338 (w), 1252 (m), 1237 (m), 1147 (s), 1041 (w), 860 (s), 844 (s), 761 (w), 639 (w);

HRMS(EI) calcd for C₁₁H₁₈O₂Si⁺ (M) 210.1076, found 210.1088;

Anal. calcd for C₁₁H₁₈O₂Si: C, 62.81; H, 8.62. Found: C, 62.83; H, 8.89.

(E)-3-Methyl-5-trimethylsilyl-pent-2-en-4-yn-1-ol¹³ (**17**)

Ester **16** (0.42 g, 2.0 mmol, 1.0 equiv) was diluted in dry Et₂O (5 mL) at 23 °C under argon and the mixture was cooled to -20 °C at which temperature DIBALH (1 M in CH₂Cl₂, 6.5 mL, 6.5 mmol, 3.2 equiv) was added dropwise. The reaction mixture was then allowed to warm to 23 °C over 2 h and stirred 1 h at 23 °C, then quenched with MeOH (0.5 mL) and H₂O (0.7 mL). After gas evolution was finished, potassium/sodium tartrate (1 g) and Et₂O (10 mL) were added and the resulting suspension stirred overnight, whereas the aluminium salts precipitated. The suspension was decanted, the solid extracted with Et₂O (3x10 mL), the combined organic layers dried over Na₂SO₄ and the solvent removed under reduced pressure. The crude product was purified by column chromatography (AcOEt/hexane 1:3) to afford enyne **17** (315 mg, 1.87 mmol, 94%).

¹H NMR (CDCl₃, 300 MHz) δ 6.03 (tq, 1H, *J* = 6.8, 1.6 Hz, alkene H), 4.19 (d, 2H, *J* = 6.8 Hz, CH₂OH), 1.80 (d, 3H, *J* = 1.6 Hz, alkene CH₃), 1.72 (bm, 1H, OH), 0.17 (s, 9H, SiCH₃);

¹³C NMR (CDCl₃, 75 MHz) δ 136.5, 120.6, 107.3, 92.1, 59.0, 17.3, -0.1;

IR ν 3325 (s), 2960 (s), 2899 (m), 2144 (s), 1631 (w), 1410 (m), 1378 (m), 1249 (s), 1199 (m), 1090 (w), 1009 (s), 842 (s), 760 (s), 699 (m), 643 (m), 580 (w), 534 (w).

(13) Roush, W. R.; Brown, B. B. *J. Am. Chem. Soc.* **1993**, *115*, 2268.

3-Vinyl-cyclopent-3-ene-1,1-dicarboxylic acid diethylester (18)

18 was prepared by metathesis of the corresponding 1,6-enyne following a previously reported procedure.¹⁴ The spectroscopic data is in agreement with the one previously reported.¹⁵

N-tosyl-1-vinyldihydropyrrole (19)

19 was prepared by metathesis of the corresponding 1,6-enyne¹⁶ following a previously reported procedure.¹⁴

¹H NMR (CDCl₃, 300 MHz, 23 °C) δ 7.73 (d, J = 8.1 Hz, 2H, Ph), 7.32 (d, J = 8.1 Hz, 2H, Ph), 6.36 (dd, 1H, J = 17.6 Hz, 10.6 Hz, -CH=CH₂), 5.58 (s, 1H, NCH₂CH=), 5.16 (d, J = 10.6 Hz, 1H, =CH₂), 5.02 (d, J = 17.6 Hz, 1H, =CH₂), 4.19 (m, 4H, CH₂NCH₂), 2.42 (s, 3H, CH₃-Ph);

IR ν 3086 (w), 2917 (w), 2893 (w), 2852 (m), 1812 (w), 1649 (w), 1596 (m), 1492 (w), 1472 (m), 1342 (s), 1303 (s), 1162 (s), 1105 (s), 999 (m), 904 (s), 831 (s), 812 (s), 709 (m), 666 (s), 589 (s), 546 (s).

Part 3: Amination of 1,3-dienes

Di-*tert*-butyl 4,5-dimethylpyridazine-1,2(3H,6H)-dicarboxylate (4)

To a solution of di-*tert*-butylazodicarboxylate (0.17 g, 0.75 mmol, 1.5 equiv) in EtOH (2.5 mL) was added 2,3-dimethyl-1,3-butadiene (**2**) (57 μ L, 0.50 mmol, 1.0 equiv) under argon. The resulting solution was stirred at 23 °C during 24 h, concentrated under vacuum and purified by column chromatography (hexane/EtOAc 9:1), to afford **4** (133 mg, 0.426 mmol, 85%) as a colorless solid.

Mp 87-88 °C;

¹H-NMR(CDCl₃, 300 MHz, 50 °C) δ 4.00-4.20 (br s, 2H, CH₂N), 3.50-3.65 (br s, 2H, CH₂N), 1.62 (s, 6H, CH₃), 1.47 (s, 18H, C(CH₃)₃);

(14) Bentz, D.; Laschat S. *Synthesis* **2000**, 12, 1766.

(15) Fürstner, A.; Stelzer, F.; Szillat, H. *J. Am. Chem. Soc.* **2001**, 123, 11863.

(16) Patel, M. C.; Livinghouse, T.; Pagenkopf, B. L. *Org. Synth.* **2003**, 80, 93.

^{13}C NMR (CDCl_3 , 75 MHz, 23 °C) δ 154.3, 122.8, 80.6, 48.1, 46.6, 28.2, 15.4;

IR ν 2977 (m), 2930 (w), 1706 (s), 1392 (s), 1366 (s), 1345 (w), 1285 (w), 1221 (s), 1175 (s), 1131 (s), 772 (s);

MS (ES) 335.1 (M+Na);

Anal. calcd for $\text{C}_{16}\text{H}_{28}\text{N}_2\text{O}_4$: C, 61.51; H, 9.03; N, 8.97. Found: C, 61.67; H, 9.09; N, 8.77.

General procedure A (amination of dienes):

Di-*tert*-butyl 1-(2,3-dimethylbut-2-enyl)hydrazine-1,2-dicarboxylate (5)

To a solution of catalyst **1** (6 mg, 0.012 mmol, 2.5 mol%) in EtOH (2.5 mL) was added TMDSO (75 μL , 0.42 mmol, 0.80 equiv) at 23 °C under argon, followed by 2,3-dimethyl-1,3-butadiene (**2**) (57 μL , 0.50 mmol, 1.0 equiv) and di-*tert*-butylazodicarboxylate (0.17 g, 0.75 mmol, 1.5 equiv). The resulting solution was stirred at 23 °C overnight (12 h) and then concentrated under vacuum. The residue was purified by column chromatography (hexane/EtOAc 9:1) yielding a colorless solid (130 mg, 0.414 mmol, 83%).

Mp (pentane) 77-78 °C;

^1H NMR (CDCl_3 , 300 MHz, 50 °C) δ 6.05 (br s, 1H, NH), 4.11 (s, 2H, CH_2N), 1.69 (m, 6H, $\text{C}=\text{C}(\text{CH}_3)_2$), 1.64 (m, 3H, $\text{C}=\text{CCH}_3$), 1.48 (s, 9H, $\text{C}(\text{CH}_3)_3$), 1.47 (s, 9H, $\text{C}(\text{CH}_3)_3$);

^{13}C NMR (CDCl_3 , 75 MHz, 23 °C) δ 156.5, 155.8, 130.6, 123.0, 81.1, 50.6, 28.4, 28.2, 21.1, 20.2, 17.0;

IR ν 3296 (m), 2980 (m), 2859 (w), 1729 (s), 1675 (s), 1519 (m), 1393 (s), 1366 (s), 1284 (s), 1254 (s), 1165 (s), 1142 (s), 1052 (m), 1012 (m), 917 (w), 859 (w), 766 (w), 617 (w);

MS (ES) 337.3 (M+Na);

Anal. calcd for $\text{C}_{16}\text{H}_{30}\text{N}_2\text{O}_4$: C, 61.12; H, 9.62; N, 8.91. Found: C, 61.35; H, 9.49; N, 8.81.

4:1 mixture of di-*tert*-butyl 1-(3-methylbut-2-enyl)hydrazine-1,2-dicarboxylate and di-*tert*-butyl 1-(2-methylbut-2-enyl)hydrazine-1,2-dicarboxylate (20a, 20b)

Following general procedure A from isoprene (50 μL , 0.50 mmol, 1.0 equiv), a 4:1 mixture (according to the ^1H -NMR) of isomers **20a** and **20b** (90 mg, 0.30 mmol, 60% combined yield) was obtained as a colorless solid after purification by column chromatography (hexane/EtOAc 9:1).

Mp (pentane) 64-65 °C;

^1H NMR (CDCl_3 , 300 MHz, 50 °C) δ 6.09 (br s, 1H, NH), 5.36 (m, 1H, $=\text{CHCH}_2\text{N}$, 20% 2. isomer), 5.23 (m, 1H, $=\text{CHCH}_2\text{N}$, 80% 2. isomer), 4.05 (d, 2H, J = 6.9 Hz, CH_2N , 80% 1.

isomer), 3.95 (s, 2H, CH₂N, 20% 2. isomer), 1.74 (d, 3H, J = 1.2 Hz, =CCH₃), 1.66 (d, 3H, J = 0.6 Hz, =CCH₃), 1.47 (s, 9H, C(CH₃)₃), 1.46 (s, 9H, C(CH₃)₃);

¹³C NMR (CDCl₃, 75 MHz, 23 °C) δ 155.4, 136.6, 119.1, 81.0, 46.7, 28.2, 25.8, 17.8;

IR ν 3304 (m), 2980 (m), 2931 (w), 1715 (s), 1508(s), 1406 (s), 1366 (s), 1285 (s), 1256 (s), 1158 (s), 1054 (m), 869 (w), 755 (w), 615 (w);

MS (ES) 323.3 (M+Na);

Anal. calcd for C₁₅H₂₈N₂O₄: C, 59.98; H, 9.39; N, 9.33. Found: C, 60.21; H, 9.24; N, 9.24.

7:1 mixture of di-*tert*-butyl 1-(2,5-dimethylhex-4-en-3-yl)hydrazine-1,2-dicarboxylate and di-*tert*-butyl 1-(2,5-dimethylhex-3-en-2-yl)hydrazine-1,2-dicarboxylate (21a, 21b)

Following general procedure **A** from 2,5-dimethyl-2,4-hexadiene (72 μ L, 0.50 mmol, 1.0 equiv), a 7:1 mixture (according to the ¹H-NMR spectra) of isomers **21a** and **21b** (121 mg, 0.355 mmol, 71% combined yield) was obtained as a colorless oil after purification by column chromatography (hexane/EtOAc 9:1). Crystallization from pentane afforded a colorless solid.

Mp (pentane) 75-77 °C;

¹H NMR (CDCl₃, 300 MHz, 50 °C) δ 6.23 (br s, 1H, 14%, 2. isomer, NH), 5.99 (br s, 1H, 86%, 1. isomer, NH), 5.60 (d, $J_{3,4}$ = 16 Hz, 1H, 14%, 2. isomer, =CH), 5.36 (dd, $J_{4,3}$ = 16 Hz, $J_{4,5}$ = 6.6 Hz, 1H, 15%, 2. isomer, =CH), 5.08 (d, $J_{3,4}$ = 8.7 Hz, 1H, 85%, 1. isomer, =CH), 4.40 (br s, 1H, 85%, 1. isomer, CHN), 2.24 (m, 1H, 15%, 2. isomer, CH(CH₃)₂), 1.66 (s, 4H, CH₃C= and CH(CH₃)₂, 1. isomer), 1.62 (s, 3H, CH₃C=, 1. isomer), 1.44 (s, 18 H, C(CH₃)₃), 1.00 (m, 3H, CH₃CH), 0.90 (d, J = 6.9 Hz, 3H, CH₃CH);

¹³C NMR (CDCl₃, 75 MHz, 23 °C) δ 156.2, 155.0, 136.0, 133.5, 122.4, 80.4, 30.7, 28.2, 26.9, 25.7, 22.5, 19.9, 19.1, 18.6;

IR ν 3263 (w), 2976 (s), 2931 (m), 1703 (br. s), 1478 (w), 1454 (w), 1392 (s), 1366 (s), 1301 (m), 1255 (m), 1168 (s), 1048 (w), 1020 (w), 947 (w), 913 (w), 744 (m);

MS (ES) 365.0 (M+Na);

Anal. calcd for C₁₈H₃₄N₂O₄: C, 63.13; H, 10.01; N, 8.18. Found: C, 62.88; H, 9.91; N, 7.95.

Di-*tert*-butyl 1-(3,7-dimethylocta-2,6-dienyl)hydrazine-1,2-dicarboxylate (22)

Following general procedure **A** from myrcene (0.19 mL, 1.0 mmol, 1.0 equiv), but using 1.25 mol% of catalyst **1** and 0.1 M solution in EtOH, **22** was obtained (241 mg, 0.655 mmol, 65%) as a colorless oil after purification by column chromatography (hexane/EtOAc 9:1).

¹H NMR (CDCl₃, 300 MHz, 50 °C) δ 6.28 (br s, 1H, NH), 5.32 (m, 1H, 22%, 2. rotamer, =CHCH₂N), 5.21 (t, *J* = 14 Hz, 1H, 78%, 1. rotamer, =CHCH₂N), 5.07 (m, 1H, CH=C(CH₃)₂), 4.05 (br s, 2H, CH₂N), 2.04 (s, 4H, CH₂CH₂), 1.72 (s, 3H, CH₃), 1.67 (s, 3H, CH₃), 1.59 (s, 3H, CH₃), 1.45 (s, 18H, (CH₃)₃C);

¹³C NMR (CDCl₃, 75 MHz, 23 °C) δ 154.9, 139.5, 131.4, 123.5, 119.6, 80.5, 46.2, 31.7, 28.0, 26.5, 25.5, 23.3, 17.5;

IR ν 3325 (w), 2976 (s), 2931 (m), 1710 (br. s), 1478 (w), 1454 (w), 1392 (s), 1366 (s), 1254 (m), 1159 (s), 1049 (w), 864 (w), 758 (m);

MS (ES) 391.2 (M+Na);

Anal. calcd for C₂₀H₃₆N₂O₄: C, 65.19; H, 9.85; N, 7.60. Found: C, 65.42; H, 10.10; N, 7.55.

Di-*tert*-butyl 3-(4-methylpent-3-enyl)pyridazine-1,2(3H,6H)-dicarboxylate (23)

Procedure A: Compound **23** was obtained as a side product in the hydrohydrazidation of myrcene (see above for experimental details). After purification of the reaction mixture by column chromatography (hexane/EtOAc 9:1), **23** was isolated as a colorless oil (11 mg, 0.03 mmol, 3%).

Procedure B: To a solution of di-*tert*-butylazodicarboxylate (0.17 g, 0.75 mmol) in EtOH (2.5 mL) was added myrcene (95 μL, 0.50 mmol) under argon. The resulting solution was stirred at 23 °C during 48 h, concentrated under vacuum and purified by column chromatography (hexane/EtOAc 9:1), to afford **23** (154 mg, 0.410 mmol, 84%) as a colorless oil.

¹H NMR (CDCl₃, 300 MHz, 23 °C) δ 5.41 (br s, 1H, =CHCH₂N), 5.04 (t, *J* = 6.0 Hz, 1H, Me₂C=CH-), 4.00-4.40 (m, 2H, CH₂N), 3.40-3.80 (m, 2H, CH₂N), 1.94-2.06 (m, 4H, CH₂CH₂), 1.64 (s, 3H, CH₃), 1.55 (s, 3H, CH₃), 1.43 (s, 18H, C(CH₃)₃);

¹³C NMR (CDCl₃, 75 MHz, 23 °C) δ 154.4, 135.1, 132.1, 123.2, 117.2, 80.8, 45.7, 42.9, 34.2, 28.3, 26.1, 25.7, 17.8;

IR ν 2976 (s), 2929 (m), 2854 (m), 1707 (br s), 1478 (w), 1451 (w), 1392 (s), 1366 (s), 1252 (m), 1231 (s), 1175 (s), 1138 (s), 1052 (w), 949 (w), 888 (w), 861 (m), 758 (m);

MS (ES) 389.2 (M+Na);

Anal. calcd for C₂₀H₃₄N₂O₄: C, 65.54; H, 9.35; N, 7.64. Found: C, 65.79; H, 9.12; N, 7.57.

Di-*tert*-butyl 1-(cyclohex-2-enyl)hydrazine-1,2-dicarboxylate (24)

Following general procedure **A** from cyclohexadiene (48 μL, 0.50 mmol, 1.0 equiv), **24** (114 mg, 0.365 mmol, 73%) was obtained as a colorless solid after purification by column chromatography (hexane/EtOAc 9:1).

Mp 128°C;

¹H-NMR(CDCl₃, **300 MHz**, **50 °C**) δ 6.04 (br s, 1H, NH), 5.86-5.77 (m, 1H, C=CH), 5.59 (m, 1H, C=CH), 4.75 (m, 1H, CHN), 2.32-1.6 (m, 6H), 1.49 (s, 9H, C(CH₃)₃), 1.47 (s, 9H, C(CH₃)₃);

¹³C NMR (CDCl₃, **75 MHz**, **23 °C**) δ 155.5, 154.9, 130.6, 127.4, 81.1, 80.8, 53.7, 28.3, 28.2, 27.1, 24.5, 21.0;

IR ν 3318 (m), 2978 (m), 2934 (m), 1702 (s), 1519 (m), 1455 (m), 1402 (s), 1386 (s), 1366 (s), 1315 (s), 1287 (s), 1245 (s), 1162 (s), 1124 (m), 1063 (m), 992 (m), 878 (m), 857 (m), 759 (m);

MS (ES) 335.4 (M+Na);

Anal. calcd for C₁₆H₂₈N₂O₄: C, 61.51; H, 9.03; N, 8.97. Found: C, 61.69; H, 9.06; N, 8.85.

Di-*tert*-butyl 1-(cyclohept-2-enyl)hydrazine-1,2-dicarboxylate (25)

Following general procedure **A** from cycloheptadiene (57 μL, 0.50 mmol, 1.0 equiv), **25** (137 mg, 0.420 mmol, 84%) was obtained as a colorless solid after purification by column chromatography (hexane/EtOAc 9:1).

Mp 124 °C;

¹H NMR (CDCl₃, **300 MHz**, **50 °C**) δ 6.18 (br s, 1H, C=CH), 5.80-5.65 (m, 2H, C=CH and NH), 4.91 (m, 1H, CHN), 2.32-2.22 (m, 1H, C=CH), 2.15-2.06 (m, 1H, C=CH), 2.30-1.89 (m, 4H), 1.71-1.55 (m, 4H), 1.47 (s, 9H, C(CH₃)₃), 1.46 (s, 9H, C(CH₃)₃);

¹³C NMR (CDCl₃, **75 MHz**, **23 °C**) δ 155.7, 154.3, 133.7, 131.0, 81.1, 80.8, 58.7, 32.2, 28.6, 28.3, 28.2, 28.0, 26.7;

IR ν 3277 (m), 2979 (m), 2929 (m), 2858 (w), 1744 (s), 1673 (s), 1519 (m), 1480 (w), 1457 (w), 1412 (s), 1394 (s), 1364 (s), 1319 (s), 1284 (w), 1238 (s), 1153 (s), 1124 (m), 1049 (w), 1021 (w), 952 (w), 886 (m), 777 (m), 759 (m), 683 (m);

MS (ES) 349.2 (M +Na);

Anal. calcd for C₁₇H₃₀N₂O₄: C, 62.55; H, 9.26; N, 8.58. Found: C, 62.53; H, 9.24; N, 8.76.

Di-*tert*-butyl 1-(cyclooct-2-enyl)hydrazine-1,2-dicarboxylate (26)

Following general procedure **A** from cyclooctadiene (64 μL, 0.50 mmol, 1.0 equiv), **26** (138 mg, 0.406 mmol, 81%) was obtained in 6 h as a colorless solid after purification by column chromatography (hexane/EtOAc 9:1).

Mp 123-125 °C;

¹H NMR (CDCl₃, **300 MHz**, **50 °C**) δ 6.06 (br s, 1H, NH), 5.73-5.67 (m, 1H, C=CH), 5.59 (m, 1H, C=CH), 5.05 (m, 1H, CHN), 2.32-2.22 (m, 1H, CH₂C=), 2.15-2.06 (m, 1H, CH₂C=), 1.79-1.53 (m, 8H), 1.49 (s, 9H, C(CH₃)₃), 1.47 (s, 9H, C(CH₃)₃);

¹³C NMR (CDCl₃, 75 MHz, 23 °C) δ 156.0, 154.7, 129.5, 128.8, 81.1, 55.3, 33.3, 29.2, 28.3, 28.2, 26.7, 26.5, 24.4;

IR ν 3322 (m), 2980 (m), 2930 (m), 2858 (w), 1698 (s), 1503 (m), 1455 (m), 1415 (s), 1392 (s), 1366 (s), 1320 (s), 1256 (s), 1161 (s), 1053 (m), 1012 (m), 936 (m), 853 (m), 783 (m), 758 (s), 617 (w);

MS (ES) 363.4 (M+Na);

Anal. calcd for C₁₈H₃₂N₂O₄: C, 63.50; H, 9.47; N, 8.23. Found: C, 63.54; H, 9.54; N, 8.09.

Di-*tert*-butyl 1-(cyclopent-2-enyl)hydrazine-1,2-dicarboxylate (27)

Following general procedure **A** with cyclopentadiene (freshly distilled, 90 μL, 1.0 mmol, 1.0 equiv), TMDSO (150 μL, 0.849 mmol, 0.85 equiv), di-*tert*-butylazodicarboxylate (340 mg, 1.50 mmol, 1.50 equiv) and 2.5 mol% of catalyst **1** in 5 mL of EtOH, **27** (134 mg, 0.450 mmol, 45%) was obtained as a colorless solid after purification by column chromatography (hexane/EtOAc 9:1).

¹H NMR (CDCl₃, 300 MHz, 50 °C) δ 5.94-6.15 (m, 1H, NH), 5.65 (m, 2H, C=CH), 5.26 (m, 1H, 50%, 1. rotamer, CHN), 4.88 (m, 1H, 50%, 2. rotamer, CHN), 2.35-2.60 (m, 3H), 2.05-2.15 (m, 1H), 1.46 (s, 18H, C(CH₃)₃);

¹³C NMR (CDCl₃, 75 MHz, 23 °C) δ 154.2, 155.3, 135 (br s), 129.0, 80.8, 80.7, 63.0 (br s), 56.5 (br s), 35.7, 31.2, 28.1, 28.0;

IR ν 3315 (m), 3058 (w), 2979 (m), 2934 (m), 1702 (br s), 1519 (m), 1457 (w), 1402 (s), 1367 (s), 1327 (m), 1288 (s), 1250 (s), 1159 (s), 1130 (m), 1057 (w), 1031 (w), 1014 (w), 944 (w), 858 (w), 757 (m), 691 (w), 609 (w);

MS (ES, -Q1MS) 297.2 (M-1);

Anal. calcd for C₁₅H₂₆N₂O₄: C, 60.38; H, 8.78; N, 9.39. Found: C, 60.59; H, 8.75; N, 9.27.

Di-*tert*-butyl 2,3-diaza-bicyclo[2.2.1]hept-5-ene-2,3-dicarboxylate (28)

28¹⁷ was a side product in the hydrohydrazination of cyclopentadiene (see above) and after purification of the reaction mixture by column chromatography (hexane/EtOAc 9:1) **28** was obtained as colorless solid (65 mg, 0.23 mmol, 23%).

¹H NMR (CDCl₃, 75 MHz, 50 °C) δ 6.85 (s, 2H, C=CH), 5.05 (m, 2H, CH-NBoc), 1.69 (m, 2H, CH₂), 1.48 (s, 18H, C(CH₃)₃);

IR ν 3003 (m), 2978 (s), 2935 (m), 1701 (br s), 1480 (w), 1456 (w), 1391 (m), 1368 (s), 1342 (m), 1255 (s), 1154 (s), 1129 (m), 1110 (m), 1093 (m), 1046 (w), 1011 (w), 956 (w), 914 (w), 889 (m), 858 (m), 821 (w), 781 (m), 760 (m), 723 (m), 640 (w), 582 (w), 535 (w);

(17) Stout, D. M., Takaya, T., and Meyers, A. I.; *J. Org. Chem.*, **1975**, *40*, 563.

5:1 mixture of di-*tert*-butyl 1-(1-(4,4-bis(ethoxycarbonyl)cyclopent-1-enyl)ethyl)hydrazine-1,2-dicarboxylate (29a) and di-*tert*-butyl 1-(4,4-bis(ethoxycarbonyl)-2-ethylidenecyclopentyl)hydrazine-1,2-dicarboxylate (29b).

Following the general procedure **A** from 3-vinyl-cyclopent-3-ene-1,1-dicarboxylic acid diethylester (**18**) (119 mg, 0.500 mmol, 1.00 equiv), **29** (176 mg, 0.375 mmol, 75%, 5:1 mixture of isomers **29a** and **29b**) was obtained in 2 h, as a colorless oil after purification by column chromatography (hexane/EtOAc 9:1).

¹H NMR (CDCl₃, 300 MHz, 50 °C) δ 6.10 (br s, 1H, NH), 5.44 (s, 1H, =CH), 4.78 (br s, 1H, CHN), 4.20 (m, 4H, CH₃CH₂O rotamers), 2.85-3.20 (m, 4H, CH₂C(CO₂Et)₂CH₂), 1.65 (m, 3H, 17% minor isomers *Z/E*, CH₃C=), 1.47 (s, 18H, C(CH₃)₃), 1.22-1.31 (m, 9H, CH₃-CHN and CH₃CH₂O);

¹³C NMR (CDCl₃, 75 MHz, 23 °C) δ 161.5, 162.3, 155.2, 154.8, 141.7, 141.2, 124.3, 123.6, 80.8, 80.3, 61.5, 59.2, 51.3, 51.0, 40.7, 40.1, 35.3, 28.0, 27.8, 21.4, 15.2, 14.3, 13.8;

IR ν 3300 (w), 2924 (m), 2932 (m), 1733 (br s), 1458 (m), 1367 (s), 1250 (s), 1160 (s), 912 (s);

MS (ES) 493.2 (M+Na);

Di-*tert*-butyl 1-(1-(1-tosyl-2,5-dihydro-1H-pyrrol-3-yl)ethyl)hydrazine-1,2-dicarboxylate (30)

Following the general procedure **A** from N-tosyl-1-vinyldihydropyrrole (**19**) (125 mg, 0.500 mmol, 1.00 equiv), **30** (218 mg, 0.453 mmol, 90%) was obtained in 7 h, as a colorless oil after purification by column chromatography (hexane/EtOAc 8:2). Crystallization from Et₂O afforded a colorless solid.

Mp (Et₂O) 128-130 °C;

¹H NMR (CDCl₃, 300 MHz, 50 °C) δ 7.67 (d, *J* = 8.1 Hz, 2H, Ph), 7.26 (d, *J* = 8.1 Hz, 2H, Ph), 5.90 (br s, 1H, NH), 5.44 (s, 1H, =CH), 4.65 (br s, 1H, CHN), 4.05 (m, 4H, CH₂NCH₂), 2.37 (s, 3H, CH₃-Ph), 1.40 (s, 18H, C(CH₃)₃), 1.20 (d, *J* = 7.2 Hz, 3H, CH₃-CHN);

¹³C NMR (CDCl₃, 75 MHz, 23 °C) δ 155.2, 154.4, 143.2, 139.4, 133.9, 129.6, 127.4, 120.7, 81.4, 80.7, 55.2, 54.8, 49.7, 28.0, 27.9, 21.4, 15.2;

IR ν 3324 (m), 2978 (m), 2932 (m), 1743 (br s), 1704 (br s), 1598 (w), 1478 (m), 1456 (m), 1392 (s), 1368 (s), 1344 (s), 1244 (s), 1164 (s), 1104 (w), 1053 (w), 1066 (w), 1017 (w), 914 (w), 814 (w), 734 (s), 671 (s), 597 (m), 549 (m);

MS (ES) 504.2 (M+Na);

Anal. calcd for C₂₃H₃₅N₃O₆S: C, 57.36; H, 7.32; N, 8.72. Found: C, 57.07; H, 7.60; N, 8.83.

Part 4: Amination of enynes

General procedure B (amination of enynes):

N-(1-Methyl-3-trimethylsilylprop-2-ynyl)-*N'*-(*tert*-butoxycarbonyl)hydrazinecarboxylic acid *tert*-butyl ester (**31**)

The cobalt catalyst **1** (10 mg, 0.025 mmol, 0.05 equiv) was dissolved in ethanol (2.5 mL, degassed) at 23 °C under N₂. Enyne **8** (62 mg, 0.50 mmol, 1.0 equiv) and phenylsilane (93 µL, 0.75 mmol, 1.5 equiv) or TMDSO (0.22 mL, 1.2 mmol, 2.5 equiv) were added to the brown-red solution, followed by di-*tert*-butyl azodicarboxylate (0.23 g, 1.0 mmol, 1.5 equiv) in one portion. The resulting solution was stirred at 23 °C and monitored by TLC (AcOEt/hexane 1:5). After completion (2 h) the reaction was quenched with H₂O (1 mL), brine (5 mL) was added and the reaction mixture was extracted with AcOEt (3x10 mL). The combined organic layers were dried over Na₂SO₄, filtered and the solvents were removed under reduced pressure. The isolated product was purified by column chromatography (AcOEt/hexane 1:10) to afford **31** (148 mg, 0.415 mmol, 83% with phenylsilane and 134 mg, 0.376 mmol, 75% with TMDSO) as a colorless solid.

R_f (AcOEt/hexane 1:5) 0.30;

Mp 82-83 °C;

¹H NMR (CDCl₃, 300 MHz, 52 °C) δ 6.14 (br s, 1H, NH), 5.02 (br m, 1H, CHN), 1.47 (s, 9H, CCH₃), 1.46 (s, 9H, CCH₃), 1.38 (d, *J* = 6.9 Hz, 3H, CNCH₃), 0.15 (s, 9H, SiCH₃);

¹³C NMR (CDCl₃, 75 MHz, 52 °C) δ 155.2, 154.0, 104.3, 87.7, 81.7, 80.9, 46.9, 28.3, 19.6, 0.1;

IR ν 3322 (w), 2980 (m), 2935 (m), 2903 (w), 2174 (w), 1713 (s), 1480 (m), 1456 (m), 1393 (s), 1368 (s), 1320 (s), 1250 (s), 1161 (s), 1112 (s), 1070 (m), 1031 (m), 1028 (w), 1006 (w), 933 (m), 901 (m), 844 (s), 760 (m), 700 (w), 643 (w);

MS (ESI) 357.0 (M+H);

Anal. calcd for C₁₇H₃₂N₂O₄Si: C, 57.27; H, 9.05; N, 7.86. Found: C, 57.53; H, 9.09; N, 7.65.

4-Methoxy-benzoic acid 4-(*N,N'*-bis(*tert*-butoxycarbonyl)-hydrazino)-1,1-dimethylpent-2-ynyl ester (**32**)

32 (145 mg, 0.304 mmol, 61%) was obtained as a gel following the general procedure **B** with enyne **7** (122 mg, 0.500 mmol, 1.00 equiv) and phenylsilane (93 μ l, 0.75 mmol, 1.5 equiv) in 1.5 h with 5 mol% catalyst after purification by column chromatography (AcOEt/hexane 1:4).

R_f (AcOEt/hexane 1:5) 0.15;

¹H NMR (CDCl₃, 300 MHz, 52 °C) δ 7.96 (d, J = 9.0 Hz, 2H, Ar H), 6.90 (d, J = 9.0 Hz, 2H, Ar H), 6.21 (br s, 1H, NH), 5.09 (br m, 1H, CHN), 3.85 (s, 3H, OCH₃), 1.75 (s, 6H, propargylic CH₃), 1.46 (s, 9H, OCCH₃), 1.45 (s, 9H, OCCH₃), 1.36 (d, J = 6.8 Hz, 3H, CNCH₃);

¹³C NMR (CDCl₃, 75 MHz, 52 °C) δ 164.5, 163.2, 155.2, 154.2, 131.4, 123.5, 113.4, 108.7, 84.8, 83.3, 81.4, 80.4, 72.2, 55.3, 46.2, 29.2, 29.0, 28.3, 28.2, 27.1, 19.1;

IR ν 3334 (m), 2980 (m), 2937 (m), 2842 (w), 2254 (w), 1710 (s), 1607 (s), 1511 (m), 1480 (m), 1457 (m), 1393 (s), 1368 (s), 1317 (m), 1284 (s), 1257 (s), 1170 (s), 1100 (s), 1030 (m), 971 (w), 917 (w), 849 (m), 772 (m), 734 (m), 698 (w), 621 (w);

MS (ESI) 499.2 (M+Na);

Anal. calcd for C₂₅H₃₆N₂O₇: C, 63.01; H, 7.61; N, 5.88. Found: C, 63.00; H, 7.69; N, 5.73.

***N*-(1-Methyl-3-phenyl-prop-2-ynyl)- *N'*-(*tert*-butoxycarbonyl)-hydrazinecarboxylic acid *tert*-butyl ester (**33**)**

33 (68 mg, 0.19 mmol, 38%, with phenylsilane or 101 mg, 0.280 mmol, 56%, with TMSO) was obtained as a gel following the general procedure **B** with enyne **9** (64 mg, 0.50 mmol, 1.0 equiv) in 30 min with 5 mol% catalyst after purification by column chromatography (AcOEt/hexane 1:10). Quenching of the reaction as soon as the starting material was consumed proved to be essential to prevent partial reduction of the alkyne triple bond.

R_f (AcOEt/hexane 1:10) 0.2;

¹H NMR (CDCl₃, 300 MHz, 52 °C) δ 7.42-7.39 (m, 2H, Ar H), 7.29-7.27 (m, 3H, Ar H), 6.28 (br s, 1H, NH), 5.28 (br m, 1H, CHN), 1.53-1.39 (m, 21H, OCCH₃ and CHNCH₃);

¹³C NMR (CDCl₃, 75 MHz, 52 °C) δ 155.5, 154.2, 131.8, 128.2, 122.9, 87.9, 83.3, 81.7, 81.0, 46.5, 28.2, 28.2, 19.6;

IR ν 3316 (w), 2979 (m), 2933 (w), 1747 (s), 1705 (s), 1598 (w), 1490 (m), 1479 (w), 1455 (w), 1392 (s), 1368 (s), 1305 (m), 1253 (m), 1160 (s), 1098 (m), 1068 (w), 1050 (w), 1024 (w), 1003 (w), 916 (w), 898 (w), 855 (w), 838 (w), 757 (m), 692 (m), 554 (w), 460 (w), 428 (w);

MS (ESI) 361.1 (M+H), 383.1 (M+Na), 743.2 (2M+Na);

Anal. calcd for C₂₀H₂₈N₂O₄: C, 66.64; H, 7.83; N, 7.77. Found: C, 66.61; H, 8.07; N, 7.62.

***N*-(1-Methyl-hept-2-ynyl)- *N'*-(*tert*-butoxycarbonyl)-hydrazinecarboxylic acid *tert*-butyl ester (34)**

34 (132 mg, 0.389 mmol, 78%, with phenylsilane or 119 mg, 0.350 mmol, 70%, with TMDSO) was obtained as a colorless solid following the general procedure **B** with enyne **10** (54 mg, 0.50 mmol, 1.0 equiv) in 45 min with 5 mol% catalyst after purification by column chromatography (AcOEt/hexane 1:12). Quenching of the reaction as soon as the starting material was consumed proved to be essential to prevent partial reduction of the alkyne triple bond.

R_f (AcOEt/hexane 1:10) 0.25;

Mp 79-80 °C;

¹H NMR (CDCl₃, 300 MHz, 52 °C) δ 6.14 (br s, 1H, NH), 4.98 (br m, 1H, CHN), 2.19-2.10 (m, 2H, alkyne-CH₂), 1.51-1.27 (m, 4H, CH₂), 1.45 (s, 9H, OCCH₃), 1.44 (s, 9H, OCCH₃), 1.32 (d, *J* = 6.9 Hz, 3H, CHNCH₃), 0.88 (t, *J* = 7.1 Hz, 3H, CH₂CH₃);

¹³C NMR (CDCl₃, 75 MHz, 52 °C) δ 155.4, 154.2, 83.7, 81.4, 80.7, 78.5, 46.2, 30.8, 28.2, 28.2, 21.8, 19.8, 18.3, 13.4;

IR ν 3320 (m), 2978 (s), 2933 (s), 2873 (m), 1747 (s), 1705 (s), 1479 (m), 1456 (m), 1392 (s), 1367 (s), 1321 (s), 1240 (s), 1160 (s), 1081 (m), 1055 (m), 1019 (m), 992 (w), 959 (w), 910 (w), 858 (w), 756 (m), 700 (w), 603 (w), 533 (w), 461 (w), 426 (w);

MS (ESI) 341.3 (M+H), 363.2 (M+Na), 703.3 (2M+Na);

Anal. calcd for C₁₈H₃₂N₂O₄: C, 63.50; H, 9.47; N, 8.23. Found: C, 63.27; H, 9.65; N, 8.06.

***N*-(1-Ethyl-3-trimethylsilanyl-prop-2-ynyl)- *N'*-(*tert*-butoxycarbonyl)hydrazinecarboxylic acid *tert*-butyl ester (35)**

35 (80 mg, 0.22 mmol, 55%, with phenylsilane or 77 mg, 0.21 mmol, 52%, with TMDSO) was obtained as a colorless solid following the general procedure **B** with enyne **11** (55 mg, 0.40 mmol, 1.0 equiv) in 3 h with 5 mol% catalyst after purification by column chromatography (AcOEt/hexane 1:20).

R_f (AcOEt/hexane 1:5) 0.4;

Mp 84-86 °C;

¹H NMR (CDCl₃, 300 MHz, 52 °C) δ 6.15 (br s, 1H, NH), 4.78 (br m, 1H, CHN), 1.82-1.62 (m, 2H, CH₂), 1.47 (s, 9H, OCCH₃), 1.46 (s, 9H, OCCH₃), 0.99 (t, *J* = 7.5 Hz, 3H, CH₂CH₃), 0.15 (s, 9H, SiCH₃);

¹³C NMR (CDCl₃, 75 MHz, 52 °C) δ 155.2, 154.3, 103.1, 88.9, 81.6, 80.9, 53.0, 28.3, 27.0, 10.8, -0.1;

IR ν 3322 (m), 2977 (s), 2935 (m), 2255 (w), 2177 (w), 1941 (w), 1706 (s), 1480 (m), 1456 (m), 1392 (s), 1367 (s), 1250 (s), 1162 (s), 1087 (m), 1053 (m), 1017 (m), 979 (m), 844 (s), 760 (s), 700 (w), 643 (w);

MS (ESI) 393.4 (M+Na);

Anal. calcd for $C_{18}H_{34}N_2O_4Si$: C, 58.34; H, 9.25; N, 7.56. Found: C, 58.18; H, 9.28; N, 7.63.

3-(*N,N'*-bis(*tert*-butoxycarbonyl)-hydrazino)-5-trimethylsilyl-pent-4-ynoic acid methyl ester (36)

36 (49 mg, 0.12 mmol, 47%, with phenylsilane) was obtained as a gel following the general procedure **A** with enyne **12** (46 mg, 0.25 mmol, 1.0 equiv) in 4 h with 5 mol% catalyst after purification by column chromatography (AcOEt/hexane 1:5).

R_f (AcOEt/hexane 1:5) 0.2;

¹H NMR ($CDCl_3$, 300 MHz, 52 °C) δ 6.20 (br s, 1H, NH), 5.35 (br m, 1H, CHN), 3.67 (s, 3H, OCH₃), 2.85 (dd, J = 15.3, 7.2 Hz, 1H, CH₂), 2.69 (dd, J = 15.6, 7.5 Hz, 1H, CH₂), 1.47 (s, 18H, OCCH₃), 0.14 (s, 9H, SiCH₃);

¹³C NMR ($CDCl_3$, 75 MHz, 52 °C) δ 169.9, 155.1, 153.7, 101.5, 89.6, 82.1, 81.2, 51.7, 48.3, 38.8, 28.3, -0.1;

IR ν 3327 (w), 2978 (m), 2178 (w), 1743 (s), 1716 (s), 1479 (w), 1458 (w), 1438 (w), 1391 (m), 1368 (s), 1250 (s), 1162 (s), 1024 (w), 903 (w), 844 (s), 760 (m), 700 (w), 644 (w);

MS (ESI) 437.2 (M+Na);

Anal. calcd for $C_{19}H_{34}N_2O_6Si$: C, 55.05; H, 8.27; N, 6.76. Found: C, 54.92; H, 8.23; N, 6.60.

***N*-(1-(2-Hydroxy-ethyl)-3-trimethylsilyl-prop-2-ynyl)-*N'*-(*tert*-butoxycarbonyl)hydrazinecarboxylic acid *tert*-butyl ester (37)**

37 (88 mg, 0.23 mmol, 77%, with phenylsilane and 86 mg, 0.22 mmol, 73%, with TMSO) was obtained as a gel following the general procedure **B** with enyne **13** (46 mg, 0.30 mmol, 1.0 equiv) in 2 h with 5 mol% catalyst after purification by column chromatography (AcOEt/hexane 1:4).

R_f (AcOEt/hexane 1:3) 0.25;

¹H NMR ($CDCl_3$, 300 MHz, 52 °C) δ 6.40 (br s, 1H, NH), 5.03 (br m, 1H, CHN), 3.74-3.58 (m, 2H, CH₂OH), 2.79 (s, 1H, OH), 1.89-1.83 (m, 2H, CHNCH₂), 1.42 (s, 9H, OCCH₃), 1.41 (s, 9H, OCCH₃), 0.10 (s, 9H, SiCH₃);

^{13}C NMR (CDCl_3 , 75 MHz, 52 °C) δ 155.4, 154.7, 102.7, 89.0, 82.0, 81.1, 58.9, 48.5, 36.6, 28.1, 28.0, -0.2;

IR ν 3405 (m), 3321 (m), 2977 (m), 2934 (m), 2252 (w), 2177 (w), 1714 (s), 1479 (m), 1455 (m), 1392 (s), 1368 (s), 1324 (m), 1250 (s), 1159 (s), 1089 (m), 1050 (m), 985 (w), 918 (m), 845 (s), 760 (m), 735 (m), 700 (w), 647 (w);

MS (ESI) 387.2 (M+H), 409.1 (M+Na);

Anal. calcd for $\text{C}_{18}\text{H}_{34}\text{N}_2\text{O}_5\text{Si}$: C, 55.93; H, 8.86; N, 7.25. Found: C, 56.00; H, 8.97; N, 7.31.

***N*-{1-[2-(*tert*-Butyl-dimethyl-silanyloxy)-ethyl]-3-trimethylsilanyl-prop-2-ynyl}-*N'*-(*tert*-butoxycarbonyl)hydrazinecarboxylic acid *tert*-butyl ester (38)**

38 (96 mg, 0.19 mmol, 63%, with phenylsilane and 93 mg, 0.19 mmol, 63%, with TMDSO) was obtained as a gel following the general procedure **B** with enyne **14** (81 mg, 0.30 mmol, 1.0 equiv) in 2 h with 5 mol% catalyst after purification by column chromatography (AcOEt/hexane 1:20).

R_f (AcOEt/hexane 1:5) 0.6;

^1H NMR (CDCl_3 , 300 MHz, 52 °C) δ 6.23 (br s, 1H, NH), 5.08 (m, 1H, CHN), 3.83-3.66 (m, 2H, CH_2OTBS), 2.04-1.93 (m, 1H, CHNCH_2), 1.85-1.74 (m, 1H, CHNCH_2), 1.46 (s, 9H, OCCH_3), 1.46 (s, 9H, OCCH_3), 0.89 (s, 9H, SiCCH_3), 0.14 (s, 9H, TMS SiCH_3), 0.05 (s, 6H, TBS SiCH_3);

^{13}C NMR (CDCl_3 , 75 MHz, 52 °C) δ 155.3, 154.4, 103.0, 88.9, 81.6, 80.8, 59.6, 48.8, 36.4, 28.2, 26.0, 18.3, -0.1, -5.4;

IR ν 3281 (w), 2958 (m), 2931 (m), 2899 (m), 2858 (m), 2174 (w), 1755 (m), 1711 (s), 1473 (m), 1391 (s), 1368 (s), 1319 (m), 1301 (m), 1251 (s), 1158 (s), 1106 (m), 1049 (w), 1018 (w), 951 (w), 840 (s), 777 (m), 760 (m), 700 (w), 665 (w);

MS (ESI) 501.2 (M+H), 523.1 (M+Na);

Anal. calcd for $\text{C}_{24}\text{H}_{48}\text{N}_2\text{O}_5\text{Si}_2$: C, 57.56; H, 9.66; N, 5.59. Found: C, 57.73; H, 9.68; N, 5.61.

***N*-(1,1-Dimethyl-3-trimethylsilanyl-prop-2-ynyl)-*N'*-(*tert*-butoxycarbonyl)hydrazine carboxylic acid *tert*-butyl ester (39)**

39 (62 mg, 0.17 mmol, 42%, with phenylsilane and 54 mg, 0.15 mmol, 36%, with TMDSO) was obtained as a gel following the general procedure **B** with enyne **15** (55 mg, 0.40 mmol, 1.0 equiv) in 4 h with 5 mol% catalyst after purification by column chromatography (AcOEt/hexane 1:10).

R_f (AcOEt/hexane 1:5) 0.35;

¹H NMR (CDCl₃, 300 MHz, 52 °C) δ 6.29 (br s, 1H, NH), 1.67 (s, 3H, CNCH₃), 1.55 (s, 3H, CNCH₃), 1.45 (s, 9H, OCCH₃), 1.44 (s, 9H, OCCH₃), 0.10 (s, 9H, SiCH₃);

¹³C NMR (CDCl₃, 75 MHz, 52 °C) δ 155.7, 154.1, 109.2, 85.6, 81.4, 80.8, 57.0, 0.0;

IR ν 3340 (m), 2978 (m), 2933 (m), 2902 (w), 2166 (w), 1716 (s), 1479 (m), 1456 (m), 1390 (s), 1367 (s), 1250 (s), 1165 (s), 1103 (w), 1078 (m), 1047 (w), 1020 (w), 966 (w), 912 (w), 874 (s), 843 (s), 760 (m), 698 (w), 640 (w);

MS (ESI) 371.0 (M+H), 393.2 (M+Na);

Anal. calcd for C₁₈H₃₄N₂O₄Si: C, 58.34; H, 9.25; N, 7.56. Found: C, 58.27; H, 9.24; N, 7.69.

***N*-[1-(2-Hydroxy-ethyl)-1-methyl-3-trimethylsilanyl-prop-2-ynyl]-*N'*-(*tert*-butoxycarbonyl)hydrazine carboxylic acid *tert*-butyl ester (**40**)**

40 (32 mg, 0.080 mmol, 27%, with TMDSO) was obtained as a colorless solid following the general procedure **B** with enyne **17** (51 mg, 0.30 mmol, 1.0 equiv) in 4 h with 5 mol% catalyst after purification by column chromatography (AcOEt/hexane 1:3).

R_f (AcOEt/hexane 1:5) 0.35;

Mp 84-85 °C;

¹H NMR (CDCl₃, 300 MHz, 52 °C) (partially in coalescence) δ 6.41 (br s, 1H, NH), 4.01-3.67 (m, 3H, CH₂OH and OH), 3.10-2.25 (m, 2H, CNCH₂), 1.66 (m, 3H, CNCH₃), 1.49 (s, 9H, OCCH₃), 1.48 (s, 9H, OCCH₃), 0.16 (s, 9H, SiCH₃);

¹³C NMR (CDCl₃, 75 MHz, 23 °C) (mixture of rotamers) δ 156.7, 156.3, 154.5, 153.7, 108.3, 106.1, 90.2, 86.9, 82.2, 81.9, 81.5, 81.4, 62.6, 59.7, 59.1, 42.8, 42.2, 28.2, 28.2, 27.6, 27.5, -0.1, -0.1;

IR ν 3328 (w), 2978 (m), 2934 (w), 2171 (w), 1714 (s), 1479 (w), 1455 (w), 1392 (m), 1367 (s), 1251 (s), 1160 (s), 1076 (m), 1021 (w), 948 (w), 913 (w), 843 (s), 761 (m), 734 (w), 699 (w), 640 (w), 452 (w), 441 (w), 426 (w);

MS (ESI) 423.1 (M+Na);

Anal. calcd for C₁₉H₃₆N₂O₅Si: C, 56.97; H, 9.06; N, 6.99. Found: C, 56.93; H, 9.15; N, 6.89.

***N*-(1-Methyl-3-trimethylsilanyl-prop-2-ynyl)-*N'*-(ethoxycarbonyl)hydrazinecarboxylic acid ethyl ester (**41**)**

41 (108 mg, 0.359 mmol, 73%) was obtained as a colorless solid following the general procedure **B** with enyne **8** (61 mg, 0.49 mmol, 1.0 equiv), phenylsilane (65 μ l, 0.52, 1.1 equiv) and diethyl azodicarboxylate (0.12 mL, 0.76 mmol, 1.5 equiv) in 1 h with 5 mol% catalyst after purification by column chromatography (AcOEt/hexane 1:5).

R_f (AcOEt/hexane 1:5) 0.30;

Mp 100-102 °C;

¹H NMR (CDCl₃, 300 MHz) δ 6.48 (br s, 1H, NH), 5.07 (m, 1H, CHN), 4.23-4.07 (m, 4H, OCH₂CH₃), 1.37 (d, J = 6.9 Hz, 3H, CHNCH₃), 1.28-1.15 (m, 6H, OCH₂CH₃), 0.12 (s, 9H, SiCH₃);

¹³C NMR (CDCl₃, 75 MHz) δ 156.1, 155.0, 103.4, 88.1, 62.8, 62.0, 47.1, 19.5, 14.5, 0.0;

IR (KBr pellets) ν 3287 (m), 2982 (m), 2912 (w), 2178 (w), 1748 (s), 1685 (s), 1527 (m), 1482 (m), 1430 (m), 1387 (m), 1323 (s), 1281 (m), 1246 (s), 1143 (m), 1112 (m), 1080 (m), 1062 (m), 1024 (w), 926 (m), 843 (s), 761 (m), 702 (w), 638 (w), 592 (w);

MS (ESI) 301.0 (M+H);

Anal. calcd for C₁₃H₂₄N₂O₄Si: C, 51.97; H, 8.05; N, 9.32. Found: C, 52.15; H, 8.04; N, 9.08.

4-Methoxy-benzoic acid 4-(*N,N'*-bis(ethoxycarbonyl)-hydrazino)-1,1-dimethyl-pent-2-ynyl ester (42)

42 (79 mg, 0.19 mmol, 75%) was obtained as a gel following the general procedure **B** with enyne **7** (61 mg, 0.25 mmol, 1.0 equiv), phenylsilane (65 μ l, 0.52, 2.0 equiv) and diethyl azodicarboxylate (60 μ l, 0.38 mmol, 1.5 equiv) in 1 h with 5 mol% catalyst after purification by column chromatography (AcOEt/hexane 1:2).

R_f (AcOEt/hexane 1:5) 0.1;

¹H NMR (CDCl₃, 300 MHz) δ 7.97 (d, J = 9.0 Hz, 2H, Ar H), 6.90 (d, J = 9.0 Hz, 2H, Ar H), 5.12 (br m, 1H, CHN), 4.21-4.10 (m, 4H, OCH₂CH₃), 3.86 (s, 3H, OCH₃), 1.74 (s, 6H, propargylic CH₃), 1.36 (d, J = 6.9 Hz, 3H, CHNCH₃), 1.27-1.22 (m, 6H, OCH₂CH₃);

¹³C NMR (CDCl₃, 75 MHz) δ 164.8, 163.2, 156.1, 155.2, 131.5, 123.1, 113.3, 85.4, 82.9, 72.1, 62.6, 61.6, 55.4, 46.7, 29.1, 28.8, 19.2, 14.5;

IR ν 3315 (w), 2986 (m), 2938 (w), 2841 (w), 2253 (w), 1754 (s), 1715 (s), 1606 (s), 1512 (m), 1467 (m), 1419 (m), 1383 (m), 1316 (s), 1285 (s), 1258 (s), 1171 (m), 1140 (s), 1099 (s), 1061 (m), 1028 (m), 968 (w), 918 (w), 848 (w), 772 (m), 734 (w), 698 (w), 622 (w);

MS (ESI) 443.1 (M+Na);

Anal. calcd for C₂₁H₂₈N₂O₇: C, 59.99; H, 6.71; N, 6.66. Found: C, 59.74; H, 6.82; N, 6.50.

***N*-(1-Methyl-3-phenyl-prop-2-ynyl)- *N'*-(ethoxycarbonyl)-hydrazinecarboxylic acid ethyl ester (43)**

43 (68 mg, 0.22 mmol, 46%) was obtained as a gel following the general procedure **B** with enyne **9** (61 mg, 0.48 mmol, 1.0 equiv), phenylsilane (65 μ L, 0.52, 1.1 equiv) and diethyl azodicarboxylate (0.12 mL, 0.75 mmol, 1.6 equiv) in 1.5 h with 5 mol% catalyst after purification by column chromatography (AcOEt/hexane 1:5).

R_f (AcOEt/hexane 1:5) 0.2;

¹H NMR (CDCl₃, 300 MHz, 52 °C) δ 7.41-7.38 (m, 2H, Ar H), 7.29-7.22 (m, 3H, Ar H), 6.49 (br s, 1H, NH), 5.34-5.28 (m, 1H, CHN), 4.25-4.19 (m, 4H, OCH₂CH₃), 1.51 (d, J = 6.9 Hz, 3H, CHNCH₃), 1.31-1.24 (m, 6H, OCH₂CH₃);

¹³C NMR (CDCl₃, 75 MHz, 52 °C) δ 156.5, 155.3, 131.7, 128.3, 128.2, 122.6, 87.2, 83.8, 62.8, 62.0, 47.1, 19.5, 14.4, 14.4;

IR ν 3295 (m), 2984 (m), 2937 (w), 1712 (s), 1491 (m), 1413 (s), 1383 (s), 1301 (s), 1226 (s), 1174 (m), 1136 (m), 1097 (m), 1060 (s), 1019 (m), 919 (w), 759 (s), 693 (m), 610 (w);

MS (ESI) 305.1 (M+H), 327.0 (M+Na);

Anal. calcd for C₁₆H₂₀N₂O₄: C, 63.14; H, 6.62; N, 9.20. Found: C, 62.86; H, 6.64; N, 8.94.

***N*-(1-Methyl-hept-2-ynyl)- *N'*-(ethoxycarbonyl)-hydrazinecarboxylic acid ethyl ester (44)**

44 (93 mg, 0.33 mmol, 72%) was obtained as a gel following the general procedure **B** with enyne **10** (50 mg, 0.46 mmol, 1.0 equiv), phenylsilane (65 μ L, 0.52, 1.1 equiv) and diethyl azodicarboxylate (0.12 mL, 0.75 mmol, 1.6 equiv) in 1.5 h with 5 mol% catalyst after purification by column chromatography (AcOEt/hexane 1:5).

R_f (AcOEt/hexane 1:5) 0.2;

¹H NMR (CDCl₃, 300 MHz) δ 6.48 (br s, 1H, NH), 5.04 (br m, 1H, CHN), 4.21-4.07 (m, 4H, OCH₂CH₃), 2.15-2.10 (m, 2H, alkyne-CH₂), 1.48-1.21 (m, 10H, 2CH₂ and OCH₂CH₃) 1.34 (d, J = 6.9 Hz, 3H, CHNCH₃), 0.87 (t, J = 7.2 Hz, 3H, CH₂CH₂CH₃);

¹³C NMR (CDCl₃, 75 MHz) δ 156.2, 155.2, 84.2, 77.9, 62.7, 62.0, 46.7, 30.8, 22.0, 20.0, 18.4, 14.5, 13.7;

IR ν 3298 (m), 2983 (m), 2960 (m), 2933 (m), 2873 (w), 2246 (w), 1713 (s), 1517 (m), 1468 (m), 1414 (s), 1383 (s), 1302 (s), 1225 (s), 1174 (w), 1132 (m), 1063 (s), 760 (m), 611 (w);

MS (ESI) 307.4 (M+Na);

Anal. calcd for C₁₄H₂₄N₂O₄: C, 59.13; H, 8.51; N, 9.85. Found: C, 59.26; H, 8.24; N, 9.78.

***N*-(1-Ethyl-3-trimethylsilanyl-prop-2-ynyl)- *N'*-(ethoxycarbonyl) hydrazinecarboxylic acid ethyl ester (45)**

45 (63 mg, 0.21 mmol, 45%) was obtained as a gel following the general procedure **B** with enyne **11** (65 mg, 0.47 mmol, 1.0 equiv) with phenylsilane (65 μ l, 0.52, 1.1 equiv) and diethyl azodicarboxylate (0.12 mL, 0.75 mmol, 1.6 equiv) in 1.5 h with 5 mol% catalyst after purification by column chromatography (AcOEt/hexane 1:7).

R_f (AcOEt/hexane 1:5) 0.2;

Mp 76-77 °C;

¹H NMR (CDCl₃, 300 MHz, 52 °C) δ 6.33 (br s, 1H, NH), 4.86-4.80 (m, 1H, CHN), 4.24-4.12 (m, 4H, OCH₂CH₃), 1.81-1.63 (m, 2H, CHNCH₂), 1.29-1.23 (m, 6H, OCH₂CH₃), 0.97 (t, *J* = 7.5 Hz, 3H, CHNCH₂CH₃), 0.14 (s, 9H, SiCH₃);

¹³C NMR (CDCl₃, 75 MHz, 52 °C) δ 156.3, 155.6, 102.4, 89.6, 62.7, 61.9, 53.5, 26.8, 14.4, 10.5, -0.1;

IR ν 3298 (m), 2968 (m), 2178 (w), 1714 (s), 1517 (w), 1467 (m), 1413 (s), 1382 (s), 1294 (s), 1250 (s), 1221 (s), 1096 (m), 1064 (m), 1025 (m), 977 (w), 927 (w), 845 (s), 760 (m), 700 (w), 646 (w);

HRMS (ESI) calcd for C₁₄H₂₆N₂O₄SiNa⁺ (M+Na): 337.1560, found 337.1550.

Anal. calcd for C₁₄H₂₆N₂O₄Si: C, 53.47; H, 8.33; N, 8.91. Found: C, 53.64; H, 8.17; N, 8.72.

***N*-(1,1-Dimethyl-3-trimethylsilanyl-prop-2-ynyl)- *N'*-(ethoxycarbonyl)hydrazine carboxylic acid ethyl ester (46)**

46 (84 mg, 0.27 mmol, 67%, with phenylsilane and 84 mg, 0.27 mmol, 67%, with TMDSO) was obtained as a gel following the general procedure **B** with enyne **15** (55 mg, 0.40 mmol, 1.0 equiv) in 2 h with 1 mol% catalyst after purification by column chromatography (AcOEt/hexane 1:7).

R_f (AcOEt/hexane 1:5) 0.3;

¹H NMR (CDCl₃, 300 MHz, 52 °C) δ 6.46 (br s, 1H, NH), 4.25-4.14 (m, 4H, OCH₂CH₃), 1.74 (s, 3H, CNCH₃), 1.63 (s, 3H, CNCH₃), 1.30-1.20 (m, 6H, OCH₂CH₃), 0.14 (s, 9H, SiCH₃);

¹³C NMR (CDCl₃, 75 MHz, 52 °C) δ 156.5, 108.2, 86.8, 62.1, 61.9, 58.1, 49.8, 28.7, 14.4, -0.1;

IR ν 3307 (m), 2983 (m), 2961 (m), 2902 (w), 2166 (w), 1719 (s), 1518 (w), 1468 (w), 1446 (w), 1403 (m), 1376 (s), 1341 (s), 1251 (s), 1186 (w), 1097 (s), 1062 (m), 965 (w), 910 (w), 874 (m), 844 (s), 761 (m), 699 (w), 641 (w), 607 (w);

MS (ESI) 337.0 (M+Na);

Anal. calcd for C₁₄H₂₆N₂O₄Si: C, 53.47; H, 8.33; N, 8.91. Found: C, 53.53; H, 8.31; N, 8.97.

***N*-[1-(2-Hydroxy-ethyl)-1-methyl-3-trimethylsilyl-prop-2-ynyl]-*N'*-(ethoxycarbonyl)hydrazine carboxylic acid ethyl ester (**47**)**

47 (68 mg, 0.20 mmol, 66%, with phenylsilane and 73 mg, 0.21 mmol, 71%, with TMSO) was obtained as a gel following the general procedure **B** with enyne **17** (51 mg, 0.30 mmol, 1.0 equiv) in 3 h with 1 mol% catalyst after purification by column chromatography (AcOEt/hexane 1:2).

R_f (AcOEt/hexane 1:1) 0.3;

¹H NMR (CDCl₃, 300 MHz) (mixture of rotamers) δ 7.12-6.90 (m, 1H, NH), 4.24-4.03 (m, 5H, OH and OCH₂CH₃), 4.00-3.72 (m, 2H, CH₂OH), 3.04-2.08 (m, 2H, CNCH₂), 1.85-1.58 (m, 3H, CNCH₃), 1.27-1.20 (m, 6H, OCH₂CH₃), 0.11 (s, 9H, SiCH₃);

¹³C NMR (CDCl₃, 75 MHz) (mixture of rotamers) δ 157.0, 156.6, 156.2, 155.2, 154.7, 106.3, 105.2, 90.6, 89.1, 62.9, 62.3, 62.2, 62.1, 61.9, 61.7, 61.0, 60.8, 59.5, 59.3, 41.8, 41.6, 40.5, 28.4, 27.9, 27.2, 26.5, 14.5, 14.4, -0.1, -0.2;

IR ν 3450 (w), 3304 (m), 2982 (m), 2172 (w), 1945 (w), 1720 (s), 1522 (m), 1467 (m), 1446 (m), 1406 (m), 1377 (s), 1342 (s), 1251 (s), 1172 (m), 1077 (s), 971 (w), 947 (w), 914 (w), 844 (s), 761 (m), 700 (w), 641 (w);

MS (ESI) 345.2 (M+H), 367.1 (M+Na);

Anal. calcd for C₁₅H₂₈N₂O₅Si: C, 52.30; H, 8.19; N, 8.13. Found: C, 52.25; H, 8.28; N, 7.99.