Supporting Information

Cobalt-mediated Cross-coupling Reactions of Primary and Secondary Alkyl Halides with 1-(Trimethylsilyl)ethenyl- and 2-Trimethylsilylethynylmagnesium reagents.

Hirohisa Ohmiya, Hideki Yorimitsu*, and Koichiro Oshima*

Department of Material Chemistry, Graduate School of Engineering, Kyoto University,

Kyoto-daigaku Katsura, Nishikyo-ku, Kyoto 615-8510, Japan

Table of Contents

Instrumentation and Chemicals	S2
Experimental Procedure	S3–S4
Characterization Data	S5–S11
NMR Spectra of All the Products	S12–S39

Instrumentation and Chemicals

^1H NMR (300 MHz and 500 MHz) and ^{13}C NMR (125.7 MHz) spectra were taken on Varian Mercury 300 and UNITY INOVA 500 spectrometers and were recorded in CDCl_3 . Chemical shifts (δ) are in parts per million relative to tetramethylsilane at 0.00 ppm for ^1H and relative to CDCl_3 at 77.2 ppm for ^{13}C unless otherwise noted. IR spectra were determined on a SHIMADZU FTIR-8200PC spectrometer. Mass spectra (EI unless otherwise noted) were determined on a JEOL Mstation 700 spectrometer. TLC analyses were performed on commercial glass plates bearing 0.25-mm layer of Merck Silica gel 60F₂₅₄. Silica gel (Wakogel 200 mesh) was used for column chromatography. Elemental analyses were carried out at the Elemental Analysis Center of Kyoto University.

Unless otherwise noted, materials obtained from commercial suppliers were used without further purification. Anhydrous $\text{Co}(\text{acac})_3$ was purchased from Wako Pure Chemicals. TMEDA was purchased from Nacalai tesque. THF was purchased from Kanto Chemical Co., stored under nitrogen, and used as it is. All reactions were carried out under argon atmosphere.

Experimental Section

Preparation of 1-trimethylsilylethenylmagnesium bromide: Magnesium turnings (1.2 g, 50 mmol) were expeditiously ground down in a mortar under air. The magnesium and anhydrous THF (10 mL) were placed in a 50-mL reaction flask. 1,2-Dibromoethane (0.1 mL) was then added under argon. A solution of 1-bromoethenyltrimethylsilane (4.48 g, 25 mmol) in THF (15 mL) was slowly added at room temperature. The reaction started within a few minutes. After the addition, the mixture was stirred for 30 min at room temperature. 1-Trimethylsilylethenylmagnesium bromide was obtained in a concentration of 0.90–1.0 M. The concentration (0.90–1.0 M) is crucial for the success of the reaction. Lower concentrations resulted in lower yields.

Preparation of 2-trimethylsilylethynylmagnesium bromide: Isopropylmagnesium bromide (1.0 M THF solution, 2.0 mL, 2.0 mmol) was placed in a 30-mL reaction flask under argon. Ethynyltrimethylsilane (196 mg, 2.0 mmol) was added, and the reaction mixture was stirred for 2 h at room temperature.

Typical procedure for cobalt-mediated coupling reaction: The reaction of bromocyclohexane with 1-(trimethylsilyl)ethenylmagnesium bromide is representative (Table 1, entry 2). 1-Trimethylsilylethenylmagnesium bromide (1.0 M THF solution, 2.0 mL, 2.0 mmol) was placed in a 50-mL flask under argon. THF was evaporated under a reduced pressure for 5 min to give a solid. Immediately, TMEDA (1.5 mL) was added to the flask to furnish a suspension. Cobalt(III) acetylacetonate (71.2 mg, 0.20 mmol) was

placed in a 20-mL reaction flask and was heated with a hair dryer in vacuo for 2 min. Anhydrous TMEDA (1 mL) was charged under argon. The mixture was stirred for 3 min at room temperature. Bromocyclohexane (82 mg, 0.50 mmol) was added. The suspension of 1-(trimethylsilyl)ethenylmagnesium bromide was then added over 3 s to the reaction mixture at room temperature via a syringe. While being stirred for 15 min at 25 °C, the mixture turned brown. The reaction mixture was poured into saturated ammonium chloride solution. The products were extracted with hexane (20 mL × 2). The combined organic layer was dried over Na₂SO₄ and concentrated. Silica gel column purification (hexane) of the crude product provided the corresponding product **10** (78 mg, 0.42 mmol) in 85% isolated yield.

Characterization Data

Compounds **11**¹, **13**², **16**³ and **25**⁴ are found in the literature.

Halo acetals **6**, **7**, **8**, **23**, **24** and **26** were prepared according to the literature.⁵

[1-(1-Ethylpropyl)ethenyl]trimethylsilane (14): oil. ¹H NMR (CDCl₃) δ 0.08 (s, 9H), 0.80 (m, *J* = 7.2 Hz, 6H), 1.35–1.45 (m, 4H), 1.98 (quintet, *J* = 6.9, 0.9 Hz, 1H), 5.42 (d, *J* = 2.7 Hz, 1H), 5.54 (dd, *J* = 2.7, 0.9 Hz, 1H); ¹³C NMR (CDCl₃) δ –0.60, 12.33, 27.60, 48.94, 124.09, 155.71.

{1-[2-(2,5-Dioxacyclopentyl)ethyl]ethenyl}trimethylsilane (15): oil. IR (neat) 838, 1248, 1733, 2954 cm^{–1}; ¹H NMR (CDCl₃) δ 0.09 (s, 9H), 1.78 (m, 2H), 2.25 (t, *J* = 7.8 Hz, 2H), 3.83–3.91 (m, 2H), 3.92–4.00 (m, 2H), 4.87 (t, *J* = 4.5 Hz, 1H), 5.33 (m, 1H), 5.58 (m, 1H); ¹³C NMR (CDCl₃) δ –1.36, 30.03, 33.32, 65.05, 104.55, 123.95, 151.54; Found: C, 59.74; H, 9.80%. Calcd for C₁₀H₂₀O₂Si: C, 59.95; H, 10.06%.

1-(*p*-Toluenesulfonyl)-4-(1-trimethylsilylethenyl)piperidine (17): solid. IR (nujol) 741, 1164, 1265, 3055 cm^{–1}; ¹H NMR (CDCl₃) δ 0.04 (s, 9H), 1.51–1.68 (m, 4H), 1.92 (m, 1H), 2.21 (td, *J* = 12.3, 3.6 Hz, 2H), 2.43 (s, 3H), 3.83–3.87 (m, 2H), 5.36 (d, *J* = 2.1 Hz, 1H), 5.57 (m, 1H), 7.33 (d, *J* = 8.1 Hz, 2H), 7.65 (d, *J* = 8.1 Hz, 2H); ¹³C NMR (CDCl₃) δ –0.97, 21.73, 31.89, 40.80, 47.14, 123.94, 127.93, 129.77, 133.25, 143.58, 155.26; Found: C, 60.42; H, 7.88%. Calcd for C₁₇H₂₇NO₂SSi: C, 60.49; H, 8.06%. m.p. 64–66 °C.

1-(*p*-Toluenesulfonyl)-4-(trimethylsilylethynyl)piperidine (18): solid. IR (nujol) 1265,

2304, 3054 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.02 (s, 9H), 1.65–1.76 (m, 2H), 1.80–1.90 (m, 2H), 2.42 (s, 3H), 2.48 (m, 1H), 2.90–2.98 (m, 2H), 3.08–3.18 (m, 2H), 7.31–7.34 (m, 2H), 7.62–7.65 (m, 2H); ^{13}C NMR (CDCl_3) δ 0.09, 21.69, 26.81, 30.66, 44.15, 87.23, 107.91, 127.93, 129.74, 132.79, 143.69; HRMS (EI) Found: 335.1374 [M^+]; $\text{C}_{17}\text{H}_{25}\text{NO}_2\text{SSi}$ Calcd: 335.1376. m.p. 171–173 $^\circ\text{C}$.

[1-(2-Isopropoxy-3-oxacyclopentyl)ethenyl]trimethylsilane (19): oil. IR (neat) 838, 1248, 1454, 2937 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.11 (s, 9H), 1.12 (d, $J = 6.3$ Hz, 3H), 1.18 (d, $J = 6.3$ Hz, 3H), 1.72 (m, 1H), 2.20 (m, 1H), 2.85 (m, 1H), 3.84 (quintet, $J = 6.3$ Hz, 1H), 3.90–4.01 (m, 2H), 4.94 (d, $J = 2.1$ Hz, 1H), 5.42 (d, $J = 2.1$ Hz, 1H), 5.63 (dd, $J = 2.1, 0.9$ Hz, 1H); ^{13}C NMR (CDCl_3) δ -1.31, 22.00, 23.86, 31.47, 49.41, 67.04, 68.89, 106.79, 123.90, 152.62; Found: C, 62.82; H, 10.31%. Calcd for $\text{C}_{12}\text{H}_{24}\text{O}_2\text{Si}$: C, 63.10; H, 10.59%. The *trans* configuration was assigned by the coupling patterns for the protons at C1 and C2.^{5b}

***tert*-Butyldimethyl[2-(1-trimethylsilylethenyl)cyclopentyloxy]silane (20):** oil. IR (neat) 836, 1247, 1472, 2956 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.00 (s, 3H), 0.01 (s, 3H), 0.09 (s, 9H), 0.85 (s, 9H), 1.34 (m, 1H), 1.50–2.00 (m, 5H), 2.52 (m, 1H), 4.07 (dt, $J = 6.3, 5.4$ Hz, 1H), 5.36 (d, $J = 2.4$ Hz, 1H), 5.58 (dd, $J = 2.4, 0.9$ Hz, 1H); ^{13}C NMR (CDCl_3) δ -4.32, -4.21, -0.88, 18.26, 22.38, 26.08, 32.14, 34.99, 53.79, 78.74, 123.38, 154.61. Found: C, 64.05; H, 11.48%. Calcd for $\text{C}_{16}\text{H}_{34}\text{OSi}_2$: C, 64.35; H, 11.48%.

(1-[(1*R*,2*R*)-2-[(+)-Isomenthoxy]-3-oxacyclopentyl]ethenyl)trimethylsilane (21): oil.

IR (neat) 856, 1027, 1248, 2955 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.11 (s, 9H), 0.84 (d, J = 6.6 Hz, 3H), 0.90 (d, J = 7.2 Hz, 3H), 0.91 (d, J = 7.2 Hz, 3H), 1.21–1.63 (m, 7H), 1.72 (m, 1H), 1.90–2.01 (m, 2H), 2.25 (m, 1H), 2.85 (dd, J = 8.1, 3.6 Hz, 1H), 3.71 (dt, J = 7.8, 3.6 Hz, 1H), 3.88–3.99 (m, 2H), 4.94 (d, J = 1.5 Hz, 1H), 5.41 (d, J = 2.1 Hz, 1H), 5.63 (dd, J = 2.1, 1.2 Hz, 1H); ^{13}C NMR (CDCl_3) δ –1.31, 17.68, 19.70, 20.00, 21.28, 25.86, 27.81, 30.78, 31.33, 36.10, 48.02, 49.11, 67.13, 71.40, 105.20, 123.81, 152.77; Found: C, 70.36; H, 11.34%. Calcd for $\text{C}_{19}\text{H}_{36}\text{O}_2\text{Si}$: C, 70.31; H, 11.18%. The absolute configuration was determined by the coupling patterns for the protons at C1 and C2.^{5b}

{1-(2-Bicyclo[2.2.1]heptanyl)ethenyl}trimethylsilane (22): oil. IR (neat) 836, 1247, 2953 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.08 (s, 9H), 1.06 (m, 1H), 1.11–1.53 (m, 7H), 2.15–2.24 (m, 3H), 5.28 (dm, J = 1.8 Hz 1H), 5.51 (m, J = 1.8 Hz 1H); ^{13}C NMR (CDCl_3) δ –0.81, 29.14, 30.48, 35.67, 37.01, 37.52, 41.03, 45.82, 120.22, 156.92; HRMS (EI) Found: 194.1490 [M^+]; $\text{C}_{12}\text{H}_{22}\text{Si}$ Calcd: 194.1491.

Compound 26 (1:1 mixture of diastereomers): oil. IR (neat) 837, 1103, 2956 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.08 (s, 6H), 0.89 (s, 9H), 0.92 (t, J = 7.5 Hz, 3H), 1.34–1.46 (m, 2H), 1.53–1.70 (m, 3H), 2.68 (m, 1H), 3.17–3.27 (m, 2H), 3.43–3.63 (m, 2H), 4.57 (m, 1H), 4.65 (m, 1H), 4.73 (q, J = 5.4 Hz, 1H), 5.87–5.92 (m, 2H); ^{13}C NMR (CDCl_3) δ –4.44, –4.38, 6.01, 6.08, 14.07 (x2C), 18.34, 18.35, 19.53 (x2C), 26.06 (x2C), 31.88, 31.93, 42.06, 42.71, 65.60, 66.00, 74.89, 74.99, 79.11, 79.13, 101.18, 101.38, 132.88, 133.28, 137.72, 137.84; Found: C, 46.20; H, 7.30%. Calcd for $\text{C}_{17}\text{H}_{33}\text{IO}_3\text{Si}$: C, 46.36; H, 7.55%.

1-((2,9-Dioxa-4-bicyclo[4.3.0]nonanyl)methyl)ethenyltrimethylsilane (27): oil. IR (neat) 837, 1023, 1247, 2946 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.09 (s, 9H), 1.34–1.66 (m, 4H), 1.96 (m, 1H), 2.10 (dd, $J = 15.0, 6.9$ Hz, 1H), 2.28 (dd, $J = 15.0, 6.9$ Hz, 1H), 2.57 (m, 1H), 3.61–3.67 (m, 2H), 3.75 (m, 1H), 3.91 (t, $J = 8.1$ Hz, 1H), 5.29 (d, $J = 3.6$ Hz, 1H), 5.34 (m, 1H), 5.58 (m, 1H); ^{13}C NMR (CDCl_3) δ -1.25, 19.47, 23.39, 33.53, 36.66, 39.30, 61.20, 70.21, 102.21, 124.91, 150.14; Found: C, 64.76; H, 9.99%. Calcd for $\text{C}_{13}\text{H}_{24}\text{O}_2\text{Si}$: C, 64.95; H, 10.06%.

2-((2,9-Dioxa-4-bicyclo[4.3.0]nonanyl)methyl)ethynyltrimethylsilane (28): oil. IR (neat) 841, 1250, 2175, 2945 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.13 (s, 9H), 1.36–1.65 (m, 3H), 1.74 (m, 1H), 2.05 (m, 1H), 2.21–2.36 (m, 2H), 2.58 (m, 1H), 3.60–3.80 (m, 3H), 4.01 (t, $J = 8.1$ Hz, 1H), 5.26 (d, $J = 3.9$ Hz, 1H); ^{13}C NMR (CDCl_3) δ 0.21, 18.60, 19.39, 23.13, 36.80, 40.42, 61.41, 69.90, 85.60, 102.04, 104.83; Found: C, 65.51; H, 9.18%. Calcd for $\text{C}_{13}\text{H}_{22}\text{O}_2\text{Si}$: C, 65.50; H, 9.30%.

5,5-Dimethyl-4-[2-(trimethylsilyl)-2-propenyl]-4,5-dihydro-2(3H)-furanone (29): oil. IR (neat) 1120, 1767, 2901 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.10 (s, 9H), 1.28 (s, 3H), 1.46 (s, 3H), 2.03 (dd, $J = 14.4, 11.7$ Hz, 1H), 2.24 (dd, $J = 16.2, 10.2$ Hz, 1H), 2.33–2.44 (m, 2H), 2.56 (dd, $J = 16.2, 6.9$ Hz, 1H), 5.41 (d, $J = 2.4$ Hz, 1H), 5.58 (s, 1H); ^{13}C NMR (CDCl_3) δ -1.14, 22.12, 27.62, 35.06, 36.50, 44.61, 86.69, 126.72, 149.35, 175.84; Found: C, 63.69; H, 10.00%. Calcd for $\text{C}_{12}\text{H}_{22}\text{O}_2\text{Si}$: C, 63.66; H, 9.80%.

5,5-Dimethyl-4-[3-(trimethylsilyl)-2-propynyl]-4,5-dihydro-2(3H)-furanone (30): oil.

IR (neat) 842, 1250, 1775, 2932 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.14 (s, 9H), 1.33 (s, 3H), 1.51 (s, 3H), 2.31–2.54 (m, 4H), 2.72 (m, 1H); ^{13}C NMR (CDCl_3) δ 0.04, 20.68, 22.02, 28.25, 35.03, 44.14, 86.39, 87.40, 103.45, 175.07; Found: C, 64.25; H, 8.99%. Calcd for $\text{C}_{12}\text{H}_{20}\text{O}_2\text{Si}$: C, 64.24; H, 8.98%.

1-(*p*-Toluenesulfonyl)-3-[2-(trimethylsilyl)-2-propenyl]pyrrolidine (31): solid. IR (nujol) 838, 1162, 1597, 2359, 2954 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.02 (s, 9H), 1.40 (m, 1H), 1.87 (m, 1H), 2.00–2.06 (m, 2H), 2.18 (m, 1H), 2.43 (s, 3H), 2.78 (dd, $J = 9.9, 7.2$ Hz, 1H), 3.20 (dt, $J = 14.7, 8.1$ Hz, 1H), 3.30 (m, 1H), 3.37 (dd, $J = 10.2, 7.2$ Hz, 1H), 5.31 (d, $J = 2.7$ Hz, 1H), 5.43 (m, 1H), 7.31 (d, $J = 8.1$ Hz, 2H), 7.71 (d, $J = 8.1$ Hz, 2H); ^{13}C NMR (CDCl_3) δ -1.31, 21.71, 31.41, 37.51, 39.98, 47.60, 53.30, 125.81, 127.73, 129.79, 133.96, 143.48, 150.00; Found: C, 60.44; H, 8.12%. Calcd for $\text{C}_{17}\text{H}_{27}\text{NO}_2\text{SSi}$: C, 60.49; H, 8.06%. m.p. 34–37 $^{\circ}\text{C}$.

1-(*p*-Toluenesulfonyl)-3-[3-(trimethylsilyl)-2-propynyl]pyrrolidine (32): solid. IR (nujol) 841, 1159, 1339, 2174, 2959 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.09 (s, 9H), 1.58 (m, 1H), 1.91 (m, 1H), 2.13–2.16 (m, 2H), 2.24 (m, 1H), 2.42 (s, 3H), 2.95 (dd, $J = 9.9, 7.5$ Hz, 1H), 3.19–3.33 (m, 2H), 3.42 (dd, $J = 9.9, 6.9$ Hz, 1H), 7.32 (d, $J = 8.1$ Hz, 2H), 7.71 (d, $J = 8.1$ Hz, 2H); ^{13}C NMR (CDCl_3) δ 0.15, 21.70, 23.05, 30.45, 37.67, 47.49, 52.48, 86.37, 104.07, 127.71, 129.82, 133.69, 143.58; Found: C, 60.69; H, 7.39%. Calcd for $\text{C}_{17}\text{H}_{25}\text{NO}_2\text{SSi}$: C, 60.85; H, 7.51%. m.p. 80–83 $^{\circ}\text{C}$.

Lactone 33: solid. IR (nujol) 839, 1265, 1771, 2956 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.03 (s,

3H), 0.04 (s, 3H), 0.12 (s, 9H), 0.86 (s, 9H), 1.96 (m, 1H), 2.18 (ddd, $J = 15.0, 7.0, 5.0$ Hz, 1H), 2.55 (dd, $J = 18.0, 4.0$ Hz, 1H), 2.70 (t, $J = 4.0$ Hz, 1H), 2.78 (dd, $J = 18.0, 11.0$ Hz, 1H), 2.90 (m, 1H), 4.05 (dt, $J = 5.0, 3.5$ Hz, 1H), 5.03 (td, $J = 7.5, 1.5$ Hz, 1H), 5.40 (t, $J = 1.5$ Hz, 1H), 5.45 (d, $J = 1.5$ Hz, 1H); ^{13}C NMR (CDCl_3) δ -4.62, -4.32, -0.83, 18.06, 25.88, 36.40, 40.09, 43.00, 60.12, 78.30, 84.86, 125.41, 152.26, 177.62; Found: C, 60.66; H, 9.48%. Calcd for $\text{C}_{18}\text{H}_{34}\text{O}_3\text{Si}_2$: C, 60.96; H, 9.66%. m.p. 71–74 °C. $[\alpha]_{\text{D}}^{25} = -31$ ($c = 1$, CHCl_3).

Lactone 34: solid. IR (neat) 839, 1785, 2926 cm^{-1} ; ^1H NMR (CDCl_3) δ 0.08 (s, 3H), 0.09 (s, 3H), 0.13 (s, 9H), 0.87 (s, 9H), 2.06 (m, 1H), 2.30 (ddd, $J = 14.7, 6.6, 4.8$ Hz, 1H), 2.54 (dd, $J = 18.6, 3.6$ Hz, 1H), 2.71 (m, 1H), 2.81 (dd, $J = 18.6, 11.1$ Hz, 1H), 3.03 (m, 1H), 4.26 (m, 1H), 5.06 (tm, $J = 6.6$ Hz, 1H); ^{13}C NMR (CDCl_3) δ -4.80, -4.72, 0.14, 18.14, 25.83, 35.85, 40.88, 44.76, 47.47, 78.86, 84.08, 88.23, 105.96, 176.91. m.p. 82–85 °C. $[\alpha]_{\text{D}}^{25} = -38$ ($c = 0.6$, CHCl_3).

References

- (1) Brown, H. C.; Mahindroo, V. K.; Bhat, N. G.; Singaram, B. *J. Org. Chem.* **1991**, *56*, 1500–1505.
- (2) Kunishima, M.; Nakata, D.; Tanaka, S.; Hioki, K.; Tani, S. *Tetrahedron* **2000**, *56*, 9927–9935.
- (3) Trost, B. M.; Runge, T. A. *J. Am. Chem. Soc.* **1981**, *103*, 7559–7572.

- (4) Wakabayashi, K.; Yorimitsu, H.; Oshima, K. *J. Am. Chem. Soc.* **2001**, *123*, 5374–5375.
- (5) (a) Inoue, R.; Nakao, J.; Shinokubo, H.; Oshima, K. *Bull. Chem. Soc. Jpn.* **1997**, *70*, 2039–2049. (b) Ohmiya, H.; Yorimitsu, H.; Oshima, K. *J. Am. Chem. Soc.* **2006**, *128*, 1886–1889.

7

(4S)-4-bromobutanoic acid

8

17

CH₂Br₂
internal standard

178

12

135

mgH17 --E--TMS

B

CH₂Br₂
internal standard

179

CO₂ 15

6a

16

22

23

4

25

129

129