Supplementary material for Pavlov V, et al. Combining intraoperative carmustine wafers and Stupp regimen in multimodal first-line treatment of primary glioblastomas. *Br J Neurosurg* 2015;29:524–31.

Supplemental Table I. Postoperative surgical and medical complications.

Patient N	Surgical	WHO Toxicity Grade	Others postoperative complications	WHO Toxicity Grade
2	Meningitis	3	Pneumonia	3
3	Cerebral abscess with death 2.5 months after surgery	4		
12	Neurological worsening	3		
20	Cerebral abscess, meningitis	4, 3		
27	Wound effusion without CSF leak, meningitis	3, 3	Thrombophlebitis with pulmonary embolism	4
33	Postoperative hematoma with neurological worsening, treated surgically	4		
41	Intraoperative gas embolism	4	Acute pulmonary edema	4
45	Neurological worsening	3		
53	Neurological worsening	3		
62	Wound scarring disorder, Postoperative hematoma with minimal neurological worsening	2, 3	Pyelonephritis with septicemia	4
66	Neurological worsening	3		
67	Neurological worsening	3		
74	Neurological worsening	3		

Supplemental Table II. Adverse events during non-surgical first-line oncological treatments.

	WHO toxicity grade	n (%)
RT + CT		
Fatigue	2	13 (16.5)
Local reactions	1	9 (11.4)
Intracranial hypertension	3	2 (2.5)
Cognitive disorders	2	3 (3.8)
Neurological worsening	3	3 (3.8)
Adjuvant CT		
Fatigue	2	15 (27.7)
Hematological disorders	2	8 (14.8)
Kaposi sarcoma	4	1 (1.8)
Digestive disorders	2	3 (5.7)