**Control of Diastereoselectivity in Tandem Asymmetric Reactions
Generating Nonadjacent Stereocenters with Bifunctional Catalysis by
Cinchona Alkaloids**

Baomin Wang, Fanghui Wu, Yi Wang, Xiaofeng Liu and Li Deng*

Department of Chemistry

Brandeis University

Waltham, Massachusetts 02454-9110

Supporting Information

General Information. ^1H and ^{13}C NMR spectra were recorded on a Varian instrument (400 MHz and 100 MHz, respectively) and internally referenced to tetramethylsilane signal or residual protio solvent signals. Data for ^1H NMR are recorded as follows: chemical shift (δ , ppm), multiplicity (s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet), integration, coupling constant (Hz). Data for ^{13}C NMR are reported in terms of chemical shift (δ , ppm). Infrared spectra were recorded on a Perkin Elmer FT-IR Spectrometer and are reported in frequency of absorption. Low resolution and high resolution mass spectra were recorded on either a Micromass 70-VSE-B instrument (EI, CI) or a Micromass Q-Tof instrument (ESI). Specific rotations were measured on a Jasco Digital Polarimeter.

High performance liquid chromatography (HPLC) analysis was performed on a Hewlett-Packard 1100 Series instrument equipped with a quaternary pump, using a Daicel Chiralcel OJ, OD Column (250 x 4.6 mm) or Chiralpak AD, AS Column (250 x 4.6 mm). UV absorption was monitored at 220 nm, 254 nm or 280 nm. Chiral GC was performed on Hewlett-Packard 6890 GC system, using HP chiral (20% Permethyated β -Cyclodextrin) column.

Materials: **3a**,¹ **3b**,² **3c**,³ **3d**,⁴ **3f-k**,⁵ **3l**,⁶ **3m**,⁷ were prepared according to literature methods. **3e** was purchased from Alfa Aesar and used without purification. α -Chloroacrylonitrile **4** and acrylonitrile **6** were purchased from Aldrich and used without further purification. (*Caution: Acrylonitrile 6 is a regulated carcinogen and should be used with careful precaution in hood.*)

Quinine, quinidine, and cinchonidine were purchased from Aldrich Inc. and used without further purification. C9-*epi*-Q was prepared according to literature procedure.⁸ Q-**1a**, QD-**1a**, Q-**1b**, and QD-**1b** were prepared according to our reported methods.⁹ Q-**2a**, Q-**2b**, Q-**2c**, QD-**2c** were prepared according to literature procedures.¹⁰

S-Table 1. Asymmetric Conjugate Addition-Protonation of Michael donors (**3a-e**, **3h**, **3l**) with 2-Chloroacrylonitrile catalyzed by catalysts **2c** and **1**.

3	cat.	5			ent-5			5'			ent-5'		
		dr	ee/%	yield/%	dr	ee/%	yield/%	dr	ee/%	yield/%	dr	ee/%	yield/%
3a	Q-2c	10:1	97	99	9:1	97	99	20:1	93	95	20:1	96	95
	QD-2c												
	Q-1a												
	QD-1a												
3b	Q-2c	16:1	99	94	12:1	97	98	8:1	87	92	11:1	91	87
	QD-2c												
	Q-1a												
	QD-1a												
3c	Q-2c	9:1	97	98	9:1	96	96	25:1	94	92	25:1	95	88
	QD-2c												
	Q-1a												
	QD-1a												
3d	Q-2c	14:1	94	98	10:1	98	97	5:1	85	92	7:1	91	87
	QD-2c												
	Q-1a												
	QD-1a												
3e	Q-2c	15:1	95	98	16:1	95	99	5:1	86	45	4:1	88	85
	QD-2c												
	Q-1b												
	QD-1b												
3h	Q-2c	12:1	95	98	12:1	94	98	3:1	80	75	4:1	86	98
	QD-2c												
	Q-1b												
	QD-1b												
3l	Q-2c	9:1	96	98	9:1	96	98	7:1	88	44	10:1	93	71
	QD-2c												
	Q-1b												
	QD-1b												

^a These data were documented in our previous report.¹¹

General procedure for the asymmetric conjugate addition of α -cyano carbonyl compounds to acrylonitrile or α -chloroacrylonitrile

To a solution of α -cyano carbonyl compound **3** (0.2 mmol or 0.3 mmol) and catalyst **Q-2c** (10-20 mol%) in toluene (2.0 mL or 3.0 mL) was added α -chloroacrylonitrile **4** (3.0 equiv.) or acrylonitrile **6** (3.0 equiv.). The reaction mixture was kept at room temperature until α -cyano carbonyl compound **3** was completely consumed. The crude reaction mixture was filtered through a short plug of silica gel for the removal of the catalyst and washed with diethyl ether (5.0 mL). The filtrate was concentrated *in vacuo*, and the residue was subjected to flash chromatography on silica gel to give the pure adduct **5** or **7**.

Q-2c (12.0 mg, 10 mol%) catalyzed reaction of **3a** (31.4 mg, 0.2 mmol) and **4** (52.5 mg, 0.6 mmol) in toluene (2.0 mL) was run at room temperature for 1 h to furnish the crude product [dr = 10:1, determined by ^1H NMR analysis of the crude product]. The crude product was purified by flash chromatography on silica gel (hexanes/ethyl acetate = 3/1) to give adducts **5a** and **5'a** (dr = 10:1) as a white solid in 99% yield. The ee of the major diastereomer was determined to be 97% ee [determined by HPLC, Chiralpak AD $\times$ 2, hexanes: isopropanol = 80: 20, 1.0 mL/min, λ 220 nm, t (major) = 18.16 min, t (minor) = 23.75 min]. ^1H NMR (400 MHz, CDCl_3) δ 7.87 (d, J = 8.0 Hz, 1H), 7.78 (t, J = 7.6 Hz, 1H), 7.57-7.51 (m, 2H), 4.93 (dd, J = 5.2 Hz, 10.4 Hz, 1H), 3.83 (d, J = 17.6 Hz, 1H), 3.63 (d, J = 17.6 Hz, 1H), 2.87 (dd, J = 5.2 Hz, 14.8 Hz, 1H), 2.45 (dd, J = 10.4 Hz, 14.0 Hz, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 195.0, 150.4, 137.3, 132.1, 129.2, 126.8, 126.0, 117.3, 116.2, 45.9, 41.4, 38.4; HRMS (EI/[M^+]) m/z calcd. for $\text{C}_{13}\text{H}_9\text{N}_2\text{OCl}$: 244.0403, found: 244.0400; IR (neat) ν 1717 cm^{-1} .

ent-**5a** was obtained as a white solid in 99% yield and 97% ee with 9:1 dr from a reaction catalyzed by QD-**2c** (10 mol%) at room temperature for 1 h.

Q-**2c** (18.0 mg, 10 mol%) catalyzed reaction of **3b** (51.3 mg, 0.3 mmol) and **4** (78.8 mg, 0.9 mmol) in toluene (3.0 mL) was run at room temperature for 11 h to furnish the crude product [dr = 16:1, determined by ^1H NMR analysis of the crude product]. The crude product was purified by flash chromatography on silica gel (hexanes/ethyl acetate = 3/1) to give adducts **5b** and **5'b** (dr = 17:1) as a colorless oil in 94% yield. The ee of the major diastereomer was determined to be 99% ee [determined by HPLC, (R, R)-Whelk-O 1, hexanes: isopropanol = 98: 2, 0.6 mL/min, λ 220 nm, t (major) = 60.28 min, t (minor) = 46.04 min]. ^1H NMR (400 MHz, CDCl_3) δ 8.03 (d, J = 7.6 Hz, 1H), 7.61 (t, J = 7.2 Hz, 1H), 7.40 (t, J = 7.2 Hz, 1H), 7.31 (d, J = 7.2 Hz, 1H), 4.92 (dd, J = 5.2 Hz, 8.4 Hz, 1H), 3.45-3.37 (m, 1H), 3.11 (dt, J = 4.4 Hz, 17.6 Hz, 1H), 2.90 (dd, J = 5.6 Hz, 14.8 Hz, 1H), 2.73-2.63 (m, 2H), 2.38-2.31 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 188.3, 142.3, 135.3, 129.1, 129.0, 127.7, 116.4, 116.3, 45.9, 39.8, 38.4, 32.0, 25.5; HRMS (EI/[M^+]) m/z calcd. for $\text{C}_{14}\text{H}_{11}\text{N}_2\text{OCl}$: 258.0560, found: 258.0558; IR (neat) ν 1700 cm^{-1} .

ent-**5b** was obtained as a colorless oil in 98% yield and 97% ee with 12:1 dr from a reaction catalyzed by QD-**2c** (10 mol%) at room temperature for 11 h.

Q-**2c** (12.0 mg, 10 mol%) catalyzed reaction of **3c** (37.0 mg, 0.2 mmol) and **4** (52.5 mg, 0.6 mmol) in toluene (2.0 mL) was run at room temperature for 10 h to furnish the crude product [dr = 9:1, determined by ^1H NMR analysis of the crude product]. The crude product was purified by flash chromatography on silica gel (hexanes/ethyl acetate = 3/1) to give adducts **5c** and **5'c** (dr = 9:1) as a colorless oil in 98% yield. The ee of the major diastereomer was determined to be 97% ee [determined by HPLC, Chiralcel OD, hexanes: isopropanol = 90: 10, 1.0 mL/min, λ 220 nm, t (major) = 43.67 min, t (minor) = 33.04 min]. ^1H NMR (400 MHz, CDCl_3) δ 7.49-7.25 (m, 4H), 4.76 (t, J = 7.2 Hz, 1H), 3.21 (m, 1H), 3.04 (dd, J = 7.2 Hz, 14.4 Hz, 1H), 2.95 (m, 1H), 2.57 (dd, J = 7.2 Hz, 14.4 Hz, 1H), 2.42-2.26 (m,

2H), 2.07-1.95 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 197.9, 138.7, 137.2, 132.5, 129.8, 128.7, 127.1, 117.8, 116.1, 50.8, 40.9, 38.7, 36.3, 32.9, 24.1; HRMS (EI/[M^+]) m/z calcd. for $\text{C}_{15}\text{H}_{13}\text{N}_2\text{OCl}$: 272.0716, found: 272.0714; IR (neat) ν 1700 cm^{-1} .

ent-**5c** was obtained as a colorless oil in 96% yield and 96% ee with 9:1 dr from a reaction catalyzed by QD-**2c** (10 mol%) at room temperature for 11 h.

5d Q-**2c** (18.0 mg, 10 mol%) catalyzed reaction of **3d** (41.1 mg, 0.3 mmol) and **4** (78.8 mg, 0.9 mmol) in toluene (3.0 mL) was run room temperature for 5 h to furnish the crude product [dr = 14:1, determined by ^1H NMR analysis of the crude product]. The crude product was purified by flash chromatography on silica gel (hexanes/ethyl acetate = 4/1) to give adducts **5d** and **5'd** (dr = 16:1) as a white solid in 98% yield. The ee of the major diastereomer was determined to be 94% ee [determined by GC, β -ICD, 100 $^\circ\text{C}$, 30 min, then 100 $^\circ\text{C}$ -200 $^\circ\text{C}$ (0.5 $^\circ\text{C}/\text{min}$), t (major) = 181.60 min, t (minor) = 180.52 min]. ^1H NMR (400 MHz, CDCl_3) δ 4.75 (dd, J = 6.4 Hz, 8.8 Hz, 1H), 3.00 (m, 1H), 2.93 (dd, J = 8.4 Hz, 14.4 Hz, 1H), 2.70-2.63 (m, 1H), 2.44 (dd, J = 6.8 Hz, 14.8 Hz, 1H), 2.17 (dt, J = 2.4 Hz, 13.2 Hz, 1H), 1.97-1.88 (m, 4H), 1.83-1.64 (m, 2H), 1.33-1.22 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 202.8, 118.0, 115.9, 52.0, 41.2, 40.7, 38.8, 36.9, 27.1, 25.9, 22.8; HRMS (EI/[M^+]) m/z calcd. for $\text{C}_{11}\text{H}_{13}\text{N}_2\text{OCl}$: 224.0716, found: 224.0719; IR (neat) ν 1717 cm^{-1} .

ent-**5d** was obtained as a white solid in 97% yield and 98% ee with 10:1 dr from a reaction catalyzed by QD-**2c** (10 mol%) at room temperature for 12 h.

5'd **5'd** was obtained as a colorless oil in 92% yield and 85% ee (The ee was determined by converting **5'd** to compound **8** in conc. HCl at 80 $^\circ\text{C}$ for 1 h.) with 5:1 dr from a reaction catalyzed by Q-**1a** (20 mol%) at room temperature for 96 h. ^1H NMR (400 MHz, CDCl_3) δ 4.72 (t, J = 7.2 Hz, 1H), 3.03 (m, 1H), 2.90 (dd, J = 7.2 Hz, 14.4 Hz, 1H), 2.67 (m, 1H), 2.39 (dd, J = 7.2 Hz, 14.8 Hz, 1H), 2.15 (m, 1H), 1.97 (m, 4H), 1.82 (m, 1H), 1.66 (m, 1H), 1.25 (m,

1H); ^{13}C NMR (100 MHz, CDCl_3) δ 203.0, 118.3, 115.9, 51.5, 41.6, 41.3, 38.4, 37.6, 27.1, 26.1, 22.5; HRMS (ESI/[M+H] $^+$) m/z calcd. for $\text{C}_{11}\text{H}_{14}\text{N}_2\text{OCl}$: 225.0795, found: 225.0801; IR (neat) ν 1715 cm^{-1} .

Compound **8**. 85% ee [determined by HPLC, Daicel chiralcel OJ, hexanes: isopropanol = 70: 30, 1.0 mL/min, λ 220 nm, t (major) = 21.14 min, t (minor) = 25.24 min]. ^1H NMR (400 MHz, CDCl_3) δ 7.96 (brs, 1H), 4.59 (dd, J = 5.6 Hz, 10.4 Hz, 1H), 2.93 (m, 2H), 2.71 (m, 1H), 2.37-2.25 (m, 2H), 1.94-1.83 (m, 3H), 1.73-1.52 (m, 4H); ^{13}C NMR (100 MHz, CDCl_3) δ 208.3, 172.0, 166.6, 58.6, 51.3, 41.9, 35.6, 32.7, 30.4, 26.5, 24.7; HRMS (ESI/[M+H] $^+$) m/z calcd. for $\text{C}_{11}\text{H}_{15}\text{NO}_3\text{Cl}$: 244.0740, found: 244.0738; IR (neat) ν 1715, 1689 cm^{-1} .

ent-**5'd** was obtained as a colorless oil in 87% yield and 91% ee with 7:1 dr from a reaction catalyzed by QD-**1a** (20 mol%) at room temperature for 96 h.

Q-**2c** (18.0 mg, 10 mol%) catalyzed reaction of **3e** (56.7 mg, 0.3 mmol) and **4** (78.8 mg, 0.9 mmol) in toluene (3.0 mL) was run at room temperature for 12 h to furnish the crude product [dr = 15:1, determined by ^1H NMR analysis of the crude product]. The crude product was purified by flash chromatography on silica gel (hexanes/ethyl acetate = 5/1) to give adducts **5e** and **5'e** (dr = 17:1) as a colorless oil in 98% yield. The ee of the major diastereomer was determined to be 95% ee [determined by HPLC, Chiralcel OD, hexanes: isopropanol = 80: 20, 1.0 mL/min, λ 220 nm, t (major) = 70.07 min, t (minor) = 12.04 min]. ^1H NMR (400 MHz, CDCl_3) δ 7.56 (m, 2H), 7.47 (m, 3H), 4.58 (dd, J = 5.6 Hz, 8.8 Hz, 1H), 4.34-4.22 (m, 2H), 3.21 (dd, J = 9.2 Hz, 14.8 Hz, 1H), 2.96 (dd, J = 5.6 Hz, 14.8 Hz, 1H), 1.26 (t, J = 7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.0, 132.0, 130.0, 129.8, 125.9, 116.2, 115.4, 64.2, 51.4, 42.6, 38.8, 13.6; HRMS (EI/[M $^+$]) m/z calcd. for $\text{C}_{14}\text{H}_{13}\text{N}_2\text{O}_2\text{Cl}$: 276.0666, found: 276.0670; IR (neat) ν 1734 cm^{-1} .

ent-**5e** was obtained as a colorless oil in 99% yield and 95% ee with 16:1 dr from a reaction catalyzed by QD-**2c** (10 mol%) at room temperature for 11 h.

5'e was obtained as a colorless oil in 45% yield and 86% ee with 5:1 dr from a reaction catalyzed by **Q-1b** (20 mol%) at room temperature for 152 h.

Q-2c (18.0 mg, 10 mol%) catalyzed reaction of **3f** (62.1 mg, 0.3 mmol) and **4** (78.8 mg, 0.9 mmol) in toluene (3.0 mL) was run at room temperature for 12 h to furnish the crude product [dr = 13:1, determined by ^1H NMR analysis of the crude product]. The crude product was purified by flash chromatography on silica gel (hexanes/ethyl acetate = 6/1) to give adducts **5f** and **5'f** (dr = 17:1) as a colorless oil in 98% yield. The ee of the major diastereomer was determined to be 95% ee [determined by HPLC, Chiralcel OD, hexanes: isopropanol = 90: 10, 1.0 mL/min, λ 220 nm, t (major) = 18.79 min, t (minor) = 15.83 min]. ^1H NMR (400 MHz, CDCl_3) δ 7.55 (q, 2H), 7.17 (t, J = 8.8 Hz, 2H), 4.61 (dd, J = 6.0 Hz, 8.4 Hz, 1H), 4.32-4.22 (m, 2H), 3.21 (dd, J = 9.2 Hz, 15.2 Hz, 1H), 2.92 (dd, J = 5.6 Hz, 14.8 Hz, 1H), 1.27 (t, J = 7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 165.9, 163.4 (d, J = 249.7 Hz), 128.1 (d, J = 9.1 Hz), 127.8 (d, J = 3.8 Hz), 116.9 (d, J = 22 Hz), 116.0, 115.3, 64.4, 50.9, 42.7, 38.7, 13.6; HRMS (EI/[M^+]) m/z calcd. for $\text{C}_{14}\text{H}_{12}\text{N}_2\text{O}_2\text{ClF}$: 294.0571, found: 294.0578; IR (neat) ν 1734 cm^{-1} .

ent-**5f** was obtained as a colorless oil in 98% yield and 95% ee with 12:1 dr from a reaction catalyzed by **QD-2c** (10 mol%) at room temperature for 12 h.

Q-2c (12.0 mg, 10 mol%) catalyzed reaction of **3g** (44.7 mg, 0.2 mmol) and **4** (52.5 mg, 0.6 mmol) in toluene (2.0 mL) was run at room temperature for 12 h to furnish the crude product [dr = 13:1, determined by ^1H NMR analysis of the crude product]. The crude product was purified by flash chromatography on silica gel (hexanes/ethyl acetate = 6/1) to give adducts **5g** and **5'g** (dr = 14:1) as a colorless oil in 98% yield. The ee of the major diastereomer was determined to be 94% ee [determined by HPLC, Chiralcel OD,

hexanes: isopropanol = 80: 20, 1.0 mL/min, λ 220 nm, t (major) = 21.45 min, t (minor) = 11.89 min]. ^1H NMR (400 MHz, CDCl_3) δ 7.50 (d, J = 8.8 Hz, 2H), 7.44 (d, J = 8.4 Hz, 2H), 4.61 (dd, J = 5.6 Hz, 9.2 Hz, 1H), 4.35-4.20 (m, 2H), 3.22 (dd, J = 9.2 Hz, 15.2 Hz, 1H), 2.90 (dd, J = 6.4 Hz, 15.6 Hz), 1.27 (t, J = 7.2 Hz, 3 H); ^{13}C NMR (100 MHz, CDCl_3) δ 165.7, 136.3, 130.5, 129.9, 127.4, 115.8, 115.2, 64.5, 51.0, 42.4, 38.7, 13.6; HRMS (ESI/[M+H] $^+$) m/z calcd. for $\text{C}_{14}\text{H}_{13}\text{N}_2\text{O}_2\text{Cl}_2$: 311.0354, found: 311.0365; IR (neat) ν 1747 cm^{-1} .

Q-2c (18.0 mg, 10 mol%) catalyzed reaction of **3h** (80.4 mg, 0.3 mmol) and **4** (78.8 mg, 0.9 mmol) in toluene (3.0 mL) was run at room temperature for 12 h to furnish the crude product [dr = 12:1, determined by ^1H NMR analysis of the crude product]. The crude product was purified by flash chromatography on silica gel (hexanes/ethyl acetate = 5/1) to give adducts **5h** and **5'h** (dr = 14:1) as a colorless oil in 98% yield. The ee of the major diastereomer was determined to be 95% ee [determined by HPLC, Chiralcel OD, hexanes: isopropanol = 90: 10, 1.0 mL/min, λ 220 nm, t (major) = 63.06 min, t (minor) = 22.78 min]. ^1H NMR (400 MHz, CDCl_3) δ 7.60 (d, J = 8.8 Hz, 2H), 7.43 (d, J = 8.8 Hz, 2H), 4.61 (dd, J = 5.2 Hz, 8.8 Hz, 1H), 4.35-4.20 (m, 2H), 3.22 (dd, J = 9.2 Hz, 14.4 Hz, 1H), 2.90 (dd, J = 5.6 Hz, 15.2 Hz, 1H), 1.27 (t, J = 7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 165.7, 132.9, 131.1, 127.6, 124.4, 115.8, 115.2, 64.5, 51.1, 42.3, 38.7, 13.6; HRMS (EI/[M $^+$]) m/z calcd. for $\text{C}_{14}\text{H}_{12}\text{N}_2\text{O}_2\text{ClBr}$: 353.9771, found: 353.9762; IR (neat) ν 1750 cm^{-1} .

ent-5h was obtained as a colorless oil in 98% yield and 94% ee with 12:1 dr from a reaction catalyzed by **QD-2c** (10 mol%) at room temperature for 12 h.

5'h was obtained as a colorless oil in 75% yield and 80% ee [determined by HPLC, Chiralcel OD, hexanes: isopropanol = 90: 10, 1.0 mL/min, λ 220 nm, t (major) = 16.07 min, t (minor) = 19.38 min] with 3:1 dr from a reaction catalyzed by **Q-1b** (20 mol%) at room temperature for 152 h. ^1H NMR (400 MHz, CDCl_3) δ 7.59 (d, J = 9.2 Hz, 2H), 7.43 (d, J = 8.8 Hz,

2H), 4.67 (dd, $J = 6.4$ Hz, 8.8 Hz, 1H), 4.93-4.22 (m, 2H), 3.26 (dd, $J = 9.2$ Hz, 12.0 Hz, 1H), 2.76 (dd, $J = 6.4$ Hz, 14.8 Hz, 1H), 1.29 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 165.4, 132.8, 131.7, 127.5, 124.3, 115.8, 115.1, 64.6, 51.1, 42.9, 38.2, 13.6; HRMS (EI/[M^+]) m/z calcd. for $\text{C}_{14}\text{H}_{12}\text{N}_2\text{O}_2\text{ClBr}$: 353.9771, found: 353.9762; IR (neat) ν 1747 cm^{-1} .

ent-**5'h** was obtained as a colorless oil in 98% yield and 86% ee with 4:1 dr from a reaction catalyzed by QD-**1b** (20 mol%) at room temperature for 96 h.

Q-**2c** (12.0 mg, 10 mol%) catalyzed reaction of **3i** (40.6 mg, 0.2 mmol) and **4** (52.5 mg, 0.6 mmol) in toluene (2.0 mL) was run at room temperature for 12 h to furnish the crude product [dr = 14:1, determined by ^1H NMR analysis of the crude product]. The crude product was purified by flash chromatography on silica gel (hexanes/ethyl acetate = 6/1) to give adducts **5i** and **5'i** (dr = 17:1) as a colorless oil in 100% yield. The ee of the major diastereomer was determined to be 95% ee [determined by HPLC, Chiralcel OJ, hexanes: isopropanol = 60: 40, 1.0 mL/min, λ 220 nm, t (major) = 58.70 min, t (minor) = 12.59 min]. ^1H NMR (400 MHz, CDCl_3) δ 7.41 (d, $J = 7.6$ Hz, 2H), 7.25 (d, $J = 8.0$ Hz, 2H), 4.56 (dd, $J = 5.6$ Hz, 9.2 Hz, 1H), 4.34-4.19 (m, 2H), 3.18 (dd, $J = 8.4$ Hz, 14.8 Hz, 1H), 2.94 (dd, $J = 5.2$ Hz, 14.4 Hz, 1H), 2.38 (s, 3H), 1.26 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.2, 140.1, 130.4, 128.9, 125.7, 116.4, 115.5, 64.1, 51.1, 42.5, 38.8, 21.0, 13.6; HRMS (ESI/[$\text{M}+\text{H}$] $^+$) m/z calcd. for $\text{C}_{15}\text{H}_{16}\text{N}_2\text{O}_2\text{Cl}$: 291.0900, found: 291.0911; IR (neat) ν 1747 cm^{-1} .

Q-**2c** (18.0 mg, 10 mol%) catalyzed reaction of **3j** (65.7 mg, 0.3 mmol) and **4** (78.8 mg, 0.9 mmol) in toluene (3.0 mL) was run at room temperature for 12 h to furnish the crude product [dr = 13:1, determined by ^1H NMR analysis of the crude product]. The crude product was purified by flash chromatography on silica gel (hexanes/ethyl acetate = 5/1) to give adducts **5j** and **5'j** (dr = 20:1) as a colorless oil in 98% yield. The ee of the major diastereomer was determined to be 95% ee [determined

by HPLC, Chiralcel OD, hexanes: isopropanol = 80: 20, 1.0 mL/min, λ 220 nm, t (major) = 20.60 min, t (minor) = 13.05 min]. ^1H NMR (400 MHz, CDCl_3) δ 7.44 (d, J = 8.4 Hz, 2H), 6.95 (d, J = 8.4 Hz, 2H), 4.57 (dd, J = 5.6 Hz, 8.8 Hz, 1H), 4.34-4.19 (m, 2H), 3.83 (s, 3H), 3.16 (dd, J = 9.6 Hz, 14.4 Hz, 1H), 2.94 (dd, J = 6.0 Hz, 15.2 Hz, 1H), 1.26 (t, J = 7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.2, 160.6, 127.2, 123.6, 116.5, 115.5, 115.0, 64.1, 55.4, 50.8, 42.6, 38.8, 13.6; HRMS (EI/[M^+]) m/z calcd. for $\text{C}_{15}\text{H}_{15}\text{N}_2\text{O}_3\text{Cl}$: 306.0771, found: 306.0773; IR (neat) ν 1748 cm^{-1} .

ent-**5j** was obtained as a colorless oil in 99% yield and 95% ee with 15:1 dr from a reaction catalyzed by QD-**2c** (10 mol%) at room temperature for 12 h.

5k Q-**2c** (18.0 mg, 10 mol%) catalyzed reaction of **3k** (71.7 mg, 0.3 mmol) and **4** (78.8 mg, 0.9 mmol) in toluene (3.0 mL) was run at room temperature for 12 h to furnish the crude product [dr = 14:1, determined by ^1H NMR analysis of the crude product]. The crude product was purified by flash chromatography on silica gel (hexanes/ethyl acetate = 5/1) to give adducts **5k** and **5'k** (dr = 17:1) as a colorless oil in 98% yield. The ee of the major diastereomer was determined to be 95% ee [determined by HPLC, Chiralcel OD, hexanes: isopropanol = 80: 20, 1.0 mL/min, λ 254 nm, t (major) = 90.77 min, t (minor) = 16.58 min]. ^1H NMR (400 MHz, CDCl_3) δ 8.10 (s, 1H), 7.95-7.86 (m, 3H), 7.60-7.51 (m, 3H), 4.58 (dd, J = 5.2 Hz, 8.8 Hz, 1H), 4.36-4.19 (m, 2H), 3.30 (dd, J = 9.2 Hz, 15.6 Hz, 1H), 3.06 (dd, J = 2.8 Hz, 15.2 Hz, 1H), 1.24 (t, J = 7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.0, 133.3, 133.0, 130.0, 129.0, 128.4, 127.7, 127.6, 127.4, 126.2, 122.0, 116.4, 115.5, 64.3, 51.5, 42.3, 38.9, 13.6; HRMS (EI/[M^+]) m/z calcd. for $\text{C}_{18}\text{H}_{15}\text{N}_2\text{O}_2\text{Cl}$: 326.0822, found: 326.0815; IR (neat) ν 1749 cm^{-1} .

5l Q-**2c** (18.0 mg, 10 mol%) catalyzed reaction of **3l** (38.7 mg, 0.3 mmol) and **4** (78.8 mg, 0.9 mmol) in toluene (3.0 mL) was run at room temperature for 12 h to furnish the crude product [dr = 9:1, determined by ^1H NMR analysis of the crude product]. The crude product was purified by flash

chromatography on silica gel (hexanes/ethyl acetate = 8/1) to give adducts **5l** and **5'l** (dr = 11:1) as a colorless oil in 98% yield. The ee of the major diastereomer was determined to be 96% ee [determined by HPLC, Chiralcel OD, hexanes: isopropanol = 90: 10, 1.0 mL/min, λ 220 nm, t (major) = 22.86 min, t (minor) = 20.01 min]. ^1H NMR (400 MHz, CDCl_3) δ 4.71 (dd, J = 6.4 Hz, 8.8 Hz, 1H), 2.89 (dd, J = 8.4 Hz, 14.4 Hz, 1H), 2.55 (dd, J = 6.0 Hz, 14.8 Hz, 1H), 2.44 (s, 3H), 1.73 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 194.8, 117.9, 115.6, 48.5, 42.2, 38.6, 25.2, 12.7; HRMS (EI/[M^+]) m/z calcd. for $\text{C}_8\text{H}_9\text{N}_2\text{OClS}$: 216.0124, found: 216.0120; IR (neat) ν 1684 cm^{-1} .

ent-**5l** was obtained as a colorless oil in 98% yield and 96% ee with 9:1 dr from a reaction catalyzed by QD-**2c** (10 mol%) at room temperature for 12 h.

5'l was obtained as a colorless oil in 44% yield and 88% ee with 7:1 dr from a reaction catalyzed by Q-**1b** (20 mol%) at room temperature for 96 h.

5m Q-**2c** (18.0 mg, 10 mol%) catalyzed reaction of **3m** (46.5 mg, 0.3 mmol) and **4** (78.8 mg, 0.9 mmol) in toluene (3.0 mL) was run at room temperature for 14 h to furnish the crude product [dr = 9:1, determined by ^1H NMR analysis of the crude product]. The crude product was purified by flash chromatography on silica gel (hexanes/ethyl acetate = 6/1) to give adduct **5m** and **5'm** (dr = 12:1) as a colorless oil in 99% yield. The ee of the major diastereomer was determined to be 95% ee [determined by HPLC, Chiralcel OJ, hexanes: isopropanol = 96: 4, 1.0 mL/min, λ 220 nm, t (major) = 65.75 min, t (minor) = 80.75 min]. ^1H NMR (400 MHz, CDCl_3) δ 5.84-5.74 (m, 1H), 5.34 (d, J = 9.6 Hz, 1H), 5.28 (d, J = 17.2 Hz, 1H), 4.70 (dd, J = 5.6 Hz, 9.2 Hz, 1H), 2.87 (dd, J = 9.2 Hz, 14.8 Hz, 1H), 2.75 (dd, J = 6.8 Hz, 14.0 Hz, 1H), 2.55 (m, 2H), 2.43 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 194.3, 128.7, 122.8, 116.7, 115.5, 53.3, 42.6, 40.6, 38.8, 12.8; HRMS (EI/[M^+]) m/z calcd. for $\text{C}_{10}\text{H}_{11}\text{N}_2\text{OClS}$: 242.0281, found: 242.0290; IR (neat) ν 1684 cm^{-1} .

ent-**5m** was obtained as a colorless oil in 98% yield and 95% ee with 10:1 dr from a reaction catalyzed by QD-**2c** (10 mol%) at room temperature for 12 h.

Q-**2c** (24.0 mg, 20 mol%) catalyzed reaction of **3a** (31.4 mg, 0.2 mmol) and **6** (32.0 mg, 0.6 mmol) in toluene was run at room temperature for 12 h to furnish the adduct (-)-**7a** as a white solid in 100% yield after flash chromatography (elution gradient: hexanes/ethyl acetate=2/1) and 93% ee as determined by HPLC analysis [Daicel chiralcel OD, hexanes: isopropanol = 60:40, 1.0 mL/min, λ 220 nm, *t* (major) = 21.50 min, *t* (minor) = 15.66 min]. $[\alpha]_D^{25} = -45.2$ (*c* 0.92, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.84 (d, *J* = 8 Hz, 1H), 7.75 (t, *J* = 7.2 Hz, 1H), 7.52 (m, 2H), 3.76 (d, *J* = 17.6 Hz, 1H), 3.37 (d, *J* = 17.6 Hz, 1H), 2.90-2.82 (m, 1H), 2.78-2.69 (m, 1H), 2.38-2.31 (m, 1H), 2.24-2.17 (m, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 196.3, 150.2, 137.0, 132.4, 129.1, 126.6, 125.8, 118.0, 117.9, 46.5, 38.4, 31.9, 13.6; HRMS (EI/[M⁺]) *m/z* calcd. for C₁₃H₁₀N₂O: 210.0793, found: 210.0789; IR (neat) ν 1734 cm⁻¹.

Q-**2c** (24.0 mg, 20 mol%) catalyzed reaction of **3b** (34.2 mg, 0.2 mmol) and **6** (32.0 mg, 0.6 mmol) in toluene was run at room temperature for 36 h to furnish the adduct (+)-**7b** as a colorless oil in 92% yield after flash chromatography (elution gradient: hexanes/ethyl acetate=4/1) and 94% ee as determined by HPLC analysis [Daicel chiralcel OD, hexanes: isopropanol = 80:20, 1.0 mL/min, λ 220 nm, *t* (major) = 30.63 min, *t* (minor) = 23.87 min]. $[\alpha]_D^{25} = +35.6$ (*c* 0.96, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 8.03 (d, *J* = 7.6 Hz, 1H), 7.59 (t, *J* = 7.6 Hz, 1H), 7.39 (t, *J* = 7.6 Hz, 1H), 7.30 (d, *J* = 8.0 Hz, 1H), 3.39-3.32 (m, 1H), 3.15-3.08 (m, 1H), 2.84-2.76 (m, 1H), 2.73-2.65 (m, 1H), 2.61-2.47 (m, 2H), 2.33-2.26 (m, 1H), 2.23-2.16 (m, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 189.2, 142.3, 135.1, 129.4, 129.0, 128.9, 127.6, 118.3, 117.2, 46.9, 32.3, 29.9, 25.5, 13.7; HRMS (EI/[M⁺]) *m/z* calcd. for C₁₄H₁₂N₂O: 224.0950, found: 224.0947; IR (neat) ν 1700 cm⁻¹.

 Q-2c (24.0 mg, 20 mol%) catalyzed reaction of **3e** (38.0 mg, 0.2 mmol) and **6** (32.0 mg, 0.6 mmol) in toluene was run at room temperature for 96 h to furnish the adduct (-)-**7e** as a colorless oil in 95% yield after flash chromatography (elution gradient: hexanes/ethyl acetate=5/1) and 89% ee as determined by HPLC analysis [Daicel chiralcel OD, hexanes: isopropanol = 80:20, 1.0 mL/min, λ 220 nm, t (major) = 17.64 min, t (minor) = 14.04 min]. $[\alpha]_D^{25} = -23.6$ (c 1.16, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.52-7.43 (m, 5H), 4.26 (m, 2H), 2.76 (m, 1H), 2.55 (m, 2H), 2.39 (m, 1H), 1.25 (t, J = 7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.2, 132.5, 129.6, 125.7, 117.5, 116.8, 63.9, 52.8, 33.6, 13.8, 13.7; HRMS (EI/[M^+]) m/z calcd. for $\text{C}_{14}\text{H}_{14}\text{N}_2\text{O}_2$: 242.1055, found: 242.1063; IR (neat) ν 1734 cm^{-1} .

(+)-**7e** was obtained as a colorless oil in 93% yield and 90% ee from a reaction catalyzed by QD-2c (20 mol%) at room temperature for 96 h.

 Q-2c (24.0 mg, 20 mol%) catalyzed reaction of **3f** (41.4 mg, 0.2 mmol) and **6** (32.0 mg, 0.6 mmol) in toluene was run at room temperature for 120 h to furnish the adduct (-)-**7f** as a colorless oil in 84% yield after flash chromatography (elution gradient: hexanes/ethyl acetate=5/1) and 88% ee as determined by HPLC analysis [Daicel chiralpak AD, hexanes: isopropanol = 98:2, 1.0 mL/min, λ 220 nm, t (major) = 22.86 min, t (minor) = 20.00 min]. $[\alpha]_D^{25} = -22.4$ (c 1.09, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.51 (dd, J = 4.8 Hz, 9.2 Hz, 2H), 7.16 (t, J = 8.0 Hz, 2H), 4.34-4.21 (m, 2H), 2.80-2.71 (m, 1H), 2.60-2.39 (m, 3H), 1.26 (t, J = 7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.1, 163.2 (d, J = 248.9 Hz), 128.4 (d, J = 3.8 Hz), 127.8 (d, J = 8.4 Hz), 117.3, 116.7 (d, J = 22.1 Hz), 116.7, 64.0, 52.3, 33.7, 13.8, 13.7; HRMS (EI/[M^+]) m/z calcd. for $\text{C}_{14}\text{H}_{13}\text{N}_2\text{O}_2\text{F}$: 260.0961, found: 260.0966; IR (neat) ν 1747 cm^{-1} .

(+)-**7f** was obtained as a colorless oil in 87% yield and 90% ee from a reaction catalyzed by QD-2c (20 mol%) at room temperature for 72 h.

Q-2c (24.0 mg, 20 mol%) catalyzed reaction of **3g** (44.7 mg, 0.2 mmol) and **6** (32.0 mg, 0.6 mmol) in toluene was run at room temperature for 52 h to furnish the adduct (-)-**7g** as a colorless oil in 96% yield after flash chromatography (elution gradient: hexanes/ethyl acetate=5/1) and 88% ee as determined by HPLC analysis [Daicel chiralcel OJ, hexanes: isopropanol = 60:40, 0.8 mL/min, λ 220 nm, t (major) = 38.01 min, t (minor) = 28.57 min]. $[\alpha]_D^{25} = -15.0$ (c 1.07, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.44 (m, 4H), 4.32-4.22 (m, 2H), 2.79-2.71 (m, 1H), 2.60-2.38 (m, 3H), 1.26 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 165.9, 135.9, 131.1, 129.8, 127.2, 117.3, 116.5, 64.1, 52.4, 33.5, 13.8, 13.7; HRMS (EI/[M^+]) m/z calcd. for $\text{C}_{14}\text{H}_{13}\text{N}_2\text{O}_2\text{Cl}$: 276.0666, found: 276.0671; IR (neat) ν 1733 cm^{-1} .

(+)-**7g** was obtained as a colorless oil in 96% yield and 89% ee from a reaction catalyzed by **QD-2c** (20 mol%) at room temperature for 52 h.

Q-2c (24.0 mg, 20 mol%) catalyzed reaction of **3h** (53.6 mg, 0.2 mmol) and **6** (32.0 mg, 0.6 mmol) in toluene was run at room temperature for 72 h to furnish the adduct (-)-**7h** as a colorless oil in 89% yield after flash chromatography (elution gradient: hexanes/ethyl acetate=5/1) and 89% ee as determined by HPLC analysis [(R, R)-Whelk-O 1, hexanes: isopropanol = 97:3, 1.0 mL/min, λ 220 nm, t (major) = 18.39 min, t (minor) = 22.44 min]. $[\alpha]_D^{25} = -11.2$ (c 0.98, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.59 (d, $J = 8.8$ Hz, 2H), 7.39 (d, $J = 8.8$ Hz, 2H), 4.34-4.20 (m, 2H), 2.79-2.71 (m, 1H), 2.61-2.38 (m, 3H), 1.26 (t, $J = 7.2$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 165.8, 132.8, 131.7, 127.5, 124.1, 117.3, 116.4, 64.2, 52.5, 33.5, 13.8, 13.7; HRMS (EI/[M^+]) m/z calcd. for $\text{C}_{14}\text{H}_{13}\text{N}_2\text{O}_2\text{Br}$: 320.0160, found: 320.0167; IR (neat) ν 1747 cm^{-1} .

(+)-**7h** was obtained as a colorless oil in 82% yield and 89% ee from a reaction catalyzed by **QD-2c** (20 mol%) at room temperature for 72 h.

Q-**2c** (24.0 mg, 20 mol%) catalyzed reaction of **3i** (40.6 mg, 0.2 mmol) and **6** (32.0 mg, 0.6 mmol) in toluene was run at room temperature for 114 h to furnish the adduct (-)-**7i** as a colorless oil in 92% yield after flash chromatography (elution gradient: hexanes/ethyl acetate=5/1) and 90% ee as determined by HPLC analysis [Daicel chiralcel OJ, hexanes: isopropanol = 60:40, 1.0 mL/min, λ 220 nm, t (major) = 30.98 min, t (minor) = 21.24 min]. $[\alpha]_D^{25} = -19.5$ (c 0.91, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.37 (d, J = 8.0 Hz, 2H), 7.24 (d, J = 8.8 Hz, 2H), 4.31-4.20 (m 2H), 2.76-2.70 (m, 1H), 2.55-2.47 (m, 2H), 2.41-2.35 (m, 1H), 2.37 (s, 3H), 1.25 (t, J = 7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.3, 139.7, 130.2, 129.5, 125.6, 117.6, 117.0, 63.8, 52.5, 33.6, 21.0, 13.70, 13.68; HRMS (EI/[M^+]) m/z calcd. for $\text{C}_{15}\text{H}_{16}\text{N}_2\text{O}_2$: 256.1212, found: 256.1227; IR (neat) ν 1749 cm^{-1} .

Q-**2c** (24.0 mg, 20 mol%) catalyzed reaction of **3j** (44.0 mg, 0.2 mmol) and **6** (32.0 mg, 0.6 mmol) in toluene was run at room temperature for 117 h to furnish the adduct (-)-**7j** as a colorless oil in 85% yield after flash chromatography (elution gradient: hexanes/ethyl acetate=5/1) and 90% ee as determined by HPLC analysis [Daicel chiralcel OD, hexanes: isopropanol = 80:20, 1.0 mL/min, λ 220 nm, t (major) = 17.34 min, t (minor) = 23.57 min]. $[\alpha]_D^{25} = -18.3$ (c 0.98, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.41 (d, J = 8.4 Hz, 2H), 6.94 (d, J = 8.4 Hz, 2H), 4.28-4.23 (m, 2H), 3.83 (s, 3H), 2.72 (m, 1H), 2.55-2.50 (m, 2H), 2.42-2.37 (m, 1H), 1.25 (t, J = 7.2 Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 166.4, 160.3, 127.1, 124.2, 117.6, 117.1, 114.8, 63.8, 55.4, 52.2, 33.6, 13.7; HRMS (EI/[M^+]) m/z calcd. for $\text{C}_{15}\text{H}_{16}\text{N}_2\text{O}_3$: 272.1161, found: 272.1163; IR (neat) ν 1734 cm^{-1} .

Q-**2c** (24.0 mg, 20 mol%) catalyzed reaction of **3k** (47.8 mg, 0.2 mmol) and **6** (32.0 mg, 0.6 mmol) in toluene was run at room temperature for 90 h to furnish the adduct (-)-**7k** as a colorless oil in 89% yield after flash chromatography (elution gradient: hexanes/ethyl acetate=5/1) and 89% ee as determined by HPLC analysis [(R, R)-Whelk-O 1, hexanes: isopropanol = 80:20, 1.0 mL/min, λ 280 nm, t (major) = 9.57 min, t (minor) = 12.38 min]. $[\alpha]_D^{25} = -$

27.0 (c 1.07, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 8.04 (s, 1H), 7.93-7.86 (m, 3H), 7.59-7.49 (m, 3H), 4.33-4.21 (m, 2H), 2.89-2.82 (m, 1H), 2.68-2.54 (m, 2H), 2.42-2.35 (m, 1H), 1.24 (t, *J* = 7.2 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 166.2, 133.2, 133.0, 129.9, 129.6, 128.3, 127.7, 127.5, 127.3, 126.0, 122.0, 117.6, 116.9, 64.0, 53.0, 33.5, 13.8, 13.7; HRMS (EI/[M⁺]) *m/z* calcd. for C₁₈H₁₆N₂O₂: 292.1212, found: 292.1208; IR (neat) ν 1749 cm⁻¹.

Q-2c (24.0 mg, 20 mol%) catalyzed reaction of **3l** (25.8 mg, 0.2 mmol) and **6** (32.0 mg, 0.6 mmol) in toluene was run at room temperature for 72 h to furnish the adduct (-)-**7l** as a colorless oil in 80% yield after flash chromatography (elution gradient: hexanes/ethyl acetate=4/1) and 93% ee as determined by HPLC analysis [Daicel chiralpak AS, hexanes: isopropanol = 80:20, 1.0 mL/min, λ 220 nm, *t* (major) = 19.79 min, *t* (minor) = 24.11 min]. [α]_D²⁵ = -5.1 (c 1.07, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 2.64-2.39 (m, 3H), 2.43 (s, 3H), 2.19-2.11 (m, 1H), 1.68 (s, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 195.1, 118.6, 117.5, 49.8, 33.5, 24.2, 13.7, 12.6; HRMS (ESI/[M+H]⁺) *m/z* calcd. for C₈H₁₁N₂OS: 183.0592, found: 183.0597; IR (neat) ν 1684 cm⁻¹.

Q-2c (24.0 mg, 20 mol%) catalyzed reaction of **3m** (31.0 mg, 0.2 mmol) and **6** (32.0 mg, 0.6 mmol) in toluene was run at room temperature for 88 h to furnish the adduct (-)-**7m** as a colorless oil in 82% yield after flash chromatography (elution gradient: hexanes/ethyl acetate=5/1) and 91% ee as determined by HPLC analysis [Daicel chiralpak AS, hexanes: isopropanol = 85:15, 1.0 mL/min, λ 220 nm, *t* (major) = 21.38 min, *t* (minor) = 18.55 min]. [α]_D²⁵ = -35.5 (c 0.93, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 5.81-5.72(m, 1H), 5.30 (d, *J* = 10.4 Hz, 1H), 5.26 (d, *J* = 16.8 Hz, 1H), 2.73 (m, 1H), 2.64-2.34 (m, 4H), 2.43 (s, 3H), 2.20-2.12 (m, 1H); ¹³C NMR (100 MHz, CDCl₃) δ 194.4, 129.1, 122.2, 117.6, 117.5, 54.9, 41.8, 32.0, 13.6, 12.7; HRMS (EI/[M⁺]) *m/z* calcd. for C₁₀H₁₂N₂OS: 208.0670, found: 208.0680; IR (neat) ν 1684 cm⁻¹.

Absolute configuration determination of compounds **5a**, **5l**, **7a**, **7l**.

As determined previously by us,¹¹ compound **ent-5'a** (obtained from **QD-1a** catalyzed reaction of **3a** and **4**) has the absolute configuration of (1*R*, 3*S*) as shown above. Compound **ent-5'l** (obtained from **QD-1b** catalyzed reaction of **3l** and **4**) has the absolute configuration of (1*R*, 3*S*) as shown above.¹¹

To a solution of **ent-5'a** (24.5 mg, 0.1 mmol) in diethyl ether (0.5 mL) was added acetic acid (0.5 mL) and zinc powder (26.2 mg, 0.4 mmol). The resulting mixture was stirred at room temperature for 20 h. TLC analysis showed **ent-5'a** was completely consumed. The reaction mixture was diluted with diethyl ether (5 mL), washed with water (5 mL × 3) and brine, dried over anhydrous Na₂SO₄. Purification by flash chromatography on silica gel (hexanes/ethyl acetate = 2/1) afforded dechlorination product **(-)-7a** in quantitative yield. Thus **(-)-7a** has the absolute configuration of *R*. According to the same procedure, the dechlorination of **5a** (obtained from **Q-2c** catalyzed reaction of **3a** and **4**) was performed. The dechlorination product was found to be **(-)-7a** as shown by comparison with the

dechlorination product obtained from *ent*-**5'a** via HPLC analysis, thus **5a** was determined to have the absolute configuration of (1*S*, 3*S*). The respective dechlorination of *ent*-**5'1** and **51** afforded the same product (-)-**71** by the same procedures, thus (-)-**71** has the absolute configuration of *S*, and **51** has the absolute configuration of (1*S*, 3*S*).

Formal synthesis of Manzacidin C.^{12,13}

To a solution of (1R,3R)-**5I** (1.07 g, 4.94 mmol, dr = 90:10) in DMSO (25 mL) was added NaN_3 (482 mg, 7.41 mmol) at room temperature. After stirring at the same temperature for 42 h, the reaction mixture was diluted with water (50 mL), and extracted with diethyl ether (2×50 mL). The combined organic layers were washed with water (4×50 mL) and brine, dried over anhydrous Na_2SO_4 . After removal of the solvent under reduced

pressure, the residue was subjected to flash chromatography on silica gel (hexanes /ethyl acetate=5/1) to afford **9** as a colorless oil (573 mg, 52% yield) in 89:11 dr. ^1H NMR (400 MHz, CDCl_3) δ 4.42 (dd, J = 5.6 Hz, 8.0 Hz, 1H), 2.52 (dd, J = 6.0 Hz, 14.8 Hz, 1H), 2.41 (s, 3H), 2.24 (dd, J = 8.4 Hz, 14.4 Hz, 1H), 1.69 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 195.0, 118.3, 114.8, 47.9, 47.7, 39.6, 25.4, 12.7; HRMS ($\text{CI}/[\text{M}+\text{H}]^+$) m/z calcd. for $\text{C}_8\text{H}_{10}\text{ON}_5\text{S}$: 224.0606, found: 224.0605; IR (neat) ν 1682, 2117 cm^{-1} .

A solution of **9** (705 mg, 3.16 mmol) and TMSCl (1.45 mL) in anhydrous MeOH (6.6 mL) was kept at 0 $^\circ\text{C}$ without stirring. After 48 h, the reaction mixture was diluted with water (30 mL), extracted with ethyl acetate (3 $\times$ 30 mL). The combined organic layers were washed with brine, dried over anhydrous Na_2SO_4 . After removal of the solvent under reduced pressure, the residue was subjected to flash chromatography on silica gel (hexanes /ethyl acetate=5/1) to afford **10** as a colorless oil (753 mg, 93% yield) in 83:17 dr. ^1H NMR (400 MHz, CDCl_3) δ 4.07 (dd, J = 4.0 Hz, 9.6 Hz, 1H), 3.83 (s, 3H), 2.55 (dd, 4.0 Hz, 14.4 Hz, 1H), 2.42 (s, 3H), 2.15 (dd, J = 9.2 Hz, 14.8 Hz, 1H), 1.71 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 195.8, 169.3, 119.0, 58.7, 53.0, 48.2, 38.2, 25.3, 12.5; HRMS ($\text{CI}/[\text{M}+\text{H}]^+$) m/z calcd. for $\text{C}_9\text{H}_{13}\text{O}_3\text{N}_4\text{S}$: 257.0708, found: 257.0711; IR (neat) ν 1682, 1747, 2114 cm^{-1} .

To a solution of **10** (666 mg, 2.60 mmol) in EtOH was added $\text{Hg}(\text{OAc})_2$ (828.5 mg, 2.60 mmol) at 0 $^\circ\text{C}$. The resulting mixture was stirred for 10 mins, and NaBH_4 (786 mg, 20.8

mmol) was added in portions. After 20 min, the reaction was quenched with saturated aq. NH_4Cl solution, and the resulting mixture was extracted with ethyl acetate (4×50 mL). The combined organic layers were washed brine, dried over anhydrous Na_2SO_4 . After removal of the solvent under reduced pressure, the residue was subjected to flash chromatography on silica gel (hexanes /ethyl acetate=1/1) to afford **11** as a colorless oil (461 mg, 96% yield) in 86:14 dr. ^1H NMR (400 MHz, CD_3OD) δ 3.72-3.51 (m, 5H), 1.82 (dd, J = 2.4 Hz, 14.4 Hz, 1H), 1.49 (dd, J = 9.6 Hz, 14.8 Hz, 1H), 1.38 (s, 3H); ^{13}C NMR (100 MHz, CD_3OD) δ 125.5, 69.8, 67.9, 64.0, 40.9, 38.5, 23.7; HRMS ($\text{CI}/[\text{M}+\text{H}]^+$) m/z calcd. for $\text{C}_7\text{H}_{13}\text{O}_2\text{N}_4$: 185.1039, found: 185.1040; IR (neat) ν 2115, 3418 cm^{-1} .

To a solution of **11** (396 mg, 2.15 mmol) and imidazole (442 mg, 6.46 mmol) in anhydrous DMF (3.0 mL) was added *tert*-butyl chlorodiphenylsilane (1.22 mL, 4.73 mmol) at room temperature. After stirring at the same temperature for 12 h, the reaction was quenched with water (30 mL). The reaction mixture was extracted with diethyl ether (3×30 mL). The combined organic layers were washed with water (3×30 mL) and brine, dried over anhydrous Na_2SO_4 . After removal of the solvent under reduced pressure, the residue was subjected to flash chromatography on silica gel (hexanes /ethyl acetate=20/1) to afford **12** as a colorless oil (1.41 g, 99% yield) in 86:14 dr. ^1H NMR (400 MHz, CDCl_3) δ 7.68-7.63 (m, 8H), 7.46-7.38 (m, 12H), 3.74-3.52 (m, 5H), 1.81 (d, J = 14.8 Hz, 1H), 1.54-1.48 (m, 1H), 1.41 (s, 3H), 1.07 (s, 18H); ^{13}C NMR (100 MHz, CDCl_3) δ 139.9, 135.61, 135.57, 135.54, 132.6, 132.4, 132.3, 130.04, 130.01, 129.9, 127.88, 127.86, 122.5, 68.9, 67.6, 60.6, 38.5, 35.9, 26.71, 26.66, 21.5, 19.3, 19.1; HRMS ($\text{ESI}/[\text{M}+\text{H}]^+$) m/z calcd. for $\text{C}_{39}\text{H}_{49}\text{O}_2\text{N}_4\text{Si}_2$: 661.3394, found: 661.3371; IR (neat) ν 2116 cm^{-1} .

To a solution of **12** (465 mg, 0.7 mmol) in EtOH (2.0 mL) was added [PtH(PMe₂OH)(PMe₂O)₂H] (15.7 mg, 0.035 mmol). The resulting mixture was stirred at 80 °C for 12 h. After cooled to room temperature, the reaction mixture was diluted with water (20 mL), and extracted with EtOAc (3×20 mL). The combined organic layers were washed with brine, dried over anhydrous Na₂SO₄. After removal of the solvent under reduced pressure, the residue was subjected to flash chromatography on silica gel (hexanes /ethyl acetate=4/1) to afford **13** as a colorless oil (460 mg, 97% yield) in 86:14 dr. ¹H NMR (400 MHz, CDCl₃) δ 7.68-7.63 (m, 8H), 7.43-7.35 (m, 12H), 6.57 (brs, 1H), 6.00 (brs, 1H), 3.71-3.53 (m, 5H), 1.78 (d, *J* = 14.0 Hz, 1H), 1.39 (dd, *J* = 9.2 Hz, 14.4 Hz, 1H), 1.23 (s, 3H), 1.07 (s, 18H); ¹³C NMR (100 MHz, CDCl₃) δ 178.2, 135.53, 135.50, 132.80, 132.76, 132.4, 132.3, 129.9, 129.7, 127.8, 127.7, 69.3, 68.0, 60.4, 46.3, 35.7, 26.8, 26.6, 19.6, 19.2, 19.0; HRMS (ESI/[M+H]⁺) *m/z* calcd. for C₃₉H₅₁O₃N₄Si₂: 679.3500, found: 679.3472; IR (neat) ν 1676 cm⁻¹.

To a suspension of **13** (1.172 g, 1.73 mmol), 10% palladium on carbon (150 mg) in EtOH (20 mL) was added Boc₂O (755 mg, 3.46 mmol). The resulting mixture was stirred under H₂ atmosphere (1 atm) for 23 at room temperature, followed by filtration on Cellite[®] 521. The filtrate was concentrated in vacuo, and then subjected to flash chromatography on silica gel (hexanes /ethyl acetate=5/1) to afford **14** as a colorless oil (900 mg, 69% yield). The dr of **14** can not be determined by ¹H NMR. ¹H NMR (400 MHz, CDCl₃) δ 7.66-7.60

(m, 8H), 7.44-7.34 (m, 12H), 6.88 (brs, 1H), 5.50 (brs, 1H), 4.73 (d, $J = 9.2$ Hz, 1H), 3.98 (d, $J = 10.4$ Hz, 1H), 3.73 (m, 1H), 3.51 (m, 3H), 2.00-1.93 (m, 1H), 1.60 (d, $J = 12.4$ Hz, 1H), 1.41 (s, 9H), 1.21 (s, 3H), 1.08 (s, 9H), 1.03 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ 179.2, 155.2, 135.6, 135.5, 133.3, 133.2, 132.5, 129.9, 129.7, 129.6, 127.8, 127.8, 127.7, 127.6, 78.8, 48.6, 45.8, 37.5, 28.4, 28.2, 26.9, 26.8, 21.9, 19.3, 19.2; HRMS (ESI/[$\text{M}+\text{H}$] $^+$) m/z calcd. for $\text{C}_{44}\text{H}_{61}\text{O}_5\text{N}_2\text{Si}_2$: 753.4119, found: 753.4099; IR (neat) ν 1678, 1710 cm^{-1} .

To a solution of **14** (247 mg, 0.33 mmol) in $^t\text{BuOH}$ (8.0 mL) was added Pb(OAc)_4 (291 mg, 0.66 mmol) at room temperature. The resulting suspension was heated to reflux for 3 h. After cooled to room temperature, the reaction mixture was diluted with diethyl ether (30 mL), and quenched with saturated aq. NaHCO_3 solution (20 mL). The aqueous phase was extracted with diethyl ether (3 $\times$ 20 mL). The combined organic layers were washed with brine and dried over anhydrous Na_2SO_4 . After removal of the solvent under reduced pressure, the residue was subjected to flash chromatography on silica gel (hexanes /ethyl acetate=15/1) to afford **15** as a white amorphous solid (152 mg, 56% yield) in 80:20 dr. ^1H NMR (400 MHz, CDCl_3) δ 7.64 (m, 8H), 7.42-7.35 (m, 10H), 4.96 (brs, 1H), 4.67 (d, $J = 8.4$ Hz, 1H), 3.82 (m, 1H), 3.73-3.57 (m, 4H), 2.08 (d, $J = 14.0$ Hz, 1H), 1.76-1.65 (m, 1H), 1.39 (s, 9H), 1.38 (s, 9H), 1.29 (s, 3H), 1.07 (s, 9H), 1.06 (s, 9H); ^{13}C NMR (100 MHz, CDCl_3) δ 155.1, 154.8, 135.6, 133.32, 133.26, 133.20, 129.7, 127.7, 78.9, 68.4, 66.9, 55.6, 48.5, 37.5, 28.4, 26.92, 26.88, 19.4, 19.3; HRMS (ESI/[$\text{M}+\text{H}$] $^+$) m/z calcd. for $\text{C}_{48}\text{H}_{69}\text{O}_6\text{N}_2\text{Si}_2$: 825.4694, found: 825.4672; IR (neat) ν 1714 cm^{-1} .

To a solution of **15** (254 mg, 0.31 mmol, dr = 80:20) in THF was added TBAF (330 mg, 1.26 mmol) at room temperature. After stirring for 1 h, the reaction was quenched with saturated aq. NH_4Cl solution. The resulting mixture was diluted with brine (30 mL), and extracted with EtOAc (3×30 mL). The combined organic layers were washed with brine and dried over anhydrous Na_2SO_4 . After removal of the solvent under reduced pressure, the residue was subjected to flash chromatography on silica gel (hexanes /ethyl acetate=1/1) to afford **16** in **diastereomerically pure form** as a colorless oil (70 mg, 65% yield). The ee of **16** was determined by transforming the diol to diester **17**. $[\alpha]_{\text{D}}^{25} = +1.8$ (c 1.07, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 5.41 (s, 1H), 5.16 (s, 1H), 4.64 (brs, 1H), 3.70–3.62 (m 1H), 3.60–3.50 (m, 4H), 3.29 (brs, 1H), 1.98 (dd, $J = 3.6$ Hz, 14.8 Hz, 1H), 1.81 (dd, $J = 7.6$ Hz, 14.4 Hz, 1H), 1.40 (s, 9H), 1.39 (s, 9H), 1.19 (s, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 156.53, 156.24, 79.87, 79.71, 69.19, 66.40, 55.99, 49.14, 36.68, 28.37, 28.32, 22.73; HRMS (ESI/[$\text{M}+\text{H}$] $^+$) m/z calcd. for $\text{C}_{16}\text{H}_{33}\text{O}_6\text{N}_2$: 349.2339, found: 349.2337; IR (neat) ν 1694, 3352 cm^{-1} . The analytic data of **16** was consistent with the reported data of the intermediate for the synthesis of manzacidin C¹² (S-Table 2).

To a solution of **16** (20 mg, 0.057 mmol) in CH_2Cl_2 (0.5 mL) was added pyridine (23 mg, 0.29 mmol), DMAP (2.0 mg), and benzoyl chloride (32 mg, 0.23 mmol) sequentially. The resulting solution was stirred at room temperature for 1 h. The reaction mixture was diluted with EtOAc (20 mL), washed with water (10 mL) and brine, dried over anhydrous

Na₂SO₄. After removal of the solvent under reduced pressure, the residue was subjected to flash chromatography on silica gel (hexanes /ethyl acetate=5/1) to afford **17** as a white amorphous solid (27.6 mg, 86% yield). The ee of **17** was determined to be 94% ee [determined by HPLC, Chiralcel OD, hexanes: isopropanol = 97: 3, 1.0 mL/min, λ 220 nm, t (major) = 16.79 min, t (minor) = 13.68 min]. $[\alpha]_D^{25} = -1.1$ (*c* 1.10, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 8.03 (m, 4H), 7.58-7.54 (m, 2H), 7.47-7.40 (m, 4H), 4.90 (s, 1H), 4.78 (d, *J* = 8.0 Hz, 1H), 4.54 (d, *J* = 10.8 Hz, 1H), 4.37 (d, *J* = 11.2 Hz, 1H), 4.29 (m, 3H), 2.34 (d, *J* = 14.0 Hz, 1H), 1.88 (dd, *J* = 8.0 Hz, 14.8 Hz, 1H), 1.43 (s, 3H), 1.41 (s, 9H), 1.39 (s, 9H); ¹³C NMR (100 MHz, CDCl₃) δ 166.4, 166.1, 155.2, 154.5, 133.13, 133.08, 129.69, 129.61, 128.42, 128.33, 79.7, 68.5, 67.7, 54.4, 46.0, 37.2, 28.32, 28.27, 22.8; HRMS (ESI/[M+H]⁺) *m/z* calcd. for C₃₀H₄₁O₈N₂: 557.2863, found: 557.2851; IR (neat) ν 1714 cm⁻¹.

S-Table 2. Comparison of characteristic data of **16** with that of literature reported¹².

16		Manzacidin C intermediate, ref.12
Optical rotation	$[\alpha]_D^{25} = + 1.8$ (<i>c</i> 1.07, CHCl ₃)	$[\alpha]_D^{20} = + 3.9$ (<i>c</i> 1.30, CHCl ₃)
¹ H NMR (400 MHz, CDCl ₃)	5.41 (s, 1H) 5.16 (s, 1H) 4.64 (brs, 1H) 3.70-3.62 (m, 1H) 3.60-3.50 (m, 4H) 3.29 (brs, 1H) 1.98 (dd, <i>J</i> = 3.6 Hz, 14.8 Hz, 1H) 1.81 (dd, <i>J</i> = 7.6 Hz, 14.4 Hz, 1H) 1.40 (s, 9H) 1.39 (s, 9H) 1.19 (s, 3H)	5.40 (s, 1H) 5.23 (s, 1H) 3.66 (m, 1H) 3.60-3.50 (m, 5H) 1.98 (dd, <i>J</i> = 3.8 Hz, 14.5 Hz, 1H) 1.81 (dd, <i>J</i> = 7.8 Hz, 14.5 Hz, 1H) 1.41 (s, 9H) 1.35 (s, 9H) 1.20 (s, 3H)
¹³ C NMR (100 MHz, CDCl ₃)	156.53 156.24 79.87 79.71 69.19 66.40 55.99 49.14 36.68 28.37 28.32 22.73	156.43 156.18 79.78 79.69 69.05 66.21 55.93 49.11 36.80 28.37 28.33 22.65

References:

- 1) Johnson, W. S.; Anderson, J. M.; Sheler, W. E. *J. Am. Chem. Soc.* **1944**, *66*, 218.
- 2) Johnson, W. S.; Sheler, W. E. *J. Am. Chem. Soc.* **1945**, *67*, 1745
- 3) Hirota, T.; Ieno, K.; Sasaki, K. *J. Heterocyclic Chem.* **1986**, *23*, 1685.
- 4) Liu, H.-J.; Ly, T.-W.; Tai, C.-L.; Wu, J.-D.; Liang, J.-K.; Guo, J.-C.; Tseng, N.-W.; Shia, K.-S. *Tetrahedron* **2003**, *59*, 1209.
- 5) (a) Kaiser, E. M.; Solter, L. E.; Schwarz, R. A.; Beard, R. D.; Hause, C. R. *J. Am. Chem. Soc.* **1971**, *93*, 4237; (b) Albarella, J. M. *J. Org. Chem.* **1977**, *42*, 2009.
- 6) Yokoyama, M.; Hatanaka, H.; Sasaki, A.; Shiraishi, T.; Kumata, K.; Sakamoto, K.; Ogata, K. *J. Chem. Soc., Perkin I* **1986**, 1197.
- 7) (a) Samii, Z. K. M. A. E.; Ashmawy, M. I. A.; Mellor, J. M. *J. Chem. Soc., Perkin I* **1988**, 2523. (b) Jevric, M.; Taylor, D. K.; Greatrex, B. W.; Tiekink, E. R. T. *Tetrahedron* **2005**, *61*, 1885. (c) Heaney, F.; Fenlon, J.; McArdle, P.; Cunningham, D. *Org. Biomol. Chem.* **2003**, *1*, 1122.
- 8) Hiratake, J.; Inagaki, M.; Yamamoto, Y.; Oda, J. *J. Chem. Soc., Perkin I* **1987**, 1053.
- 9) (a) Li, H.; Wang, Y.; Tang, L.; Deng, L. *J. Am. Chem. Soc.* **2004**, *126*, 9906. (b) Li, H.; Wang, Y.; Tang, L.; Wu, F.; Liu, X.; Guo, C.; Foxman, B. M.; Deng, L. *Angew. Chem. Int. Ed.* **2005**, *44*, 105.
- 10) (a) Vakulya, B.; Varga, S.; Csámpai, A.; Soós, T. *Org. Lett.* **2005**, *7*, 1967. (b) Li, B.; Jiang, L.; Liu, M.; Chen, Y.; Ding, L.; Wu, Y. *Synlett* **2005**, *4*, 603.
- 11) Wang, Y.; Liu, X.; Deng, L. *J. Am. Chem. Soc.* **2006**, *128*, 3928.
- 12) Namba, K.; Shinada, T.; Teramoto, T.; Ohfun, Y. *J. Am. Chem. Soc.* **2000**, *122*, 10708.
- 13) For other asymmetric total syntheses of manzacidins, see: (a) Wehn, P. M.; Du Bois, J. *J. Am. Chem. Soc.* **2002**, *124*, 12950. (b) Lanter, J. C.; Chen, H.; Zhang, X.; Sui, Z. *Org. Lett.* **2005**, *7*, 5905. (c) Kano, T.; Hashimoto, T.; Maruoka, K. *J. Am. Chem. Soc.* **2006**, *128*, 2174.

HPLC conditions for ee determination: Chiralpak AD $\times$ 2, hexanes: isopropanol = 80: 20, 1.0 mL/min, 220 nm.

97% ee, product obtained with Q-2c

97% ee, product obtained with QD-2c

HPLC conditions for ee determination: (R, R)- Whelk-O 1, hexanes: isopropanol = 98: 2, 0.6 mL/min, 220 nm.

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU	Area %	Height [mAU]	Area %
1	53.111	BV	1.8676	7330.52148	56.33213	20.8914	
2	58.030	VV	2.1332	9827.24512	64.27335	28.0069	
3	63.513	VV	2.0080	6946.03174	52.54915	19.7956	
4	66.780	VB	2.7595	1.09849e4	54.76469	31.3061	
Totals :				3.50887e4	227.91932

99% ee, product obtained with Q-2c

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU	Area %	Height [mAU]	Area %
1	46.042	BB	1.2665	252.45657	2.98434	0.5965	
2	56.110	BV	1.5212	554.62823	5.41448	1.3104	
3	60.277	VV	3.0400	3.96997e4	176.73721	93.7944	
4	70.104	VB	2.5725	1819.52954	9.96659	4.2988	
Totals :				4.23263e4	195.10262

97% ee, product obtained With QD-2c

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU	Area %	Height [mAU]	Area %
1	47.316	BV	2.8348	7.46090e4	360.36267	91.1398	
2	57.177	VB	2.2946	4025.65603	25.04466	4.9176	
3	64.157	BV	1.7517	1084.33948	9.49452	1.3246	
4	67.129	VB	2.4347	2143.19824	12.70024	2.6181	
Totals :				8.18622e4	407.60209		

HPLC conditions for ee determination: Chiralcel OD, hexanes: isopropanol = 90: 10, 1.0 mL/min, 220 nm.

Racemic-5c

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	25.464	BV	0.8619	1.61880e4	294.80591	28.0968
2	28.377	VV	0.9922	1.62526e4	253.64578	28.2089
3	32.668	VB	1.1247	1.25671e4	171.12521	21.8122
4	43.994	BB	1.5200	1.26074e4	128.21075	21.8821
Totals				5.76151e4	847.78766

5c

97% ee, product obtained
with Q-2c

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	25.676	BV	0.8301	3289.41357	61.85086	4.7249
2	28.681	VB	0.9756	4136.66162	64.98447	5.9419
3	33.037	BB	1.0952	1042.00769	14.69447	1.4967
4	43.665	BB	1.5924	6.11507e4	596.32397	87.8365
Totals				6.96187e4	737.85377

ent-5c

96% ee, product obtained
with QD-2c

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	25.622	BV	0.8348	3787.75879	70.49281	7.6077
2	28.611	VV	0.9393	2300.93311	38.00307	4.6214
3	32.103	VB	1.1983	4.27195e4	542.48987	85.8023
4	43.884	BB	1.4574	980.09027	10.32047	1.9685
Totals				4.97883e4	661.50622	

GC conditions for ee determination: β -CD, 100 °C, 30 min, then 100 °C-200 °C (0.5 °C/min).

Racemic-**5d**

Error storing signal to M:SIGNAL .BNA
 DIRECTORY FULL
 Storing processed peaks to M:SIGNAL .PRA

RUN# 6922 SEP 18, 2006 20:39:40

AREA#	RT	AREA TYPE	WIDTH	AREAX
179.895	8884	UU	.494	49.32816
180.556	4594	UU	.316	25.50806
181.157	4532	UP	.305	25.16380

TOTAL AREA= 18010
 MUL FACTOR=1.0000E+00

5d

94% ee, product obtained
 with Q-2c

Error storing signal to M:SIGNAL .BNA
 DIRECTORY FULL
 Storing processed peaks to M:SIGNAL .PRA

RUN# 6928 SEP 23, 2006 20:06:57

AREA#	RT	AREA TYPE	WIDTH	AREAX
179.845	3280	BU	.441	8.66899
180.520	1104	UU	.347	2.91786
181.602	33452	UP	.435	88.41315

TOTAL AREA= 37836
 MUL FACTOR=1.0000E+00

ent-**5d**

98% ee, product obtained
 with QD-2c

Error storing signal to M:SIGNAL .BNA
 DIRECTORY FULL
 Storing processed peaks to M:SIGNAL .PRA

RUN# 6926 SEP 23, 2006 15:52:06

AREA#	RT	AREA TYPE	WIDTH	AREAX
3.664	13	P8	.054	.16018
88.461	93	UP	.310	1.14580
179.685	150	P8	.270	1.04020
180.640	7777	BU	.307	95.82307
181.082	83	UP	.198	1.02267

TOTAL AREA= 8116
 MUL FACTOR=1.0000E+00

HPLC conditions for ee determination: Chiralcel OJ, hexanes: isopropanol = 70: 30, 1.0 mL/min, 220 nm.

85% ee, product from 5'd

91% ee, product from ent-5'd

HPLC conditions for ee determination: Chiralcel OD, hexanes: isopropanol = 80: 20, 1.0 mL/min, 220 nm.

Racemic-5e

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU	*s	Height [mAU]	Area %
1	9.421	VV	0.3308	9397.56152		439.80234	19.2999
2	11.935	VV	0.4375	1.48072e4		519.28290	30.4003
3	15.270	VV	0.5577	9431.16016		256.11746	19.3629
4	68.830	BB	2.6325	1.50715e4		88.79630	30.9430
Totals :				4.87074e4		1303.99900

5e

95% ee, product obtained with Q-2c

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU	*s	Height [mAU]	Area %
1	9.433	BB	0.3030	872.06958		43.64915	2.4430
2	12.037	BB	0.4117	841.08429		31.97898	2.3562
3	15.439	BB	0.5415	757.05945		21.98342	2.1208
4	70.071	BB	2.7259	3.32263e4		176.99734	93.0800
Totals :				3.56966e4		274.60890

ent-5e

95% ee, product obtained
with QD-2c

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU *s	Height [mAU]	Area %
1	9.624	BB	0.3425	862.20917	40.35199	2.7957
2	12.227	BB	0.5066	2.85182e4	852.91779	92.4681
3	16.064	BB	0.5761	751.52026	20.09534	2.4367
4	70.999	BB	2.5018	709.18103	4.20853	2.2995

Totals : 3.08411e4 917.57365

5'e

86% ee, product obtained
with Q-1b

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU *s	Height [mAU]	Area %
1	9.488	VB	0.3335	1.96862e4	922.07776	75.0192
2	12.412	BB	0.4283	2400.42480	87.35324	9.1474
3	15.901	BB	0.5574	1505.89087	42.07033	5.7386
4	73.286	BB	2.3532	2649.02002	15.20248	10.0948

Totals : 2.62415e4 1066.70381

Racemic-**5f**

95% ee, product obtained
with Q-2c

95% ee, product obtained
with QD-2c

HPLC conditions for ee determination: Chiralcel OD, hexanes: isopropanol = 80: 20, 1 mL/min, 220 nm.

94% ee, product obtained with Q-2c

HPLC conditions: Chiralcel OD, hexanes: isopropanol = 90: 10, 1.0 mL/min, 220 nm.

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area [mAU]	Height [mAU]	Area %
1	16.039	BB	0.6003	5.18682e4	1313.87708	20.5331
2	19.105	BV	0.6862	5.15922e4	1150.54395	20.4239
3	21.324	VV	0.9865	7.41756e4	1081.48792	29.3640
4	61.484	BB	2.6479	7.49715e4	443.47781	29.6790
Totals :				2.52608e5	3989.38675

95% ee, product obtained with Q-2c

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area [mAU]	Height [mAU]	Area %
1	16.608	BB	0.5487	359.07928	10.24379	3.3334
2	19.618	BB	0.6640	345.99854	8.05895	3.2120
3	22.776	BB	0.7721	292.57236	5.82713	2.7160
4	63.058	BB	2.2684	9774.50781	66.71517	90.7386
Totals :				1.07722e4	90.84505

94% ee, product obtained With QD-2c

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area [mAU]	Height [mAU]	Area %
1	16.556	BB	0.5441	3979.43774	114.81927	3.7225
2	19.428	BV	0.6223	3378.78540	85.84633	3.1607
3	21.326	VB	1.0316	9.63756e4	1330.25891	90.1536
4	63.423	BB	2.2044	3167.73804	22.14810	2.9632
Totals :				1.06902e5	1553.07260

80% ee, product obtained
with **Q-1b**

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU *s	Height [mAU]	Area %
1	16.069	BB	0.5982	4.02204e4	1050.36060	68.1202
2	19.378	BB	0.6489	4351.36377	104.48685	7.3698
3	22.162	BB	0.7936	7335.71436	143.68570	12.4243
4	63.310	BB	2.2260	7135.74902	49.59743	12.0856
Totals :				5.90432e4	1348.13057

86% ee, product obtained
with **QD-1b**

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU *s	Height [mAU]	Area %
1	16.594	BB	0.5430	4222.16211	122.17205	5.5088
2	19.293	BB	0.7218	5.65461e4	1232.14624	73.7771
3	22.484	BV	0.8303	9585.53027	180.18568	12.5065
4	63.708	BB	2.2514	6290.76855	43.36925	8.2077
Totals :				7.66446e4	1577.87323	

HPLC conditions for ee determination: Chiralcel OJ, hexanes: isopropanol = 60: 40, 1.0 mL/min, 220 nm.

95% ee, product obtained with Q-2c

HPLC conditions for ee determination: Chiralcel OD, hexanes: isopropanol = 80: 20, 1.0 mL/min, 220 nm.

=====
Area Percent Report
=====

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU	*s	Height [mAU]	Area %
1	10.342	BB	0.4326	1.67588e4		618.50116	41.0593
2	12.960	BB	0.4990	1.20011e4		389.81458	29.4028
3	20.586	BB	0.7467	1.20562e4		251.04202	29.5379

Totals : 4.08160e4 1259.35776

95% ee, product obtained with Q-2c

=====
Area Percent Report
=====

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU	*s	Height [mAU]	Area %
1	10.345	BB	0.4068	9070.26855		350.40521	5.5471
2	13.046	VB	0.4865	4166.71924		131.36748	2.5482
3	20.599	BB	0.9038	1.50276e5		2659.81860	91.9046

Totals : 1.63513e5 3141.59129

95% ee, product obtained with QD-2c

=====
Area Percent Report
=====

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU	*s	Height [mAU]	Area %
1	10.981	BB	0.4969	1886.10144		61.60828	5.0526
2	13.590	BB	0.5569	3.46348e4		942.37213	92.7816
3	22.634	BB	0.8557	808.50049		14.59995	2.1659

Totals : 3.73294e4 1018.58036

HPLC conditions for ee determination: Chiralcel OD, hexanes: isopropanol = 80: 20, 1.0 mL/min, 254 nm.

=====
Area Percent Report
=====

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=254 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU *s	Height [mAU]	Area %
1	13.274	BB	0.4863	1835.55566	61.79380	21.7461
2	16.454	BB	0.6053	2378.25220	59.59758	28.1755
3	24.801	BB	0.9067	1851.55505	30.98944	21.9356
4	92.852	BB	3.5749	2375.48755	10.21559	28.1428

Totals : 8440.85046 162.59641

95% ee, product obtained
with Q-2c

=====
Area Percent Report
=====

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=254 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU *s	Height [mAU]	Area %
1	13.306	BB	0.4963	251.94394	8.24400	2.9792
2	16.581	BB	0.6105	207.61452	5.41361	2.4550
3	24.772	BB	0.9043	208.52855	3.62368	2.4658
4	90.766	BB	3.9090	7788.72803	31.25552	92.1000

Totals : 8456.81503 48.53681

Racemic-5l

51

96% ee, product obtained
with Q-2c

ent-5l

96% ee, product obtained
with QD-2c

HPLC conditions for ee determination: Chiralcel OJ, hexanes: isopropanol = 70: 30, 1.0 mL/min, 220 nm.

Racemic-**5I**

88% ee, product obtained
with **Q-1b**

HPLC conditions for ee determination: Chiralcel OJ, hexanes: isopropanol = 96: 4, 1.0 mL/min, 220 nm.

Racemic-5m

=====
Area Percent Report
=====

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU	*s	Height [mAU]	Area %
1	41.979	BV	1.9133	2654.05176		21.26474	21.3301
2	44.951	VB	2.4206	3460.73315		20.41280	27.8133
3	67.336	BB	3.3759	3203.34839		12.99369	25.7608
4	79.315	BB	3.2430	3122.61572		11.47034	25.0959

Totals : 1.24427e4 66.14156

5m

95% ee, product obtained
with Q-2c

=====
Area Percent Report
=====

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU	*s	Height [mAU]	Area %
1	42.471	BV	1.7020	540.08264		4.39367	2.9730
2	45.718	VB	2.0686	759.49011		4.92069	4.1807
3	65.749	BB	4.2378	1.64789e4		55.88909	90.7102
4	80.751	BB	2.7720	388.06070		1.68093	2.1361

Totals : 1.81665e4 66.88438

ent-5m

95% ee, product obtained
with QD-2c

=====
Area Percent Report
=====

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU	*s	Height [mAU]	Area %
1	42.846	BV	1.6845	503.40369		4.11205	3.1555
2	46.043	VB	2.0217	730.60315		4.56455	4.5797
3	68.783	BB	3.3775	400.98837		2.01712	2.5135
4	77.559	BB	4.5428	1.43181e4		42.80846	89.7513

Totals : 1.59531e4 53.50218

HPLC conditions for ee determination: Chiralcel OD, hexanes: isopropanol = 60: 40, 1.0 mL/min, 220 nm.

=====
Area Percent Report
=====

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU *s	Height [mAU]	Area %
1	15.569	BB	0.6344	9983.01465	247.11082	49.8219
2	21.042	BB	0.8398	1.00544e4	186.16653	50.1781

Totals : 2.00374e4 433.27736

93% ee, product obtained
with Q-2c

=====
Area Percent Report
=====

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VWD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU *s	Height [mAU]	Area %
1	15.661	BB	0.6188	1084.90442	27.77279	3.7283
2	21.504	BB	1.0132	2.80142e4	445.50516	96.2717

Totals : 2.90991e4 473.27795

HPLC conditions for ee determination: Chiralcel OD, hexanes: isopropanol = 80: 20, 1.0 mL/min, 220 nm.

Racemic-7b

(+)-7b

94% ee, product obtained
with Q-2c

HPLC conditions for ee determination: Chiralcel OD, hexanes: isopropanol = 80: 20, 1.0 mL/min, 220 nm.

HPLC conditions for ee determination: Chiralpak AD, hexanes: isopropanol = 98: 2, 1.0 mL/min, 220 nm.

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VMD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU	Area %	Height [mAU]	Area %
1	19.222	BV	0.9359	6627.45020	108.20633	49.8789	
2	21.989	VB	1.0485	6659.62939	96.64352	50.1211	
Totals :				1.32871e4	204.84985

88% ee, product obtained with Q-2c

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VMD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU	Area %	Height [mAU]	Area %
1	19.996	BV	1.0088	2725.59692	40.37809	5.9155	
2	22.857	VB	1.0606	4.33499e4	619.81958	94.0845	
Totals :				4.60755e4	660.19767

90% ee, product obtained with QD-2c

Area Percent Report

Sorted By : Signal
Multiplier : 1.0000
Dilution : 1.0000
Sample Amount : 1.00000 [ng/ul] (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs

Signal 1: VMD1 A, Wavelength=220 nm

Peak #	RetTime [min]	Type	Width [min]	Area mAU	Area %	Height [mAU]	Area %
1	19.897	BV	0.9288	2.27236e4	368.63620	94.8896	
2	22.632	VB	1.0961	1223.81543	17.48987	5.1104	
Totals :				2.39474e4	386.12207		

HPLC conditions for ee determination: Chiralcel OJ, hexanes: isopropanol = 60: 40, 0.8 mL/min, 220 nm.

88% ee, product obtained with Q-2c

89% ee, product obtained with QD-2c

HPLC conditions for ee determination: (R, R)-Whelk-O 1, hexanes: isopropanol = 97: 3, 1.0 mL/min, 220 nm.

89% ee, product obtained with Q-2c

89% ee, product obtained with QD-2c

HPLC conditions for ee determination: Chiralcel OJ, hexanes: isopropanol = 60: 40, 1.0 mL/min, 220 nm.

90% ee, product obtained with Q-2c

HPLC conditions for ee determination: Chiralcel OD, hexanes: isopropanol = 80: 20, 1.0 mL/min, 220 nm.

90% ee, product obtained with Q-2c

HPLC conditions for ee determination: (R, R)-whelk-O 1, hexanes: isopropanol = 80: 20, 1.0 mL/min, 280 nm.

Racemic-7k

(-)-7k

89% ee, product obtained with Q-2c

HPLC conditions for ee determination: Chiralpak AS, hexanes: isopropanol = 80: 20, 1.0 mL/min, 220 nm.

93% ee, product obtained
 with Q-2c

HPLC conditions for ee determination: Chiralpak AS, hexanes: isopropanol = 85: 15, 1.0 mL/min, 220 nm.

91% ee, product obtained
with Q-2c

7m

7m

HPLC conditions for ee determination: Chiralcel OD, hexanes: isopropanol = 97: 3, 1.0 mL/min, 220 nm.

```

=====
                          Area Percent Report
=====
Sorted By      :      Signal
Multiplier     :      1.0000
Dilution       :      1.0000
Sample Amount  :      1.00000 [ng/ul]   (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs
  
```

```

Signal 1: VWD1 A, Wavelength=220 nm

Peak RetTime Type Width Area Height Area
# [min] [min] [min] mAU *s [mAU] %
-----|-----|-----|-----|-----|-----|
1 13.549 BB 0.7704 2.10434e4 420.38342 50.2603
2 16.976 BB 0.9728 2.08254e4 318.38510 49.7397
Totals :                      4.18688e4 738.76852
  
```

```

=====
                          Area Percent Report
=====
Sorted By      :      Signal
Multiplier     :      1.0000
Dilution       :      1.0000
Sample Amount  :      1.00000 [ng/ul]   (not used in calc.)
Use Multiplier & Dilution Factor with ISTDs
  
```

```

Signal 1: VWD1 A, Wavelength=220 nm

Peak RetTime Type Width Area Height Area
# [min] [min] [min] mAU *s [mAU] %
-----|-----|-----|-----|-----|-----|
1 13.683 VV 0.8177 1492.45349 28.49780 2.9641
2 16.790 VB 0.9641 4.88587e4 752.65070 97.0359
Totals :                      5.03512e4 781.14849
  
```