Supporting Information

Convergent Stereoselective Synthesis of the Visual Pigment A2E

Cristina Sicre, M. Magdalena Cid*

*Departamento de Química Orgánica, Universidade de Vigo,
Campus Lagoas-Marcosende, 36310 Vigo (Spain)*

mcid@uvigo.es

General. Reagents and solvents were purchased as reagent-grade and used without further purification unless otherwise stated. Solvents were dried according to published methods¹ and distilled before use. All reactions were performed in oven-dried or flame-dried glassware under an inert atmosphere of Ar unless otherwise stated. Chromatography refers to flash chromatography (FC) on SiO₂ 60 (230-400 mesh) from *Merck* or neutral Al₂O₃ 58 (~150 mesh) from Aldrich activated Brockmann III, head pressure of ca 0.2 bar. TLC: UV₂₅₄ SiO₂-coated plates from *Merck*, visualization by UV light (254 nm) or by coloring with 15% ethanolic phosphomolybdic acid solution. NMR spectra were recorded in a Bruker AMX400 (400.13 MHz and 100.61 MHz for proton and carbon respectively) spectrometer at 298 K with residual solvent peaks as internal reference and the chemical shifts are reported in δ [ppm], coupling constants J are given in [Hz] and the multiplicities assigned with DEPT experiments and expressed as follows: s = singlet, d = doublet, t = triplet, q = cuartet, m = multiplet. COSY, HMBC and HSQC methods were used to stablish atom connectivities. Electronic impact ionization (EI) and Fast atom bombardment (FAB) mass spectra were recorded on a VG-Autospec M instrument. Electrospray spectra were performed on a Bruker Daltonics APEX III FT-ICR (Fourier Transform Ion Cyclotron Resonance), equipped with a 7T actively shielded magnet, using an Apollo API electrospray ionization source. Infrared spectra (IR) were obtained on a JASCO FT/IR-4200 infrared spectrometer. Peaks are quoted in wave numbers (cm⁻¹) and their relative intensities are reported as follows: s = strong, m = medium, w = weak. UV/VIS spectra were recorded on a Varian-CARY 100 Bio Scan using MeOH as solvent and a 1 cm cell. Absorption maxima are reported in λ_{max} [nm] and ϵ [M⁻¹ cm⁻¹]. Reactions carried out under dark were performed using amber glassware. Melting points (mp) were taken on a Stuart Scientific apparatus. Elemental analysis was performed using a Fisons EA-1108 analyzer. High-performance liquid chromatography (HPLC) was performed with a Waters 510 machine with a dual-wave detector (300 and 290 nm).

¹ Perrin, D.; Amarego, W. *Purification of Laboratory Chemicals*; Pergamon Press: Oxford 1998.

2,4-Dibromopyridine (4)

2,4-Dihydroxypyridine (2.7 g, 24.3 mmol) and POBr₃ (23.0 g, 80.2 mmol) were heated in a Schlenk tube at 125 °C for 4.5 h. After cooling to 25 °C, the reaction mixture was carefully poured into water, basified to pH = 9 with aqueous NaHCO₃ (sat) and extracted with CH₂Cl₂. The combined extracts were dried over Na₂SO₄ (anh) and the solvent removed *in vacuo*. The residue was purified by flash chromatography (40:60, CH₂Cl₂/hexane) to give 4.9 g (85 % yield) of 2,4-dibromopyridine (**4**) as a white solid (mp = 36 °C, EtOH, lit² 38.0-38.5 °C) and 0.9 g (12%) of 2,3,4-tribromopyridine as a white solid (mp = 86 °C, EtOH, lit² 84-85 °C). **4**: ¹H-NMR (CDCl₃) δ 7.43 (dd, *J* = 5.3, 1.6 Hz, 1H, H⁵), 7.70 (d, *J* = 1.6 Hz, 1H, H³), 8.21 (d, *J* = 5.3 Hz, 1H, H⁶) ppm. ¹³C-NMR (CDCl₃) δ 121.5 (CH), 125.8 (CH), 128.6 (C), 136.3 (C), 143.5 (CH) ppm. MS (EI+) *m/z* 239 (M⁺ ⁸¹[Br]⁸¹[Br], 37), 237 (M⁺ ⁸¹[Br]⁷⁹[Br], 75), 235 (M⁺ ⁷⁹[Br]⁷⁹[Br], 40), 158 (97), 156 (100). **2,3,4-Tribromopyridine**: ¹H-NMR (CDCl₃) δ 7.52 (d, *J* = 5.1 Hz, 1H, H⁵), 8.11 (d, *J* = 5.1 Hz, 1H, H⁶) ppm. ¹³C-NMR (CDCl₃) δ 126.9 (C), 127.8 (CH), 136.7 (C), 144.8 (C), 147.7 (CH) ppm. MS (EI+) *m/z* (%) 319 (M⁺ ⁸¹[Br]⁸¹[Br]⁸¹[Br], 33), 317 (M⁺ ⁸¹[Br]⁸¹[Br]⁷⁹[Br], 100), 315 (M⁺ ⁸¹[Br]⁷⁹[Br]⁷⁹[Br], 85), 313 (M⁺ ⁷⁹[Br]⁷⁹[Br]⁷⁹[Br], 35), 238 (45), 236 (91).

2-[(1*E*)-3-*tert*-Butyldimethylsilyloxy-2-methylprop-1-en-1-yl]-4-halopyridine

Method A: Suzuki conditions.

X=Br. To a cold (-78 °C) solution of (*E*)-*tert*-butyl(3-iodo-2-methylallyloxy)dimethylsilane (0.45 g, 1.44 mmol) in THF (4.0 mL) in a Schlenk flask was added dropwise ^tBuLi (1.8 mL, 1.7 M in pentane, 3.03 mmol) and the mixture was stirred at -78 °C for 30 min. Then, B(OiPr)₃ (0.7 mL, 2.88 mmol) was added dropwise and the mixture was stirred at 0 °C for 2 h. At this time, Pd(PPh₃)₄ (78 mg, 0.07 mmol), a solution of **4a** (0.19 g, 0.80 mmol) in THF (4.0 mL) and a 10% aqueous TIOH (7.0 mL, 3.2 mmol) solution were added sequentially and the mixture was stirred at 25 °C until complete disappearance of dibromopyridine as observed by TLC and NMR (10 h). The crude mixture was diluted with Et₂O, filtered through a Celite® plug and washed with an aqueous saturated NaHCO₃ solution (2×). The organic layer was dried over Na₂SO₄ (anh) and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel (96:4 hexane/EtOAc) affording 0.19 g (70%) of **6a** as a

² Den Hertog, H. J. Rec. Trav. Chim., **1944**, 63, 85-101.

yellow oil and 12 mg (4%) of **7** as a yellow oil. **6a**: $^1\text{H-NMR}$ (CD_2Cl_2) δ 0.12 (s, 6H, $\text{Si}(\text{CH}_3)_2$), 0.95 (s, 9H, *t*Bu), 2.05 (br. s, 3H, CH_3), 4.19 (d, $J = 0.7$ Hz, 2H, $\text{H}^{3'}$), 6.51 (m, 1H, $\text{H}^{1'}$), 7.26 (dd, $J = 5.3$ Hz, 1.8 Hz, 1H, H^5), 7.39 (d, $J = 1.8$ Hz, 1H, H^3), 8.38 (d, $J = 5.3$ Hz, 1H, H^6) ppm. $^{13}\text{C-NMR}$ (CD_2Cl_2) δ -5.1 (CH_3 , $\text{Si}(\text{CH}_3)_2$), 15.8 (CH_3 , $\text{CH}_3\text{-C}^{2'}$), 18.9 (C, C-*t*Bu), 26.3 (CH_3 , *t*Bu), 68.5 (CH_2 , $\text{C}^{3'}$), 121.7 (CH, $\text{C}^{1'}$), 124.3 (CH, C^5), 127.6 (CH, C^3), 132.9 (C, C^4), 145.1 (C, $\text{C}^{2'}$), 150.4 (CH, C^6), 159.2 (C, C^2) ppm. IR (neat, cm^{-1}) 2929 (w, C-H), 2855 (w, C-H), 1693 (w), 1660 (w), 1564 (m), 1541 (m), 1465 (m), 1366 (w), 1252 (m), 1167 (w), 1107 (m), 1084 (m), 1004 (w), 883 (w), 835 (s), 775 (s), 710 (m), 670 (m). MS (EI+) m/z (%) 343 ($\text{M}^+ {}^{81}\text{Br}$], 21), 341 ($\text{M}^+ {}^{79}\text{Br}$], 21), 286 ($[\text{M-}t\text{Bu}]^+ {}^{81}\text{Br}$], 100), 284 ($[\text{M-}t\text{Bu}]^+ {}^{79}\text{Br}$], 95), 212 (25), 210 (25), 75 (37), 73 (23). High resolution MS calc for $\text{C}_{15}\text{H}_{24}\text{NOSi}^{81}\text{Br}$ 343.0790 and $\text{C}_{15}\text{H}_{24}\text{NOSi}^{79}\text{Br}$ 341.0810; found 343.0794 and 341.0810. **7**: $^1\text{H-NMR}$ (CDCl_3) δ 0.11 (s, 6H, $\text{Si}(\text{CH}_3)_2$), 0.95 (s, 9H, *t*Bu), 1.83 (s, 3H, CH_3), 4.17 (s, 2H, $\text{H}^{3'}$), 6.44 (s, 1H, $\text{H}^{1'}$), 7.12 (dd, $J = 5.1$ Hz, 1.2 Hz, 1H, H^5), 7.35 (br. s, 1H, H^3), 8.28 (d, $J = 5.1$ Hz, 1H, H^6) ppm. $^{13}\text{C-NMR}$ (CDCl_3) δ -5.4 (CH_3 , $\text{Si}(\text{CH}_3)_2$), 15.2 (CH_3 , $\text{CH}_3\text{-C}^{2'}$), 18.4 (C, C-*t*Bu), 25.9 (CH_3 , *t*Bu), 67.5 (CH_2 , $\text{C}^{3'}$), 119.5 (CH, $\text{C}^{1'}$), 122.7 (CH, C^5), 127.5 (CH, C^3), 142.3 (C, C^2), 143.7 (C, C^2), 148.7 (C, C^4), 149.7 (CH, C^6) ppm. IR (neat, cm^{-1}) 2951 (w, C-H), 2929 (w, C-H), 2855 (w, C-H), 1660 (w), 1582 (m), 1524 (m), 1462 (m), 1366 (m), 1252 (m), 1170 (w), 1108 (s), 1078 (s), 1005 (w), 989 (w), 834 (s), 775 (s). MS (EI+) m/z (%) 343 ($\text{M}^+ {}^{81}\text{Br}$], 2), 342 ($[\text{M-1}]^+ {}^{81}\text{Br}$], 11), 341 ($\text{M}^+ {}^{79}\text{Br}$], 2), 340 ($[\text{M-1}]^+ {}^{79}\text{Br}$], 11), 300 (100), 298 (98), 286 ($[\text{M-}t\text{Bu}]^+ {}^{81}\text{Br}$], 25), 284 ($[\text{M-}t\text{Bu}]^+ {}^{79}\text{Br}$], 25), 117 (24), 116 (20), 75 (79), 73 (39). High resolution MS calc for $\text{C}_{15}\text{H}_{24}\text{NOSi}^{81}\text{Br}$ 343.0790 and $\text{C}_{15}\text{H}_{24}\text{NOSi}^{79}\text{Br}$ 341.0810; found 343.0781 and 341.0801.

X=Cl. 58 mg (0.29 mmol) of **4b** and 0.10 g (0.32 mmol) of (*E*)-*tert*-butyl(3-iodo-2-methylallyloxy)dimethylsilane afforded, after 30 min at 25 °C and further purification by column chromatography on silica gel (90:10 hexane/EtOAc), 51 mg (59%) of **6b** as a yellow oil. **6b**: $^1\text{H-NMR}$ (CDCl_3) δ 0.11 (s, 12H, $\text{Si}(\text{CH}_3)_2$), 0.95 (s, 18H, *t*Bu), 2.04 (s, 3H, $\text{CH}_3\text{-C}^{2'}$), 4.19 (s, 2H, $\text{H}^{3'}$), 6.55 (s, 1H, $\text{H}^{1'}$), 7.09 (dd, $J = 5.3$ Hz, 1.9 Hz, 1H, H^5), 7.22 (d, $J = 1.9$ Hz, 1H, H^3), 8.48 (d, $J = 5.3$ Hz, 1H, H^6) ppm. $^{13}\text{C-NMR}$ (CDCl_3) δ -5.3 (CH_3), 15.3 (CH_3), 18.4 (C), 25.9 (CH_3), 67.9 (CH_2), 120.9 (CH), 121.4 (CH), 124.1 (CH), 143.7 (C), 144.2 (C), 149.9 (CH), 158.7 (C) ppm. IR (neat, cm^{-1}) ν 2952 (m, C-H), 2929 (m, C-H), 2855 (m, C-H), 1662 (w), 1571 (s), 1546 (m), 1465 (m), 1366 (w), 1252 (m), 1107 (s), 1086 (s), 1005 (w), 835 (s), 774 (s), 738 (m), 667 (w). MS (EI+) m/z (%) 299 ($\text{M}^+ {}^{37}\text{Cl}$], 9), 298 ($[\text{M+1}]^+ {}^{35}\text{Cl}$], 6), 297 ($\text{M}^+ {}^{35}\text{Cl}$], 26), 242 ($[\text{M-}t\text{Bu}]^+ {}^{37}\text{Cl}$], 37), 240 ($[\text{M-}t\text{Bu}]^+ {}^{35}\text{Cl}$], 100), 166 ($[\text{M-OTBS}]^+ {}^{35}\text{Cl}$], 42), 151 (22), 71 (36). High resolution MS calc for $\text{C}_{15}\text{H}_{24}{}^{37}\text{ClNOSi}$ 299.1286 and $\text{C}_{15}\text{H}_{24}{}^{35}\text{ClNOSi}$ 297.1316; found 299.1272 and 297.1302.

Method B: Negishi conditions.

X=Br. To a cold (-78 °C) solution of (*E*)-*tert*-butyl(3-iodo-2-methylallyloxy)dimethylsilane (3.6 g, 11.66 mmol) in THF (27.0 mL) was added dropwise $^t\text{BuLi}$ (13.7 mL, 1.7 M in pentane, 23.34 mmol) and the mixture was stirred at -78 °C for 30 min. Then, a solution of ZnCl_2 (2.5 g, 18.34 mmol) in THF (18.0 mL), drying ZnCl_2 previously by heating at 120 °C under vacuum for 2 h, was added dropwise and the mixture was allowed to warm to 25 °C for 1 h. This solution was cannulated to a solution of $\text{Pd}(\text{PPh}_3)_4$ (0.45 g, 0.39 mmol) and **4a** (1.53 g, 6.40 mmol) in

THF (17.0 mL) in a Schlenk flask. Upon stirring for 20 h, the reaction was quenched with an aqueous saturated NH_4Cl solution. The aqueous phase was extracted with Et_2O (3 $\times$) and the combined organic extracts were washed with brine (1 $\times$), dried over Na_2SO_4 (anh.) and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel (96:4 hexane/ EtOAc) affording 1.6 g (73%) of **6a** as a yellow oil, 0.21 g (10%) of **7** as a yellow oil.

4-Bromo-2-[(1E)-3-hidroxy-2-methyl-prop-1-en-1-yl]pyridine (**8**)

A solution of silylether **6** (0.30 g, 0.87 mmol) in THF (6.0 mL) was treated with $n\text{Bu}_4\text{NF}$ (1.0 mL, 1 M in THF, 1.04 mmol) and the mixture was stirred for 30 min at 25 °C. It was diluted with EtOAc and washed with an aqueous saturated NaHCO_3 solution (2 $\times$). The combined aqueous layers were extracted with EtOAc (3 $\times$) and the resulting organic extracts washed with brine (2 $\times$), dried over Na_2SO_4 (anh.) and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel (60:40 hexane/ EtOAc) affording 0.20 g (100%) of **8** as a yellow oil. ^1H -NMR (CD_2Cl_2) δ 1.56 (br. s, 1H, OH), 2.10 (d, $J = 1.3$ Hz, 3H, CH_3), 4.18 (s, 2H, $\text{H}^{3'}$), 6.50 (q, $J = 1.3$ Hz, 1H, $\text{H}^{1'}$), 7.28 (dd, $J = 5.3$ Hz, 1.8 Hz, 1H, H^5), 7.42 (d, $J = 1.8$ Hz, 1H, H^3), 8.39 (d, $J = 5.3$ Hz, 1H, H^6) ppm. ^{13}C -NMR (CD_2Cl_2) δ 16.0 (CH_3 , $\text{CH}_3\text{-C}^{2'}$), 68.9 (CH_2 , $\text{C}^{3'}$), 122.3 (CH , $\text{C}^{1'}$), 124.6 (CH , C^5), 127.7 (CH , C^3), 133.0 (C , C^4), 145.3 (C , $\text{C}^{2'}$), 150.4 (CH , C^6), 158.9 (C , C^2) ppm. IR (NaCl, cm^{-1}) 3500-3100 (br., O-H), 2912 (m, C-H), 2854 (m, C-H), 1683 (m), 1659 (m), 1567 (s), 1542 (s), 1463 (m), 1386 (m), 1076 (m), 1018 (m), 885 (m), 823 (m). MS (EI+) m/z (%) 230 ($[\text{M}+1]^+ ^{81}\text{Br}$, 17), 229 ($\text{M}^+ ^{81}\text{Br}$, 100), 228 ($[\text{M}+1]^+ ^{79}\text{Br}$, 76), 227 ($\text{M}^+ ^{79}\text{Br}$, 74), 226 (46), 210 (77), 208 (95), 198 (64), 196 (61), 117 (98). High resolution MS calc for $\text{C}_9\text{H}_{10}\text{NO}^{81}\text{Br}$ 228.9925 and $\text{C}_9\text{H}_{10}\text{NO}^{79}\text{Br}$ 226.9946; found, 228.9932 y 226.9951.

General procedure for Sonogashira reaction of 4-bromopyridines.

To a Schlenk type flask charged with $\text{Pd}(\text{PPh}_3)_2\text{Cl}_2$ (4 mol%), CuI (4 mol%) and a solution of the bromopyridine (100 mol%) in Et_3N (0.3 M) was added (trimethylsilyl)acetylene (150 mol%). The solution was stirred at 105 °C for 4 h.

Finally, triethylamine was removed *in vacuo* and the residue was purified by column chromatography on silica gel.

For R=TBS: 2-[(1E)-3-*tert*-Butyldimethylsilyloxy-2-methylprop-1-en-1-yl]-4-[2-trimethylsilyleth-1-yn-1-yl]pyridine (10a)

161 mg (0.47 mmol) of bromopyridine **6** afforded, after purification by column chromatography on silica gel (95:5 hexane/EtOAc), 163 mg (97%) of **10a** as an oil. $^1\text{H-NMR}$ (CDCl_3) δ 0.11 (s, 6H, $\text{Si}(\text{CH}_3)_2$), 0.26 (s, 9H, $\text{Si}(\text{CH}_3)_3$), 0.95 (s, 9H, *t*Bu), 2.03 (br. s, 3H, $\text{CH}_3\text{-C}^2$), 4.19 (d, $J = 0.7$ Hz, 2H, H^3), 6.56 (m, 1H, H^1), 7.09 (dd, $J = 5.1$ Hz, 1.4 Hz, 1H, H^5), 7.23 (br. s, 1H, H^3), 8.53 (d, $J = 5.1$ Hz, 1H, H^6) ppm. $^{13}\text{C-NMR}$ (CDCl_3) δ -5.3 (CH_3), -0.3 (CH_3), 15.3 (CH_3), 18.4 (C), 25.9 (CH_3), 68.0 (CH_2), 99.0 (C), 102.5 (C), 122.0 (CH), 122.7 (CH), 126.1 (CH), 131.0 (C), 143.1 (C), 149.0 (CH), 157.3 (C) ppm. MS (EI+) m/z (%) 360 ($[\text{M}+1]^+$, 20), 359 (M^+ , 60), 344 (33), 303 (36), 302 ($[\text{M}-t\text{Bu}]^+$, 100), 228 ($[\text{M}-\text{OTBS}]^+$, 31), 73 (22). High resolution MS calc $\text{C}_{20}\text{H}_{33}\text{NOSi}_2$ 359.2101; found 359.2091.

For R=H: 2-[(1E)-3-Hidroxy-2-methylprop-1-en-1-yl]-4-[2-trimethylsilyleth-1-yn-1-yl]pyridine (10c)

1.5 g (6.71 mmol) of bromopyridine **8** afforded, after purification by column chromatography on silica gel (60:38:2 EtOAc/hexane/ Et_3N), 1.5 g (93%) of **10c** as an oil. $^1\text{H-NMR}$ (CDCl_3) δ 0.25 (s, 9H, $\text{Si}(\text{CH}_3)_3$), 2.00 (s, 3H, CH_3), 4.16 (s, 2H, H^3), 6.57 (s, 1H, H^1), 7.10 (d, $J = 5.0$ Hz, 1H, H^5), 7.25 (s, 1H, H^3), 8.48 (d, $J = 5.0$ Hz, 1H, H^6) ppm. $^{13}\text{C-NMR}$ (CDCl_3) δ -0.3 (CH_3), 15.4 (CH_3), 67.8 (CH_2), 99.4 (C), 102.2 (C), 122.5 (CH), 123.0 (CH), 126.1 (CH), 131.3 (C), 143.8 (C), 148.8 (CH), 156.9 (C) ppm. IR (neat, cm^{-1}) 3500-3000 (br, OH), 2958 (w, C-H), 2901 (w, C-H), 2857 (w, C-H), 2162 (w), 1588 (m), 1534 (w), 1468 (w), 1250 (m), 1170 (w), 1075 (w), 946 (m), 839 (s), 758 (m), 702 (w), 647 (w). MS (EI+) m/z (%) 246 ($[\text{M}+1]^+$, 20), 245 (M^+ , 100), 244 ($[\text{M}-1]^+$, 75), 230 (19), 228 (14), 227 (18), 226 (35), 216 (23), 214 (61). High resolution MS calc for $\text{C}_{14}\text{H}_{19}\text{NOSi}$ 245.1236; found 245.1235.

Sonogashira reaction of 4-chloropyridine.

To a solution of (trimethylsilyl)acetylene (0.011 mL, 0.08 mmol) in THF (0.5 mL) in a Schlenk flask at -78°C was added dropwise *n*-BuLi (0.049 mL, 1.48 M in hexane, 0.07 mmol) and the mixture was stirred at -78°C for 30 min. Then, Me_3SnCl (14 mg, 0.07 mmol) was added and the mixture was stirred at -78°C for 2 h. At this time, the solvent was removed *in vacuo* and a solution of **6b** (19 mg, 0.06 mmol) in toluene (0.5 mL) followed by $\text{Pd}(\text{PPh}_3)_4$ (7 mg, 0.006 mmol) were added. The reaction mixture was

stirred at 100 °C overnight. Finally, it was allowed to cool to 25 °C and filtered through a Celita® plug washing with EtOAc. The solvent was removed *in vacuo* and the residue was purified by column chromatography on silica gel (95:5 hexane/EtOAc) affording 11 mg (48%) of **10a** and 9.9 mg (52%) of **6b**.

4-Ethynyl-2-[(1E)-3-hidroxy-2-methylprop-1-en-1-yl]pyridine (**10b**)

A solution of silylether **10a** (69 mg, 0.19 mmol) in THF (1.3 mL) was treated with *n*Bu₄NF (0.46 mL, 1 M in THF, 0.46 mmol) and the mixture was stirred for 2 h at 25 °C. It was diluted with EtOAc and washed with an aqueous saturated NaHCO₃ solution (2×). The combined aqueous layers were extracted with EtOAc (3×) and the resulting organic extracts washed with brine (2×), dried over Na₂SO₄ and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel (60:40 hexane/EtOAc) affording 25 mg (76%) of **10b** as an oil. ¹H-NMR (CDCl₃) δ 2.02 (br s, 3H, CH₃-C^{2'}), 3.14 (br s, 1H, OH), 3.26 (s, 1H, H^{2''}), 4.18 (s, 2H, H^{3'}), 6.59 (d, *J* = 1.3 Hz, 1H, H^{1'}), 7.15 (dd, *J* = 5.0 Hz, 1.4 Hz, 1H, H⁵), 7.29 (br s, 1H, H³), 8.53 (d, *J* = 5.0 Hz, 1H, H⁶) ppm. ¹³C-NMR (CDCl₃) δ 15.5 (CH₃), 68.1 (CH₂), 81.2 (C), 81.4 (CH), 122.6 (CH), 123.2 (CH), 126.4 (CH), 130.4 (C), 143.8 (C), 149.1 (CH), 157.0 (C) ppm. MS (EI+) *m/z* (%) 174 ([M+1]⁺, 13), 173 (M⁺, 100), 172 ([M-1]⁺, 63), 156 ([M-OH], 19), 155 (23), 154 (85), 144 (44), 142 (86), 141 (53). High resolution MS calc for C₁₁H₁₁NO 173.0841; found 173.0837.

General procedure for Dess-Martin oxidation

To a solution of Dess-Martin periodinane (1.5 mol equiv) and pyridine (2.0 mol equiv) in CH₂Cl₂ (0.3 M) at 0 °C was added, *via cannula*, a solution of the alcohol (1.0 mol equiv) in CH₂Cl₂ (1 M) and the reaction mixture was stirred at 0 °C for 2 h. The reaction mixture was diluted with EtOAc and washed with saturated NaHCO₃ aqueous solution (2×) and saturated Na₂S₂O₃ aqueous solution (2×). The combined aqueous layers were extracted with EtOAc and the resulting organic extracts dried over Na₂SO₄ (anh), and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel.

For R=H : 4-Ethynyl-2-[(1E)-3-formil-2-methylprop-1-en-1-yl]pyridine (11a)

25 mg (0.14 mmol) of alcohol **10b** afforded, after purification by column chromatography on silica gel (85:15 hexane/EtOAc), 20 mg (79%) of **11a** as a white solid (mp 107-108 °C, hexane/CH₂Cl₂). ¹H-NMR (CDCl₃) δ 2.24 (d, *J* = 1.4 Hz, 3H, CH₃-C²), 3.34 (s, 1H, H^{2''}), 7.21 (m, 1H, H^{1'}), 7.31 (dd, *J* = 5.0 Hz, 1.4 Hz, 1H, H⁵), 7.54 (br. s, 1H, H³), 8.70 (d, *J* = 5.0 Hz, 1H, H⁶), 9.64 (s, 1H, CHO) ppm. ¹³C-NMR (CDCl₃) δ 11.0 (CH₃), 80.6 (C), 82.4 (CH), 125.4 (CH), 128.1 (CH), 130.9 (C), 142.2 (C), 146.3 (CH), 150.0 (CH), 154.7 (C), 195.6 (CH) ppm. IR (neat, cm⁻¹) 3228 (s, C≡C), 2843 (w, C-H), 1663 (s, C=O), 1583 (w), 1360 (m), 1183 (m), 1024 (m), 916 (m), 828 (m), 725 (m), 688 (m). MS (EI+) *m/z* (%) 172 ([M+1]⁺, 9), 171 (M⁺, 73), 143 (11), 142 (100), 141 (30), 115 (10). High resolution MS calc for C₁₁H₉NO 171.0684; found 171.0685.

For R=TMS: 2-[(1E)-3-Formil-2-methylprop-1-en-1-yl]-4-[2-trimethylsilyleth-1-yn-1-yl]pyridine (11b)

1.5 g (6.24 mmol) of alcohol **10c** afforded, after purification by column chromatography on neutral alumina Act III (85:15 hexane/EtOAc), 1.4 g (92%) of **11b** as a white solid (mp 66-67 °C, hexane/CH₂Cl₂). ¹H-NMR (CD₂Cl₂) δ 0.27 (s, 9H, Si(CH₃)₃), 2.22 (d, *J* = 1.2 Hz, 3H, CH₃-C²), 7.19 (m, 1H, H^{1'}), 7.27 (dd, *J* = 4.9 Hz, 1.3 Hz, 1H, H⁵), 7.51 (br s, 1H, H³), 8.66 (d, *J* = 4.9 Hz, 1H, H⁶), 9.61 (s, 1H, CHO) ppm. ¹³C-NMR (CD₂Cl₂) δ -0.1 (CH₃), 11.3 (CH₃), 101.0 (C), 102.0 (C), 125.6 (CH), 128.5 (CH), 132.3 (C), 142.6 (C), 146.9 (CH), 150.4 (CH), 155.3 (C), 196.0 (CH) ppm. IR (neat, cm⁻¹) 1676 (s, C=O), 1577 (m), 1533 (w), 1465 (w), 1415 (w), 1364 (w), 1297 (w), 1250 (m), 1178 (m), 1026 (m), 939 (m), 894 (w), 839 (s), 761 (m), 701 (m), 646 (m). MS (EI+) *m/z* (%) 244 ([M+1]⁺, 16), 243 (M⁺, 75), 228 (11), 215 (19), 214 (100), 154 (8), 99 (9), 85 (6), 73 (13). High resolution MS calc for C₁₄H₁₇NOSi 243.1079; found 243.1081. Anal. calcd (%) C 69.09, H 7.04, N 5.76; found C 68.90, H 7.07, N 5.76.

2-[(1E)-2-Methylbut-1-en-3-yn-1-yl]-4-[2-trimethylsilyleth-1-yn-1-yl]pyridine (12a)

Trimethylsilyldiazomethane (0.03 mL, 2 M in hexane, 0.19 mmol) was added dropwise to a solution of LDA, prepared from diisopropylamine (0.027 mL, 0.19 mmol) and *n*-BuLi (0.19 mL, 1 M in hexane, 0.19 mmol) in THF (0.5 mL), at -78 °C and the mixture was stirred at this temperature for 30 min. A solution of the aldehyde (36 mg, 0.15 mmol) in THF (0.3 mL) was then added dropwise at -78 °C and the mixture was stirred

under dark for 5 h. After being quenched with water, the mixture was allowed to warm to 25 °C and extracted with EtOAc. The organic extracts were dried over Na₂SO₄ (anh), and concentrated *in vacuo*. The residue was purified by column chromatography on silica gel (96:2:2 hexane/EtOAc/Et₃N) affording 23 mg (65%) of **12a** as a white solid. ¹H-NMR (CDCl₃) δ 0.26 (s, 9H, Si(CH₃)₃), 2.31 (s, 3H, CH₃-C^{2'}), 3.05 (s, 1H, H^{4'}), 6.81 (s, 1H, H^{1'}), 7.13 (d, *J* = 5.0 Hz, 1H, H⁵), 7.22 (s, 1H, H³), 8.55 (d, *J* = 5.0 Hz, 1H, H⁶) ppm. ¹³C-NMR (CDCl₃) δ -0.3 (CH₃), 19.4 (CH₃), 77.4 (CH), 87.2 (C), 99.7 (C), 102.0 (C), 123.6 (CH), 124.3 (C), 126.7 (CH), 131.3 (C), 135.0 (CH), 149.2 (CH), 155.7 (C) ppm. IR (NaCl, cm⁻¹) 3297 (m, C≡C-H), 2958 (m, C-H), 2925 (m, C-H), 2854 (w, C-H), 1585 (m), 1534 (w), 1467 (w), 1251 (m), 942 (w), 847 (s), 760 (w). MS (EI+) *m/z* (%) 240 ([M+1]⁺, 60), 239 (M⁺, 100), 238 ([M-H]⁺, 99), 224 (49), 180 (23), 166 ([M-TMS]⁺, 14). High resolution MS (FAB⁺) calc for C₁₅H₁₈NSi 240.1209; found 240.1213. UV (MeOH) λ_{max} = 226 nm (ε 14900), 248 nm (ε 20600), 258 nm (ε 23200), 309 nm (ε 5600).

4-Ethynyl-2-[(1*E*)-2-methylbut-1-en-3-yn-1-yl]pyridine (**12b**)

A solution of triphenylphosphine (1.1 g, 4.11 mmol) in CH₂Cl₂ (5.0 mL) was added to a cooled (0 °C) solution of carbon tetrabromide (0.68 g, 2.06 mmol) in CH₂Cl₂ (7.6 mL) under dark. This solution was stirred for 30 min at 0 °C and then a solution of aldehyde **11b** (0.25 g, 1.03 mmol) in CH₂Cl₂ (2.6 mL) was added dropwise. The reaction mixture darkens on addition. After stirring for 1 h the solvent was practically evaporated and filtered through a silica plug washing with 78:20:2 hexane/EtOAc/Et₃N. The resulting filtrate was concentrated under reduced pressure to give the corresponding dibromodiene, which was used without further purification.

n-Butyllithium (4.4 mL, 1.41 M in hexane, 6.17 mmol) was added dropwise to a solution of diisopropylamine (0.9 mL, 6.48 mmol) in THF (13.0 mL) at 0 °C. After 1 h, the resulting LDA solution was cooled to -78 °C and transferred *via cannula* to a solution of the dibromodiene in THF (11.4 mL) at -78 °C under dark. The reaction was stirred at this temperature for 15 min and the resulting dark solution was quenched with MeOH (5.0 mL) and allowed to warm to room temperature. The reaction mixture was filtered through a Celite® plug, washed with EtOAc and concentrated under reduced pressure. Purification was performed by flash chromatography on silica gel (90:8:2 hexane/EtOAc/Et₃N) affording 0.14 g (80%) of **12b** as a white solid. ¹H-NMR (C₆D₆) δ 2.43 (d, *J* = 1.4 Hz, 3H, CH₃-C^{2'}), 2.73 (s, 1H, H^{2''}), 2.76 (s, 1H, H^{4'}), 6.69 (dd, *J* = 4.9 Hz, 1.3 Hz, 1H, H⁵), 6.73 (br s, 1H, H^{1'}), 6.82 (br s, 1H, H³), 8.23 (d, *J* = 4.9 Hz, 1H, H⁶) ppm. ¹³C-NMR (C₆D₆) δ 19.8 (CH₃, CH₃-C^{2'}), 78.1 (CH, C^{4'}), 81.3 (C, C^{1''}), 81.6 (CH, C^{2''}), 87.7 (C, C^{3'}), 123.6 (CH, C⁵), 125.3 (C, C^{2'}), 127.3 (CH, C³), 130.4 (C, C⁴), 134.9 (CH, C^{1'}), 149.5 (CH, C⁶), 156.4 (C, C²) ppm. IR (neat, cm⁻¹) 3270 (s, C≡C-H), 2920 (w, C-H), 2113 (w), 1618 (m), 1582 (s), 1533 (s), 1476 (m), 1436 (w), 1393 (m), 1306 (w), 1225 (m), 1148 (m), 1104 (m), 926 (w), 901 (s), 839 (s), 771 (m). MS (EI+)

m/z (%) 167 (M^+ , 7), 166 ($[M-1]^+$, 10), 76 (33), 75 (36), 73 (33), 71 (34), 69 (100). High resolution MS calc for $C_{12}H_9N$ 167.0735; found 167.0738. UV (MeOH) $\lambda_{\max}$ = 235 nm (ϵ 14800), 244 nm (ϵ 15900), 265 nm (ϵ 12200), 307 nm (ϵ 6100).

2-[(1*E*,3*E*)-4-Tributylstannyl-2-methylbuta-1,3-dien-1-yl]-4-[(1*E*)-2-(tributylstannyl)eth-1-en-1-yl]pyridine (3**)**

To a solution of hexabutylditin (0.73 mL, 1.44 mmol) in THF (2.9 mL) was added *n*-BuLi (0.91 mL, 1.58 M in hexane, 1.44 mmol) at -78°C . The solution was stirred for 30 min at -40°C . Then the mixture was added *via cannula* to a suspension of CuCN (64 mg, 0.72 mmol) in THF (0.1 mL) at -78°C under dark. The solution was stirred at -40°C for 5 min. Dried MeOH (1.6 mL) was added at -78°C and the temperature was allowed to warm to -40°C . After 15 min, a solution of the alkyne **12b** (0.03 g, 0.18 mmol) in THF (0.2 mL) was added dropwise. The dark solution was stirred for 5 min and then quenched with dried MeOH (2 mL). After 15 min water (2 mL) was added and the reaction mixture allowed to warm to 25°C over 15 min. Then, it was extracted with diethyl ether and the organic layer was washed with brine, dried over Na_2SO_4 , filtered, and concentrated *in vacuo*. Purification was performed by flash chromatography on silica gel (98:2 hexane/ Et_3N , then hexane) affording 0.10 g (76%) of a 2.8:1 mixture of 2-[(1*E*,3*E*)-4-tributylstannyl-2-(methylbuta-1,3-dienyl)]-4-[(*E*)-2-(tributylstannyl)vinyl]pyridine (**3**) and 2-[(1*E*)-3-tributylstannyl-2-methylbuta-1,3-dien-1-yl]-4-[(1*E*)-2-(tributylstannyl)eth-1-en-1-yl]pyridine (**13**) as a yellow oil. The mixture was separated by HPLC [Waters Spherisorb s5 NH_2 , 10×250 mm, 70:30 hexane/ CH_2Cl_2 , flow: 1.8 mL/min, detector: UV 300 nm].

3: $^1\text{H-NMR}$ (CD_2Cl_2) δ 0.91 (t, $J = 7.2$ Hz, 18H, $6 \times \text{SnCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 0.95-1.03 (m, 12H, $6 \times \text{SnCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.30-1.39 (m, 12H, $6 \times \text{SnCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 1.52-1.60 (m, 12H, $6 \times \text{SnCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 2.27 (d, $J = 0.9$ Hz, 3H, $\text{CH}_3\text{-C}^{2'}$), 6.48 (br s, 1H, H^1), 6.53 (d, $J = 19.2$ Hz, $^2J^1\text{H-}^{119}\text{Sn} = 67.6$ Hz, $^2J^1\text{H-}^{117}\text{Sn} = 64.9$ Hz, 1H, H^4), 6.77 (d, $J = 19.2$ Hz, $^3J^1\text{H-}^{119}\text{Sn} \approx ^3J^1\text{H-}^{117}\text{Sn} \approx 57.3$ Hz, 1H, H^3), 6.85 (d, $J = 19.5$ Hz, 1H, $\text{H}^{1''}$), 7.09 (dd, $J = 5.1$ Hz, 1.5 Hz, 1H, H^5), 7.20 (d, $J = 19.5$ Hz, $^2J^1\text{H-}^{119}\text{Sn} \approx ^2J^1\text{H-}^{117}\text{Sn} \approx 62.7$ Hz, 1H, $\text{H}^{2''}$), 7.23 (br s, $^5J^1\text{H-}^{119}\text{Sn} = 32.3$ Hz, $^5J^1\text{H-}^{117}\text{Sn} = 30.4$ Hz, 1H, H^3), 8.52 (d, $J = 5.1$ Hz, 1H, H^6) ppm. $^{13}\text{C-NMR}$ (CD_2Cl_2) δ 10.1 (CH_2 , $3 \times \text{SnCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 10.2 (CH_2 , $3 \times \text{SnCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 13.6 (CH_3 , $\text{CH}_3\text{-C}^{2'}$), 14.1 (CH_3 , $6 \times \text{SnCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 27.89 (CH_2 , $3 \times \text{SnCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 27.91 (CH_2 , $3 \times \text{SnCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 29.67 (CH_2 , $3 \times \text{SnCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 29.72 (CH_2 , $3 \times \text{SnCH}_2\text{CH}_2\text{CH}_2\text{CH}_3$), 118.1 (CH, C^5), 122.3 (CH, C^3), 130.7 (CH, C^1), 131.3 (CH, $\text{C}^{4'}$), 137.1 (CH, $\text{C}^{2''}$), 141.4 (C, $\text{C}^{2'}$), 144.3 (CH, $\text{C}^{1''}$), 146.0 (C, C^4), 150.0 (CH, C^6), 151.8 (CH, $\text{C}^{3'}$), 158.3 (C, C^2) ppm. IR (NaCl, cm^{-1}) 2956 (s, C-H), 2924 (s, C-H), 2851 (m, C-H), 1590 (m), 1538 (w), 1460 (m), 1376 (w), 1178 (w), 1155 (w), 1072 (w), 984 (m), 867 (w), 775 (w), 672 (m). MS (FAB $^+$) m/z (%) 752 (20), 750 (25), 748 (22), 460 (32),

458 (23), 179 (54), 177 (52), 175 (34), 172 (29), 170 (100). High resolution MS (FAB⁺) calc for C₃₆H₆₆N¹²⁰Sn¹²⁰Sn 752.3239; found 752.3228. UV (MeOH) λ_{max} = 272 nm (ϵ 19000), 320 nm (ϵ 6000). **13**: ¹H-NMR (CD₂Cl₂) δ 0.87-0.92 (m, 18H, 6 $\times$ SnCH₂CH₂CH₂CH₃), 0.99-1.05 (m, 12H, 6 $\times$ SnCH₂CH₂CH₂CH₃), 1.29-1.37 (m, 12H, 6 $\times$ SnCH₂CH₂CH₂CH₃), 1.52-1.58 (m, 12H, 6 $\times$ SnCH₂CH₂CH₂CH₃), 2.29 (s, 3H, CH₃-C²), 5.37 (d, J = 1.9 Hz, ³ J ¹H-¹¹⁹Sn $\approx$ ³ J ¹H-¹¹⁷Sn $\approx$ 62.7 Hz, 1H, H⁴-*cis*), 6.05 (d, J = 1.9 Hz, ³ J ¹H-¹¹⁹Sn = 132.9 Hz, ³ J ¹H-¹¹⁷Sn = 129.3 Hz, 1H, H⁴-*trans*), 6.32 (s, 1H, H¹), 6.84 (d, J = 19.6 Hz, 1H, H^{1'}), 7.08 (d, J = 5.1 Hz, 1H, H⁵), 7.14 (s, 1H, H³), 7.19 (d, J = 19.6 Hz, 1H, H^{2'}), 8.51 (d, J = 5.1 Hz, 1H, H⁶) ppm. ¹³C-NMR (CD₂Cl₂) δ 10.2 (CH₂, 3 $\times$ SnCH₂CH₂CH₂CH₃), 11.0 (CH₂, 3 $\times$ SnCH₂CH₂CH₂CH₃), 14.1 (CH₃, 6 $\times$ SnCH₂CH₂CH₂CH₃), 16.7 (CH₃, CH₃-C²), 27.88 (CH₂, 3 $\times$ SnCH₂CH₂CH₂CH₃), 27.96 (CH₂, 3 $\times$ SnCH₂CH₂CH₂CH₃), 29.65 (CH₂, 3 $\times$ SnCH₂CH₂CH₂CH₃), 29.69 (CH₂, 3 $\times$ SnCH₂CH₂CH₂CH₃), 117.8 (CH, C⁵), 122.1 (CH, C³), 125.8 (CH₂, C⁴), 128.5 (CH, C^{1'}), 137.0 (CH, C^{2'}), 144.3 (CH, C^{1''}), 146.0 (C, C⁴), 147.5 (C, C²), 149.9 (CH, C⁶), 158.4 (C, C²), 158.8 (C, C³) ppm. IR (NaCl, cm⁻¹) 2956 (s, C-H), 2924 (s, C-H), 2852 (m, C-H), 1591 (m), 1538 (w), 1461 (m), 1376 (w), 1072 (w), 986 (w), 910 (w), 896 (w), 864 (w), 773 (w), 686 (m), 671 (m). MS (FAB⁺) m/z (%) 752 (10), 750 (14), 748 (11), 692 (22), 690 (20), 290 (24), 288 (21), 281 (31), 235 (22), 221 (31), 207 (34), 179 (98), 177 (97), 175 (61), 172 (31), 170 (100). High resolution MS (FAB⁺) calc for C₃₆H₆₆N¹²⁰Sn¹²⁰Sn 752.3239; found 752.3218. UV (MeOH) λ_{max} = 225 nm (ϵ 13600), 267 nm (ϵ 20900), 318 nm (ϵ 3300).

2-[(1*E*,3*E*,5*E*,7*E*)-2,6-Dimethyl-8-(2,6,6-trimethylcyclohex-1-en-1-yl)octa-1,3,5,7-tetraen-1-yl]-4-[(1*E*,3*E*,5*E*)-4-methyl-6-(2,6,6-trimethylcyclohex-1-en-1-yl)hexa-1,3,5-trien-1-yl]pyridine (14)

Method A: PdCl₂(PhCN)₂ (4 mg, 0.010 mmol) was added to a solution of the iodide **2** (95 mg, 0.30 mmol) in DMF (5.0 mL) and the resulting mixture was thoroughly degassed using freeze-thaw cycles (3 $\times$). After 5 min at 25 °C, a solution of a 4.5:1 mixture of stannanes **3** and **13** (75 mg, 0.10 mmol) and (*i*Pr)₂NEt (0.10 mL, 0.60 mmol) in THF (5.0 mL), previously thoroughly degassed using freeze-thaw cycles (3 $\times$), was added followed by BHT (trace). After stirring for 15 h, another portion of PdCl₂(PhCN)₂ (2 mg, 0.006 mmol) was added and the mixture was stirred for 31 h at 25 °C. Then, an aqueous saturated KF solution was added, the layers were separated after 30 min of stirring and the aqueous layer was extracted with EtOAc (3 $\times$). The combined

organic layers were washed with brine (5×) and KF (1×), dried over Na₂SO₄ and the solvent was removed. The residue was purified by column chromatography on neutral alumina Act III (99:1 hexane/EtOAc) affording 22 mg (50%) of **14**³ as a bright yellow solid and **15**.

Method B: To a solution of **2** (53 mg, 0.17 mmol), Pd₂dba₃ (4 mg, 0.004 mmol) and AsPh₃ (11 mg, 0.04 mmol) in NMP (0.7 mL) under dark previously thoroughly degassed using freeze-thaw cycles (3×), was added a degassed solution of a 16:1 mixture of **3** and **13** (42 mg, 0.06 mmol) in NMP (0.7 mL) and BHT (trace). The reaction mixture was stirred at 25 °C for 8 h, when another portion of Pd₂dba₃ (4 mg, 0.004 mmol) and AsPh₃ (11 mg, 0.04 mmol) was added. It was stirred 23 h at 25 °C and then, an aqueous saturated KF solution was added, the layers were separated after 30 min of stirring and the aqueous layer was extracted with EtOAc (3×). The combined organic layers were washed with brine (5×) and KF (1×), dried over Na₂SO₄ (anh.) and the solvent was removed. The residue was purified by column chromatography on neutral alumina Act III (99:1 hexane/EtOAc) affording 0.02 g (69%) of **14** as a bright yellow solid and traces of **15**.

14: ¹H-NMR (CDCl₃) δ 1.04 (s, 12H, 2 × CH₃-C^{12''} and 2 × CH₃-C^{14''}), 1.47 (m, 4H, (CH₂)^{11''} and (CH₂)^{13''}), 1.63 (m, 4H, (CH₂)^{9''} and (CH₂)^{11'}), 1.73 (s, 6H, CH₃-C^{8''} and CH₃-C^{10''}), 2.00 (s, 3H, CH₃-C^{6'}), 2.03 (m, 4H, (CH₂)^{10''} and (CH₂)^{12'}), 2.05 (s, 3H, CH₃-C^{4''}), 2.36 (s, 3H, CH₃-C^{2'}), 6.12-6.23 (m, 5H, 5 × vinylic CH), 6.29 (d, *J* = 16.1 Hz, 1H, vinylic CH), 6.46 (d, *J* = 15.3 Hz, 1H, H^{1''}), 6.47 (d, *J* = 14.9 Hz, 1H, H^{3'}), 6.52 (s, 1H, H^{1'}), 6.83 (dd, *J* = 14.9 Hz, 11.4 Hz, 1H, H^{4'}), 7.08 (d, *J* = 5.1 Hz, 1H, H^{5'}), 7.20 (s, 1H, H^{3'}), 7.35 (dd, *J* = 15.3 Hz, 11.6 Hz, 1H, H^{2''}), 8.51 (d, *J* = 5.1 Hz, 1H, H^{6'}) ppm. ¹³C-NMR (CDCl₃) δ 12.8 (CH₃, CH₃-C^{6'}), 12.9 (CH₃, CH₃-C^{4''}), 14.1 (CH₃, CH₃-C^{2'}), 19.2 (CH₂, C^{9''} or C^{11'}), 19.3 (CH₂, C^{9''} or C^{11'}), 21.7 (CH₃, CH₃-C^{8''} and CH₃-C^{10''}), 29.0 (CH₃, 2 × CH₃-C^{12''} and 2 × CH₃-C^{14''}), 33.1 (CH₂, C^{10''} and C^{12'}), 34.3 (C, C^{12''} and C^{14''}), 39.60 (CH₂, C^{11''} or C^{13''}), 39.64 (CH₂, C^{11''} or C^{13''}), 117.4 (CH, C^{5'}), 122.0 (CH, C^{3'}), 126.7 (CH, C^{4'}), 127.1 (CH, C^{6''} or C^{8''}), 128.7 (CH, C^{6''} or C^{8''}), 129.1 (CH, C^{1''}), 129.2 (CH, C^{3''}), 129.4 (CH, C^{2''}), 130.0 (C, C^{8''} or C^{10''}), 130.1 (CH, C^{1'}), 130.3 (CH, C^{5'}), 136.8 (C, C^{6'}), 137.2 (CH, C^{5''}), 137.68 (CH, C^{7'}), 137.71 (C, C^{7''} or C^{9''}), 137.8 (CH, C^{3'}), 137.9 (C, C^{7''} or C^{9''}), 139.4 (C, C^{4''}), 140.5 (C, C^{2'}), 145.1 (C, C^{4'}), 149.4 (CH, C^{6'}), 157.4 (C, C^{2'}) ppm. IR (NaCl, cm⁻¹) 2928 (s, C-H), 2865 (m, C-H), 1714 (w), 1675 (w), 1590 (s), 1450 (m), 1362 (m), 1257 (w), 1169 (w), 969 (s), 899 (w), 755 (s). MS (EI+) *m/z* (%) 549 ([M+2]⁺, 11), 548 ([M+1]⁺, 45), 547 (M⁺, 100), 533 (11), 532 ([M-Me]⁺, 24), 411 (14), 410 (31), 358 (16). High resolution MS calc for C₄₀H₅₃N 547.4178; found 547.4161. UV (MeOH) λ_{max} = 368 nm (ε 49900). **15:** ¹H-NMR (CDCl₃) δ 0.89 (t, 9H, *J* = 7.3 Hz, 3 × SnCH₂CH₂CH₂CH₃), 0.99-1.06 (m, 6H, 3 × SnCH₂CH₂CH₂CH₃), 1.04 (s, 6H, 2 × CH₃-C^{12''}), 1.29-1.38 (m, 6H, 3 × SnCH₂CH₂CH₂CH₃), 1.46-1.49 (m, 2H, (CH₂)^{11''}), 1.50-1.56 (m, 6H, 3 × SnCH₂CH₂CH₂CH₃), 1.61-1.66 (m, 2H, (CH₂)^{10''}), 1.73 (s, 3H, CH₃-C^{8''}), 2.00-2.04 (m, 2H, (CH₂)^{9''}), 2.04 (s, 3H, CH₃-C^{4''}), 2.27 (d, 3H, *J* = 1.0 Hz, CH₃-C^{2'}), 5.34 (d, *J* = 2.1 Hz, ³*J* ¹H-¹¹⁹Sn ≈ ³*J* ¹H-¹¹⁷Sn ≈ 62.2 Hz, 1H, H^{4'}-*cis*), 6.04 (d, *J* = 2.1 Hz, ³*J* ¹H-¹¹⁹Sn = 133.4 Hz, ³*J* ¹H-¹¹⁷Sn = 128.7 Hz, 1H, H^{4'}-*trans*), 6.16 (d, *J* = 16.2 Hz, 1H, H^{5''}), 6.22 (d, *J* = 11.4 Hz, 1H, H^{3''}), 6.29 (d, *J* = 16.2 Hz, 1H, H^{6''}), 6.33 (br s, 1H, H^{1'}), 6.46 (d, *J* = 15.4 Hz, 1H, H^{1''}), 7.09 (dd, *J* = 5.1 Hz, 1.3 Hz, 1H, H^{5'}),

³ Ren, R. X.-F.; Sakai, N.; Nakanishi, K. *J. Am. Chem. Soc.* **1997**, *119*, 3619-3620.

7.12 (br s, 1H, H³), 7.34 (dd, $J = 15.4$ Hz, 11.4 Hz, 1H, H^{2''}), 8.51 (d, $J = 5.1$ Hz, 1H, H⁶) ppm. ¹³C-NMR (CDCl₃) δ 10.5 (CH₂, 3 $\times$ SnCH₂CH₂CH₂CH₃), 12.9 (CH₃, CH₃-C^{4''}), 13.7 (CH₃, 3 $\times$ SnCH₂CH₂CH₂CH₃), 16.5 (CH₃, CH₃-C^{2'}), 19.2 (CH₂, C^{10''}), 21.7 (CH₃, CH₃-C^{8''}), 27.3 (CH₂, 3 $\times$ SnCH₂CH₂CH₂CH₃), 29.0 (CH₃, 2 $\times$ CH₃-C^{12''}), 29.1 (CH₂, 3 $\times$ SnCH₂CH₂CH₂CH₃), 33.1 (CH₂, C^{9''}), 34.3 (C, C¹²), 39.6 (CH₂, C^{11''}), 117.4 (CH, C⁵), 121.7 (CH, C³), 125.4 (CH₂, C^{4'}), 127.5 (CH, C^{1'}), 128.6 (CH, C^{6''}), 129.2 (2 $\times$ CH, C^{1''} and C^{3''}), 129.3 (CH, C^{2''}), 129.9 (C, C^{8''}), 137.2 (CH, C^{5''}), 137.7 (C, C^{7''}), 139.3 (C, C^{4''}), 145.0 (C, C^{4'}), 147.2 (C, C^{2'}), 149.4 (CH, C⁶), 157.8 (C, C²), 158.0 (C, C^{3'}) ppm. IR (NaCl, cm⁻¹) 2955 (s, C-H), 2924 (s, C-H), 2856 (s, C-H), 1586 (s), 1538 (m), 1459 (s), 1379 (m), 1288 (w), 1260 (w), 1155 (w), 1073 (w), 1022 (w), 964 (s), 893 (w), 810 (w). MS (ESI) Calc for C₃₈H₆₀N¹²⁰Sn₂ 650.3742; found 650.3737. UV (MeOH) $\lambda_{\text{max}} = 217$ nm (ϵ 6500), 358 nm (ϵ 32800).

A2-E (1)

A mixture of **14** (25 mg, 0.05 mmol) and 2-iodoethanol (0.034 mL, 0.43 mmol) in CH₃NO₂ (1.5 mL) was heated at 100 °C in the dark for 18 h. The reaction mixture was concentrated under reduced pressure. Purification was performed by flash chromatography on silica gel in the dark (SiO₂, 92:8:0.001 CH₂Cl₂/MeOH/TFA) affording 24 mg (74%) of A2E (**1**)⁴ as a bright red syrup and 2 mg (6%) of *iso*-A2E.⁵ **A2-E (1):** ¹H-NMR (CD₃OD) δ 1.05 (s, 6H, CH₃-C^{12''} or CH₃-C^{14'}), 1.06 (s, 6H, CH₃-C^{12''} or CH₃-C^{14'}), 1.51 (m, 4H, cyclic CH₂), 1.66 (m, 4H, cyclic CH₂), 1.73 (s, 3H, CH₃-C^{8''} or CH₃-C^{10'}), 1.75 (s, 3H, CH₃-C^{8''} or CH₃-C^{10'}), 2.05 (s, 3H, CH₃), 2.08 (m, 4H, cyclic CH₂), 2.16 (s, 3H, CH₃), 2.18 (s, 3H, CH₃), 3.93 (t, $J = 4.8$ Hz, 2H, NCH₂CH₂OH), 4.55 (t, $J = 4.8$ Hz, 2H, NCH₂CH₂OH), 6.18 (d, $J = 16.0$ Hz, 1H, H^{5''} or H^{6''} or H^{7''} or H^{8''}), 6.25 (d, $J = 11.4$ Hz, 1H, H^{3''} or H^{5'}), 6.28 (d, $J = 16.0$ Hz, 1H, H^{5''} or H^{6''} or H^{7''} or H^{8''}), 6.34 (d, $J = 16.2$ Hz, 1H, H^{5''} or H^{6''} or H^{7''} or H^{8''}), 6.42 (d, $J = 11.7$ Hz, 1H, H^{3''} or H^{5'}), 6.55 (d, $J = 16.2$ Hz, 1H, H^{5''} or H^{6''} or H^{7''} or H^{8''}), 6.63 (d, $J = 15.0$ Hz, 1H, H^{1''} or H^{3'}), 6.71 (s, 1H, H^{1'}), 6.77 (d, $J = 15.3$ Hz, 1H, H^{1''} or H^{3'}), 7.12 (dd, $J = 15.0$ Hz, 11.4 Hz, 1H, H^{2''} or H^{4'}), 7.86 (s, 1H, H³), 7.94 (d, $J = 7.0$ Hz, 1H, H⁵), 7.99 (dd, $J = 15.3$ Hz, 11.7 Hz, 1H, H^{2''} or H^{4'}), 8.55 (d, $J = 7.0$ Hz, 1H, H⁶) ppm. ¹³C-NMR (CD₃OD) δ 12.9 (CH₃), 13.3 (CH₃), 15.0 (CH₃), 20.2 (CH₂), 20.3 (CH₂), 21.9 (CH₃), 22.0 (CH₃), 29.4 (CH₃), 34.0 (CH₂), 34.1 (CH₂), 35.26 (C), 35.29 (C), 40.8 (CH₂), 60.1

⁴ Sakai, N.; Decatur, J.; Nakanishi, K. *J. Am. Chem. Soc.* **1996**, *118*, 1559-1560.

⁵ Parish, C. A.; Hashimoto, M.; Nakanishi, K.; Dillon, J.; Sparrow, J. *Proc. Natl. Acad. Sci. USA* **1998**, *95*, 14609-14613.

(CH₂), 61.1 (CH₂), 120.4 (CH), 121.2 (CH), 127.0 (CH), 127.2 (CH), 129.8 (CH), 130.5 (CH), 130.8 (CH), 130.9 (C), 132.1 (CH), 132.2 (C), 133.0 (CH), 135.6 (CH), 138.4 (CH), 138.87 (CH), 138.92 (C), 139.0 (C), 139.4 (CH), 140.9 (C), 146.1 (CH), 146.9 (C), 148.9 (C), 153.4 (C), 154.7 (C) ppm.

14