SUPPORTING INFORMATION

Hetero-Allylic Asymmetric Alkylation(h-AAA): Cu-Catalyzed Asymmetric Synthesis of Chiral Allylic Esters

Koen Geurts, Stephen P. Fletcher and Ben L. Feringa*

*Department of Organic Chemistry and Molecular Inorganic Chemistry,
Stratingh Institute, University of Groningen, Nijenborgh 4, 9747 AG,
Groningen, The Netherlands*

Contents:

General information	S2
General procedure for the synthesis of racemic allylic esters	S2
General procedure for asymmetric synthesis of allylic esters	S2
Preparation of allylic esters 2	S3
Synthesis of (+)-(<i>S</i>)-5-Ethyl-2(<i>5H</i>)-furanone	S5
Synthesis of (–)-(<i>S</i>)-Benzoic acid-cyclopent-2-enyl ester	S5
Notes and references	S5
¹ H NMR and ¹³ C NMR spectra for compounds 2, 6 and 7	S6
Representative chiral HPLC chromatograms	S17

General information: All reactions were carried out under nitrogen atmosphere and in flame dried glassware unless otherwise noted. The following Grignard reagents were purchased from Aldrich as solutions in Et₂O: MeMgBr (3M), EtMgBr (3M), *i*-BuMgBr (2M), PentMgBr (2M). Other Grignard reagents were prepared from the corresponding alkyl bromides and magnesium turnings in Et₂O following standard procedures. Grignard reagents were titrated using *s*-BuOH and catalytic amounts of 1,10-phenanthroline. Ligands **4** and **5** were generously donated by Solvias (Basel) and CuBr•Me₂S and benzoyl bromide were purchased from Acros. Solvents were purified before use employing standard techniques.¹ Column chromatography was performed using ultra pure silica gel (60 Angstrom, 40–63 μm, Silicycle, Canada) as the stationary phase. TLC was performed using precoated silica gel plates (0.25 mm thick, 60F₂₅₄) and the product was observed under UV light or with either phosphomolybdic acid dip or KMnO₄ based dip. All ¹H NMR and ¹³C NMR spectra were recorded in CDCl₃ on Varian VXR-300 (300 MHz) or a Varian Mercury Plus (400 MHz) spectrometer. Chemical shifts were determined relative to the residual solvent peaks (CDCl₃, δ = 7.26 ppm for ¹H-NMR, δ = 77.0 ppm for ¹³C NMR). Data are reported as follows: chemical shift, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, br = broad and m = multiplet), coupling constants and integration. Melting points were determined on a Buchi B-545 melting point apparatus and are uncorrected. Mass spectra were recorded on an AEI-MS-902 mass spectrometer. Optical rotations were measured on a Smidt + Haensdch polarimeter (Polartronic MH8) with a 10 cm cell (c given in g/100 mL). Ee's of allylic esters were, in all cases except **2h**, determined by comparison of the HPLC traces of the corresponding racemic esters on a chiral non-racemic stationary phase (Chiracel OB-H or OD-H column) using a Shimadzu LC 10AD VP HPLC.

Synthesis of starting materials: *Trans*-**1**, **1a** and **1b** were readily prepared in a single step from the corresponding acid bromide and acrolein by simply mixing in CH₂Cl₂ for 72 h at room temperature and subsequent recrystallization according to the procedure of Trombini and Lombardo *et al.*² **1c** was prepared by heating a CH₂Cl₂ solution of methacrolein and benzoyl bromide to reflux overnight before the usual workup procedure.

General procedure for the synthesis of racemic allylic esters: Racemic allylic esters were synthesized by modification of a literature procedure.³ The Grignard reagent (1.0 eq) in Et₂O (1–3M) was added dropwise over 5 min to a stirred and cooled (–15 °C) solution of freshly distilled acrolein (1.1 eq) in THF. Stirring was continued for 1 h before the mixture was warmed to room temperature, quenched with water, partitioned between water and ether and the aqueous layer extracted with Et₂O. The combined organic layers were washed with brine, dried (MgSO₄), and the solvent carefully evaporated *in vacuo*. The crude alcohol (1 eq) was used without further purification. Benzoyl chloride (1.1 eq) was injected into a cooled (0 °C) solution of crude alcohol (1 eq) and pyridine (2 eq) in CH₂Cl₂. The reaction mixture was allowed to warm to room temperature and stirred for 1 h, before the solvent was evaporated *in vacuo*. The residue was taken up in Et₂O and washed with aq. 2N KOH, aq. 1N H₂SO₄, sat. aq. NaHCO₃ and brine. The organic layer was dried (MgSO₄), and the solvent evaporated *in vacuo*. Flash chromatography of the residue over silica gel, using pentane or Et₂O-pentane mixtures, gave racemic allylic esters as clear colorless oils.

General procedure for the Cu-catalyzed asymmetric synthesis of allylic esters with Grignard reagents

Several points deserved noting: Two general procedures (A and B, see below) were used for these experiments – to the best of our knowledge the experimental procedures give identical results. We found procedure B more convenient and used it for our later experiments. In general the yields of the reactions appear higher when done at higher concentrations. All reactions were conducted using 5 mol% catalyst, although two experiments demonstrate that much lower catalyst loading (0.8 and 0.05 mol %) can be used without deleterious effects. Reactions are typically very clean and only **2**, **3**, and the catalyst can be observed after completion: Flash chromatography (as a glorified filtration) with pentane or Et₂O-pentane mixtures served to remove the catalyst from the product(s). In the case of substrates **2g** and **2h**, more elaborate Grignard reagents were used and the reaction was accompanied by by-products which appear to arise from homo-coupling of the Grignard, purification of **2g** and **2h** therefore required more careful chromatography. Substrate **2i**, also required proper flash chromatography to provide analytically pure product. Recovery of copper-catalyst complex is easily accomplished by flushing the column with EtOAc and collecting the yellow coloured fractions.

A: CH₂Cl₂ was added to a mixture of CuBr•SMe₂ (5 mol%) and Taniaphos (6 mol%) under a nitrogen atmosphere at room temperature and the orange solution was stirred for 10 min before cooling to –75 °C. While ensuring that stirring continued, the Grignard reagent (1.3 eq in Et₂O) was added dropwise over 2 min. After an additional 15 min the allylic bromide in CH₂Cl₂ (0.7 mL) was added dropwise over 5 min. Stirring at –75 °C was continued (typically overnight) and the reaction mixture was then quenched by the addition of MeOH (*ca.* 1 mL). The reaction mixture was removed from the cooling bath and sat. aq. NH₄Cl (*ca.* 1 mL), and then water (*ca.* 2 mL) were added. The mixture was partitioned between Et₂O (5 mL) and water, and the aqueous layer extracted three times with Et₂O. The combined organic layers were dried (MgSO₄), filtered and the solvent evaporated *in vacuo*.

B: The Grignard reagent (1.3–2.0 eq in Et₂O) was added dropwise over 5 min to a homogeneous, stirred and cooled (–75 °C) solution of the allylic bromide, CuBr•SMe₂ (5 mol%) and Taniaphos (5 mol%) in CH₂Cl₂ (2.5 mL) under a nitrogen atmosphere. If the solution was not again homogeneous after 5 min, further minimal portions of CH₂Cl₂ were added to dissolve precipitates. Stirring was continued until NMR or TLC (5% Et₂O/pentane) analysis showed the reaction had reached completion (typically overnight) and the reaction was quenched with MeOH (5 mL). The reaction mixture was removed from the cooling bath and sat. aq. NH₄Cl (*ca.* 5 mL) was added. The mixture was partitioned between CH₂Cl₂ (5 mL) and water. The organic layer was dried (MgSO₄), filtered and the solvent evaporated *in vacuo*.

Purification of Crude Products (for procedures A or B): Flash chromatography of the residue over silica gel using pentane or Et₂O-pentane gave allylic esters as clear colorless oils. As stated above compounds **2g**, **2h** and **2i** required careful flash chromatography.

Preparation of allylic esters 2:

(+)-(S)-Benzoic acid-1-methyl-allyl ester: **2**⁴

Methylmagnesium bromide (3.0 M, 0.50 mL, 1.5 mmol) was added dropwise over 2 min to a stirred and cooled (-73°C) solution of (**180** mg, 0.75 mmol), $\text{CuBr}\cdot\text{SMe}_2$ (8.1 mg, 39 μmol) and (*R,S*)-Taniaphos (31.2 mg, 45.4 μmol) in CH_2Cl_2 (3 mL) under a nitrogen atmosphere. Stirring was continued for 18 h at -73°C before the reaction was quenched with MeOH (2 mL). The reaction mixture was removed from the cooling bath and sat. aq. NH_4Cl (ca. 5 mL) was added. The mixture was partitioned between CH_2Cl_2 (5 mL) and water, and the aqueous layer extracted with CH_2Cl_2 . The combined organic layers were dried (MgSO_4), filtered and the solvent evaporated *in vacuo*. Flash chromatography of the dark orange residue over silica gel, using first pentane and then 100:1 pentane-Et₂O gave (+)-(S)-benzoic acid-1-methyl-allyl ester **2** (112 mg, 85%) as a clear colorless oil: ^1H NMR (CDCl_3 , 400 MHz) δ 8.11–8.03 (m, 2 H), 7.59–7.51 (m, 1 H), 7.47–7.39 (m, 2 H), 5.97 (ddd, $J = 5.7, 10.6, 17.3$ Hz, 1 H), 5.67–5.55 (m, 1 H), 5.34 (td, $J = 1.3, 17.3$ Hz, 1 H), 5.19 (td, $J = 1.3, 10.6$ Hz, 1 H), 1.45 (d, $J = 6.5$ Hz, 3 H); ^{13}C NMR (CDCl_3 , 100.6 MHz) δ 165.8 (s), 137.7 (d), 132.8 (d), 130.6 (s), 129.5 (d, 2xC), 128.3 (d, 2xC), 115.8 (t), 71.5 (d), 20.0 (q); exact mass m/z calcd for $\text{C}_{11}\text{H}_{12}\text{O}_2$ 176.08372, found 176.08445; $[\alpha]_{\text{D}}^{27} = +43.7$ (c 0.35, CHCl_3), Lit⁵: $[\alpha]_{\text{D}}^{20} = +48.03$ (EtOH). HPLC analysis indicated an enantiomeric excess of 98% [Chiralcel OB-H column; flow: 0.5 mL/min; heptane/*i*-PrOH: 99.75:0.25; $\lambda = 220$ nm; major enantiomer (+)-**2**, $t_{\text{R}} = 13.65$ min; minor enantiomer (–)-**2**, $t_{\text{R}} = 15.25$ min]. Absolute configuration assigned by comparison of optical rotation.⁵

(+)-(E)-3-Phenyl-acrylic acid-1-methyl-allyl ester: **2a**

CH_2Cl_2 (1 mL) was added to a mixture of $\text{CuBr}\cdot\text{SMe}_2$ (3.4 mg, 17 μmol) and (*R,S*)-Taniaphos (15.4 mg, 22.5 μmol) under a nitrogen atmosphere at room temperature, and the resulting orange solution was stirred for 10 min before cooling to -77°C . While ensuring that stirring continued, MeMgBr (3M, 0.14 mL, 0.47 mmol) was added dropwise over 2 min, to provide a yellow mixture. After an additional 15 min **1a** (100.2 mg, 0.375 mmol) in CH_2Cl_2 (0.7 mL) was added dropwise over 5 min. Stirring at -77°C was continued for 13 h and the reaction mixture was then quenched by the addition of MeOH (ca. 1 mL). The reaction mixture was removed from the cooling bath and sat. aq. NH_4Cl (ca. 1 mL), and then water (ca. 2 mL) were added. The mixture was partitioned between Et₂O (5 mL) and water, and the aqueous layer extracted three times with Et₂O. The combined organic layers were dried (MgSO_4), filtered and the solvent evaporated *in vacuo*. Flash chromatography of the dark orange residue over silica gel using 100:1 pentane-Et₂O gave (+)-(E)-3-Phenyl-acrylic acid-1-methyl-allyl ester **2a** (60 mg, 80%) as a clear colorless oil: ^1H NMR (300 MHz) δ 7.70 (d, $J = 16.0$ Hz, 1 H), 7.52–7.46 (m, 2 H), 7.37–7.32 (m, 3 H), 6.46 (d, $J = 16.0$ Hz, 1 H), 5.93 (ddd, $J = 5.8, 10.5, 17.2$ Hz, 1 H), 5.55–5.46 (m, 1 H), 5.31 (td, $J = 1.2, 17.3$ Hz, 1 H), 5.18 (td, $J = 1.2, 10.5$ Hz, 1 H), 1.40 (d, $J = 6.5$ Hz, 3 H); ^{13}C NMR (CDCl_3 , 100.6 MHz) δ 165.8 (s), 144.4 (d), 137.5 (d), 134.2 (s), 129.9 (d), 128.6 (d, 2 x C), 127.8 (d, 2 x C), 118.2 (d), 115.5 (t), 70.7 (d), 19.7 (q); exact mass m/z calcd for $\text{C}_{13}\text{H}_{14}\text{O}_2$ 202.09937, found 202.10045; $[\alpha]_{\text{D}}^{20} = +50.0$ (c 0.04, CHCl_3). HPLC analysis indicated an enantiomeric excess of 98% [Chiralcel OB-H column; flow: 0.5 mL/min; heptane/*i*-PrOH: 99.5:0.5; $\lambda = 225$ nm; minor enantiomer (–)-**2a**, $t_{\text{R}} = 21.10$ min; major enantiomer (+)-**2a**, $t_{\text{R}} = 22.35$ min].

(+)-2,4,6-Trimethylbenzoic acid-1-methyl-allyl ester: **2b**

Methylmagnesium bromide (3.0 M, 0.48 mL, 1.4 mmol) was added dropwise over 3 min to a stirred and cooled (-73°C) solution of **1b** (204 mg, 0.722 mmol), $\text{CuBr}\cdot\text{SMe}_2$ (8.0 mg, 38.9 μmol) and (*R,S*)-Taniaphos (30.0 mg, 43.6 μmol) in CH_2Cl_2 (3.5 mL) under a nitrogen atmosphere. Stirring was continued for 15 h at -73°C before the reaction was quenched with MeOH (5 mL). The reaction mixture was removed from the cooling bath and sat. aq. NH_4Cl (ca. 5 mL) was added. The mixture was partitioned between CH_2Cl_2 (5 mL) and water. The organic layer was dried (MgSO_4), filtered and the solvent evaporated *in vacuo*. Flash chromatography of the dark orange residue over silica gel, using 2% Et₂O in pentane gave (+)-2,4,6-trimethylbenzoic acid-1-methyl-allyl ester **2b** (143.1 mg, 91%), as a clear colorless oil: ^1H NMR (CDCl_3 , 400 MHz) δ 6.86 (s, 2 H), 5.96 (ddd, $J = 6.2, 0.5, 17.2$ Hz, 1 H), 5.65 (td, $J = 1.2, 6.4$ Hz, 1 H), 5.37 (td, $J = 1.3, 17.3$ Hz, 1 H), 5.22 (td, $J = 1.2, 10.5$ Hz, 1 H), 2.31 (s, 6 H), 2.29 (s, 3 H), 1.46 (d, $J = 6.5$ Hz, 3 H); ^{13}C NMR (CDCl_3 , 100.6 MHz) δ 169.2 (s), 139.0 (s), 137.4 (d), 134.8 (s, 2 x C), 131.1 (s), 128.2 (d, 2 x C), 116.4 (t), 71.6 (d), 21.0 (q), 19.9 (q), 19.5 (q, 2 x C); exact mass m/z calcd for $\text{C}_{14}\text{H}_{18}\text{O}_2$ 218.13067, found 218.13034; $[\alpha]_{\text{D}}^{28} = +12.2$ (c 0.33, CHCl_3). HPLC analysis indicated an enantiomeric excess of 98% [Chiralcel OB-H column; flow: 0.5 mL/min; heptane/*i*-PrOH: 99.5:0.5; $\lambda = 230$ nm; minor enantiomer (–)-**2b**, $t_{\text{R}} = 8.73$ min; major enantiomer (+)-**2b**, $t_{\text{R}} = 9.22$ min].

(+)-Benzoic acid-1-ethyl-allyl ester: 2c

Ethylmagnesium bromide (3.0 M, 0.50 mL, 1.5 mmol) was added dropwise over 5 min to a stirred and cooled (-79°C) solution of **1** (181 mg, 0.75 mmol), $\text{CuBr}\cdot\text{SMe}_2$ (7.2 mg, 35 μmol) and (*R,S*)-Taniaphos (30.3 mg, 44 μmol) in CH_2Cl_2 (3 mL) under a nitrogen atmosphere. Stirring was continued for 16 h at -79°C before the reaction was quenched with MeOH (1 mL). The reaction mixture was removed from the cooling bath and sat. aq. NH_4Cl (ca. 1 mL) was added. The mixture was partitioned between CH_2Cl_2 (5 mL) and water and the aqueous layer extracted with CH_2Cl_2 . The combined organic layers were dried (MgSO_4), filtered and the solvent evaporated *in vacuo*. Flash chromatography of the dark orange residue over silica gel, using first pentane and then 2% Et_2O -pentane gave (+)-benzoic acid-1-ethyl-allyl ester **2c** (124 mg, 87 %) as a clear colorless oil: ^1H NMR (400 MHz) δ 8.10–8.06 (m, 2 H), 7.58–7.54 (m, 1 H), 7.47–7.41 (m, 2 H), 5.90 (ddd, J = 6.2, 10.6, 17.2 Hz, 1 H), 5.48–5.41 (m, 1 H), 5.33 (td, J = 1.3, 17.3 Hz, 1 H), 5.22 (td, J = 1.3, 10.6 Hz, 1 H), 1.87–1.72 (m, 2 H), 0.99 (t, J = 7.4 Hz, 3 H); ^{13}C NMR (CDCl_3 , 100.6 MHz) δ 165.8 (s), 136.2 (d), 130.6 (s), 129.5 (d, 2 x C), 128.3 (d, 2 x C), 116.7 (t), 76.4 (d), 27.3 (t), 9.4 (q); exact mass m/z calcd for $\text{C}_{12}\text{H}_{14}\text{O}_2$ = 190.09937, found 190.09917; $[\alpha]_{\text{D}}^{25} = +43.8$ (c 1.5, CHCl_3). HPLC analysis indicated an enantiomeric excess of 98% [Chiralcel OB-H column; flow: 0.5 mL/min; heptane/*i*-PrOH: 99.5:0.5; λ = 210 nm; major enantiomer (+)-**2c**, t_{R} = 11.62 min; minor enantiomer (–)-**2c**, t_{R} = 13.01 min]

(–)-Benzoic acid-1-vinyl-hexyl ester: 2d⁶

Pentylmagnesium bromide (2.0 M, 0.76 mL, 1.5 mmol) was added dropwise over 5 min to a stirred and cooled (-75°C) solution of **1** (184.3 mg, 0.765 mmol), $\text{CuBr}\cdot\text{SMe}_2$ (8.0 mg, 50 μmol) and (*S,R*)-Taniaphos (30.0 mg, 43.6 μmol) in CH_2Cl_2 (2.5 mL) under a nitrogen atmosphere. After 5 min a further 1 mL CH_2Cl_2 was added to dissolve the resulting precipitates and stirring was continued for 66 h at -75°C before the reaction was quenched with MeOH (5 mL). The reaction mixture was removed from the cooling bath and sat. aq. NH_4Cl (ca. 5 mL) was added. The mixture was partitioned between CH_2Cl_2 (5 mL) and water. The organic layer was dried (MgSO_4), filtered and the solvent evaporated *in vacuo*. Flash chromatography of the dark orange residue over silica gel, using first pentane and then 1% Et_2O pentane gave (–)-benzoic acid-1-vinyl-hexyl ester **2d** (164.0 mg, 99%) as a clear colorless oil: ^1H NMR (CDCl_3 , 400 MHz) δ 8.07 (apparent d, J = 8.4 Hz, 2 H), 7.59–7.51 (m, 1 H), 7.47–7.40 (m, 2 H), 5.90 (ddd, J = 6.3, 10.5, 16.9 Hz, 1 H), 5.50 (dd, J = 6.4, 12.6 Hz, 1 H), 5.33 (dd, J = 1.1, 17.2 Hz, 1 H), 5.20 (dd, J = 1.1, 10.5 Hz, 1 H), 1.88–1.62 (m, 2 H), 1.50–1.18 (m, 6 H), 0.89 (br s, 3 H); ^{13}C NMR (CDCl_3 , 100.6 MHz) δ 166.1 (s), 136.9 (d), 133.1 (d), 130.8 (s), 129.8 (d, 2 x C), 128.5 (d, 2 x C), 116.7 (t), 75.6 (d), 34.5 (t), 31.8 (t), 24.9 (t), 22.8 (t), 14.2 (q); exact mass m/z calcd for $\text{C}_{15}\text{H}_{20}\text{O}_2$ 232.14632, found 232.14714; $[\alpha]_{\text{D}}^{20} = -27.6$ (c 0.47, CHCl_3). HPLC analysis indicated an enantiomeric excess of 97% [Chiralcel OB-H column; flow: 0.5 mL/min; heptane/*i*-PrOH: 99.5:0.5; λ = 230 nm; minor enantiomer (+)-**2d**, t_{R} = 9.74 min; major enantiomer (–)-**2d**, t_{R} = 10.92 min].

(+)-(*S*)-Benzoic acid-1-vinyl-pent-4-enyl ester: 2f

Butenylmagnesium bromide (1.1M, 1.37 mL, 1.51 mmol) was added dropwise over 2 min to a stirred and cooled (-73°C) solution of **1** (182.0 mg, 0.755 mmol), $\text{CuBr}\cdot\text{SMe}_2$ (8.0 mg, 50 μmol) and (*R,S*)-Taniaphos (30.0 mg, 43.6 μmol) in CH_2Cl_2 (3.5 mL) under a nitrogen atmosphere. Stirring was continued for 20 h at -73°C before the reaction was quenched with MeOH (5 mL). The reaction mixture was removed from the cooling bath and sat. aq. NH_4Cl (ca. 5 mL) was added. The mixture was partitioned between CH_2Cl_2 (5 mL) and water. The organic layer was dried (MgSO_4), filtered and the solvent evaporated *in vacuo*. Flash chromatography of the dark orange residue over silica gel, using 2% Et_2O -pentane gave (+)-(*S*)-benzoic acid-1-vinyl-pent-4-enyl ester **2f** (156.3 mg, 96%) as a clear colorless oil: ^1H NMR (400 MHz) δ 8.11–8.06 (m, 2 H), 7.59–7.53 (m, 1 H), 7.48–7.42 (m, 2 H), 5.87 (dddd, J = 6.4, 10.4, 13.2, 27.3 Hz, 2 H), 5.57–5.51 (m, 1 H), 5.35 (td, J = 1.3, 17.2 Hz, 1 H), 5.23 (td, J = 1.2, 10.5 Hz, 1 H), 5.05 (ddd, J = 1.6, 3.4, 17.1 Hz, 1 H), 5.02–4.98 (m, 1 H), 2.20 (qdd, J = 1.4, 6.7, 13.3 Hz, 2 H), 1.98–1.88 (m, 2 H); ^{13}C NMR (CDCl_3 , 100.6 MHz) δ 165.6 (s), 137.4 (d), 136.2 (d), 132.7 (d), 130.3 (s), 129.4 (d, 2 x C), 128.2 (d, 2 x C), 116.7 (t), 115.1 (t), 74.5 (d), 33.3 (t), 29.2 (t); exact mass m/z calcd for $\text{C}_{14}\text{H}_{16}\text{O}_2$ 216.11502, found 216.11595; $[\alpha]_{\text{D}}^{20} = +27.2$ (c 0.29, CHCl_3). HPLC analysis indicated an enantiomeric excess of 97% [Chiralcel OB-H column; flow: 0.5 mL/min; heptane/*i*-PrOH: 99.75:0.25; λ = 220 nm; major enantiomer (+)-**2f**, t_{R} = 15.82 min; minor enantiomer (–)-**2f**, t_{R} = 18.76 min]. The absolute stereochemistry is based on the conversion to known **7**.

(+)-Benzoic acid-1-phenethyl-allyl ester: 2g⁷

Phenethylmagnesium bromide (1.1M, 1.37 mL, 1.51 mmol) was added dropwise over 2 min to a stirred and cooled (-73°C) solution of **1** (183.1 mg, 0.760 mmol), $\text{CuBr}\cdot\text{SMe}_2$ (8.0 mg, 50 μmol) and (*R,S*)-Taniaphos (30.0 mg, 43.6 μmol) in CH_2Cl_2 (3.5 mL) under a nitrogen atmosphere. Stirring was continued for 16 h at -73°C before the reaction was quenched with MeOH (5 mL). The reaction mixture was removed from the cooling bath and sat. aq. NH_4Cl (ca. 5 mL) was added. The mixture was partitioned between CH_2Cl_2 (5 mL) and water. The organic layer was dried (MgSO_4), filtered and the solvent evaporated *in vacuo*. Flash chromatography of the dark orange residue over silica gel, using first pentane and then 5% Et_2O -pentane gave impure (+)-benzoic acid-1-phenethyl-allyl ester **2g**. A second, more careful, round of chromatography over silica gel, using first pentane and then 2% Et_2O pentane gave (+)-benzoic acid-1-phenethyl-allyl ester **2g** (188.2 mg, 93%) which still contained a ^1H NMR detectable amount of Grignard homo-coupled impurity: ^1H NMR (400 MHz) δ 8.12–8.08 (m, 2 H), 7.62–7.56 (m, 1 H), 7.51–7.45 (m, 2 H), 7.34–7.28 (m, 2 H), 7.24–7.18 (m, 3 H), 5.96 (ddd, $J = 6.2, 10.6, 17.1$ Hz, 1 H), 5.57 (ddd, $J = 1.1, 6.0, 12.0$ Hz, 1 H), 5.39 (td, $J = 1.3, 17.2$ Hz, 1 H), 5.27 (td, $J = 1.2, 10.6$ Hz, 1 H), 2.78 (ddd, $J = 4.7, 6.4, 9.0$ Hz, 2 H), 2.23–2.04 (m, 2H); ^{13}C NMR (CDCl_3 , 100.6 MHz) δ 166.0 (s), 141.6 (s), 136.5 (d), 133.2 (d), 130.7 (s), 129.9 (d, 2 x C), 128.7 (d, 2 x C), 128.6 (d, 4 x C), 126.3 (d), 117.3 (t), 75.0 (d), 36.2 (t), 31.7 (t); exact mass m/z calcd $\text{C}_{18}\text{H}_{18}\text{O}_2$ 266.13067, found 266.12978; $[\alpha]_{589}^{20} = +8.1$ (c 0.42, CHCl_3). HPLC analysis indicated an enantiomeric excess of 93 % [Chiralcel OB-H column; flow: 0.5 mL/min; heptane/*i*-PrOH: 98:2; $\lambda = 220$ nm; major enantiomer (+)-**2g**, $t_R = 15.28$ min; minor enantiomer (–)-**2g**, $t_R = 26.58$ min].

(+)-Benzoic acid-1-vinyl-nonadecyl ester: 2h

Octadecane magnesium bromide (1.5M in ether, 0.98 mL, 1.5 mmol) was added at a fast dropwise rate over 1 min (to avoid solidification of the Grignard in the syringe) to a stirred and cooled (-70°C) solution of **1** (177.5 mg, 0.736 mmol), $\text{CuBr}\cdot\text{SMe}_2$ (8.0 mg, 50 μmol) and (*R,S*)-Taniaphos (30.0 mg, 43.6 μmol) in CH_2Cl_2 (2.5 mL) under a nitrogen atmosphere. After 5 min a further 1 mL of CH_2Cl_2 was added to dissolve the resulting precipitates and stirring was continued for 18 h at -70°C before the reaction was quenched with MeOH (5 mL). The reaction mixture was removed from the cooling bath and sat. aq. NH_4Cl (ca. 5 mL) was added. The mixture was partitioned between CH_2Cl_2 (5 mL) and water. The organic layer was dried (MgSO_4), filtered and the solvent evaporated *in vacuo*. Flash chromatography of the dark orange residue over silica gel, using first pentane and then 5% Et_2O -pentane gave impure (+)-benzoic acid-1-vinyl-nonadecyl ester **2h**. A solid by-product was removed by crystallization from hot diisopropyl ether (ca. 5 mL) and filtration. The mother liquor was concentrated and flash chromatography of the colorless residue over silica gel, using first pentane and then 2% Et_2O -pentane gave pure (+)-benzoic acid-1-vinyl-nonadecyl ester **2h** (284.3 mg, 93%) as a clear colorless oil: ^1H NMR (400 MHz) δ 8.07 (apparent d, $J = 8.4$ Hz, 2 H), 7.58–7.52 (m, 1 H), 7.47–7.41 (m, 2 H), 5.95–5.84 (m, 1 H), 5.49 (apparent q, $J = 6.5$ Hz, 1 H), 5.32 (dd, $J = 0.6, 17.2$ Hz, 1 H), 5.20 (dd, $J = 1.0, 10.5$ Hz, 1 H), 1.87–1.66 (m, 2 H), 1.45–1.20 (m, 32 H), 0.88 (t, 3 H, $J = 6.8$ Hz); ^{13}C NMR (CDCl_3 , 100.6 MHz) δ 165.9 (s), 136.6 (d), 132.8 (d), 130.6 (s), 129.6 (d, 2 x C), 128.3 (d, 2 x C), 116.5 (t), 75.4 (d), 34.3 (t), 31.9 (t), 29.7 (t, 7 x C), 29.7 (t), 29.6 (t), 29.5 (t), 29.4 (t), 25.1 (t), 22.7 (t), 14.1 (q); exact mass m/z calcd for $\text{C}_{28}\text{H}_{46}\text{O}_2$ 414.34977, found 414.35186. Optical rotation was measured on the corresponding alcohol of (+)-benzoic acid-1-vinyl-nonadecyl ester: $[\alpha]_{589}^{24} = +17.0$ (c 0.094, CHCl_3). Enantiomeric excess was measured by ^1H NMR and ^{19}F NMR analysis of the corresponding Mosher⁸ ester >95%.

(+)-(S)-(E)-3-Phenyl-acrylic acid-1-ethyl-allyl ester: 2i

Ethylmagnesium bromide (3.0 M, 2.1 mL, 6.4 mmol) was added dropwise over 5 min to a stirred and cooled (-73°C) solution of **1b** (852.7 mg, 3.194 mmol), $\text{CuBr}\cdot\text{SMe}_2$ (4.8 mg, 23 μmol) and (*R,S*)-Taniaphos (17.2 mg, 25.0 μmol) in CH_2Cl_2 (7 mL) under a nitrogen atmosphere. Stirring was continued for 17 h at -73°C before the reaction was quenched with MeOH (5 mL). The reaction mixture was removed from the cooling bath and sat. aq. NH_4Cl (ca. 5 mL) was added. The mixture was partitioned between CH_2Cl_2 (5 mL) and water and the aqueous layer extracted with CH_2Cl_2 . The combined organic layers were dried (MgSO_4), filtered and the solvent evaporated *in vacuo*. Careful flash chromatography of the dark orange residue over silica gel, using Et_2O -pentane mixtures from 2–5% Et_2O -pentane gave (*S*)-(*E*)-3-phenyl-acrylic acid-1-ethyl-allyl ester **2i** (549.8 mg, 80%) as a clear colorless oil: ^1H NMR (400 MHz) δ 7.71 (d, $J = 16.0$ Hz, 1 H), 7.55–7.51 (m, 2 H), 7.40–7.36 (m, 3 H), 6.47 (d, $J = 16.0$ Hz, 1 H), 5.85 (ddd, $J = 6.3, 10.5, 17.0$ Hz, 1 H), 5.36–5.27 (m, 2 H), 5.21 (td, $J = 1.2, 10.6$ Hz, 1 H), 1.82–1.65 (m, 2 H), 0.96 (t, $J = 7.4$ Hz, 3H); ^{13}C NMR (CDCl_3 , 100.6 MHz) δ 166.2 (s), 144.6 (d), 136.3 (d), 134.4 (s), 130.2 (d), 128.8 (d, 2 x C), 127.9 (d, 2 x C), 118.3 (d), 116.7 (t), 76.0 (d), 27.3 (t), 9.4 (q); exact mass m/z calcd for $\text{C}_{14}\text{H}_{16}\text{O}_2$ 216.11502, found 216.11604; $[\alpha]_{589}^{20} = +53.4$ (c 0.058, CHCl_3). HPLC analysis indicated an enantiomeric excess of at least 98 % [Chiralcel OB-H column; flow: 0.5 mL/min; heptane/*i*-PrOH: 99.5:0.5; $\lambda = 225$ nm; major enantiomer (+)-**2i**, $t_R = 16.83$ min; minor enantiomer, racemate, (–)-**2i**, $t_R = 17.8$ min]. The absolute stereochemistry based on the conversion to known **6**.

(+)-(S)-5-Ethyl-2(5H)-furanone: 6⁹

Hoveyda-Grubbs II catalyst¹⁰ (15.3 mg, 0.024 mmol) was tipped into a stirred, dilute (0.005M) solution of (*S*)-(*E*)-3-phenyl-acrylic acid-1-ethyl-allyl ester **2i** (55.2 mg, 0.255 mmol) in, degassed CH₂Cl₂ (50 mL) under a nitrogen atmosphere. The reaction flask was placed in a preheated oil bath (60 °C) and stirred while heating to reflux for 43 h. The reaction mixture was cooled to room temperature and the solvent carefully removed under vacuum. Flash chromatography of the residue over silica gel, using EtOAc-pentane mixtures from 10–50% EtOAc gave (+)-(*S*)-5-ethyl-2(5H)-furanone **6** (21.8 mg, 78%) as a volatile colorless oil; ¹H NMR (400 MHz) δ 7.44 (dd, *J* = 1.5, 5.7 Hz, 1 H), 6.10 (dd, *J* = 2.0, 5.7 Hz, 1 H), 4.99 (tdd, *J* = 1.8, 5.3, 1.8 Hz, 1 H), 1.90–1.68 (m, 2 H), 0.99 (t, *J* = 7.4 Hz, 3 H); ¹³C NMR (CDCl₃, 100.6 MHz) δ 173.1 (s), 155.8 (d), 121.6 (d), 84.2 (d), 26.2 (t), 8.9 (q); [α]_D²⁹ = +76.9 (*c* 0.026, CHCl₃), lit⁹: [α]_D = +105 (*c* 4.1, CH₂Cl₂). HPLC analysis indicated an enantiomeric excess of 98% [Chiralcel OB-H column; flow: 0.5 mL/min; heptane/*i*-PrOH: 90:10; λ = 210 nm; major enantiomer (+)-**6**, *t*_R = 26.93 min; minor enantiomer (–)-**6**, *t*_R = 30.42 min]. Note 3

(–)-(S)-Benzoic acid-cyclopent-2-enyl ester: 7¹¹

Grubbs II catalyst¹² (19.6 mg, 0.0231 mmol) was tipped into a stirred solution of benzoic acid 1-vinyl-pent-4-enyl ester **2f** (97.1 mg, 0.450 mmol) in degassed CH₂Cl₂ (10 mL) under a nitrogen atmosphere. The reaction flask was placed in a preheated oil bath (60° C) and stirred while heating to reflux for 18 h. The reaction mixture was cooled to room temperature and the solvent removed *in vacuo*. Flash chromatography of the residue over silica gel, using a 2% Et₂O-pentane mixture gave (*S*)-benzoic acid-cyclopent-2-enyl ester **7** (72.0 mg, 85%) as a clear colorless oil; ¹H NMR (400 MHz) δ 8.03 (m, 2 H), 7.54 (m, 1 H), 7.42 (m, 2 H), 6.16 (m, 1 H), 5.95 (m, 2 H), 2.60 (m, 1 H), 2.39 (m, 2 H), 1.98 (m, 1 H); ¹³C NMR (CDCl₃, 100.6 MHz) δ 166.5 (s), 137.6 (d), 132.6 (d), 130.6 (s), 129.5 (d), 129.3 (d, 2 x C), 128.2 (d, 2 x C), 81.0 (d), 31.1 (t), 29.8 (t); exact mass *m/z* calcd C₁₂H₁₂O₂ 188.08372, found 188.08463; [α]_D²⁸ = –195.7 (*c* 0.19, CHCl₃), lit¹¹: [α]_D²⁵ = –98.9 (*c* 2.05, CHCl₃). HPLC analysis indicated an enantiomeric excess of 97% [Chiralcel OD-H column; flow: 0.5mL/min; heptane/*i*-PrOH: 99.9:0.1; λ = 225 nm; major enantiomer (–)-**7**, *t*_R = 18.57 min; minor enantiomer (+)-**7**, *t*_R = 30.42 min, 97% ee; Chiralcel OD column; flow: 1 mL/min; heptane/*i*-PrOH: 99.5:0.5; λ = 220 nm; major enantiomer (–)-**7**, *t*_R = 6.43 min; minor enantiomer (+)-**7**, *t*_R = 7.10 min, 97% ee]. Note 4.

Benzoic acid 1-ethyl-2-methylallyl ester: 2i

O, O'-(*R*)-(1,1'-dinaphthyl)-2,2'-diyl)-*R*,*R*'-di-(*R*,*R*)-1-phenylethylphosphoramidite (26.1 mg, 47 μmol) and CuTC (8.0 mg, 43 μmol) were added to a stirred solution of β-methyl allylbromide **1c** (218 mg, 0.859 mmol) in CH₂Cl₂ (2 mL) under a nitrogen atmosphere. Stirring at room temperature was continued for 30 min before the reaction was cooled to –70 °C. A solution of EtMgBr (3.0 M in diethyl ether, 0.57 mL, 1.7 mmol) was diluted with CH₂Cl₂ (1.2 mL), and the Grignard (now *ca* 1 M in 2 mL CH₂Cl₂-ether) was then injected slowly via syringe pump over 1.5 h. Although TLC (5% EtOAc/pentane) analysis showed the reaction had reached completion after a further 30 min, stirring at –70 °C was arbitrarily continued overnight before the reaction was quenched with EtOH (5 mL). The reaction mixture was removed from the cooling bath and sat. aq. NH₄Cl (*ca*. 5 mL) was added. The mixture was partitioned between CH₂Cl₂ (5 mL) and water, and the organic layer dried (MgSO₄), filtered and the solvent evaporated *in vacuo*. Flash chromatography of the residue over silica gel, using Et₂O-pentane mixtures from 0–5% Et₂O, gave a mixture of benzoic acid 1-ethyl-2-methylallyl ester **2i** and **3i** (170 mg, 97 %) as a 2.5:1 mixture contaminated with traces (¹H NMR spectroscopy) of phosphoramidite ligand. Note: Ee determination using the below method *requires* complete removal of **3i** and chiral ligand due to overlapping retention times. Repeated flash chromatography provided a sample free of **3i** (¹H NMR spectroscopy), which was then distilled (Kugelrohr, 7 mmHg, oven *ca* 200 °C) to removed traces of phosphoramidite ligand. Pure **2i** had: ¹H NMR (400 MHz) 8.08 (m, 2 H), 7.56 (m, 1 H), 7.45 (m, 2 H), 5.37 (t, *J* = 6.6 Hz, 1 H), 5.04 (s, 1 H), 4.94 (s, 1 H), 1.80 (m, 5 H), 0.96 (t, *J* = 7.4 Hz, 1 H); ¹³C NMR (CDCl₃, 100.6 MHz) δ 165.7(s), 142.8 (s), 132.7 (d), 130.5 (s), 129.5 (d 2 x C), 128.2 (d 2 x C), 112.7 (s), 79.0 (d), 25.6 (t), 18.1 (q), 9.6 (q); exact mass *m/z* calcd for C₁₂H₁₄O₂ = 204.11502, found 204.11557; HPLC analysis indicated an enantiomeric excess of 97% [Chiralcel OB-H column; flow: 0.5 mL/min; heptane/*i*-PrOH: 99:1; λ = 207 nm; major enantiomer **2i**, *t*_R = 8.94 min; minor enantiomer **2i**, *t*_R = 11.24 min]

Benzoic acid 1-isopropenylhexyl ester: **2j**

O, O'-(*R*)-(1,1'-dinaphthyl)-2,2'-diyl)-*R,R'*-di-(*R,R*)-1-phenylethylphosphoramidite (23.2 mg, 43 μ mol) and CuTC (7.5 mg, 39 μ mol) were added to a stirred solution of β -methyl allylbromide **1c** (200 mg, 0.780 mmol) in CH₂Cl₂ (2 mL) under a nitrogen atmosphere. Stirring at room temperature was continued for 30 min before the reaction was cooled to -70°C . A solution of PentylMgBr (1.2 M in diethyl ether, 1.3 mL, 1.56 mmol) was diluted with CH₂Cl₂ (2.6 mL), and the Grignard (now *ca* 0.4 M in 3.9 mL CH₂Cl₂-ether) was then injected slowly via syringe pump over 1 h. Stirring at -70°C was arbitrarily continued overnight before the reaction was quenched with MeOH (3 mL). The reaction mixture was removed from the cooling bath and sat. aq. NH₄Cl (*ca.* 5 mL) was added. The mixture was partitioned between CH₂Cl₂ (5 mL) and water, and the organic layer dried (MgSO₄), filtered and the solvent evaporated *in vacuo*. Flash chromatography of the residue over silica gel, using Et₂O-pentane mixtures from 0-5% Et₂O, gave a mixture of benzoic acid 1-isopropenylhexyl ester **2j** and **3j** (185 mg, 96 %) as a 2:1 mixture contaminated with traces (¹H NMR spectroscopy) of phosphoramidite ligand. Note: Distillation (Kugelrohr, 7 mmHg, oven *ca* 250 $^{\circ}\text{C}$) removed the traces of phosphoramidite ligand. Flash chromatography provided a sample significantly enriched (6:1) in **2j**, so that it was suitable for HPLC analysis on a chiral non-racemic stationary phase. Signals corresponding to **2j** (from spectra of the mixture – by comparison with the pure racemate) have: ¹H NMR (400 MHz) δ 8.15–8.08 (m, 2 H), 7.61–7.53 (m, 1 H), 7.49–7.43 (m, 2 H), 5.46 (t, *J* = 6.4 Hz, 1 H), 5.06 (s, 1 H), 4.94 (s, 1 H), 1.85–1.79 (m, s H), 1.82 (s, 3 H), 1.50–1.25 (m, 6 H), 0.95–0.85 (m, 3 H); ¹³C NMR (CDCl₃, 100.6 MHz) δ 165.7 (s), 143.2 (s), 132.7 (d), 130.6 (s), 129.5 (d, 2 x C), 128.4 (d, 2 x C), 112.6 (s), 77.9 (d), 32.6 (t), 31.5 (t), 25.0 (t), 22.5 (t), 18.1 (q), 14.0 (q); HPLC analysis indicated an enantiomeric excess of 98% [Chiralcel OB-H column; flow: 0.5 mL/min; heptane/*i*-PrOH: 99.75:0.25; λ = 230 nm; major enantiomer **2j**, *t_R* = 10.49 min; minor enantiomer **2j**, *t_R* = 13.46 min]

Notes:

- 1 An error in the artwork of the following paper makes it appear that hetero allylic acetates have been used in an AAA.¹³
- 2 Secondary and sp² hybridized Grignard reagents proved to be unreactive as nucleophiles in the h-AAA.
- 3 The absolute configuration of the natural occurring butyrolactone was not determined.¹⁴ Absolute configuration assigned by comparison of the optical rotations.⁹
- 4 Absolute configuration assigned by comparison of the optical rotations.¹¹

References:

- (1) *Purification of Laboratory Chemicals 3rd Edition*; Perrin, D. D.; Armarego, W. L. F.; Butterworth-Heinemann Ltd, Oxford, Great Britain, 1988.
- (2) (a) Lombardo, M.; Morganti, S.; Trombini, C. *J. Org. Chem.* **2003**, *68*, 997–1006. (b) Lombardo, M.; Girotti, R.; Morganti, S.; Trombini, C. *Chem. Commun.* **2001**, 2310–2311.
- (3) Ihle, N. C.; Heathcock, C. H. *J. Org. Chem.* **1993**, *58*, 560–563.
- (4) Nakamura, K.; Takenaka, K. *Tetrahedron Asym.* **2002**, *13*, 415–422.
- (5) Engman, L. *J. Am. Chem. Soc.* **1984**, *106*, 3977–3984.
- (6) Andrus, M. B.; Argade, A. B.; Chen, X.; Pamment, M. G. *Tetrahedron Lett.* **1995**, *36*, 2945–2948.
- (7) Uenishi, J.; Kubo, Y. *Tetrahedron Lett.* **1994**, *35*, 6697–6700.
- (8) Dale, J. D.; Mosher, H. S. *J. Am. Chem. Soc.* **1968**, *90*, 3732–3738.
- (9) van Oeveren, A.; Feringa, B. L. *J. Org. Lett.* **1996**, *61*, 2920–2921.
- (10) Kingsbury, J. S.; Harrity, J. P. A.; Bonitatebus Jr., P. J.; Hoveyda, A. H. *J. Am. Chem. Soc.* **1999**, *121*, 791–799.
- (11) Asami, M. *Bull. Chem. Soc. Jpn.* **1990**, *33*, 721–727, and references therein. Representative methods for making cyclopent-2-enol or its derivatives: (a) Kitamura, M.; Kasahara, I.; Manabe, K.; Noyori, R.; Takaya, H. *J. Org. Chem.* **1988**, *53*, 708–710. (b) Corey, E. J.; Reichard, A. G. *Tetrahedron Lett.* **1989**, *30*, 5547–5550. (c) Andrus, M. B.; Asgari, D. *Tetrahedron* **2000**, *56*, 5775–5780. (d) Seki, A.; Asami, M. *Tetrahedron* **2002**, *58*, 4655–4665. (e) Bertilsson, K. S.; Sodergron, J. M.; Andersson, G. P. J. *J. Org. Chem.* **2002**, *67*, 1567–1573. (f) Gais, J. H.; Bondarev, O.; Hetzer, R. *Tetrahedron Lett.* **2005**, *46*, 6279–6283.
- (12) Trnka, T. M.; Grubbs, R. H. *Acc. Chem. Res.* **2001**, *34*, 18–29.
- (13) Okamoto, S.; Tominaga, S.; Saino, N.; Kase, K.; Shimoda, K. *J. Organomet. Chem.* **2005**, *690*, 6001–6007.
- (14) Tava, A.; Pecetti, L. *Phytochemistry*. **1997**, *45*, 1145–1148.

^1H NMR and ^{13}C NMR:

(+)-(S)-Benzoic acid-1-methyl-allyl ester: 2

(+)-(E)-3-Phenyl-acrylic acid-1-methyl-allyl ester: 2a

(+)-2,4,6-Trimethylbenzoic acid-1-methyl-allyl ester: 2b

(+)-Benzoic acid-1-ethyl-allyl ester: **2c**

(-)-Benzoic acid-1-vinyl-hexyl ester: 2d

(+)-(S)-Benzoic acid-1-vinyl-pent-4-enyl ester: 2f

(+)-Benzoic acid-1-phenethyl-allyl ester: **2g**

(+)-Benzoic acid-1-vinyl-nonadecyl ester: 2h

(+)-(S)-(E)-3-Phenyl-acrylic acid-1-ethyl-allyl ester: 2i

(+)-(S)-5-Ethyl-2(5H)-furanone: 6

(-)-(S)-Benzoic acid-cyclopent-2-enyl ester: **7**

Benzoic acid 1-ethyl-2-methylallyl ester: 2i

Benzoic acid 1-isopropenylhexyl ester: 2j

Representative chiral HPLC

(+)-(S)-Benzoic acid-1-methyl-allyl ester: 2