Copper-Free Sonogashira Coupling Reaction with PdCl₂ in H₂O under Aerobic Conditions

Bo Liang,[†] Mingji Dai,[†] Jiahua Chen*[†] and Zhen Yang*^{†,‡}

Key Laboratory of Bioorganic Chemistry and Molecular Engineering of Ministry of Education, College of Chemistry, Beijing 100871, China, Laboratory of Chemical Genetics, Shenzhen Graduate School of Peking University, The University Town, Shenzhen 518055, China, and VivoQuest, Inc., 711 Executive Blvd, Valley Cottage, New York 10989

Supporting Information

Table of Contents

Experimental Procedures and Spectra Data..... 2-10

¹H NMR, ¹³C NMR and MS Spectra11-46

Page 11-12: ¹H NMR, ¹³C NMR and MS of compound **1a**

Page 12-13: ¹H NMR, ¹³C NMR and MS of compound **1b**

Page 14-15: ¹H NMR, ¹³C NMR and MS of compound **1c**

Page 15-16: ¹H NMR, ¹³C NMR and MS of compound **1d**

Page 17-18: ¹H NMR, ¹³C NMR and MS of compound **1e**

Page 18-19: ¹H NMR, ¹³C NMR and MS of compound **1f**

Page 20-21: ¹H NMR, ¹³C NMR and MS of compound **1g**

Page 21-22: ¹H NMR, ¹³C NMR and MS of compound **1h**

Page 23-24: ¹H NMR, ¹³C NMR and MS of compound **1i**

Page 24-25: ¹H NMR, ¹³C NMR and MS of compound **1j**

Page 26-27: ¹H NMR, ¹³C NMR and MS of compound **1k**

Page 27-28: ¹H NMR, ¹³C NMR and MS of compound **1l**

Page 29-30: ¹H NMR, ¹³C NMR and MS of compound **2a**

Page 30-31: ¹H NMR, ¹³C NMR and MS of compound **2b**

Page 32-33: ¹H NMR, ¹³C NMR and MS of compound **2c**

Page 33-34: ¹H NMR, ¹³C NMR and MS of compound **2d**

Page 35-36: ¹H NMR, ¹³C NMR and MS of compound **2e**

Page 36-37: ¹H NMR, ¹³C NMR and MS of compound **2f**

Page 38-39: ¹H NMR, ¹³C NMR and MS of compound **2g**

Page 39-40: ¹H NMR, ¹³C NMR and MS of compound **2h**

Page 41-42: ¹H NMR, ¹³C NMR and MS of compound **2i**

Page 42-43: ¹H NMR, ¹³C NMR and MS of compound **2j**

Page 43-44: ¹H NMR, ¹³C NMR and MS of compound **2k**

Page 44-45: ¹H NMR, ¹³C NMR and MS of compound **2l**

Page 45-46: ¹H NMR, ¹³C NMR and MS of compound **2m**

General Information

All reagents were used as received and used directly without further purification. Silica gel (200-300) for purification and silica gel TLC (F₂₅₄) were purchased from Qing Dao Hai Yang Chemical Industry Co. of China.

¹H NMR spectra were recorded on a Varian Inova 300 MHz spectrometer. ¹³C NMR spectra were recorded on a Varian Inova 75 MHz spectrometer. Mass spectra were obtained on a ZAB-HS (EI, 70eV) spectrometer.

Procedure of Sonogashira reaction in water:

PdCl₂ (3.5 mg, 0.02 mmol), 1-iodo-4-nitrobenzene (498 mg, 2.0 mmol), H₂O (2.5 mL) and pyrrolidine (0.83 mL, 10 mmol) were added to a flask (25 mL), and the mixture was stirred for five minutes. To this solution, phenylacetylene (0.26 mL, 2.4 mmol) was added, and the mixture was stirred at room temperature for 24 hours. The reaction mixture was then extracted with EtOAc (3 ×10 mL), and the organic layer was dried over Na₂SO₄. The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane/EtOAc = 20/1) to give the product **1a** (433 mg, 97 %) as a yellow solid. ¹H NMR (300MHz, CDCl₃): δ 7.38-7.40 (m, 3H), 7.54-7.58 (m, 2H), 7.65 (d, *J* = 9.0 Hz, 2H), 8.21(d, *J* = 9.0 Hz, 2H); ¹³C NMR (75MHz, CDCl₃): δ87.5, 94.7, 122.1, 123.6, 128.5, 129.3, 130.2, 131.8, 132.2, 146.9; MS (EI) [C₁₄H₉NO₂]: *m/z* (M⁺): Calcd 223, found 223. (Known compound, see: Sakai, N.; Annaka, K.; Konakahara, T. *Org. Lett.* **2004**, 6, 1527.)

PdCl₂ (7.0 mg, 0.04 mmol), 1-(trifluoromethyl)-4-iodobenzene (1.09 g, mmol), H₂O (5 mL) and pyrrolidine (1.7 mL, 20 mmol) were added to a flask (25 mL), the mixture was stirred for five minutes. To this solution, phenylacetylene(0.52 mL, 4.8 mmol) was added, and the reaction was stirred at room temperature for 24 hours. The mixture was then extracted with EtOAc (3 ×10 mL), and the organic layer was dried over Na₂SO₄. The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane) to give the product **1b** (886 mg, 90%) as a white solid; ¹H NMR (300 MHz, CDCl₃): δ 7.31-7.33 (m, 3H), 7.51-7.59 (m, 6H); ¹³C NMR (75MHz, CDCl₃): δ87.9, 91.8, 122.6, 125.2, 125.2, 125.2, 125.3, 127.1, 128.4, 128.8, 131.7, 131.8; MS (EI) [C₁₅H₉F₃]: *m/z* (M⁺): Calcd 246, found 246. (Known compound, see: Gelman, D.; Hohne, G.; Schumann, H.; Blum, J. *Synthesis* **2001**, 591.)

PdCl_2 (3.5 mg, 0.02 mmol), 4-iodobenzonitrile (458 mg, 2 mmol), H_2O (2.5 mL) and pyrrolidine (0.83 mL, 10 mmol) were added to a flask (25 mL), and the mixture was stirred for five minutes. To this solution, phenylacetylene (0.26 mL, 2.4 mmol) was added, and the reaction mixture was stirred at room temperature for 24 hours. The reaction mixture was then extracted with EtOAc (3 $\times$ 10 mL), and the organic layer was dried over Na_2SO_4 . The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane/EtOAc= 20/1) to give the product **1c** (771 mg, 95 %) as a white solid; ^1H NMR (300 MHz, CDCl_3): δ 7.36-7.39 (m, 3H), 7.52-7.59 (m, 2H), 7.61-7.62 (m, 4H); ^{13}C NMR (75MHz, CDCl_3): δ 87.6, 93.7, 111.3, 118.4, 122.1, 128.1, 128.4, 129.0, 131.7, 131.9, 131.9; MS (EI) [$\text{C}_{15}\text{H}_9\text{N}$]: m/z (M^+): Calcd 203, found 203. (Known compound, see: Elangovan, A.; Wang, Y.-H.; Ho, T.-I. *Org. Lett.* 2003, 5, 1841.)

PdCl_2 (3.5 mg, 0.02 mmol), 4-iodoacetophenone (492 mg, 2 mmol), H_2O (2.5 mL) and pyrrolidine (0.83 mL, 10 mmol) were added to a flask (25 mL), the mixture was stirred for five minutes. To this solution, phenylacetylene (0.26 mL, 2.4 mmol) was added, and the reaction mixture was stirred at room temperature for 24 hours. The reaction mixture was extracted with EtOAc (3 $\times$ 10 mL), and the extract was dried over Na_2SO_4 . The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane/EtOAc= 20/1) to give the product **1d** (392 mg, 89 %) as a white solid; ^1H NMR (300MHz, CDCl_3): δ 2.61 (s, 3H), 7.36-7.38 (m, 3H), 7.54-7.57 (m, 2H), 7.59-7.63 (m, 2H), 7.92-7.95 (m, 2H); ^{13}C NMR (75MHz, CDCl_3): δ 26.6, 88.6, 92.7, 122.6, 128.2, 128.3, 128.4, 128.8, 131.7, 131.7, 136.1, 197.3; MS (EI) [$\text{C}_{15}\text{H}_{12}\text{O}$]: m/z (M^+): Calcd 220, found 220. (Known compound, see; Urgaonkar, S.; Verkade, J. G. *J. Org. Chem.* 2004, ASAP, web release data: 16-Jul-2004)

PdCl_2 (3.5 mg, 0.02 mmol), 1-bromo-4-iodobenzene (566 mg, 2 mmol), H_2O (2.5 mL) and pyrrolidine (0.83 mL, 10 mmol) were added to a flask (25 mL), and the mixture was stirred at 50 $^\circ\text{C}$ for five minutes. To this solution, phenylacetylene (0.26 mL, 2.4 mmol) was added, and reaction mixture was stirred at 50 $^\circ\text{C}$ for 24 hours. The reaction mixture was then extracted with EtOAc (3 $\times$ 10 mL), and the extract was dried over Na_2SO_4 . The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane) give the product **1e** (420 mg, 82 %) as a white solid; ^1H NMR (300MHz, CDCl_3): δ 7.23-7.30 (m, 5H), 7.35-7.44 (m, 4H); ^{13}C NMR (75MHz, CDCl_3): δ 88.3, 90.5, 122.2, 122.4, 122.9, 128.4, 128.5, 131.6, 132.9;

MS (EI) [C₁₄H₉Br]: m/z (M⁺): Calcd 256, found 256. (Known compound, see: Katritzky, A. R.; Abdel-Fattah, A. A. A.; Wang, M. *J. Org. Chem.* **2002**, 67, 7526.)

PdCl₂ (7.0 mg, 0.04 mmol), 1-bromo-2-iodobenzene (1.13 g, 4 mmol), H₂O (5 mL) and pyrrolidine (1.7 mL, 20 mmol) were added to a flask (25 mL), and the mixture was stirred at 50 °C for five minutes. To this solution, phenylacetylene (0.52 mL, 4.8 mmol) was added, and the reaction mixture was stirred at 50 °C for 24 hours. The reaction mixture was then extracted with EtOAc (3 × 10 mL), and the organic layer was dried over Na₂SO₄. The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane) to give the product **1f** (900 mg, 88 %) as a colorless liquid; ¹H NMR (300 MHz, CDCl₃): δ 7.09-7.15 (m, 1H), 7.21-7.27 (m, 1H), 7.30-7.37 (m, 3H), 7.51-7.59 (m, 4H); ¹³C NMR (75 MHz, CDCl₃): δ 87.9, 93.9, 122.8, 125.3, 125.6, 126.9, 128.3, 128.6, 129.3, 131.6, 132.4, 133.2; MS (EI) [C₁₄H₉Br]: m/z (M⁺): Calcd 256, found 256. (Known compound, see: Kamikawa, T.; Hayashi, T. *J. Org. Chem.* **1998**, 63, 8922.)

PdCl₂ (17.5 mg, 0.1 mmol), idiobenzene (2.04 g, 10 mmol), H₂O (12.5 mL) and pyrrolidine (4.2 mL, 50 mmol) were added to a flask (25 mL), and the mixture was stirred at 50 °C for five minutes. To this solution, phenylacetylene (1.3 mL, 12 mmol) was injected into the tube, and the reaction mixture was stirred at 50 °C for 24 hours. The mixture was then extracted with EtOAc (3 × 10 mL), and the extract was dried over Na₂SO₄. The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane) to give the product **1g** (1.67 g, 94 %) as a white solid; ¹H NMR (300 MHz, CDCl₃): δ 7.25-7.28 (m, 6H), 7.50-7.53 (m, 4H); ¹³C NMR (75 MHz, CDCl₃): δ 89.4, 123.2, 128.2, 128.3, 131.5; MS (EI) [C₁₄H₁₀]: m/z (M⁺): Calcd 178, found 178. (Known compound, see: Soheili, A.; Albaneze-Walker, J.; Murry, J. A.; Dormer, P. G.; Hughes, D. L. *Org. Lett.* **2003**, 5, 4191.)

PdCl₂ (17.5 mg, 0.1 mmol), 1-ido-4-methylbenzene (2.18 g, 10 mmol), H₂O (12.5 mL) and pyrrolidine (4.2 mL, 50 mmol) were added to a flask (25 mL), and the mixture was stirred at 50 °C for five minutes. To this solution, phenylacetylene (1.3 mL, 12 mmol) was added, and the mixture was stirred at 50 °C for 24 hours. The mixture was then extracted with EtOAc (3 × 10 mL), and the extract was dried over Na₂SO₄. The solvent was removed under vacuum and the residue was purified flash chromatography on

silica gel (hexane) to give the product **1h** (1.73 g, 90 %) as a white solid; ^1H NMR (300MHz, CDCl_3): δ 2.32 (s, 3H), 7.10-7.13 (m, 2H), 7.28-7.30 (m, 3H), 7.40-7.43 (m, 2H), 7.49-7.51 (m, 2H); ^{13}C NMR (75MHz, CDCl_3): δ 21.4, 88.7, 89.6, 120.1, 123.4, 128.0, 128.3, 129.1, 131.4, 131.5, 138.3; MS (EI) [$\text{C}_{15}\text{H}_{12}$]: m/z (M^+): Calcd 192, found 192. (Known compound, see: Sakai, N.; Annaka, K.; Konakahara, T. *Org. Lett.* **2004**, 6, 1527.)

PdCl_2 (3.5 mg, 0.02 mmol), 4-iodoanisole (468 mg, 2 mmol), H_2O (2.5 mL) and pyrrolidine (0.83 mL, 10 mmol) were added to a flask (25 mL), the mixture was stirred at 50°C for five minutes. To this solution, phenylacetylene (0.26 mL, 2.4 mmol) was added, and the reaction mixture was stirred at 50°C for 24 hours. The mixture was then extracted with EtOAc (3 $\times$ 10 mL), and the extract was dried over Na_2SO_4 . The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane/EtOAc = 20/1) to give the product **1i** (354 mg, 85 %) as a white solid; ^1H NMR (300MHz, CDCl_3): δ 3.75 (s, 3H), 6.82-6.85 (m, 2H), 7.28-7.31 (m, 3H), 7.44-7.52 (m, 4H); ^{13}C NMR (75MHz, CDCl_3): δ 55.1, 88.0, 89.4, 113.9, 115.3, 123.5, 127.9, 128.3, 131.4, 132.9, 159.5; MS (EI) [$\text{C}_{15}\text{H}_{12}\text{O}$]: m/z (M^+): Calcd 208, found 208. (Known compound, see; Urgaonkar, S.; Verkade, J. G. *J. Org. Chem.* **2004**, ASAP, web release data: 16-Jul-2004)

PdCl_2 (3.5 mg, 0.02 mmol), 1-iodonaphthalene (508 mg, 2 mmol), H_2O (2.5 mL) and pyrrolidine (0.83 mL, 10 mmol) were added to a flask (25 mL), and the mixture was stirred at 50°C for five minutes. To this solution, phenylacetylene (0.26 mL, 2.4 mmol) was added, and the reaction was stirred at 50°C for 24 hours. The mixture was then extracted with EtOAc (3 $\times$ 10 mL), and the extract was dried over Na_2SO_4 . The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane) to give the product **1j** (388 mg, 85 %) as a colorless liquid; ^1H NMR (300MHz, CDCl_3): δ 7.34-7.66 (m, 8H), 7.74-7.84 (m, 3H), 8.84-8.46 (m, 1H); ^{13}C NMR (75MHz, CDCl_3): δ 87.5, 94.3, 120.9, 123.4, 125.2, 126.2, 126.4, 126.7, 128.3, 128.3, 128.4, 128.7, 130.3, 131.6, 133.2, 133.2; MS (EI) [$\text{C}_{18}\text{H}_{12}$]: m/z (M^+): Calcd 228, found 228. (Known compound, see: Chow, H.-F.; Wan, C.-W.; Low, K.-H.; Yeung, Y.-Y. *J. Org. Chem.* **2001**, 66, 1910.)

PdCl_2 (3.5 mg, 0.02 mmol), 3-iodopyridine (410 mg, 2 mmol), H_2O (2.5 mL) and pyrrolidine (0.83 mL, 10 mmol) were added to a flask, the mixture was stirred at 50°C for five minutes. To this solution, phenylacetylene (0.26 mL, 2.4 mmol) was added, and the reaction mixture was stirred at 50°C for 24 hours. The mixture was extracted with EtOAc (3 $\times$ 10 mL), and the extract was dried over Na_2SO_4 . The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane/EtOAc = 8/1) to give product **1k** (340 mg, 95%) as a yellow solid; ^1H NMR (300 MHz, CDCl_3): δ 7.20-7.24 (m, 1H), 7.33-7.35 (m, 3H), 7.52-7.55 (m, 2H), 7.75-7.78 (m, 1H), 8.51-8.53 (m, 1H), 8.77 (s, 1H); ^{13}C NMR (75 MHz, CDCl_3): δ 85.8, 92.5, 120.2, 122.3, 122.8, 128.3, 128.6, 131.5, 138.2, 148.4, 152.0; MS (EI) [$\text{C}_{13}\text{H}_9\text{N}$]: m/z (M^+): Calcd 179, found 179. (Known compound, see: Elangovan, A.; Wang, Y.-H.; Ho, T.-I. *Org. Lett.* **2003**, 5, 1841.)

PdCl_2 (3.5 mg, 0.02 mmol), 4-iodobenzonitrile (458 mg, 2 mmol), H_2O (2.5 mL) and pyrrolidine (0.83 mL, 10 mmol) were added to a flask (25 mL), the mixture was stirred at room temperature for five minutes. To this solution, n-hexyne (0.28 mL, 2.4 mmol) was added, and the reaction mixture was stirred at room temperature for 24 hours. The mixture was extracted with EtOAc (3 $\times$ 10 mL), and the extract was dried over Na_2SO_4 . The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane/EtOAc = 20/1) to give the product **2a** (256 mg, 70 %) as a colorless liquid; ^1H NMR (300 MHz, CDCl_3): δ 0.95 (t, 3H), 1.44-1.51 (m, 2H), 1.55-1.62 (m, 2H), 2.43 (t, 2H), 7.44 (d, J = 8.1 Hz, 2H), 7.56 (d, J = 8.1 Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3): δ 13.5, 19.1, 21.9, 30.4, 79.4, 95.5, 110.6, 118.5, 129.0, 131.8, 131.9; MS (EI) [$\text{C}_{13}\text{H}_{13}\text{N}$]: m/z (M^+): Calcd 183, found 183. (Known compound, see: Molander, G. A.; Katona, B. W.; Machrouhi, F. *J. Org. Chem.* **2002**, 67, 8416.)

PdCl_2 (3.5 mg, 0.02 mmol), 1-iodo-4-nitrobenzene (498 mg, 2 mmol), H_2O (2.5 mL) and pyrrolidine (0.83 mL, 10 mmol) were added to a flask (25 mL), and the mixture was stirred at room temperature for five minutes. To this solution, n-hexyne (0.28 mL, 2.4 mmol) was added and the reaction mixture was stirred at room temperature for 24 hours. The mixture was extracted with EtOAc (3 $\times$ 10 mL), and the extract was dried over Na_2SO_4 . The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane/EtOAc = 20/1) to give the product **2b** (345 mg, 85 %) as a yellow liquid; ^1H NMR (300 MHz, CDCl_3): δ 0.95 (t, 3H), 1.44-1.54 (m, 2H), 1.56-1.63 (m, 2H), 2.44 (t, 2H), 7.47-7.51 (m, 2H), 8.11-8.15 (m, 2H); ^{13}C NMR (75 MHz, CDCl_3): δ 13.4, 19.1, 21.9, 30.3, 79.1, 96.6, 123.2, 131.0, 132.0, 146.4; MS (EI) [$\text{C}_{12}\text{H}_{13}\text{NO}_2$]: m/z (M^+): Calcd 203, found 203.

(Known compound, see: Molander, G. A.; Katona, B. W.; Machrouhi, F. *J. Org. Chem.* **2002**, 67, 8416.)

PdCl_2 (3.5 mg, 0.02 mmol), 3-iodopyridine (510 mg, 2 mmol), H_2O (2.5 mL) and pyrrolidine (0.83 mL, 10 mmol) were added to a flask (25 mL), and the mixture was stirred at room temperature for five minutes. To this solution, n-hexyne (0.28 mL, 2.4 mmol) was added, and the reaction was stirred at room temperature for 24 hours. The mixture was then extracted with EtOAc (3 $\times$ 10 mL), and the extract was dried over Na_2SO_4 . The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane/EtOAc = 20/1) to give the product **2c** (226 mg, 71 %) as a colorless liquid; ^1H NMR (300MHz, CDCl_3): δ 0.95 (t, 3H), 1.44-1.55 (m, 2H), 1.57-1.62 (m, 2H), 2.42 (t, 2H), 7.17-7.21 (m, 2H), 7.64-7.68 (m, 1H), 8.46-8.48 (m, 1H), 8.62-8.63 (m, 1H); ^{13}C NMR (75MHz, CDCl_3): δ 13.5, 18.9, 21.9, 30.5, 77.2, 93.9, 121.1, 122.7, 138.2, 147.7, 152.2; MS (EI) [$\text{C}_{11}\text{H}_{13}\text{N}$]: m/z (M^+): Calcd 159, found 159. (Known compound, see: Novak, Z.; Szabo, A.; Repasi, J.; Kotschy, A. *J. Org. Chem.* **2003**, 68, 3327.)

PdCl_2 (3.5 mg, 0.02 mmol), 3-iodo-pyridine (410 mg, 2 mmol), 4-N,N-dimethyl phenylacetylene (348 mg, 2.4 mmol), H_2O (2.5 mL) and pyrrolidine (0.83 mL, 10 mmol) were added to a flask, and the reaction mixture was stirred at 50 $^\circ\text{C}$ for 24 hours. The mixture was then extracted with EtOAc (3 $\times$ 10 mL), and the extract was dried over Na_2SO_4 . The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane/EtOAc = 8/1) to give the product **2d** (377 mg, 85 %) as a yellow solid; ^1H NMR (300MHz, CDCl_3): δ 2.99 (s, 6H), 6.64-6.67 (m, 2H), 7.21-7.26 (m, 1H), 7.40-7.43 (m, 2H), 7.73-7.77 (m, 1H), 8.47-8.49 (m, 1H), 8.72-8.73 (m, 1H); ^{13}C NMR (75MHz, CDCl_3): δ 40.1, 84.0, 94.1, 108.9, 111.7, 121.3, 122.9, 132.8, 137.9, 147.6, 150.3, 151.9; MS (EI) [$\text{C}_{15}\text{H}_{14}\text{N}_2$]: m/z (M^+): Calcd 222, found 222. (Known compound, see: Beletskaya, I. P.; Latyshev, G. V.; Tsvetkov, A. V.; Lukashev, N. V. *Tetrahedron Lett.* **2003**, 44, 5011.)

PdCl_2 (3.5 mg, 0.02 mmol), 3-iodo-pyridine (410 mg, 2 mmol), 4-methoxy-phenylacetylene (317 mg, 2.4 mmol), H_2O (2.5 mL) and pyrrolidine (0.83 mL, 10 mmol) were added to a flask (25 mL), and the mixture was stirred at 50 $^\circ\text{C}$ for 24 hours. The mixture was then extracted with EtOAc (3 $\times$ 10 mL), and the extract was dried

over Na₂SO₄. The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane/EtOAc = 20/1) to give the product **2e** (411 mg, 81 %) as a yellow solid; ¹H NMR (300MHz, CDCl₃): δ 3.78 (s, 3H), 6.85-6.88 (m, 2H), 7.20-7.24 (m, 1H), 7.46-7.49 (m, 2H), 7.73-7.77 (m, 1H), 8.49-8.51 (m, 1H), 8.74-8.75 (m, 1H); ¹³C NMR (75MHz, CDCl₃): δ 55.1, 84.6, 92.6, 113.9, 114.3, 120.6, 122.8, 132.9, 137.9, 148.0, 151.9, 159.8; ; MS (EI) [C₁₄H₁₁NO]: m/z (M⁺): Calcd 209, found 209.

PdCl₂ (3.5 mg, 0.02 mmol), idiobenzene (408 mg, 2 mmol), 4-N,N-dimethylphenylacetylene (348 mg, 2.4 mmol), H₂O (2.5 mL) and pyrrolidine (0.83 mL, 10 mmol) were added to a flask (25 mL), and the reaction mixture was stirred at 50 °C for 24 hours. The mixture was then extracted with EtOAc (3 × 10 mL), and the extract was dried over Na₂SO₄. The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane/EtOAc = 20/1) to give the product **2f** (336 mg, 76 %) as a white solid; ¹H NMR (300MHz, CDCl₃): δ 2.92 (s, 6H), 6.61 (d, *J* = 9.3 Hz, 2H), 7.26-7.29 (m, 3H), 7.40 (d, *J* = 9.0 Hz, 2H), 7.47-7.50 (m, 2H); ¹³C NMR (75MHz, CDCl₃): δ 40.1, 87.3, 90.6, 109.9, 111.7, 124.1, 127.4, 127.5, 128.2, 131.2, 132.6, 149.9; MS (EI) [C₁₆H₁₅N]: m/z (M⁺): Calcd 221, found 221. (Known compound, see: Batey, R. A.; Shen, M.; Lough, A. J. *Org. Lett.* **2002**, 4, 1411.)

PdCl₂ (3.5 mg, 0.02 mmol), 1-ido-4-methylbenzene (436 mg, 2 mmol), 4-N,N-dimethylphenylacetylene (348 mg, 2.4 mmol), H₂O (2.5 mL) and pyrrolidine (0.83 mL, 10 mmol) were added to a flask (25 mL), and the reaction mixture was stirred at 50 °C for 24 hours. The mixture was extracted with EtOAc (3 × 10 mL), and the extract was dried over Na₂SO₄. The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane/EtOAc = 20/1) to give the product **2g** (329 mg, 70 %) as a white solid; ¹H NMR (300MHz, CDCl₃): δ 2.33 (s, 3H), 2.94 (s, 6H), 6.63 (d, *J* = 9.0 Hz, 2H), 7.11 (d, *J* = 8.1 Hz, 2H), 7.37-7.40 (m, 4H); ¹³C NMR (75MHz, CDCl₃): δ 21.4, 40.1, 87.4, 89.8, 110.2, 111.8, 121.0, 128.9, 131.1, 132.6, 137.4, 149.9; MS (EI) [C₁₇H₁₇N]: m/z (M⁺): Calcd 235, found 235. (Known compound, see: Van Koten, G.; Ten H., Richard W. M.; Noltes, J. G. *J. Org. Chem.* **1977**, 42, 2705.)

PdCl₂ (3.5 mg, 0.02 mmol), 4-idiobenzotrifluoride (544 mg, 2 mmol), 4-N,N-dimethylphenylacetylene (348 mg, 2.4 mmol), H₂O (2.5 mL) and pyrrolidine (0.83 mL,

10 mmol) were added to a flask (25 mL), and the reaction mixture was stirred at room temperature for 24 hours. The mixture was then extracted with EtOAc (3 ×10 mL), and the extract was dried over Na₂SO₄. The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane/EtOAc = 20/1) to give the product **2h** (474 mg, 82 %) as a white solid; ¹H NMR (300MHz, CDCl₃): δ 2.97 (s, 6H), 6.64 (d, *J* = 9.0 Hz, 2H), 7.41 (d, *J* = 9.0 Hz, 2H), 7.55 (s, 4H); ¹³C NMR (75MHz, CDCl₃): δ40.1, 86.3, 93.4, 109.1, 111.7, 125.1, 125.2, 128.1, 131.3, 132.9, 150.4; ; MS (EI) [C₁₇H₁₄F₃N]: m/z (M⁺): Calcd 289, found 289.

PdCl₂ (3.5 mg, 0.02 mmol), 4-iodoacetophenone (492 mg, 2 mmol), 4-methoxyphenylacetylene (317 mg, 2.4 mmol), H₂O (2.5 mL) and pyrrolidine (0.83 mL, 10 mmol) were added to a flask (25 mL), and the reaction mixture was stirred at room temperature for 24 hours. The mixture was then extracted with EtOAc (3 ×10 mL), and the extract was dried over Na₂SO₄. The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane/EtOAc = 20/1) to give the product **2i** (400 mg, 80 %) as a white solid; ¹H NMR (300MHz, CDCl₃): δ 2.59 (s, 3H), 3.82 (s, 3H), 6.88 (d, *J* = 8.4 Hz, 2H), 7.47-7.58 (m, 4H), 7.91 (d, *J* = 8.4 Hz, 2H); ¹³C NMR (75MHz, CDCl₃): δ26.5, 55.2, 87.5, 92.9, 114.0, 114.6, 128.2, 128.5, 131.4, 133.2, 135.8, 159.9, 197.3; MS (EI) [C₁₇H₁₄O₂]: m/z (M⁺): calcd 250, found 250. (Known compound, see: Nishihara, Y.; Ikegashira, K.; Hirabayashi, K.; Ando, J.; Mori, A.; Hiyama, T. *J. Org. Chem.* **2000**, 65, 1780.)

PdCl₂ (3.5 mg, 0.02 mmol), 1-iodo-4-methylbenzene (516 mg, 2 mmol), 4-methoxyphenylacetylene (317 mg, 2.4 mmol), H₂O (2.5 mL) and pyrrolidine (0.83 mL, 10 mmol) were added to a flask, and the mixture was stirred at 50 °C for 24 hours. The mixture was then extracted with EtOAc (3 ×10 mL), and the extract was dried over Na₂SO₄. The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane/EtOAc = 20/1) to give the product **2j** (333 mg, 75 %) as a white solid; ¹H NMR (300MHz, CDCl₃): δ 2.35 (s, 3H), 3.80 (s, 3H), 6.86 (d, *J* = 9.0 Hz, 2H), 7.12-7.14 (m, 2H), 7.39-7.47 (m, 4H); ¹³C NMR (75MHz, CDCl₃): δ21.4, 55.2, 88.2, 88.6, 113.9, 115.5, 120.5, 129.0, 131.3, 132.9, 137.9, 159.4; MS (EI) [C₁₆H₁₄O₂]: m/z (M⁺): Calcd 238, found 238. (Known compound, see: Mio, M. J.; Kopel, L. C.; Braun, J. B.; Gadzikwa, T. L.; Hull, K. L.; Brisbois, R. G.; Markworth, C. J.; Grieco, P. A. *Org. Lett.* **2002**, 4, 3199.)

PdCl_2 (3.5 mg, 0.02 mmol), 1-iodo-4-methylbenzene (516 mg, 2 mmol), 4-bromophenylacetylene (434 mg, 2.4 mmol), H_2O (2.5 mL) and pyrrolidine (0.83 mL, 10 mmol) were added to a flask, and the reaction mixture was stirred at 50°C for 24 hours. The mixture was then extracted with EtOAc (3 $\times$ 10 mL), and the extract was dried over Na_2SO_4 . The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane) to give the product **2k** (369 mg, 68 %) as a white solid; ^1H NMR (300MHz, CDCl_3): δ 2.35 (s, 3H), 7.14 (d, $J = 7.8$ Hz, 2H), 7.34-7.47 (m, 6H); ^{13}C NMR (75MHz, CDCl_3): δ 21.5, 87.7, 90.7, 119.8, 122.2, 122.4, 129.1, 131.4, 131.5, 132.9, 138.7; MS (EI) [$\text{C}_{15}\text{H}_{11}\text{Br}$]: m/z (M^+): Calcd 270, found 270. (Known compound, see: Gray, G. W.; Mosley, A. *Molecular Crystals and Liquid Crystals* **1976**, 37, 213.)

PdCl_2 (3.5 mg, 0.02 mmol), 1,4-diiodobenzene (330 mg, 1 mmol), H_2O (2.5 mL) and pyrrolidine (0.83 mL, 10 mmol) were added to a flask (25 mL), the mixture was stirred at 50°C for five minutes. To this solution, phenylacetylene (0.26 mL, 2.4 mmol) was added, and the reaction mixture was stirred at 50°C for 24 hours. The mixture was then extracted with EtOAc (3 $\times$ 10 mL), and the extract was dried over Na_2SO_4 . The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane) to give the product **2l** (222 mg, 80 %) as a white solid; ^1H NMR (300MHz, CDCl_3): δ 7.33-7.35 (m, 6H), 7.50-7.55 (m, 8H); ^{13}C NMR (75MHz, CDCl_3): δ 89.1, 91.2, 122.9, 123.1, 128.4, 128.4, 131.5, 131.6; MS (EI) [$\text{C}_{22}\text{H}_{14}$]: m/z (M^+): Calcd 278, found 278. (Known compound, see: Lee, P. H.; Lee, S. W.; Seomoon, D. *Org. Lett.* **2003**, 5, 4963.)

PdCl_2 (17.5 mg, 0.1 mmol), 1,4-dibromo-2,5-diiodobenzene (2.4 g, 5 mmol), H_2O (12.5 mL) and pyrrolidine (4.2 mL, 50 mmol) were added to a flask, the mixture was stirred at 50°C for five minutes. To this solution, phenylacetylene (1.3 mL, 12 mmol) was added, and the reaction mixture was stirred at 50°C for 24 hours. The mixture was then extracted with EtOAc (3 $\times$ 10 mL), and the extract was dried over Na_2SO_4 . The solvent was removed under vacuum and the residue was purified flash chromatography on silica gel (hexane/EtOAc = 80/1) to give the product **2m** (1.42 g, 65%) as a white solid; ^1H NMR (300MHz, CDCl_3): δ 7.37-7.38 (m, 6H), 7.56-7.59 (m, 4H), 7.77 (s, 2H); ^{13}C NMR (75MHz, CDCl_3): δ 86.8, 96.7, 122.3, 123.7, 126.4, 128.4, 129.1, 131.8, 135.9; MS (EI) [$\text{C}_{22}\text{H}_{12}\text{Br}_2$]: m/z (M^+): Calcd 436, found 436. (Known compound, see: Goldfinger, M. B.; Crawford, K. B.; Swager, T. M. *J. Am. Chem. Soc.* **1997**, 119, 4578.)

File Name : d:\maspec2\data\40764.ms2
 Creation Date/Time : 04-5-11 at 15:27:03
 File Type : Lo-Res Data - Ctd (Magnet)
 File Source : Acquired on MASPEC II system [msw/9888]
 Operator : Peking University
 Instrument : ZAB-HS

SCAN GRAPH. Flagging=Nom.M/z.
 Scan 22#1:55. Sub.=6#0:31. "T=100 C". Entries=247. Base M/z=223. 100% Int.=6.79936. EI. POS.

sonogeshira
 p-CF3
 Archive directory: /export/home/vnmr1/vnmrsys/data
 Sample directory:
 File: PROTON
 Pulse Sequence: szp

sonogashira
p-CF₃
Pulse Sequence: s2pul

File Name : d:\mspec2\data\40061.ms2
Creation Date/Time : 04-5-16 at 17:06:50
File Type : Lo-Res Data - Ctd (Magnet)
File Source : Acquired on MASPPEC II system [msw0606]
Operator : Peking University
Instrument : ZAB-4IS

SCAN GRAPH, Flagger=Nom.M/z
Scan 790:36, Sub=390:16, T=80 C, Entries=144, Base M/z=245.7, 100% Int.=0.78272, EI, POS.

```
sonogashira-1-9
Archive directory: /export/home/vnmri/vnmrsys/data
Sample directory:
File: PROTON
```

sonogashira
p-Oh-hexyne
Pulse Sequence: szpu1

sonogashira-Ph-Ph-Ph
 Archive directory: /export/home/vnmr1/vnmrsys/data
 Sample directory:
 File: PROTON
 Pulse Sequence: zgpg30

sonogashira-Ph-Ph-Ph
 Pulse Sequence: s2pul