SUPPORTING INFORMATION AVAILABLE

Cross-Coupling and Dehalogenation Reactions Catalyzed by (NHC)Pd(allyl)Cl (NHC = N-Heterocyclic Carbene) Complexes

*Oscar Navarro, Harneet Kaur, Parisa Mahjoor, and Steven P. Nolan**

Department of Chemistry, University of New Orleans, New Orleans, Louisiana 70148

snolan@uno.edu

Table of Contents

General Considerations	S1
Conventional and Microwaved-Assisted Suzuki-Miyaura Reactions	S2
Conventional and Microwave-Assisted Dehalogenation of Aryl Chlorides	S7
Catalytic Amination of Aryl Triflates	S8

General Considerations

- All aryl halides, boronic acids, amines and aryl triflates were used as received. Technical grade 2-butanol was also used as received. 1,4-Dioxane (anhydrous) and toluene were distilled under argon from sodium benzophenone ketyl prior to use. Cesium carbonate, potassium and cesium fluoride, potassium phosphate, potassium and sodium *tert*-butoxide, barium hydroxide were stored under argon in an inert atmosphere glovebox or in desiccators over anhydrous calcium carbonate.
- (IMes)Pd(allyl)Cl (**1**) (IMes = (*N,N'*-bis(2,4,6-trimethylphenyl)imidazol-2-ylidene)), (SIMes)Pd(allyl)Cl (**2**) (SIMes = (*N,N'*-bis(2,4,6-trimethylphenyl)-4,5-dihydroimidazol-2-ylidene)), (IPr)Pd(allyl)Cl (**3**) (IPr = (*N,N'*-bis(2,6-diisopropylphenyl)imidazol-2-ylidene)), (SIPr)Pd(allyl)Cl (**4**) (SIPr = (*N,N'*-bis(2,6-

diisopropylphenyl)-4,5-dihydroimidazol-2-ylidene)), (ItBu)Pd(allyl)Cl (**5**) (ItBu = (*N,N*-bis(2,6-*tert*-butylphenyl)imidazol-2-ylidene)) were prepared according to reported procedures.¹

- All reactions were carried out under an atmosphere of argon in screw cap vials and stirred in a Lab-Line Orbit bench shaker if conventionally heated, or in sealed cap vials and placed in an Enrys Optimizer microwave reactor if microwave heated.
- Flash chromatography was performed using silica gel 60 (230-400 mesh) using hexanes:ethylacetate mixtures.
- ¹H NMR spectra were recorded on a 300 or 400 MHz spectrometer at ambient temperature in CDCl₃.

Conventional and Microwaved-Assisted Suzuki-Miyaura Cross-Coupling Reactions

Effect of the Base on the Suzuki-Miyaura Cross-Coupling. *General Procedure:* In a glovebox, **1** (2 mol %, 9.8 mg), base (1.5 mmol) and phenylboronic acid (1 mmol, 122 mg) were added in turn to a vial, and closed with a screw cap fitted with a septum. Outside the glovebox, 4-chlorotoluene (1 mmol, 118 µL) and 1,4-dioxane (3 mL) were injected into the vial in that order. The vial was then placed in a Lab-Line Orbit bench shaker set at 80 °C for 4 h. The mixture was then analyzed by gas chromatography.

Effect of the Different NHC on the Suzuki-Miyaura Cross-Coupling. *General Procedure:* In a glovebox, catalyst (2 mol %), NaO^tBu (3 mmol, 288 mg) and phenylboronic acid (1.1 mmol, 134 mg) were added in turn to a vial, and closed with a screw cap fitted with a septum. Outside the glovebox, 4-chlorotoluene (1 mmol, 118 µL) and 1,4-dioxane (1 mL) were injected into the vial in that order. The vial was then placed in a Lab-Line Orbit bench shaker set at 80 °C for 20 min. The mixture was then analyzed by gas chromatography.

(1) Viciu, M. S.; Germaneau, R. F.; Navarro-Fernandez, O; Stevens, E. D.; Nolan, S. P. *Organometallics*, **2002**, *21*, 5470-5472.

Catalyzed Suzuki-Miyaura Cross Coupling of Aryl Chlorides with Phenylboronic

Acid. *General Procedure:* In a glovebox, **3** (2 mol %, 11.4 mg), NaO^tBu (3 mmol, 288 mg) and phenylboronic acid (1.2 mmol, 146 mg) were added in turn to a vial, and closed with a screw cap fitted with a septum. Outside the glovebox, the aryl chloride (1 mmol) and 1,4-dioxane (1.5 mL) were injected into the vial in that order. The vial was then placed in a Lab-Line Orbit bench shaker set at 60 °C for 3 h, and subsequently allowed to cool to room temperature. After analysis by gas chromatography, silica gel was added to the vial, the solvent was evaporated *in vacuo* and the product isolated by flash chromatography. The amounts of isolated product expressed below are the average of two runs.

4-Cyanobiphenyl (Table 3, entry 1).² The procedure afforded 315 mg (88%) of the title compound.

4-Trifluoromethylbiphenyl (Table 3, entry 2).³ The procedure afforded 417.5 mg (94 %) of the title compound.

2-Pyridinylphenyl (Table 3, entry 3).⁴ The procedure afforded 289 mg (93 %) of the title compound.

3-Pyridinylphenyl (Table 3, entry 4).⁵ The procedure afforded 282 mg (91 %) of the title compound.

(2) Klement, I.; Rohlander, M.; Tucker, C. E.; Majid, T. N.; Knochel, P.; Venegas, P.; Cahiez, G.

Tetrahedron **1996**, *52*, 7201-7220.

(3) Molander, G. A.; Biolatto, B. *Org. Lett.* **2002**, *4*, 1867-1870.

(4) Ellames, G. J.; Gibson, J. S.; Herbert, J. M.; McNeill, A. H. *Tetrahedron* **2001**, *57*, 9487-9498.

(5) Sakamoto, T.; Kondo, Y.; Murata, N.; Yamanaka, H. *Tetrahedron* **1993**, *49*, 9713-9720.

Biphenyl (Table 3, entry 5).⁶ The procedure afforded 305 mg (99 %) of the title compound.

4-Methylbiphenyl (Table 3, entry 6).⁷ The procedure afforded 313 mg (93 %) of the title compound.

4-Methoxybiphenyl (Table 3, entry 7).⁷ The procedure afforded 342 mg (93%) of the title compound.

3-Methoxybiphenyl (Table 3, entry 8).⁸ The procedure afforded 339 mg (92 %) of the title compound.

2-Methoxybiphenyl (Table 3, entry 9).⁹ The procedure afforded 342 mg (93 %) of the title compound.

Catalyzed Suzuki-Miyaura Cross Coupling of Aryl Bromides with Phenylboronic Acid. *General Procedure:* In a glovebox, **3** (1 mol %, 5.7 mg), NaO^tBu (3 mmol, 288 mg) and phenylboronic acid (1.2 mmol, 146 mg) were added in turn to a vial, and closed with a screw cap fitted with a septum. Outside the glovebox, the aryl bromide (1 mmol) and 1,4-dioxane (1.5 mL) were injected into the vial in that order. The vial was then placed in a Lab-Line Orbit bench shaker set at 60 °C for 3 h, and subsequently allowed to cool to room temperature. After analysis by gas chromatography, silica gel was added to the vial, the solvent was evaporated *in vacuo* and the product isolated by flash

(6) Litke, A. F.; Dai, D.; Fu, G. C. *J. Am. Chem. Soc.* **2000**, *122*, 4020-4028.

(7) Rao, M. S. C.; Rao, G. S. K. *Synthesis* **1987**, 231-233.

(8) Zapf, A.; Ehrentraut, A.; Beller, M. *Angew. Chem. Int. Ed.* **2000**, *39*, 4153-4155.

(9) Li, G. Y. *Angew. Chem. Int. Ed.* **2001**, *40*, 1513-1516.

chromatography. The amounts of isolated product expressed below are the average of two runs:

2-Pyridinylphenyl (Table 4, entry 1).⁴ The procedure afforded 279 mg (90%) of the title compound.

Biphenyl (Table 4, entry 2).⁶ The procedure afforded 287 mg (93 %) of the title compound.

4-Methylbiphenyl (Table 4, entry 3).⁷ The procedure afforded 306 mg (91 %) of the title compound.

4-Methoxybiphenyl (Table 4, entry 4).⁷ The procedure afforded 328 mg (89%) of the title compound.

Catalyzed Suzuki-Miyaura Cross Coupling of 4-Chlorotoluene with Different Boronic Acids. *General Procedure:* In a glovebox, **3** (2 mol %, 11.4 mg), NaO^tBu (3 mmol, 288 mg) and the boronic acid (1.2 mmol) were added in turn to a vial, and closed with a screw cap fitted with a septum. Outside the glovebox, 4-Cl-toluene (1 mmol, 120 mL) and 1,4-dioxane (1.5 mL) were injected into the vial in that order. The vial was then placed in a Lab-Line Orbit bench shaker set at 60 °C for 3 h, and subsequently allowed to cool to room temperature. After analysis by gas chromatography, silica gel was added to the vial, the solvent was evaporated *in vacuo* and the product isolated by flash chromatography. The amounts of isolated product expressed below are the average of two runs:

4-Methoxy-4'-methylbiphenyl (Table 5, entry 1).⁷ The procedure afforded 380 mg (96 %) of the title compound.

3-Methoxy-4'-methylbiphenyl (Table 5, entry 2).¹⁰ The procedure afforded 364 mg (92 %) of the title compound.

2,4'-Dimethylbiphenyl (Table 5, entry 3).¹¹ The procedure afforded 273 mg (75 %) of the title compound.

3-Cyano-4'-methylbiphenyl (Table 5, entry 4).¹² The procedure afforded 279 mg (72 %) of the title compound.

Microwave-Assisted Catalyzed Suzuki-Miyaura Cross Coupling of Aryl Chlorides with Phenylboronic Acid. *General procedure:* In a glovebox, **3** (2 mol %, 11.4 mg), NaO^tBu (3 mmol, 288 mg) and phenylboronic acid (1.2 mmol, 146 mg) were added in turn to a microwave vial containing a magnetic stirring bar, and sealed with a cap fitted with a septum. Outside the glovebox, the aryl chloride (1 mmol) and 1,4-dioxane (2 mL) were injected into the vial in that order. The vial was then placed in a Enrys optimizer microwave reactor set at 120 °C for 90 sec, and subsequently allowed to cool to room temperature. The mixture was then analyzed by gas chromatography and compared with a real sample.

Conventional and Microwave-Assisted Catalytic Dehalogenation of Aryl Chlorides

Effect of the Base on the Microwave-Assisted Dehalogenation. *General Procedure:* In a glovebox, base (1.05 mmol) was added to a microwave vial containing a magnetic stirring bar, and sealed with a cap fitted with a septum. Outside the glovebox, 4-chlorotoluene (1 mmol, 118 μ L) and 2 mL of a stock solution of **3** in technical grade

(10) Hatanake, Y.; Goda, K.; Okahara, Y. *Tetrahedron* **1994**, 59, 8301-8316.

(11) Kosugi, M.; Shikawa, T.; Nogami, T.; Migito, T. *Nippon Kagaku Kaishi* **1985**, 3, 520-526.

(12) Kageyama, H.; Miyazaki, T.; Kimura, Y. *Synlett*. **1994**, 5, 371-372.

isopropanol (1.5 mg in 20 mL, 0.025 mol % in 2 mL) were injected into the vial in that order. The vial was then placed in an Enrys Optimizer microwave reactor set at 120 °C for 120 sec, and subsequently allowed to cool to room temperature. The mixture was analyzed by gas chromatography and compared with an authentic sample of product.

Microwave-assisted Catalyzed Dehalogenation of Aryl Chlorides. *General Procedure:* In a glovebox, NaO^tBu (1.05 mmol, 101 mg) was added to a microwave vial containing a magnetic stirring bar, and sealed with a cap fitted with a septum. Outside the glovebox, aryl chloride (1 mmol) and 2 mL of a stock solution of **3** in technical grade isopropanol (1.5 mg in 20 mL, 0.025 mol % in 2 mL) were injected into the vial in that order. The vial was then placed in an Enrys Optimizer microwave reactor set at 120 °C for 120 sec, and subsequently allowed to cool to room temperature. The mixture was analyzed by gas chromatography and compared with an authentic sample of product.

Dehalogenation of Aryl Chlorides. *General Procedure:* In a glovebox, NaO^tBu (1.05 mmol, 101 mg) was added to a vial, and closed with a screw cap fitted with a septum. Outside the glovebox, the aryl chloride (1 mmol) and 1.5 mL of a stock solution of **3** in technical grade isopropanol (57 mg in 30 mL, 0.5 mol % in 1.5 mL) were injected into the vial in that order. The vial was then placed in a Lab-Line Orbit bench shaker set at 60 °C for 1.75 h, and subsequently allowed to cool to room temperature. The mixture was analyzed by gas chromatography and compared with an authentic sample of the product.

Catalytic Amination of Aryl Triflates

Effect of the Different NHC on the Catalytic Amination of Aryl triflates. In a glovebox, catalyst (1 mol %), NaO^tBu (1.05 mmol, 101 mg) and dry toluene (4 mL) were added in turn to a vial which was closed with a screw cap fitted with a septum. Outside the glovebox, p-tolyl triflate (1 mmol, 179 µL) and morpholine (1.2 mmol, 105 µL) were injected into the vial in that order. The vial was then placed in a Lab-Line Orbit bench shaker set at 70 °C and the reaction was monitored by gas chromatography.

Catalyzed Amination of Aryl Triflates. *General Procedure:* In a glovebox, **3** (1 mol %), NaO^tBu (1.05 mmol, 101 mg) and dry toluene (4 mL) were added in turn to a vial, which was closed with a screw cap fitted with a septum. Outside the glovebox, aryl triflate (1 mmol) and amine (1.2 mmol) were injected into the vial in that order. The vial was then placed in a Lab-Line Orbit bench shaker set at 70 °C. After analysis by gas chromatography, silica gel was added to the vial, the solvent was evaporated *in vacuo* and the product isolated by flash chromatography. The amounts of isolated product reported below are the average of two runs:

N-Cyclohexyl-4-methylaniline (Table 12, entry 1).¹³ The general procedure afforded 302 mg (80 %) of the title compound.

N-(4-Methylphenyl)benzylamine (Table 12, entry 2).¹⁴ The general procedure afforded 280 mg (71 %) of the title compound.

N-(4-Methylphenyl)morpholine (Table 12, entry 3).¹⁵ The general procedure afforded 318 mg (90 %) of the title compound.

1-(4-Methylphenyl)piperidine (Table 12, entry 4).¹⁶ The general procedure afforded 262 mg (75 %) of the title compound.

N,N-Dibutyl-4-methylaniline (Table 12, entry 5).¹⁴ The general procedure afforded 372 mg (85 %) of the title compound.

(13) Antilla, J. C.; Buchwald, S. L. *Org. Lett.* **2001**, 3, 2077-2079.

(14) Kataoka, N.; Shelby, Q.; Stambuli, J. P.; Hartwig, J. F. *J. Org. Chem.* **2002**, 67, 5553-5566.

(15) Kuwano, R.; Utsunomiya, M.; Hartwig, J. F. *J. Org. Chem.* **2002**, 67, 6479-6486.

(16) Hamann, B. C.; Hartwig, J. F. *J. Am. Chem. Soc.* **1998**, 120, 7369-7370.

N-Methyl-N-phenyl-*p*-toluidine (Table 12, entry 6).¹⁵ The general procedure afforded 354 mg (90 %) of the title compound.

N-(4-Methoxyphenyl)morpholine (Table 12, entry 7).¹⁵ The general procedure afforded 356 mg (90 %) of the title compound.

N-Methyl-N-phenyl-*p*-anisidine (Table 12, entry 8).¹⁵ The general procedure afforded 376 mg (88 %) of the title compound.

N-Phenylmorpholine (Table 12, entry 9).¹⁵ The general procedure afforded 250 mg (77 %) of the title compound.

N-Methyldiphenylamine (Table 12, entry 10).¹⁵ The general procedure afforded 330 mg (90 %) of the title compound.