Crystal chemistry and electronic structure of the metallic lithium ion conductor,

LiNiN

Zlatka Stoeva,^a Bernd Jäger,^b Ruben Gomez,^a Sabri Messaoudi,^c Mouna Ben Yahia,^c Xavier Rocquefelte,^c Gary B. Hix,^d Walter Wolf,^e Jeremy J. Titman,^a Régis Gautier,^c Peter Herzig^b and Duncan H. Gregory.^{f}*

^a School of Chemistry, University of Nottingham, University Park, Nottingham, NG7 2RD, UK.

^b Institut für Physikalische Chemie, Universität Wien, Währinger Straße 42, 1090 Vienna, Austria.

^c Laboratoire de Chimie du Solide et Inorganique Moléculaire, UMR 6511, CNRS - ENSC Rennes, Institut de Chimie de Rennes, Campus de Beaulieu, 35700 Rennes, France.

^d School of Biomedical and Natural Science, Nottingham Trent University, Clifton Lane, Nottingham, NG11 8NS, UK.

^e Materials Design s.a.r.l., 44 Avenue F-A. Bartholdi, 72000 Le Mans, France.

^f WestCHEM, Department of Chemistry, Joseph Black Building, University of Glasgow, Glasgow G12 8QQ, UK.

Author E-mail: D.Gregory@chem.gla.ac.uk

Supporting Information

1. Simulation of ^7Li MAS-NMR spectrum for LiNiN.

Figure S1. Simulation of CQ =145 kHz lithium-7 MAS spectrum with MAS rate 15 kHz and Larmor frequency 116 MHz.