**Cu and Pd-Catalyzed Asymmetric One-Pot Tandem Addition-Cyclization
Reaction of 2-(2',3'-Alkadienyl)- α -ketoesters, Organic Halides, and Dibenzyl
Azodicarboxylate: An Effective Protocol for The Enantioselective Synthesis of
Pyrrolidine Derivatives**

**Shengming Ma^{a,b*}, Ning Jiao^a, Zilong Zheng^a, Zhichao Ma^b, Zhan Lu^b, Longwu
Ye^b, Youquan Deng^b and Guofei Chen^b**

^a State Key Laboratory of Organometallic Chemistry, Shanghai Institute of Organic
Chemistry, Chinese Academy of Sciences, 354 Fenglin Lu, Shanghai 200032, P. R. China

^b Department of Chemistry, Zhejiang University, Hangzhou 310027, P. R. China

E-mail: masm@mail.sioc.ac.cn Fax: (+86)21-64167510

Supporting Information

Table of Contents

Analytical data for compounds 1	S2
¹ H NMR and ¹³ C NMR spectra of those compounds	S16

Experimental section:

1) **Typical procedure: Synthesis of 1,2-Dibenzoxycarbonyl-3-ethoxycarbonyl-3-acetyl-5-(1'-phenylethenyl)pyrazolidine (1aa) ((3R, 5R)-1aa and (3R, 5S)-1aa).** A solution of Cu(OTf)₂ (5.4 mg, 0.015 mmol) and (S)-(-)-2,2'-isopropylidenebis(4-phenyl-2-oxazoline) (5.4 mg, 0.165 mmol) in dry CH₂Cl₂ (2 mL) was stirred at room temperature for 1 h. 2-(2',3'-Dienyl)-ketoester **5a** (27 mg, 0.15 mmol) and dibenzyl azodicarboxylate (DBAD) **6** (54 mg, 0.18 mmol) were added to the solution at 0 °C. The mixture was stirred at this temperature for 3 h as monitored by TLC. Then, 1,4-dioxane (4 mL) was added directly to the mixture followed by the subsequent addition of potassium carbonate (41mg, 0.30 mmol), Pd(PPh₃)₄ (9 mg, 0.0075 mmol), and iodobenzene (37 mg, 0.18 mmol) subsequently under nitrogen. The resulting mixture was heated at 100 °C for 4 h as monitored by TLC. After evaporation, the residue was purified by flash chromatography on silica gel (eluent: petroleum ether/ether = 3:1) to afford 26 mg (31 %) of (3R, 5R)-**1aa** and 51 mg (61 %) (3R, 5S)-**1aa**.

(3R, 5R)-**1aa**: 98% ee as determined by HPLC analysis (Chiralcel AD, 15% *i*-PrOH in hexane, 0.7 mL/min, 230 nm), *t*_r 15.5 (minor), 18.8 (major); [α]_D²⁰ = +54.9 (*c* = 0.79, CHCl₃); viscous liquid; ¹H NMR (CDCl₃, 300 MHz): δ = 7.21-7.39 (m, 15 H), 5.63 (s, 1 H), 5.43 (d, *J* = 8.60 Hz, 1 H), 5.07-5.33 (m, 5 H), 3.92-4.16 (m, 2 H), 3.26 (dd, *J* = 13.43, 9.15 Hz, 1 H), 2.36 (s, 3 H), 2.25 (dd, *J* = 13.43, 3.00 Hz, 1 H), 1.10 (t, *J* = 7.15 Hz, 3 H); ¹³C NMR (CDCl₃, 75.4 MHz): δ = 201.76, 166.76, 156.71, 153.55, 144.59, 138.56, 135.29, 135.09, 128.56, 128.47, 128.45, 128.43, 128.32,

128.15, 127.95, 127.87, 126.61, 112.99, 75.41, 68.79, 68.48, 62.31, 62.04, 38.76, 26.61, 13.58; MS (70 eV): m/z (%): 556 (0.97) [M^+], 469 (53.05) [$M^+ + 1\text{-COOEt-CH}_3$], 289 (100); IR (neat): $\nu = 1734, 1709, 1629, 1458, 1400\text{ cm}^{-1}$; Anal. calcd for $C_{32}H_{32}N_2O_7$ (%): C 69.05, H 5.79, N 5.03; Found C 68.80, H 5.64, N 5.01.

(3R, 5S)-**1aa**: 98% ee as determined by HPLC analysis (Chiralcel AD, 15% *i*-PrOH in hexane, 0.7 mL/min, 230 nm), t_r 20.9 (major), 28.1 (minor); $[\alpha]_D^{20} = +14.33$ ($c = 3.855$, CHCl_3); viscous liquid; ^1H NMR (CDCl_3 , 300 MHz): $\delta = 7.22\text{--}7.38$ (m, 15 H), 5.62 (s, 1 H), 5.44 (d, $J = 6.00$ Hz, 1 H), 5.32 (s, 1 H), 5.26 (d, $J = 12.00$ Hz, 1 H), 5.20 (s, 2 H), 5.11 (d, $J = 12.00$ Hz, 1 H), 3.95–4.22 (m, 2 H), 3.07 (dd, $J = 13.78, 8.85$ Hz, 1 H), 2.30 (dd, $J = 13.78, 3.20$ Hz, 1 H), 2.09 (s, 3 H), 1.12 (t, $J = 7.10$ Hz, 3 H); ^{13}C NMR (CDCl_3 , 75.4 MHz): $\delta = 196.70, 168.72, 156.82, 153.61, 144.74, 138.21, 135.40, 135.18, 128.58, 128.41, 128.37, 128.22, 128.15, 128.12, 128.06, 127.68, 126.55, 113.36, 77.13, 68.52, 68.36, 62.46, 61.32, 39.65, 25.84, 13.68$; MS (70 eV): m/z (%): 556 (0.25) [M^+], 469 (9.69) [$M^+ + 1\text{-COOEt-CH}_3$], 91 (100); IR (neat): $\nu = 1728, 1628, 1498, 1410\text{ cm}^{-1}$; Anal. calcd for $C_{32}H_{32}N_2O_7$ (%): C 69.05, H 5.79, N 5.03; Found C 68.88, H 5.95, N 5.07.

2) **1,2-Dibenzoxycarbonyl-3-ethoxycarbonyl-3-acetyl-5-(1'-phenylethenyl)pyrazolidine (1aa)** ((3S, 5S)-**1aa** and (3S, 5R)-**1aa**). The reaction of **5a** (46 mg, 0.25 mmol), **6** (89 mg, 0.30 mmol), and PhI (61 mg, 0.3 mmol) afforded 120 mg (86%) of **1aa** ((3S, 5S)-**1aa** : (3S, 5R)-**1aa** = 35 : 65) using (R)-(+)-2,2'-isopropylidenebis(4-phenyl-2-oxazoline) as a ligand. (3S, 5S)-**1aa**: 97%

ee as determined by HPLC analysis (Chiralcel AD, 15% *i*-PrOH in hexane, 0.7 mL/min, 230 nm), t_r 17.9 (major), 22.4 (minor); $[\alpha]_D^{20} = -57.7$ ($c = 0.755$, CHCl_3);

(3S, 5R)-**1aa**: 98% ee as determined by HPLC analysis (Chiralcel AD, 15% *i*-PrOH in hexane, 0.7 mL/min, 230 nm), t_r 24.6 (minor), 33.9 (major); $[\alpha]_D^{20} = -14.8$ ($c = 0.95$, CHCl_3).

3) **1,2-Dibenzoxycarbonyl-3-ethoxycarbonyl-3-acetyl-5-(1'-(4''-methylphenyl)ethenyl)pyrazolidine (1ab) ((3R, 5R)-1ab and (3R, 5S)-1ab)**. The reaction of **5a** (46 mg, 0.25 mmol), **6** (89 mg, 0.30 mmol), and 4-iodotoluene (65 mg, 0.3 mmol) afforded 135 mg (95%) of **1ab** ((3R, 5R)-**1ab** : (3R, 5S)-**1ab** = 34 : 66) (S)-(-)-2,2'-isopropylidenebis(4-phenyl-2-oxazoline) as a ligand. (3R, 5R)-**1ab**: 98% ee as determined by HPLC analysis (Chiralcel AD, 10% *i*-PrOH in hexane, 0.7 mL/min, 230 nm), t_r 20.6 (minor), 22.0 (major); $[\alpha]_D^{20} = +45.3$ ($c = 0.96$, CHCl_3); viscous liquid; ^1H NMR (CDCl_3 , 300 MHz): $\delta = 7.16\text{--}7.30$ (m, 10 H), 7.12 (d, $J = 8.15$ Hz, 2 H), 7.04 (d, $J = 8.15$ Hz, 2 H), 5.51 (s, 1 H), 5.35 (d, $J = 7.20$ Hz, 1 H), 4.97–5.24 (m, 5 H), 3.83–4.03 (m, 2 H), 3.17 (dd, $J = 13.58, 9.40$ Hz, 1 H), 2.27 (s, 3 H), 2.24 (s, 3 H), 2.17 (dd, $J = 13.58, 3.05$ Hz, 1 H), 1.01 (t, $J = 7.25$ Hz, 3 H); ^{13}C NMR (CDCl_3 , 75.4 MHz): $\delta = 201.77, 166.78, 156.72, 153.55, 144.36, 135.60, 137.69, 135.31, 135.12, 129.13, 128.55, 128.46, 128.40, 128.31, 128.14, 127.93, 126.44, 112.15, 75.40, 68.77, 68.47, 62.30, 62.05, 38.79, 26.62, 21.06, 13.58$; MS (70 eV): m/z (%): 483 (6.55) [$\text{M}^+ + 1\text{-COOEt-CH}_3$], 91 (100); IR (neat): $\nu = 1723, 1632, 1585, 1409, 1262\text{ cm}^{-1}$; Anal. calcd for $\text{C}_{33}\text{H}_{34}\text{N}_2\text{O}_7$ (%): C 69.46, H 6.01, N 4.91; Found C 69.41, H 5.98, N 4.83.

(3R, 5S)-**1ab**: 98% ee as determined by HPLC analysis (Chiralcel AD, 10% *i*-PrOH in hexane, 0.7 mL/min, 230 nm), t_r 29.8 (major), 33.1 (minor); $[\alpha]_D^{20} = +17.3$ ($c = 1.14$, CHCl₃); viscous liquid; ¹H NMR (CDCl₃, 300 MHz): $\delta = 7.26$ -7.38 (m, 10 H), 7.22 (d, $J = 8.15$ Hz, 2 H), 7.14 (d, $J = 8.15$ Hz, 2 H), 5.60 (s, 1 H), 5.45 (d, $J = 5.70$ Hz, 1 H), 5.08-5.34 (m, 5 H), 3.96-4.21 (m, 2 H), 3.08 (dd, $J = 13.60, 9.20$ Hz, 1 H), 2.24-2.38 (m, 4 H), 2.08 (s, 3 H), 1.12 (t, $J = 7.15$ Hz, 3 H); ¹³C NMR (CDCl₃, 75.4 MHz): $\delta = 196.81, 168.77, 156.87, 153.58, 144.58, 137.96, 135.45, 135.30, 135.24, 129.29, 128.43, 128.40, 128.24, 128.16, 128.14, 127.71, 126.42, 112.57, 77.14, 68.53, 68.36, 62.45, 61.39, 39.66, 25.73, 21.03, 13.70$; MS (70 eV): m/z (%): 483 (1.95) [M⁺+1-COOEt-CH₃], 91 (100); IR (neat): $\nu = 1732, 1633, 1499, 1411$ cm⁻¹; Anal. calcd for C₃₃H₃₄N₂O₇ (%): C 69.46, H 6.01, N 4.91; Found C 69.39, H 6.04, N 4.87.

4) **1,2-Dibenzoxycarbonyl-3-ethoxycarbonyl-3-acetyl-5-(1'-(1'-(4''-methoxyphenyl)ethenyl)pyrazolidine (1ac) ((3R, 5R)-1ac and (3R, 5S)-1ac)**. The reaction of **5a** (46 mg, 0.25 mmol), **6** (89 mg, 0.30 mmol), and 4-iodoanisole (70 mg, 0.3 mmol) afforded 135 mg (92%) of **1ac** ((3R, 5R)-**1ac** : (3R, 5S)-**1ac** = 33 : 67) using (S)-(-)-2,2'-isopropylidenebis(4-phenyl-2-oxazoline) as a ligand. (3R, 5R)-**1ac**: 98% ee as determined by HPLC analysis (Chiralcel AD, 15% *i*-PrOH in hexane, 0.7 mL/min, 230 nm), t_r 24.8 (minor), 26.6 (major); $[\alpha]_D^{20} = +43.5$ ($c = 0.76$, CHCl₃); viscous liquid; ¹H NMR (CDCl₃, 300 MHz): $\delta = 7.29$ -7.41 (m, 8 H), 7.20-7.28 (m, 4 H), 6.84 (d, $J = 9.10$ Hz, 2 H), 5.55 (s, 1 H), 5.39 (d, $J = 6.80$ Hz, 1 H), 5.08-5.32 (m, 5 H), 3.94-4.11 (m, 2 H), 3.80 (s, 3 H), 3.26 (dd, $J = 13.53, 9.40$ Hz, 1 H), 2.35 (s, 3

H), 2.26 (dd, $J = 13.53, 3.15$ Hz, 1 H), 1.10 (t, $J = 7.05$ Hz, 3 H); ^{13}C NMR (CDCl_3 , 75.4 MHz): $\delta = 201.80, 166.77, 159.26, 156.71, 153.53, 143.93, 135.30, 135.11, 130.92, 128.54, 128.46, 128.41, 128.13, 127.93, 127.70, 126.60, 113.79, 111.54, 75.41, 68.76, 68.47, 62.30, 62.08, 55.21, 38.80, 26.62, 13.60$; MS (70 eV): m/z (%): 586 (0.68) $[\text{M}^+]$, 91 (100); IR (neat): $\nu = 1716, 1608, 1513, 1407, 1249$ cm^{-1} ; Anal. calcd for $\text{C}_{33}\text{H}_{34}\text{N}_2\text{O}_8$ (%): C 67.56, H 5.84, N 4.78; Found C 67.90, H 6.00, N 4.72.

(3R, 5S)-**1ac**: 99% ee as determined by HPLC analysis (Chiralcel AD, 15% *i*-PrOH in hexane, 0.7 mL/min, 230 nm), t_r 33.7 (major), 40.8 (minor); $[\alpha]_D^{20} = +24.6$ ($c = 0.965$, CHCl_3); viscous liquid; ^1H NMR (CDCl_3 , 300 MHz): $\delta = 7.28\text{--}7.41$ (m, 10 H), 7.26 (d, $J = 8.55$ Hz, 2 H), 6.85 (d, $J = 8.55$ Hz, 2 H), 5.55 (s, 1 H), 5.41 (d, $J = 6.10$ Hz, 1 H), 5.04–5.28 (m, 5 H), 3.92–4.21 (m, 2 H), 3.80 (s, 3 H), 3.09 (dd, $J = 12.30, 9.00$ Hz, 1 H), 2.30 (d, $J = 12.30$ Hz, 1 H), 2.07 (s, 3 H), 1.11 (t, $J = 7.30$ Hz, 3 H); ^{13}C NMR (CDCl_3 , 75.4 MHz): $\delta = 196.78, 168.78, 159.43, 156.87, 153.58, 144.15, 135.45, 135.24, 130.59, 128.43, 128.40, 128.25, 128.17, 128.14, 127.72, 113.94, 111.92, 77.16, 68.54, 68.37, 62.47, 61.43, 55.22, 39.70, 25.77, 13.72$; MS (70 eV): m/z (%): 586 (0.11) $[\text{M}^+]$, 91 (100); IR (neat): $\nu = 1728, 1608, 1513, 1411, 1248$ cm^{-1} ; Anal. calcd for $\text{C}_{33}\text{H}_{34}\text{N}_2\text{O}_8$ (%): C 67.56, H 5.84, N 4.78; Found C 67.92, H 5.91, N 4.72.

5) **1,2-Dibenzoxycarbonyl-3-ethoxycarbonyl-3-acetyl-5-(1'-(4''-methoxycarbonylphenyl)ethenyl)pyrazolidine (1ad) ((3R, 5R)-1ad and (3R, 5S)-1ad).**

The reaction of **5a** (46 mg, 0.25 mmol), **6** (89 mg, 0.30 mmol), and 4-iodobenzoic acid methyl ester (79 mg, 0.3 mmol) afforded 147 mg (96%) of **1ad** ((3R, 5R)-**1ad** :

(3R, 5S)-**1ad** = 38 : 62) using (S)-(-)-2,2'-isopropylidenebis(4-phenyl-2-oxazoline) as a ligand. (3R, 5R)-**1ad**: 97% ee as determined by HPLC analysis (Chiralcel AD, 15% *i*-PrOH in hexane, 1 mL/min, 230 nm), t_r 23.6 (minor), 27.9 (major); $[\alpha]_D^{20} = +40.6$ ($c = 0.85$, CHCl₃); viscous liquid; ¹H NMR (CDCl₃, 300 MHz): $\delta = 7.98$ (d, $J = 8.10$ Hz, 2 H), 7.21-7.42 (m, 12 H), 5.72 (d, $J = 1.90$ Hz, 1 H), 5.45 (d, $J = 7.30$ Hz, 1 H), 5.39 (s, 1 H), 5.04-5.33 (m, 4 H), 3.96-4.14 (m, 2 H), 3.91 (s, 3 H), 3.28 (dd, $J = 13.58, 9.35$ Hz, 1 H), 2.35 (s, 3 H), 2.20 (dd, $J = 13.58, 2.80$ Hz, 1 H), 1.10 (t, $J = 7.25$ Hz, 3 H); ¹³C NMR (CDCl₃, 75.4 MHz): $\delta = 201.69, 166.70, 166.62, 156.65, 153.51, 144.02, 143.16, 135.21, 135.00, 129.75, 129.47, 128.60, 128.51, 128.41, 128.20, 128.02, 126.64, 114.98, 75.41, 68.93, 68.58, 62.43, 61.83, 52.13, 38.72, 26.64, 13.61$; MS (70 eV): m/z (%): 527 (9.59) [M⁺+1-COOEt-CH₃], 91 (100); IR (neat): $\nu = 1720, 1608, 1406, 1282$ cm⁻¹; Anal. calcd for C₃₄H₃₄N₂O₉ (%): C 66.44, H 5.58, N 4.56; Found C 66.50, H 5.61, N 4.55.

(3R, 5S)-**1ad**: 98% ee as determined by HPLC analysis (Chiralcel AD, 15% *i*-PrOH in hexane, 1 mL/min, 230 nm), t_r 31.7 (major), 45.2 (minor); $[\alpha]_D^{20} = +19.0$ ($c = 0.38$, CHCl₃); viscous liquid; ¹H NMR (CDCl₃, 300 MHz): $\delta = 7.99$ (d, $J = 8.40$ Hz, 2 H), 7.39 (d, $J = 8.40$ Hz, 2 H), 7.25-7.37 (m, 10 H), 5.69 (s, 1 H), 5.05-5.41 (m, 6 H), 3.98-4.21 (m, 2 H), 3.92 (s, 3 H), 3.03 (dd, $J = 13.73, 9.15$ Hz, 1 H), 2.29 (d, $J = 13.73$ Hz, 1 H), 2.10 (s, 3 H), 1.12 (t, $J = 7.20$ Hz, 3 H); ¹³C NMR (CDCl₃, 75.4 MHz): $\delta = 196.47, 168.76, 166.58, 156.82, 153.85, 144.18, 142.95, 135.43, 135.17, 129.91, 129.69, 128.51, 128.48, 128.36, 128.28, 128.21, 127.81, 126.64, 115.36, 77.20, 68.67, 68.53, 62.64, 61.13, 52.19, 39.97, 26.40, 13.76$; MS (70 eV): m/z (%):

527 (18.01) [$M^+ + 1$ -COOEt-CH₃], 91 (100); IR (neat): ν = 1720, 1609, 1396, 1279 cm⁻¹; Anal. calcd for C₃₄H₃₄N₂O₉ (%): C 66.44, H 5.58, N 4.56; Found C 66.18, H 5.64, N 4.51.

6) **1,2-Dibenzoxycarbonyl-3-ethoxycarbonyl-3-acetyl-5-(1'-(4''-bromo-phenyl)-ethenyl)pyrazolidine (1ae) ((3R, 5R)-1ae and (3R, 5S)-1ae).** The reaction of **5a** (46 mg, 0.25 mmol), **6** (89 mg, 0.30 mmol), and 1-bromo-4-iodobenzene (85 mg, 0.3 mmol) afforded 150 mg (94%) of **1ae** ((3R, 5R)-**1ae** : (3R, 5S)-**1ae** = 36 : 64) using (S)-(-)-2,2'-isopropylidenebis(4-phenyl-2-oxazoline) as a ligand. (3R, 5R)-**1ae**: 97% ee as determined by HPLC analysis (Chiralcel AD, 10% *i*-PrOH in hexane, 0.7 mL/min, 230 nm), t_r 31.0 (minor), 33.55 (major); $[\alpha]_D^{20}$ = +45.4 (c = 0.71, CHCl₃); white solid, Mp: 105-107 °C (*n*-hexane/ethyl acetate); ¹H NMR (CDCl₃, 300 MHz): δ = 7.41 (d, J = 8.70 Hz, 2 H), 7.19-7.39 (m, 10 H), 7.17 (d, J = 8.70 Hz, 2 H), 5.64 (s, 1 H), 5.36 (d, J = 8.50 Hz, 1 H), 5.04-5.31 (m, 5 H), 3.92-4.14 (m, 2 H), 3.26 (dd, J = 13.55, 9.25 Hz, 1 H), 2.35 (s, 3 H), 2.21 (dd, J = 13.55, 2.95 Hz, 1 H), 1.10 (t, J = 7.05 Hz, 3 H); ¹³C NMR (CDCl₃, 75.4 MHz): δ = 201.67, 166.67, 156.62, 153.47, 143.76, 137.54, 135.20, 134.99, 131.55, 128.57, 128.50, 128.49, 128.39, 128.32, 128.18, 127.99, 121.88, 113.92, 75.41, 68.88, 68.54, 62.42, 61.88, 38.64, 26.64, 13.62; MS (70 eV): m/z (%): 635 (0.39) [M^+], 636 (0.28) [$M^+ + 1$], 91 (100); IR (KBr): ν = 1716, 1590, 1406, 1352, 1272 cm⁻¹; Anal. calcd for C₃₂H₃₁BrN₂O₇ (%): C 60.48, H 4.92, N 4.41; Found C 60.45, H 4.90, N 4.32.

(3R, 5R)-**1ae**: >98% ee as determined by HPLC analysis (Chiralcel AD, 10% *i*-PrOH in hexane, 0.7 mL/min, 230 nm), t_r 48.8 (major); $[\alpha]_D^{20}$ = +8.5 (c = 0.995, CHCl₃);

white solid, Mp: 101-103 °C (*n*-hexane/ethyl acetate); ¹H NMR (CDCl₃, 300 MHz): δ = 7.43 (d, *J* = 8.25 Hz, 2 H), 7.21-7.39 (m, 10 H), 7.17 (d, *J* = 8.25 Hz, 2 H), 5.61 (s, 1 H), 5.12-5.41 (m, 5 H), 5.09 (d, *J* = 11.70 Hz, 1 H), 3.95-4.21 (m, 2 H), 3.02 (dd, *J* = 13.23, 9.05 Hz, 1 H), 2.29 (d, *J* = 13.23 Hz, 1 H), 2.10 (s, 3 H), 1.12 (t, *J* = 7.15 Hz, 3 H); ¹³C NMR (CDCl₃, 75.4 MHz): δ = 196.42, 168.68, 156.74, 153.73, 143.86, 137.28, 135.36, 135.11, 131.66, 128.42, 128.40, 128.28, 128.23, 128.18, 128.12, 127.72, 122.08, 114.13, 77.10, 68.56, 68.42, 62.54, 61.12, 39.81, 26.29, 13.70; MS (70 eV): *m/z* (%): 635 (0.55) [M⁺], 636 (0.31) [M⁺+1], 91 (100); IR (KBr): ν = 1728, 1587, 1394, 1304 cm⁻¹; Anal. calcd for C₃₂H₃₁BrN₂O₇ (%): C 60.48, H 4.92, N 4.41; Found C 60.44, H 4.82, N 4.41

7) **1,2-Dibenzoxycarbonyl-3-ethoxycarbonyl-3-acetyl-5-(1'-(4''-nitrophenyl)-ethenyl)pyrazolidine (1af) ((3R, 5R)-1af and (3R, 5S)-1af)**. The reaction of **5a** (46 mg, 0.25 mmol), **6** (89 mg, 0.30 mmol), and 1-nitro-4-iodobenzene (75 mg, 0.3 mmol) afforded 145 mg (97%) of **1af** ((3R, 5R)-**1af** : (3R, 5S)-**1af** = 40 : 60) using (S)-(-)-2,2'-isopropylidenebis(4-phenyl-2-oxazoline) as a ligand. (3R, 5R)-**1af**: >98% ee as determined by HPLC analysis (Chiralcel AD, 40% *i*-PrOH in hexane, 0.7 mL/min, 230 nm), *t_r* 22.3 (major); [α]_D²⁰ = +45.6 (*c* = 0.87, CHCl₃); viscous liquid; ¹H NMR (CDCl₃, 300 MHz): δ = 8.12 (d, *J* = 8.75 Hz, 2 H), 7.46 (d, *J* = 8.75 Hz, 2 H), 7.20-7.41 (m, 10 H), 5.77 (d, *J* = 2.00 Hz, 1 H), 5.42 (s, 2 H), 5.04-5.28 (m, 4 H), 3.94-4.16 (m, 2 H), 3.32 (dd, *J* = 13.48, 9.25 Hz, 1 H), 2.36 (s, 3 H), 2.22 (dd, *J* = 13.48, 3.10 Hz, 1 H), 1.12 (t, *J* = 7.25 Hz, 3 H); ¹³C NMR (CDCl₃, 75.4 MHz): δ = 201.59, 166.60, 156.52, 153.41, 147.18, 145.34, 143.48, 135.05, 134.84, 128.60,

128.52, 128.49, 128.36, 128.20, 128.07, 127.59, 123.64, 116.92, 75.45, 69.03, 68.63, 62.56, 61.72, 38.56, 26.66, 13.62; MS (70 eV): m/z (%): 514 (10.96) [$M^+ + 1$ -COOEt-CH₃], 91 (100); IR (neat): ν = 1715, 1595, 1520, 1346 cm⁻¹; Anal. calcd for C₃₂H₃₁N₃O₉ (%): C 63.89, H 5.19, N 6.98; Found C 64.12, H 5.43, N 6.74.

(3R, 5S)-**1af**: 99% ee as determined by HPLC analysis (Chiralcel AD, 40% *i*-PrOH in hexane, 0.7 mL/min, 230 nm), t_r 23.2 (major), 28.4 (minor); $[\alpha]_D^{20}$ = +17.1 (c = 0.855, CHCl₃); viscous liquid; ¹H NMR (CDCl₃, 300 MHz): δ = 8.14 (d, J = 8.70 Hz, 2 H), 7.47 (d, J = 8.70 Hz, 2 H), 7.21-7.41 (m, 10 H), 5.72 (s, 1 H), 5.30-5.44 (m, 2 H), 5.11-5.28 (m, 3 H), 5.07 (d, J = 12.10 Hz, 1 H), 4.00-4.22 (m, 2 H), 2.90-3.06 (m, 1 H), 2.23-2.40 (m, 1 H), 2.16 (s, 3 H), 1.15 (t, J = 7.25 Hz, 3 H); ¹³C NMR (CDCl₃, 75.4 MHz): δ = 196.21, 168.72, 156.72, 153.96, 147.30, 145.16, 143.61, 135.33, 135.00, 128.52, 128.49, 128.43, 128.31, 128.19, 127.83, 127.58, 123.77, 116.95, 68.70, 68.60, 67.84, 62.73, 60.97, 40.20, 27.03, 13.75; MS (70 eV): m/z (%): 514 (8.79) [$M^+ + 1$ -COOEt-CH₃], 91 (100); IR (neat): ν = 1728, 1597, 1519, 1345 cm⁻¹; Anal. calcd for C₃₂H₃₁N₃O₉ (%): C 63.89, H 5.19, N 6.98; Found C 63.61, H 5.26, N 6.84.

8) **1,2-Dibenzoxycarbonyl-3-ethoxycarbonyl-3-acetyl-5-(1'-methylene-3'-methoxycarbonyl-2'-propenyl)pyrazolidine (1ag) ((3R, 5R)-1ag and (3R, 5S)-1ag)**. The reaction of **5a** (46 mg, 0.25 mmol), **6** (89 mg, 0.30 mmol), and (*E*)-1-iodo-2-methoxycarbonyl-ethylene (64 mg, 0.3 mmol) afforded 138 mg (98%) of **1ag** ((3R, 5R)-**1ag** : (3R, 5S)-**1ag** = 45 : 55) using (*S*)-(-)-2,2'-isopropylidenebis(4-phenyl-2-oxazoline) as a ligand. (3R, 5R)-**1ag**: 97%

ee as determined by HPLC analysis (Chiralcel AD, 15% *i*-PrOH in hexane, 1 mL/min, 230 nm), t_r 18.4 (minor), 20.1 (major); $[\alpha]_D^{20} = +44.3$ ($c = 1.18$, CHCl_3); viscous liquid; ^1H NMR (CDCl_3 , 300 MHz): $\delta = 7.19\text{--}7.40$ (m, 11 H), 5.86 (s, 1 H), 5.79 (d, $J = 16.40$ Hz, 1 H), 5.43 (s, 1 H), 5.28 (d, $J = 12.20$ Hz, 1 H), 5.00–5.21 (m, 4 H), 3.94–4.14 (m, 2 H), 3.76 (s, 3 H), 3.46 (dd, $J = 13.30, 9.50$ Hz, 1 H), 2.26–2.40 (m, 4 H), 1.10 (t, $J = 7.10$ Hz, 3 H); ^{13}C NMR (CDCl_3 , 75.4 MHz): $\delta = 201.54, 166.81, 166.65, 156.36, 153.28, 143.57, 141.34, 135.11, 134.90, 128.60, 128.54, 128.47, 128.26, 128.04, 123.25, 118.66, 75.34, 68.97, 68.67, 62.56, 59.78, 51.77, 39.30, 26.65, 13.57$; MS (70 eV): m/z (%): 477 (6.17) [$\text{M}^+ + 1\text{-COOEt-CH}_3$], 91 (100); IR (neat): $\nu = 1721, 1634, 1607, 1406, 1279\text{ cm}^{-1}$; Anal. calcd for $\text{C}_{30}\text{H}_{32}\text{N}_2\text{O}_9$ (%): C 63.82, H 5.71, N 4.96; Found C 63.87, H 5.78, N 4.96.

(3R, 5R)-**1ag**: 98% ee as determined by HPLC analysis (Chiralcel AD, 15% *i*-PrOH in hexane, 1 mL/min, 230 nm), t_r 25.1 (major), 28.3 (minor); $[\alpha]_D^{20} = -23.5$ ($c = 0.75$, CHCl_3); viscous liquid; ^1H NMR (CDCl_3 , 300 MHz): $\delta = 7.22\text{--}7.40$ (m, 11 H), 5.74–5.84 (m, 2 H), 5.45 (s, 1 H), 5.24 (d, $J = 12.00$ Hz, 1 H), 5.16 (s, 2 H), 4.87–5.12 (m, 2 H), 4.00–4.22 (m, 2 H), 3.76 (s, 3 H), 3.14–3.26 (m, 1 H), 2.35–2.46 (m, 1 H), 2.16 (s, 3 H), 1.15 (t, $J = 7.20$ Hz, 3 H); ^{13}C NMR (CDCl_3 , 75.4 MHz): $\delta = 195.99, 168.62, 166.68, 156.43, 153.78, 143.32, 141.42, 135.35, 135.00, 128.47, 128.39, 128.26, 128.22, 127.81, 123.30, 118.89, 77.20, 68.62, 62.74, 58.90, 51.81, 40.45, 40.42, 26.61, 13.77$; MS (70 eV): m/z (%): 477 (4.94) [$\text{M}^+ + 1\text{-COOEt-CH}_3$], 91 (100); IR (neat): $\nu = 1732, 1609, 1587, 1396, 1166\text{ cm}^{-1}$; Anal. calcd for $\text{C}_{30}\text{H}_{32}\text{N}_2\text{O}_9$ (%): C 63.82, H 5.71, N 4.96; Found C 63.61, H 5.77, N 4.89.

9) **1,2-Dibenzoxycarbonyl-3-ethoxycarbonyl-3-acetyl-5-(1'-methylene-3'-phenyl-2'(*E*)-propenyl)pyrazolidine (1ah) ((3R, 5R)-1ah and (3R, 5S)-1ah).** The reaction of **5a** (46 mg, 0.25 mmol), **6** (89 mg, 0.30 mmol), and (*E*)-1-iodo-2-phenylethylene (69 mg, 0.3 mmol) afforded 125 mg (86%) of **1ah** ((3R, 5R)-**1ah**: (3R, 5S)-**1ah** = 37 : 63) using (S)-(-)-2,2'-isopropylidenebis(4-phenyl-2-oxazoline) as a ligand. (3R, 5R)-**1ah**: 99% ee as determined by HPLC analysis (Chiralcel AD, 10% *i*-PrOH in hexane, 1 mL/min, 230 nm), t_r 16.1 (minor), 18.1 (major); $[\alpha]_D^{20} = +74.1$ ($c = 1.605$, CHCl₃); viscous liquid; ¹H NMR (CDCl₃, 300 MHz): $\delta = 7.20$ -7.43 (m, 15 H), 6.80 (d, $J = 16.90$ Hz, 1 H), 6.43 (d, $J = 16.90$ Hz, 1 H), 5.60 (s, 1 H), 5.30 (d, $J = 12.40$ Hz, 1 H), 5.12-5.24 (m, 4 H), 5.08 (d, $J = 12.40$ Hz, 1 H), 3.96-4.18 (m, 2 H), 3.54 (dd, $J = 13.05, 9.75$ Hz, 1 H), 2.48 (dd, $J = 13.05, 3.55$ Hz, 1 H), 2.39 (s, 3 H), 1.11 (t, $J = 7.10$ Hz, 3 H); ¹³C NMR (CDCl₃, 75.4 MHz): $\delta = 201.91, 166.76, 156.52, 153.34, 142.75, 136.62, 135.25, 135.10, 129.15, 128.64, 128.57, 128.50, 128.44, 128.36, 128.19, 127.97, 127.86, 127.80, 126.36, 115.75, 75.45, 68.79, 68.57, 62.47, 60.40, 39.70, 26.71, 13.59$; MS (70 eV): m/z (%): 495 (3.27) [$M^+ + 1$ -COOEt-CH₃], 91 (100); IR (neat): $\nu = 1716, 1602, 1498, 1355, 1272$ cm⁻¹; Anal. calcd for C₃₄H₃₄N₂O₇ (%): C 70.09, H 5.88, N 4.81; Found C 69.71, H 5.94, N 4.82.

(3R, 5S)-**1ah**: 99% ee as determined by HPLC analysis (Chiralcel AD, 10% *i*-PrOH in hexane, 1 mL/min, 230 nm), t_r 22.6 (major), 34.5 (minor); $[\alpha]_D^{20} = +2.9$ ($c = 0.49$, CHCl₃); viscous liquid; ¹H NMR (CDCl₃, 300 MHz): $\delta = 7.22$ -7.45 (m, 15 H), 6.82 (d, $J = 16.55$ Hz, 1 H), 6.46 (d, $J = 16.55$ Hz, 1 H), 5.60 (s, 1 H), 5.08-5.31 (m, 6 H),

4.00-4.26 (m, 2 H), 3.38 (dd, $J = 11.58, 11.58$ Hz, 1 H), 2.54 (dd, $J = 11.58, 2.70$ Hz, 1 H), 2.17 (s, 3 H), 1.17 (t, $J = 7.10$ Hz, 3 H); ^{13}C NMR (CDCl_3 , 75.4 MHz): $\delta = 196.60, 168.87, 156.65, 153.49, 142.88, 136.43, 135.46, 135.18, 129.55, 128.66, 128.45, 128.44, 128.31, 128.20, 128.19, 127.99, 127.78, 127.50, 126.42, 115.87, 77.19, 68.56, 68.48, 62.61, 59.62, 40.51, 25.99, 13.78$; MS (70 eV): m/z (%): 495 (0.96) [$\text{M}^+ + 1\text{-COOEt-CH}_3$], 91 (100); IR (neat): $\nu = 1725, 1601, 1498, 1353, 1159$ cm^{-1} ; Anal. calcd for $\text{C}_{34}\text{H}_{34}\text{N}_2\text{O}_7$ (%): C 70.09, H 5.88, N 4.81; Found C 70.17, H 5.98, N 4.99.

10) **1,2-Dibenzoxycarbonyl-3-ethoxycarbonyl-3-acetyl-5-(1'-methylene-2'(E)-heptenyl)pyrazolidine (1ai) ((3R, 5R)-1ai and (3R, 5S)-1ai).** The reaction of **5a** (46 mg, 0.25 mmol), **6** (89 mg, 0.30 mmol), and (*E*)-1-iodo-1-hexene (63 mg, 0.3 mmol) afforded 106 mg (75%) of **1ai** ((3R, 5R)-**1ai** : (3R, 5S)-**1ai** = 34 : 66) using (*S*)-(-)-2,2'-isopropylidenebis(4-phenyl-2-oxazoline) as a ligand. (3R, 5R)-**1ai**: 97% ee as determined by HPLC analysis (Chiralcel AD, 15% *i*-PrOH in hexane, 0.7 mL/min, 230 nm), t_r 9.54 (major), 13.9 (minor); $[\alpha]_D^{20} = +79.9$ ($c = 1.215, \text{CHCl}_3$); viscous liquid; ^1H NMR (CDCl_3 , 300 MHz): $\delta = 7.19\text{-}7.41$ (m, 10 H), 6.04 (d, $J = 16.30$ Hz, 1 H), 5.54 (dt, $J = 16.30, 6.85$ Hz, 1 H), 5.38 (s, 1 H), 5.28 (d, $J = 12.30$ Hz, 1 H), 4.98-5.21 (m, 4 H), 4.94 (s, 1 H), 3.94-4.18 (m, 2 H), 3.38 (dd, $J = 13.20, 9.70$ Hz, 1 H), 2.39 (dd, $J = 13.20, 3.50$ Hz, 1 H), 2.36 (s, 3 H), 2.08 (q, $J = 6.80$ Hz, 2 H), 1.22-1.41 (m, 4 H), 1.12 (t, $J = 7.05$ Hz, 3 H), 0.89 (t, $J = 7.10$ Hz, 3 H); ^{13}C NMR (CDCl_3 , 75.4 MHz): $\delta = 202.13, 166.84, 156.54, 153.41, 142.61, 135.32, 135.19, 131.67, 129.15, 128.55, 128.48, 128.40, 128.33, 128.16, 127.93, 112.84,$

75.49, 68.69, 68.49, 62.39, 60.51, 39.65, 32.79, 31.32, 26.68, 22.21, 13.90, 13.58; MS (70 eV): m/z (%): 562 (0.09) [M^+], 91 (100); IR (neat): ν = 1716, 1610, 1499, 1409, 1264 cm^{-1} ; Anal. calcd for $\text{C}_{32}\text{H}_{38}\text{N}_2\text{O}_7$ (%): C 68.31, H 6.81, N 4.98; Found C 68.60, H 6.91, N 4.96.

(3R, 5S)-**1ai**: 98% ee as determined by HPLC analysis (Chiralcel AD, 15% *i*-PrOH in hexane, 0.7 mL/min, 230 nm), t_r 13.2 (major), 15.5 (minor); $[\alpha]_D^{20} = -9.4$ ($c = 0.795$, CHCl_3); viscous liquid; ^1H NMR (CDCl_3 , 300 MHz): δ = 7.25-7.38 (m, 10 H), 6.06 (d, $J = 16.20$ Hz, 1 H), 5.60 (dt, $J = 16.20, 7.05$ Hz, 1 H), 5.38 (s, 1 H), 5.26 (d, $J = 12.00$ Hz, 1 H), 4.94-5.19 (m, 5 H), 3.98-4.23 (m, 2 H), 3.26 (dd, $J = 13.13, 9.45$ Hz, 1 H), 2.43 (dd, $J = 13.13, 3.90$ Hz, 1 H), 2.03-2.18 (m, 5 H), 1.20-1.43 (m, 4 H), 1.14 (t, $J = 7.05$ Hz, 3 H), 0.91 (t, $J = 7.20$ Hz, 3 H); MS (70 eV): m/z (%): 475 (2.50) [$M^+ + 1\text{-COOEt-CH}_3$], 91 (100); IR (neat): ν = 1728, 1610, 1588, 1456, 1352 cm^{-1} ; Anal. calcd for $\text{C}_{32}\text{H}_{38}\text{N}_2\text{O}_7$ (%): C 68.31, H 6.81, N 4.98; Found C 68.59, H 6.88, N 4.91.

11) **1,2-Dibenzoxycarbonyl-3-ethoxycarbonyl-3-acetyl-5-(1'-(2''-thienyl)-ethenyl)pyrazolidine (1aj)** ((3R, 5R)-**1aj** and (3R, 5S)-**1aj**). The reaction of **5a** (46 mg, 0.25 mmol), **6** (89 mg, 0.30 mmol), and 2-iodothiophene (63 mg, 0.3 mmol) afforded 104 mg (71%) of **1aj** ((3R, 5R)-**1aj** : (3R, 5S)-**1aj** = 36 : 64) using (S)-(-)-2,2'-isopropylidenebis(4-phenyl-2-oxazoline) as a ligand. (3R, 5R)-**1aj**: 91% ee as determined by HPLC analysis (Chiralcel AD, 10% *i*-PrOH in hexane, 1 mL/min, 230 nm), t_r 21.8 (minor), 25.7 (major); $[\alpha]_D^{20} = +43.8$ ($c = 1.05$, CHCl_3); viscous liquid; ^1H NMR (CDCl_3 , 300 MHz): δ = 7.22-7.41 (m, 10 H), 7.19 (dd, $J =$

5.10, 1.18 Hz, 1 H), 6.97 (dd, $J = 5.10, 3.68$ Hz, 1 H), 6.92 (dd, $J = 3.68, 1.18$ Hz, 1 H), 5.55 (s, 1 H), 5.39 (s, 1 H), 5.31 (d, $J = 11.80$ Hz, 2 H), 5.04-5.24 (m, 3 H), 3.92-4.13 (m, 2 H), 3.42 (dd, $J = 13.60, 9.35$ Hz, 1 H), 2.44 (dd, $J = 13.60, 3.35$ Hz, 1 H), 2.37 (s, 3 H), 1.09 (t, $J = 7.15$ Hz, 3 H); ^{13}C NMR (CDCl_3 , 75.4 MHz): $\delta = 201.77, 166.76, 156.52, 153.44, 141.71, 138.38, 135.23, 135.05, 128.58, 128.52, 128.47, 128.41, 128.25, 127.99, 127.42, 124.62, 123.82, 111.64, 75.39, 68.87, 68.59, 62.43, 61.83, 39.40, 26.66, 13.54$; MS (70 eV): m/z (%): 475 (3.15) [$\text{M}^+ + 1 - \text{COOEt} - \text{CH}_3$], 91 (100); IR (neat): $\nu = 1717, 1626, 1587, 1408, 1265\text{ cm}^{-1}$; Anal. calcd for $\text{C}_{30}\text{H}_{30}\text{N}_2\text{O}_7\text{S}$ (%): C 64.04, H 5.37, N 4.98; Found C 64.02, H 5.46, N 4.80.

(3R, 5R)-**1aj**: 99% ee as determined by HPLC analysis (Chiralcel AD, 10% *i*-PrOH in hexane, 1 mL/min, 230 nm), t_r 38.3 (major), 55.3 (minor); $[\alpha]_D^{20} = +4.4$ ($c = 1.125, \text{CHCl}_3$); viscous liquid; ^1H NMR (CDCl_3 , 300 MHz): $\delta = 7.26\text{--}7.41$ (m, 10 H), 7.21 (d, $J = 5.10$ Hz, 1 H), 6.91-7.02 (m, 2 H), 5.55 (s, 1 H), 5.42 (s, 1 H), 5.16-5.38 (m, 4 H), 5.12 (d, $J = 12.30$ Hz, 1 H), 3.96-4.23 (m, 2 H), 3.27 (dd, $J = 13.50, 9.15$ Hz, 1 H), 2.49 (dd, $J = 13.50, 3.45$ Hz, 1 H), 2.13 (s, 3 H), 1.14 (t, $J = 6.90$ Hz, 3 H); ^{13}C NMR (CDCl_3 , 75.4 MHz): $\delta = 196.47, 168.68, 156.56, 153.49, 141.24, 138.57, 135.35, 135.08, 128.60, 128.40, 128.32, 128.29, 128.16, 127.70, 127.45, 124.93, 124.04, 111.70, 77.21, 68.56, 68.45, 67.73, 62.54, 40.31, 25.86, 13.71$; MS (70 eV): m/z (%): 475 (3.28) [$\text{M}^+ + 1 - \text{COOEt} - \text{CH}_3$], 91 (100); IR (neat): $\nu = 1717, 1623, 1587, 1413, 1198\text{ cm}^{-1}$; Anal. calcd for $\text{C}_{30}\text{H}_{30}\text{N}_2\text{O}_7\text{S}$ (%): C 64.04, H 5.37, N 4.98; Found C 63.83, H 5.54, N 5.21.

jn-11-162-1

1aa (3R, 5R)

jn-11-162-1

Archive directory: /export/home/masm/vnmr/sys/data
Sample directory:
File: CARBON

Pulse Sequence: s2pu1

jn-13-10-1

1ab (3R, 5R)

jn-13-10-2

2213.2
2198.5
2189.4
2181.3
2145.7
2137.5

1679.4
1637.1
1631.4
1590.8
1588.1
1575.7
1571.2
1542.7
1530.6

1252.8
1245.9
1242.3
1235.5
1235.0
1207.6
1204.4

932.3
927.9
916.7

703.5
684.9

624.5

526.0
522.8

362.7
342.7
335.5
328.4

Jn-13-10-2

Archive directory: /export/home/masm/vnmr/sys/data
Sample directory:
File: CARBON

Pulse Sequence: s2pu1

1ab (3R, 5S)

jn-11-195-1

jn-13-11-2

jn-13-16-1

1ad (3R, 5R)

Jn-13-16-1
STANDARD 1H OBSERVE

Archive directory: /export/home/ASM/vnmrsys/data
Sample directory: auto_140oct2003

Pulse Sequence: s2pu1

jn-13-14-2

1ad (3R, 5S)

jn-13-12-1

1ae (3R, 5R)

2231.5
2223.3
2212.8
2208.1
2206.9
2193.7
2189.5
2185.6
2175.6
2174.6
2155.9
2147.6

1683.7	1600.8	1586.7	1575.0	1570.5	1563.1	1555.7	1531.4	1519.7
--------	--------	--------	--------	--------	--------	--------	--------	--------

915.6	
906.1	
901.6	
893.0	

693.1
680.2

6293

526.1

377.3
342.1
334.9
327.8

24.5

jn-13-13-1

1af (3R, 5R)

jn-13-13-2

1af (3R, 5S)

lag (3R, 5R)

Jn-13-17-2
STANDARD 1H OBSERVE

Archive directory: /export/home/ASM/vnmrsys/data
Sample directory: auto 14063000

1ag (3R, 5R)

jn-13-17-3

1ag (3R, 5S)

1ah (3R, 5R)

jn-13-42-2

jn-13-43-3

1ai (3R, 5S)

JN-13-48-1

Archive directory: /export/home/ASM/vnmr/sys/data
 Sample directory:
 File: CARBON

Pulse Sequence: s2pu1

1aj (3R, 5R)