Supplementary Material For:

Cyclic Oilgofurans: One Pot Synthesis of 30π and 40π Expanded Porphyrinoids

J. Sreedhar Reddy and Venkataramanarao G. Anand

Photosciences & Photonics Unit,

Chemical Sciences & Technology Division,

Regional Research Laboratory (CSIR),

Trivandrum 695 019,

India

Sukhendu Mandal

Solid State & Structural Chemistry Unit,

Indian Institute of Science,

Bangalore 560 012,

India

Contents:

S1 - General notes and procedures

S2 - Synthesis and spectral details of **1** and **2**

S3 - FAB Mass spectrum of **1** and **2**

S4 - ^1H NMR spectrum of **1** and **2**

S5 – Variable temperature ^1H NMR of **1**

S6 - Variable temperature ^1H NMR of **2**

S7 – Cyclic Voltammograms of **1** and **2**

General notes and procedure:

^1H NMR spectra were recorded on a 300 MHz Bruker Advance DPX spectrometer. Chemical shifts (δ) are reported in parts per million (ppm) relative to residual solvent (CHCl_3 , s, δ , 7.26). Mass spectroscopic analysis was carried out on JEOL JMS 600H spectrometer. Electronic spectra were recorded on a Perkin-Elmer Lambda 20 spectrophotometer. Chromatographic separations were performed on basic alumina and silica gel (230-400) in glass columns.

Single Crystal Structure Determination:

A black colored single crystal of **I** was carefully selected and glued to a thin glass fiber. The single crystal X-ray diffraction data were collected on a Bruker AXS Smart Apex CCD diffractometer at 293(2) K. The X-ray generator was operated at 50kV and 35mA using $\text{MoK}\alpha$ ($\lambda = 0.71073\text{\AA}$) radiation. Data were collected with ω scans of width 0.3° . A total of 606 frames were collected in three different settings of φ (0° , 90° , 180°) keeping the sample-to-detector distance fixed at 6.03cm and the detector position (2θ) fixed at -25° .

The data were reduced using SAINTPLUS^[1] and an empirical absorption correction was applied using the SADABS program.^[2] The crystal structure was solved and refined using SHELX-97 present in the WinGx suit of program (Version 1.63.04a).^[3] All the hydrogen atoms were located in the difference Fourier map and for the final refinement the hydrogen atoms were placed in geometrically ideal positions and refined using the riding mode. The last cycles of refinements included atomic positions, anisotropic thermal parameters for all the non-hydrogen atoms and isotropic thermal parameters for all the hydrogen atoms. Full-matrix-least-squares structure refinement against $|F|^2$ was carried out using the WINGX^[4] package of programs.

Observation of several electron density peaks in the difference Fourier map indicated disordered solvent molecules. These could not be resolved and were therefore treated using a continuous solvent area model with the SQUEEZE/BYPASS⁵ procedure.

References:

1. SMART (V 5.628), SAINT (V 6.45a), XPREP, SHELXTL, Bruker AXS Inc. Madison, Wisconsin, USA, **2004**.
2. G.M. Sheldrick, *Siemens Area Correction Absorption Correction Program*. University of Göttingen, Göttingen, Germany, **1994**.
3. G.M. Sheldrick, *SHELXL-97 Program for Crystal Structure Solution and Refinement*. University of Göttingen, Göttingen, Germany, **1997**.
4. J.L. Farrugia, *J. App. Crystallogr*, **1999**, 32, 837 - 838.
5. P. Van der Sluis, A.L. Spek, *Acta Crystallogr., Sect. A* **1990**, 46, 194

Synthesis and spectral data for **1** and **2**:

Furan and pentafluoro benzaldehyde were purchased from Sigma-Aldrich and used as such. Synthesis was carried out under an inert atmosphere using standard Schlenk line techniques. Dry CH₂Cl₂ was used throughout. All other reagents were used as received unless otherwise specified.

Synthetic Procedure: A solution of pentafluoro benzaldehyde (1.2 ml, 10 mmol) and furan (0.73ml, 10 mmol) in 100 ml of dry dichloromethane was placed in 250 ml flask under nitrogen. BF₃.OEt₂ (1.3 ml, 10 mmol) was added under dark, and the resulting solution was stirred for 1h. After adding FeCl₃ (8.10 g, 50 mmol), solution was opened to air and stirred for two more hours. The reaction mixture was washed with water and passed through a short alumina column. This mixture was separated by silica gel column chromatography by using CH₂Cl₂/n-hexane as eluant. Two different colored fractions- light blue (first) and pink color (last) fractions were obtained. Repeated purification of pink fraction by silica gel column chromatography, with 5% CH₂Cl₂ in n-hexane as eluant, yielded two different fractions, cyclo-hexa furans **1** (0.045 g, 2%) and cyclo-octa furans **2** (0.020 g, 1%) respectively.

Cyclco-Hexa Furan (1): ¹H NMR δ (300 MHz, CDCl₃, 298K): 7.85 (s, 6H,), 2.5(s, 6H).
UV- Vis λ_{max}(CH₂Cl₂)/nm (ε): 551(138800), 663(17700). **FABMS** m/z: 1471 (100.0%, M+1)

Cyclco-Octa Furan (2): ¹H NMR δ (300 MHz, CDCl₃, 298K): 9.4 (s, 8H), 5.85(s, 8H).
UV- Vis (CH₂Cl₂). λ_{max}(CH₂Cl₂)/nm (ε):498 (174600), 535(344400),. **FABMS** m/z 1962(100.0%, M+1)

FAB Mass spectrum of 1 and 2:

FAB Mass spectrum of 1

FAB Mass spectrum of 2

^1H NMR Spectrum of **1** and **2** at Room Temperature:

^1H NMR spectrum of **1**

^1H NMR spectrum of **2**

Variable temperature ^1H NMR of **1**:

Variable temperature ^1H NMR of **2**:

Cyclic Voltammogram of **1** and **2**:

Cyclic Voltammogram of **1** and **2** in CH_2Cl_2 containing TBAPF_6 (0.01M) recorded at 100mVs^{-1} . The concentrations of **1** and **2** were $\sim 10^{-3}\text{M}$