Supporting Information

Cyclocarbopalladation of Alkynes. A Stereoselective Method for Preparing Tetra-Substituted Dibenzoxapine Derivatives

Hannah Yu,* Rachel N. Richey, Matthew W. Carson, and Michael J. Coghlan

Chemical Product Research and Development, Lilly Research Laboratories, Eli Lilly and Company, Indianapolis, Indiana 46285, and Discovery Chemistry Research, Lilly Research Laboratories, Eli Lilly and Company, Indianapolis, Indiana 46285

General Methods.

Commercially available reagents and solvents were used without further purification. All reactions were performed under a dry nitrogen atmosphere unless specified otherwise. Reaction progress was monitored by thin-layer chromatography (TLC) or HPLC analysis. TLC analyses were performed with 0.25 mm silica gel plates containing F-254 indicator and visualized using UV light (254 nm). An HPLC gradient method was applied using a 30 min gradient of 40% acetonitrile 0.1% aqueous trifluoroacetic acid/60% 0.1% aqueous trifluoroacetic acid to 70% acetonitrile 0.1% aqueous trifluoroacetic acid/30% 0.1% aqueous trifluoroacetic acid; flow rate 1.0 mL/min; column Zorbax SB-C8, 4.6 mm $\times$ 250 mm, 5 μ m; 254 nm detection. Flash column chromatography was performed with 32-63 μ m silica gel packing.

Melting and boiling points are uncorrected. ^1H NMR and ^{13}C NMR spectra were recorded as specified for each experiment with chemical shift recorded as parts per million. Elemental analysis and high-resolution mass spectrometry data was provided by the Physical Chemistry group of Lilly Research Laboratories.

General procedure for the preparation of *o*-ethynylbenzyl alcohols (4a-4c).

To a 250 mL 3 neck RBF was added 2-iodobenzyl alcohol (34.4 mmol), copper (I) iodide (6.87 mmol), bis(triphenylphosphine)palladium (II) chloride (3.44 mmol) and diethylamine (115 mL). An excess of propyne was bubbled in for 5 minutes and the reaction was stirred at room temperature under nitrogen for 12 h. The solvent was removed *in vacuo* to a black solid. The solid was dissolved in methylene chloride (500

* To whom correspondence should be addressed. Email: yu_hannah@lilly.com

mL) and water (400 mL) was added. The layers were separated and the organic layer was dried (MgSO₄), filtered through celite and concentrated.

5-Methoxy-2-(prop-1-ynyl)benzyl alcohol (4b). The product was purified by trituration from ethyl acetate/hexane (4:1) to give an off-white solid (83% yield): mp 74 – 75 °C; ¹H NMR (500 MHz, CDCl₃) δ 7.30 (d, *J* = 8.0 Hz, 1 H), 6.94 (d, *J* = 2.5 Hz, 1 H), 6.73 (m, 1 H), 4.74 (s, 2 H), 3.79 (s, 3 H), 2.41 (bs, 1 H), 2.06 (s, 3 H); ¹³C (500 MHz, CDCl₃) δ 159.3, 144.2, 133.4, 114.0, 113.0, 112.5, 89.0, 77.0, 64.0, 55.3, 4.4; IR (KBr, cm⁻¹) ν 3312, 3008, 2965, 2910, 2832, 1898, 1608; MS (EI) *m/z* (rel intensity) 177 (*M*⁺, 100); HRMS (ES⁺) exact mass calcd for C₁₁H₁₂O₂ 177.0915, found 177.0913. Anal. Calcd for C₁₁H₁₂O₂: C, 74.98; H, 6.86. Found: C, 74.54, H, 6.88.

5-Fluoro-2-(prop-1-ynyl)benzyl alcohol (4c). The product was isolated without any further purification to give a brown solid (99% yield): ¹H NMR (500 MHz, CDCl₃) δ 7.35 (m, 1 H), 7.14 (dd, *J* = 9.5 Hz, *J* = 2.5 Hz, 1 H), 6.90 (dt, *J* = 9.0 Hz, *J* = 3.0 Hz, 1 H), 4.78 (s, 2 H), 2.07 (s, 3 H); ¹³C (500 MHz, CDCl₃) δ 163.2, 161.3, 145.2, 145.1, 133.8, 117.7, 114.3, 114.1, 113.9, 90.4, 76.2, 63.5, 4.4; IR (KBr, cm⁻¹) ν 3300, 2917, 1609, 1583, 1488, 1445, 1422, 1357; MS (EI) *m/z* (rel intensity) 164 (*M*⁺, 100); HRMS (ES⁺) exact mass calcd for C₁₀H₆FO 164.0637, found 164.0648.

General procedure for the preparation of 2-(prop-1-ynyl)-benzyl-2'-iodophenyl ethers (3a-3e).

To a mixture of benzyl alcohol (51.4 mmol), phenol (51.4 mmol) and triphenylphosphine (77.0 mmol) in 260 mL of THF was added diisopropyl azodicarboxylate (77.2 mmol) dropwise at 0 °C. The reaction mixture was warmed up to rt and was stirred at rt for 12 h. 300 mL of water was added followed by 300 mL of EtOAc. The layers were separated and the aqueous layer was extracted with additional 300 mL of EtOAc. The combined organic layers were washed with brine (300 mL), dried (Na₂SO₄) and concentrated.

2-(Prop-1-ynyl)-benzyl-2'-iodophenyl ether (3a). The crude reaction mixture was purified by silica gel column chromatography (20:1 hexanes/ethyl acetate) to give a white solid (85% yield): *R_f* (hexanes/ethyl acetate = 5:1) = 0.58; ¹H NMR (500 MHz,

CDCl₃) δ 7.81 (dd, $J = 7.5$ Hz, $J = 1.5$ Hz, 1 H), 7.68 (d, $J = 7.5$ Hz, 1 H), 7.43 (d, $J = 7.5$ Hz, 1 H), 7.34 – 7.25 (m, 3 H), 6.89 (d, $J = 4.5$ Hz, 1 H), 6.73 (m, 1 H), 5.32 (s, 2 H), 2.12 (s, 3 H); ¹³C (500 MHz, CDCl₃) δ 157.2, 139.5, 138.1, 132.0, 129.5, 128.0, 127.4, 126.8, 122.8, 121.7, 112.8, 91.5, 86.7, 69.1, 4.59; IR (KBr, cm⁻¹) ν 3060, 2914, 2843, 1581, 1569, 1486, 1475, 1437, 1374; MS (EI) m/z (rel intensity) 349 (M⁺, 100); HRMS (ES⁺) exact mass calcd for C₁₆H₁₃IO 349.0089, found 349.0111. An analytical sample was prepared by crystallization from 1:4 THF/heptane: mp 81 – 83 °C. Anal. Calcd for C₁₆H₁₃IO: C, 55.19; H, 3.76. Found: C, 55.34, H, 3.87.

5-Methoxy-2-(Prop-1-ynyl)-benzyl-2'-iodophenyl ether (3b). The reaction mixture was purified by silica gel column chromatography (50:1 hexanes/ethyl acetate) to give a pale yellow oil (76% yield): R_f (hexanes/ethyl acetate = 20:1) = 0.49; ¹H NMR (500 MHz, CDCl₃) δ 7.81 (dd, $J = 8.0$ Hz, $J = 2.0$ Hz, 1 H), 7.35 – 7.28 (m, 3 H), 6.90 (dd, $J = 8.0$ Hz, $J = 1.0$ Hz, 1 H), 6.79 – 6.72 (m, 2 H), 5.29 (s, 2 H), 3.84 (s, 3 H), 2.10 (s, 3 H); ¹³C (500 MHz, CDCl₃) δ 159.4, 157.0, 139.9, 139.5, 133.2, 129.5, 122.8, 113.6, 113.5, 112.7, 111.8, 89.7, 86.7, 76.7, 68.8, 55.4, 4.5; IR (KBr, cm⁻¹) ν 3060, 3001, 2912, 2835, 1607, 1570, 1500, 1475. MS (EI) m/z (rel intensity) 379 (M⁺, 100); HRMS (ES⁺) exact mass calcd for C₁₇H₁₅IO₂ 379.0195, found 379.0171.

2-(Prop-1-ynyl)-benzyl-5'-fluoro-2'-iodophenyl ether (3c). The reaction mixture was purified by silica gel column chromatography (20:1 hexanes/ethyl acetate) to give a white solid (74% yield): R_f (hexanes/ethyl acetate = 5:1) = 0.59; ¹H NMR (500 MHz, CDCl₃) δ 7.74 – 7.71 (m, 1 H), 7.64 (d, $J = 7.5$ Hz, 1 H), 7.43 (d, $J = 7.5$ Hz, 1 H), 7.35 – 7.32 (m, 1 H), 7.27 – 7.24 (m, 1 H), 6.66 (dd, $J = 11.0$ Hz, $J = 3.0$ Hz, 1 H), 6.53 – 6.50 (m, 1 H), 5.28 (s, 2 H), 2.12 (s, 3 H); ¹³C (500 MHz, CDCl₃) δ 164.9, 162.9, 158.3, 158.2, 139.6, 139.5, 137.3, 132.1, 128.0, 127.6, 126.8, 121.9, 109.7, 109.5, 101.3, 101.1, 91.7, 79.5, 76.9, 69.4, 69.3, 4.6; IR (KBr, cm⁻¹) ν 3437, 3062, 2913, 2248, 1869, 1606, 1580; MS (EI) m/z (rel intensity) 366 (M⁺, 100); HRMS (ES⁺) exact mass calcd for C₁₆H₁₂FIO 366.9995, found 366.9985. An analytical sample was prepared by crystallization from hexanes: mp 86 – 88 °C. Anal. Calcd for C₁₆H₁₂FIO: C, 52.48; H, 3.30. Found: C, 52.27, H, 3.28.

5-Fluoro-2-(prop-1-ynyl)-benzyl-5'-fluoro-2'-iodophenyl ether (3d). The reaction mixture was purified by silica gel column chromatography (50:1 hexanes/ethyl acetate) to give a white solid (66% yield): R_f (hexanes/ethyl acetate = 20:1) = 0.48; $^1\text{H NMR}$ (500 MHz, CDCl_3) δ 7.75 – 7.72 (m, 1 H), 7.44 – 7.38 (m, 2 H), 6.97 – 6.93 (m, 1 H), 6.62 (dd, J = 11.0 Hz, J = 3.0 Hz, 1 H), 6.56 – 6.52 (m, 1 H), 5.23 (s, 2 H), 2.11 (s, 3 H); ^{13}C (500 MHz, CDCl_3) δ 164.8, 163.3, 162.9, 161.4, 157.9, 157.8, 140.1, 139.7, 139.6, 133.8, 117.4, 114.8, 114.6, 114.1, 114.0, 113.9, 109.9, 109.8, 101.2, 101.0, 91.4, 79.4, 75.8, 68.7, 4.5; IR (KBr, cm^{-1}) ν 2447, 3101, 3070, 2910, 2844, 1875, 1839, 1599; MS (EI) m/z (rel intensity) 385 (M^+ , 100); HRMS (ES+) exact mass calcd for $\text{C}_{16}\text{H}_{11}\text{F}_2\text{IO}$ 384.9901, found 384.9913. An analytical sample was prepared by crystallization from 1:1 ethyl acetate/hexanes: mp 132 – 134 °C. Anal. Calcd for $\text{C}_{16}\text{H}_{11}\text{F}_2\text{IO}$: C, 50.02; H, 2.89. Found: C, 49.88, H, 2.93.

2-Iodobenzyl-2'-(prop-1-ynyl)-phenyl ether (3e). The reaction mixture was purified by silica gel column chromatography (20:1 hexanes/methylene chloride) to give a clear oil (77% yield): R_f (hexanes/methylene chloride = 5:1) = 0.39; $^1\text{H NMR}$ (500 MHz, CDCl_3) δ 7.89 (d, J = 8.0, 1 H), 7.68 (d, J = 7.5 Hz, 1 H), 7.46 – 7.40 (m, 2 H), 7.25 (m, 1 H), 7.06 (m, 1 H), 6.96 (t, J = 7.5 Hz, 1 H), 6.90 (d, J = 8.5 Hz, 1 H), 5.16 (s, 2 H), 2.18 (s, 3 H); ^{13}C (400 MHz, CDCl_3) δ 158.7, 139.2, 139.0, 133.6, 129.2, 128.9, 128.4, 128.2, 121.2, 114.1, 112.9, 96.2, 90.3, 75.9, 74.4, 4.8; IR (KBr, cm^{-1}) ν 3062, 2906, 2844, 1599, 1575. MS (EI) m/z (rel intensity) 349 (M^+ , 100); HRMS (ES+) exact mass calcd for $\text{C}_{16}\text{H}_{13}\text{IO}$ 349.0089, found 349.0104.

General procedure for the preparation of 3-[1-(6,11-dibenzol[*b,e*]oxepino-11-ylidene)]-ethyl-nitrobenzenes (2a-2e, 8a-8f).

To a mixture of iodo alkyne (0.72 mmol), 3-nitrophenylboronic acid (0.86 mmol), sodium carbonate (2.15 mmol) was added $\text{Pd}(\text{OAc})_2$ (0.0144 mmol in 5.6 mL of dioxane) followed by water (1.4 mL). The reaction mixture was heated at 70 – 100 °C under nitrogen for 12 h. The reaction mixture was filtered through Celite. The filtrate was diluted with 10 mL of water and 10 mL of CH_2Cl_2 and the layers were separated. The aqueous layer was extracted with CH_2Cl_2 . The combined organic layers were washed with brine and dried (MgSO_4). The solvent was removed *in vacuo*.

(Z)-3-[1-(6,11-Dibenzol*b,e*]oxepino-11-ylidene]-ethyl-nitrobenzene (2a). The

reaction mixture was purified by silica gel column chromatography (30:1 hexanes/ethyl acetate) to give an off-white solid (89% yield): *R_f* (hexanes/ethyl acetate = 10:1) = 0.48; mp 169 – 170 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.21 (t, *J* = 2.1 Hz, 1 H), 8.03 – 7.99 (m, 1 H), 7.59 – 7.56 (m, 1 H), 7.43 – 7.31 (m, 5 H), 6.96 – 6.93 (m, 1 H), 6.75 – 6.72 (m, 1 H), 6.53 – 6.43 (m, 2 H), 5.87 (d, *J* = 12.0 Hz, 1 H), 4.91 (d, *J* = 12.0 Hz, 1 H), 2.16 (s, 3 H); ¹³C (300 MHz, DMSO-*d*₆) δ 154.6, 147.5, 144.5, 142.8, 136.0, 135.7, 133.7, 132.9, 132.3, 129.5, 128.9, 128.7, 128.6, 127.8, 126.9, 124.8, 123.7, 121.4, 119.9, 118.8, 68.7, 21.4; IR (KBr, cm⁻¹) ν 3065, 2916, 2880, 1523, 1476, 1347, 1106; MS (EI) *m/z* (rel intensity) 344 (M⁺, 100); HRMS (ES⁺) exact mass calcd for C₂₂H₁₇NO₃ 344.1287, found 344.1308. Anal. calcd for C₂₂H₁₇NO₃ C, 76.95, H, 4.99, N, 4.08, found C, 76.84, H, 4.67, N, 4.01.

(Z)-3-[1-(8-Methoxy-6,11-dibenzol*b,e*]oxepino-11-ylidene]-ethyl-nitrobenzene (2b).

The reaction mixture was purified by silica gel column chromatography (9:1 hexanes/methylene chloride) to give a yellow solid (70% yield): *R_f* (hexanes/methylene chloride = 1:1) = 0.74; mp 152 – 154 °C; ¹H NMR (300 MHz, CDCl₃) δ 8.20 (m, 1 H), 8.02 – 7.98 (m, 1 H), 7.56 (d, *J* = 7.8 Hz, 1 H), 7.35 – 7.26 (m, 2 H), 7.98 – 6.91 (m, 3 H), 6.76-6.73 (m, 1 H), 6.53 – 6.43 (m, 2 H), 5.84 (d, *J* = 12.5 Hz, 1 H), 4.87 (d, *J* = 12.5 Hz, 1 H), 3.84 (s, 3 H), 2.16 (s, 3 H); ¹³C (300 MHz, CDCl₃) δ 159.2, 155.0, 148.1, 145.0, 136.8, 135.8, 135.3, 134.2, 133.3, 132.4, 128.9, 128.8, 128.4, 125.7, 123.9, 121.4, 120.3, 119.3, 114.0, 113.9, 69.8, 55.4, 21.7; IR (KBr, cm⁻¹) ν 3432, 3008, 2957, 2871, 2825, 1723, 1618, 1525, 1474; MS (EI) *m/z* (rel intensity) 374 (M⁺, 100); HRMS (ES⁺) exact mass calcd for C₂₃H₁₉NO₄ 374.1392, found 374.1408. Anal. calcd for C₂₃H₁₉NO₄ C, 73.98, H, 5.13, N, 3.75, found C, 73.57, H, 5.07, N, 3.82.

(Z)-3-[1-(3-Fluoro-6,11-dibenzol*b,e*]oxepino-11-ylidene]-ethyl-nitrobenzene (2c).

The reaction mixture was purified by silica gel column chromatography (9:1 hexanes/methylene chloride) to give a white solid (77% yield): *R_f* (hexanes/methylene chloride = 1:1) = 0.71; mp 206 – 208 °C; ¹H NMR (500 MHz, CDCl₃) δ 8.21 (s, 1 H), 8.04 (m, 1 H), 7.57 (d, *J* = 7.5 Hz, 1 H), 7.44 – 7.35 (m, 5 H), 6.50 – 6.44 (m, 2 H), 6.16 (dt, *J* = 8.0 Hz, *J* = 3.0 Hz, 1 H), 5.87 (d, *J* = 12.0 Hz, 1 H), 4.92 (d, *J* = 12.0 Hz, 1 H),

2.15 (s, 3 H); ^{13}C (500 MHz, CDCl_3) δ 163.8, 161.8, 156.2, 148.2, 144.6, 143.0, 136.2, 135.7, 133.7, 133.6, 132.6, 129.0, 128.7, 128.0, 127.1, 123.9, 121.6, 120.9, 107.7, 107.6, 106.2, 106.0, 69.7, 21.7; IR (KBr, cm^{-1}) ν 3437, 3083, 2977, 2926, 2882, 1609, 1579; MS (EI) m/z (rel intensity) 362 (M^+ , 100); HRMS (ES+) exact mass calcd for $\text{C}_{22}\text{H}_{16}\text{FNO}_3$ 362.1192, found 362.1187. Anal. calcd for $\text{C}_{22}\text{H}_{16}\text{FNO}_3$ C, 73.12, H, 4.46, N, 3.88, found C, 72.91, H, 4.53, N, 3.94.

(Z)-3-[1-(3,8-Difluoro-6,11-dibenzo[*b,e*]oxepino-11-ylidene)-ethyl-nitrobenzene (2d).

The reaction mixture was purified by silica gel column chromatography (9:1 hexanes/methylene chloride) to give a white solid (73% yield): R_f (hexanes/methylene chloride = 1:1) = 0.64; mp 200 – 202 °C; ^1H NMR (500 MHz, $\text{DMSO-}d_6$) δ 8.17 (t, J = 2.5 Hz, 1 H), 8.04 (m, 1 H), 7.74 (m, 1 H), 7.55 – 7.44 (m, 3 H), 7.29 (t, J = 3.0 Hz, 1 H), 6.56 (m, 2 H), 6.31 (t, J = 2.5 Hz, 1 H), 5.84 (d, J = 13.0 Hz, 1 H), 5.01 (d, J = 13.0 Hz, 1 H), 3.30 (3 H), 2.07 (s, 3 H); ^{13}C (500 MHz, $\text{DMSO-}d_6$) δ 163.3, 162.8, 161.4, 160.9, 156.2, 156.1, 148.0, 144.7, 139.5, 136.5, 135.4, 135.3, 135.2, 134.4, 133.9, 130.1, 129.5, 124.1, 122.0, 121.7, 121.8, 116.3, 116.1, 115.9, 107.9, 107.7, 106.0, 105.8, 68.7, 21.9; IR (KBr, cm^{-1}) ν 3431, 3076, 2970, 2893, 1611, 1581; MS (EI) m/z (rel intensity) 380 (M^+ , 100); HRMS (ES+) exact mass calcd for $\text{C}_{22}\text{H}_{15}\text{F}_2\text{NO}_3$ 380.1098, found 380.1114. Anal. calcd for $\text{C}_{22}\text{H}_{15}\text{F}_2\text{NO}_3$ C, 69.65, H, 3.99, N, 3.69, found C, 69.45, H, 3.95, N, 3.73.

(E)-3-[1-(6,11-Dibenzo[*b,e*]oxepino-11-ylidene)-ethyl-nitrobenzene (2e). The reaction mixture was purified by silica gel column chromatography (9:1 hexanes/methylene chloride) to give a clear oil (67% yield): R_f (hexanes/methylene chloride = 1:1) = 0.54; ^1H NMR (500 MHz, CDCl_3) δ 8.01 – 7.96 (m, 2 H), 7.37 – 7.29 (m, 4 H), 7.23 – 7.18 (m, 2 H), 6.89 (m, 3 H), 6.67 (d, J = 7.5 Hz, 1 H), 5.89 (d, J = 12.5 Hz, 1 H), 4.99 (d, J = 12.5 Hz, 1 H), 2.40 (s, 3 H); ^{13}C (500 MHz, CDCl_3) δ 154.9, 147.8, 145.0, 143.2, 137.3, 134.4, 133.2, 132.8, 131.1, 129.6, 128.8, 128.6, 128.2, 127.9, 127.6, 124.9, 123.2, 121.4, 120.2, 119.7, 70.1, 22.0; IR (KBr, cm^{-1}) ν 3069, 3032, 2935, 1607, 1532; MS (EI) m/z (rel intensity) 344 (M^+ , 100); HRMS (ES+) exact mass calcd for $\text{C}_{22}\text{H}_{17}\text{NO}_3$ 344.1287, found 344.1294.

(Z)-1-(6,11-Dibenzo[*b,e*]oxepino-11-ylidene)-ethylbenzene (8a). The reaction mixture was purified by silica gel column chromatography (50:1 hexanes/ethyl acetate) to give a

clear oil (77% yield): R_f (hexanes/ethyl acetate = 20:1) = 0.66; ^1H NMR (500 MHz, CDCl_3) δ 7.42 – 7.14 (m, 9 H), 6.92 (m, 1 H), 6.72 (d, J = 1.0 Hz, 1 H), 6.63 (d, J = 1.5 Hz, 1 H), 6.45 – 6.42 (m, 1 H), 5.93 (d, J = 12.0 Hz, 1 H), 4.89 (d, J = 12.0 Hz, 1 H), 2.13 (s, 3 H); ^{13}C (500 MHz, CDCl_3) δ 154.8, 143.8, 143.0, 135.8, 134.6, 133.2, 132.8, 129.3, 128.6, 128.4, 128.3, 128.0, 127.5, 127.4, 126.4, 125.9, 119.9, 118.8, 69.5, 22.0; IR (KBr, cm^{-1}) ν 3019, 1603, 1478, 1439; MS (EI) m/z (rel intensity) 299 (M^+ , 100); HRMS (ES^+) exact mass calcd for $\text{C}_{22}\text{H}_{18}\text{O}$ 299.1436, found 299.1459.

(Z)-3-[1-(6,11-Dibenzol*b*,*e*]oxepino-11-ylidene)]-ethyl-methylbenzene (8b).** The reaction mixture was purified by silica gel column chromatography (9:1 hexanes/methylene chloride) to give a clear oil (91% yield): R_f (hexanes/methylene chloride = 1:1) = 0.65; ^1H NMR (500 MHz, CDCl_3) δ 7.41 – 7.29 (m, 4 H), 7.13 – 7.08 (m, 3 H), 6.97 – 6.91 (m, 2 H), 6.70 (dd, J = 8.0 Hz, J = 1.0 Hz, 1 H), 6.64 (dd, J = 7.5 Hz, J = 1.5 Hz, 1 H), 6.46 – 6.43 (m, 1 H), 5.92 (d, J = 12.5 Hz, 1 H), 4.88 (d, J = 12.5 Hz, 1 H), 2.28 (s, 3 H), 2.10 (s, 3 H); ^{13}C (500 MHz, CDCl_3) δ 154.7, 143.9, 142.9, 137.4, 135.9, 134.4, 133.2, 132.8, 129.8, 128.5, 128.4, 128.3, 127.8, 127.5, 127.4, 127.1, 126.4, 126.0, 119.9, 118.7, 69.5, 22.0, 21.4; IR (KBr, cm^{-1}) ν 3429, 3058, 3008, 2926, 2871; MS (EI) m/z (rel intensity) 313 (M^+ , 100); HRMS (ES^+) exact mass calcd for $\text{C}_{23}\text{H}_{20}\text{O}$ 313.1592, found 313.1613.**

(Z)-4-[1-(6,11-Dibenzol*b*,*e*]oxepino-11-ylidene)]-ethyl-fluorobenzene (8c).** The reaction mixture was purified by silica gel column chromatography (9:1 hexanes/methylene chloride) to give a clear oil (76% yield): R_f (hexanes/methylene chloride = 1:1) = 0.59; ^1H NMR (500 MHz, CDCl_3) δ 7.43 – 7.28 (m, 6 H), 6.98 – 6.93 (m, 3 H), 6.74 (dd, J = 7.5 Hz, J = 1.0 Hz, 1 H), 6.62 (dd, J = 7.5 Hz, J = 1.0 Hz, 1 H), 6.5 (m, 1 H), 5.91 (d, J = 13.0 Hz, 1 H), 4.90 (d, J = 13.0 Hz, 1 H), 2.12 (s, 3 H); ^{13}C (500 MHz, CDCl_3) δ 162.3, 160.4, 154.8, 143.7, 138.9, 135.0, 134.7, 133.1, 132.7, 130.9, 130.8, 128.6, 128.5, 128.4, 127.6, 127.4, 125.8, 120.1, 119.0, 115.0, 114.8, 69.5, 22.0; IR (KBr, cm^{-1}) ν 3066, 2973, 2918, 2860, 1595; MS (EI) m/z (rel intensity) 317 (M^+ , 100); HRMS (ES^+) exact mass calcd for $\text{C}_{22}\text{H}_{17}\text{FO}$ 317.1342, found 317.1365.**

(Z)-3-[1-(6,11-Dibenzol*b*,*e*]oxepino-11-ylidene)]-ethyl-trifluoromethyl benzene (8d).** The reaction mixture was purified by silica gel column chromatography (9:1**

hexanes/methylene chloride) to give a white solid (70% yield): R_f (hexanes/methylene chloride = 1:1) = 0.55; mp 102 – 104 °C; ^1H NMR (500 MHz, CDCl_3) δ 7.66 (s, 1 H), 7.52 – 7.32 (m, 7 H), 7.02 – 6.98 (m, 1 H), 6.81 – 6.79 (m, 1 H), 6.62 – 6.60 (m, 1 H), 6.51 – 6.48 (m, 1 H), 5.95 (d, J = 12.0 Hz, 1 H), 4.95 (d, J = 12.0 Hz, 1 H), 2.19 (s, 3 H); ^{13}C (500 MHz, CDCl_3) δ 155.0, 143.8, 143.4, 136.3, 134.2, 133.1, 132.8, 130.6, 130.8, 130.6, 130.3, 130.1, 128.9, 128.7, 128.5, 128.4, 127.8, 127.4, 127.3, 126.0, 125.9, 125.4, 125.3, 123.2, 123.2, 123.1, 120.2, 119.2, 69.6, 21.7; IR (KBr, cm^{-1}) ν 3432, 3070, 2980, 2918, 2879, 1599, 1567; MS (EI) m/z (rel intensity) 367 (M^+ , 100); HRMS (ES+) exact mass calcd for $\text{C}_{23}\text{H}_{17}\text{F}_3\text{O}$ 367.1310, found 367.1292. Anal. calcd for $\text{C}_{23}\text{H}_{17}\text{F}_3\text{O}$ C, 75.40, H, 4.68, found C, 75.01, H, 4.65.

(Z)-4-[1-(6,11-Dibenzo[*b,e*]oxepino-11-ylidene)]-ethyl-methoxybenzene (8e). The reaction mixture was purified by silica gel column chromatography (4:1 hexanes/methylene chloride) to give a white form (74% yield): R_f (hexanes/methylene chloride = 1:4) = 0.60; ^1H NMR (500 MHz, CDCl_3) δ 7.41 – 7.23 (m, 6 H), 6.93 (dt, J = 7.5 Hz, J = 2.0 Hz, 1 H), 6.77 – 6.64 (m, 4 H), 6.46 (dt, J = 8.5 Hz, J = 1.0 Hz, 1 H), 5.91 (d, J = 12.5 Hz, 1 H), 4.87 (d, J = 12.5 Hz, 1 H), 3.77 (s, 3 H), 2.10 (s, 3 H); ^{13}C (500 MHz, CDCl_3) δ 158.0, 154.8, 144.1, 135.2, 135.1, 133.9, 133.2, 132.9, 130.4, 128.6, 128.4, 128.3, 127.6, 127.4, 126.2, 120.1, 118.8, 113.4, 69.5, 55.1, 22.0; IR (KBr, cm^{-1}) ν 3432, 2992, 2953, 2930, 2829, 1610; MS (EI) m/z (rel intensity) 329 (M^+ , 100); HRMS (ES+) exact mass calcd for $\text{C}_{23}\text{H}_{20}\text{O}_2$ 329.1541, found 329.1516.

(Z)-3,4-Dichloro-[1-(6,11-dibenzo[*b,e*]oxepino-11-ylidene)]-ethylbenzene (8f). The reaction mixture was purified by silica gel column chromatography (20:1 hexanes/methylene chloride) to give a white solid (78% yield): R_f (hexanes/methylene chloride = 1:1) = 0.71; mp 153 – 155 °C; ^1H NMR (500 MHz, CDCl_3) δ 7.48 – 7.33 (m, 5 H), 7.27 (dd, J = 8.5 Hz, J = 2.5 Hz, 1 H), 7.15 – 7.12 (m, 1 H), 7.03 – 6.00 (m, 1 H), 6.79 – 6.76 (m, 1 H), 6.66 – 6.63 (m, 1 H), 6.54 (t, J = 7.0 Hz, 1 H), 5.91 (d, J = 12.5 Hz, 1 H), 4.92 (d, J = 12.5 Hz, 1 H), 2.12 (s, 3 H); ^{13}C (500 MHz, CDCl_3) δ 154.9, 143.2, 143.1, 136.5, 133.2, 133.0, 132.5, 132.1, 131.0, 130.9, 130.4, 130.0, 129.0, 128.7, 128.5, 127.8, 127.3, 125.2, 120.4, 119.2, 69.5, 21.7; IR (KBr, cm^{-1}) ν 3428, 3062, 3023, 2984, 2938, 2875, 2731, 1603; MS (EI) m/z (rel intensity) 367 (M^+ , 100); HRMS (ES+) exact

mass calcd for $C_{22}H_{16}Cl_2O$ 367.0656, found 367.0647. Anal. calcd for $C_{22}H_{16}Cl_2O$ C, 71.95, H, 4.39, found C, 71.84, H, 4.57.

(Z)-3-[1-(6,11-Dibenzol*b,e*]oxepino-11-ylidene)]-ethyl-methylsulfonylaminobenzene (1a).

To a hydrogenation vessel was added dibenzoxapine **2a** (9.71 g, 28.3 mmol), triethylamine (2.86 g, 28.3 mmol), platinum (5% wt. on carbon, sulfided, 970 mg) and THF (120 mL). The vessel was then vacuumed and charged with hydrogen three times. The reaction mixture was shaken under hydrogen at 50 psi for 2 hours. The reaction mixture was filtered through celite and rinsed with ethyl acetate. The filtrate was dried (Na_2SO_4), filtered and concentrated to give the aniline as a off-white solid (8.86 g, quantitative yield) that was used without further purification.

To a mixture of the above aniline (8.86 g, 28.3 mmol) in CH_2Cl_2 (120 mL) was added pyridine (2.46 g, 31.1 mmol) followed by methanesulfonyl chloride (3.47 g, 30.3 mmol) dropwise at rt. The reaction mixture was stirred at rt for 12 h and 150 mL of 0.1 N HCl was added. The layers were separated and the organic layer was washed with water (150 mL) followed by saturated aqueous $NaHCO_3$ (150 mL). The organic layer was dried (Na_2SO_4) and concentrated to a solid, which was further purified by re-crystallizing in 160 mL of ethyl acetate and hexanes (1:1) to give a white solid (9.26 g, 22.4 mmol, 79%): mp 199 – 200 °C; 1H NMR (500 MHz, $CDCl_3$) δ 7.42 – 7.21 (m, 6 H), 7.14 (bs, 1 H), 7.02 (m, 1 H), 6.89 (m, 1 H), 6.70 (d, J = 8.0 Hz, 1 H), 6.61 (m, 1 H), 6.44 (t, J = 7.5 Hz, 1 H), 5.90 (d, J = 12.5 Hz, 1 H), 4.90 (d, J = 12.5 Hz, 1 H), 2.64 (s, 3 H), 2.11 (s, 3 H); ^{13}C (500 MHz, $CDCl_3$) δ 155.0, 144.8, 143.3, 136.4, 135.6, 135.1, 133.0, 132.7, 129.5, 128.7, 128.5, 127.7, 127.3, 126.1, 125.9, 121.9, 119.9, 119.3, 119.1, 69.6, 38.7, 21.7; IR (KBr, cm^{-1}) ν 3440, 3227, 3060, 3024, 2982, 2950, 2925, 2874, 1603, 1583; MS (EI) m/z (rel intensity) 414 (M^+ , 100); HRMS (ES+) exact mass calcd for $C_{22}H_{21}NO_3S$ 414.1140, found 414.1121. Anal. calcd for $C_{23}H_{21}NO_3S$ C, 70.56, H, 5.41, N, 3.58, found C, 70.77, H, 5.31, N, 3.40.

S11

159.308

144.169

133.440

114.044
112.976
112.510

88.993

77.311
77.059
76.998
76.807

64.042

55.297
55.259

4.395

4.774

2.074

157.202

139.545
138.126

131.960
129.465
128.023
127.367
126.817
122.766
121.697

112.770

91.519

86.689

77.296
77.037
76.785

69.124

4.593

5.283

2.115

164.855
162.894
158.285
158.209

139.583
139.499
137.340
132.052
128.030
127.596
126.848
121.865

109.687
109.656
109.511
109.481
101.347
101.133

91.740

79.501
79.470
77.311
77.052
76.907
76.800
69.383
69.337
69.284

4.570

S23

4.778

69.795

55.404

21.670

— 2.275
— 2.102
— 1.547

21.716

21.739

S45