Supporting Information for:

Designing molecules by optimizing potentials

Mingliang Wang^a, Xiangqian Hu^a, David N. Beratan^{*a,b} and Weitao Yang^{*a}

^aDepartment of Chemistry, Duke University, Durham, NC 27708

^bDepartment of Biochemistry, Duke University, Durham, NC 27708

*To whom correspondence should be addressed. Email: david.beratan@duke.edu and weitaoyang@duke.edu

1. Molecular geometry in the LCAP scheme

The molecular fragment approach of building molecules assumes that structure can be assembled approximately from a sum of “standard” parts, molecular fragments. When steric and electronic interactions among fragments have a substantial impact on the property of interest, the optimization scheme can be adjusted to account for these effects.

A. Geometry sensitivity of properties and molecular fragmentation

In optimizing the polarizabilities and hyperpolarizabilities of the 21 small molecules associated with Fig. 2 in the text, we assumed a fixed bond length (1.53 Å) for the bond between the two fragments. This assumption produced favorable property values compared with experiment, and the correct ordering of polarizabilities (Table S1, below). The viability of this fragment approach was subjected to a more challenging test on a family of push-pull polyenes, reasoning that delocalization and charge transfer mediated by a pi-electron bridge define a worst-case for the viability of the fragment approach.

Table S1. Polarizabilities (1 a.u. = 1.4819×10^{-25} esu) and hyperpolarizabilities (1 a.u. = 8.63922×10^{-33} esu) of molecules based on combinations of functional groups.

Structure	Calculated α (10^{-25} esu)	Experimental α (10^{-25} esu)	Calculated β_{μ} (10^{-33} esu)
CH ₃ -CH ₃	45.46	44.80 (5)	0.0
CH ₃ -SH	56.92	--	641.38
^a SH-SH	65.41	--	-628.59

CH ₃ -OH	35.27	33.19 (5)	-646.21
CH ₃ -NH ₂	42.62	40.13 (8)	-589.54
CH ₃ -F	28.51	26.10 (5)	-722.58 ^a
CH ₃ -Cl	45.11	45.30 (5)	-149.63
^a H ₂ O ₂	26.05	22.87 (6)	-423.67
Cl-Cl	45.02	44.98 (7)	0.0
NH ₂ -F	32.57	--	-835.93
NH ₂ -SH	52.37	--	-177.62
HO-NH ₂	32.84	--	-879.47
HO-F	19.98	--	-349.02
HO-Cl	34.71	--	267.47
HO-SH	45.52	--	133.69
F-F	14.70	12.42(7)	0.0
F-Cl	28.48	--	-1093.73
F-SH	38.90	--	-1472.99
Cl-SH	56.52	--	405.01
H ₂ N-NH ₂	39.80	34.63 (8)	-17.62
H ₂ N-Cl	41.73	--	-305.83

^aThe H₂O₂ and H₂S₂ structures are assumed locked in non-centrosymmetric geometries with a torsion angles of 90°. ^b B3LYP/Aug-POL calculation: 728.55×10⁻³³esu from (7)

We examined the first hyperpolarizabilities of the following structures:

The library of donor/acceptor groups is indicated in the Table S2 below

A1

A2

Table S2. HF/STO-3G Optimized bond lengths (Å).

			M1	M2	M3	M4
Donors (X) ->		H	NH ₂	NBu ₂	NEt ₂	NEt ₂
Bond Lengths	a	1.315	1.328	1.329	1.326	1.326
	b	1.484	1.474	1.477	1.475	1.475
	c	1.324	1.326	1.327	1.326	1.326
	d	1.479	1.476	1.476	1.476	1.476

e	1.325	1.326	1.327	1.326	1.326
f	1.478	1.477	1.476	1.476	1.476
g	1.325	1.326	1.327	1.326	1.327
h	1.479	1.477	1.474	1.476	1.476
i	1.324	1.325	1.329	1.328	1.328
j	1.484	1.479	1.470	1.473	1.473
k	1.315	1.322	1.339	1.335	1.335

Acceptors (X1,X2) ->	H,H	H,CHO	CN,CN	A1 (struct. above)	A2 (struct. above)
-------------------------	-----	-------	-------	-----------------------	-----------------------

Table S3. Properties (esu) calculated using HF/6-31+g optimized structures.

Molecule	$ \mu (\times 10^{-18})$		$\beta(0) \cdot \mu (\times 10^{-48})$		$\beta(0) \cdot \mu / \mu (\times 10^{-30})$		$\beta_{tot}(0) (\times 10^{-30})$	
	S1	S2	S1	S2	S1	S2	S1	S2
M1	8.482	8.408	-508.9	-457.4	-60.0	-54.4	63.5	62.2
M2	11.832	11.573	-1231.7	-1219.8	-104.1	-105.4	118.1	111.2
M3	8.106	7.872	-1171.3	-1101.3	-144.5	-139.9	147.3	142.7
M4	8.815	8.533	-1414.8	-1307.3	-160.5	-153.2	163.8	156.6

For S1: the full structure is optimized at the HF/STO-3G level; For S2: the structure is built from fragments based on the separate donor, acceptor and bridge units.

In the molecular constructions (note “S2” in Table S3), the double bond nearest the donor or acceptor unit is included in its fragment. That is, we take the geometry of the four double bonds in the molecule’s center from a $\text{H}(\text{HC}=\text{CH})_6\text{H}$ calculation. Table S2 compares the geometries of the pure polyene and its substituted derivatives. The data indicate that the molecular geometry of the bridge changes little on donor/acceptor substitution. Table S3 similarly shows that hyperpolarizabilities change little when the structure of the full molecule is approximated by uniting the geometries of the fragment parts. The optimizations and comparisons are based on gas phase calculations; hyperpolarizabilities of strongly solvatochromic structures should be treated using a model that includes explicit solvation; the calculations described here apply best to low dielectric media. When such solvent effects are important, the self-consistent procedure below could be used to make corrections arising from more significant structural changes than observed in the two case studies described here.

B. Self-consistent geometry optimization in the LCAP framework

Interactions among fragments will perturb its structure through steric and electronic interactions. An approach to including the influence of electronic interactions (that cause changes in orbital hybridization and chemical bonding, for example) among molecular fragments is to optimize the geometry of the LCAP optimized structure. The geometry optimized structure could then be used to modify the geometry of the input fragments and the LCAP analysis would then be repeated. This procedure could be continued until self-consistency is reached.

C. Inclusion of multiple fragment conformers in the optimization

The self-consistent procedure described above would include electronic interactions among fragments but would not sample conformational space widely. If the molecular property has a strong conformational dependence, the LCAP sum in Eq. (1) of the text can include different conformations of the fragments. This procedure would be analogous to adopting a library of amino acid rotamers, as is often done when modeling protein structure.

D. Summary of hybrid geometry/LCAP optimization schemes

The approaches described above are illustrated in Fig. S1 and could be used to expand the scope of the LCAP optimizations described in the text.

Figure S1. Flow chart for an expanded LCAP-based optimization procedure that includes multiple fragment conformers and self-consistent geometry optimization.

2. Details of the LCAP density functional (DFT) calculations

We implement the LCAP optimization in DFT calculations using the norm-conserving pseudo-potentials (NCPP) and ultra-soft pseudopotentials (USPP) based on a plane wave basis set [1-3]. The NCPP can be regarded as a special case of the USPP with the augmentation function $Q_{nm}(r)$ equal to zero and one reference energy only being used. For clarity, only the USPP is discussed here. The USPP is [2]:

$$v_A(\mathbf{r}, \mathbf{r}') = v_A^{loc}(\mathbf{r})\delta(\mathbf{r} - \mathbf{r}') + v_A^{nl}(\mathbf{r}, \mathbf{r}') = v_A^{loc}(\mathbf{r})\delta(\mathbf{r} - \mathbf{r}') + \sum_{nm} D_{nm}^0 \left| \beta_n^A \right\rangle \left\langle \beta_m^A \right|, \quad (\text{S.1})$$

where the D_{nm}^0 and β_n^A characterize the USPP for atom A .

The LCAP potential is expanded:

$$v(\mathbf{r}, \mathbf{r}') = \sum_{\mathbf{R}, A} b_A^{\mathbf{R}} v_A^{\mathbf{R}}(\mathbf{r}, \mathbf{r}'), \quad (\text{S.2})$$

Thus, it can be written in USPP form as:

$$\begin{aligned} v(\mathbf{r}, \mathbf{r}') &= \sum_{\mathbf{R}, A} b_A^{\mathbf{R}} v_A^{\mathbf{R}, loc}(\mathbf{r})\delta(\mathbf{r} - \mathbf{r}') + \sum_{\mathbf{R}, A} b_A^{\mathbf{R}} v_A^{\mathbf{R}, nl}(\mathbf{r}, \mathbf{r}') \\ &= \sum_{\mathbf{R}, A} b_A^{\mathbf{R}} v_A^{\mathbf{R}, loc}(\mathbf{r})\delta(\mathbf{r} - \mathbf{r}') + \sum_{\mathbf{R}, A} b_A^{\mathbf{R}} \sum_{nm} D_{nm}^0 \left| \beta_n^A \right\rangle \left\langle \beta_m^A \right| \end{aligned} \quad (\text{S.3})$$

The first term is the total local potential $v^{loc}(\mathbf{r})$ and the second term is the total non-local potential $v^{nl}(\mathbf{r}, \mathbf{r}')$.

The total energy of the system in an external electric field is:

$$E(F_x) = \sum_i \left\langle \phi_i \left| -\frac{\hbar^2}{2m} \nabla^2 + v^{nl}(\mathbf{r}, \mathbf{r}') \right| \phi_i \right\rangle + E_H[n(\mathbf{r})] + E_{xc}[n(\mathbf{r})] + \int d\mathbf{r} (v^{loc}(\mathbf{r}) - xF) n(\mathbf{r}) \quad (\text{S.4})$$

The derivative of energy with respect to the $b_A^{\mathbf{R}}$ coefficients is computed with a fixed number of electrons N to be:

$$\begin{aligned}
\left. \frac{\partial E(F_x)}{\partial b_A^{\mathbf{R}}} \right|_N &= \frac{\partial}{\partial b_A^{\mathbf{R}}} \left\{ \sum_i \left\langle \phi_i \left| -\frac{\hbar^2}{2m} \nabla^2 + v^{nl}(\mathbf{r}, \mathbf{r}') \right| \phi_i \right\rangle + E_H[n(\mathbf{r})] + E_{xc}[n(\mathbf{r})] + \int d\mathbf{r} (v^{loc}(\mathbf{r}) - xF) n(\mathbf{r}) \right\} \\
&= \sum_i \left[\left\langle \frac{\partial \phi_i}{\partial b_A^{\mathbf{R}}} \left| -\frac{\hbar^2}{2m} \nabla^2 + v^{nl}(\mathbf{r}, \mathbf{r}') \right| \phi_i \right\rangle + c.c. \right] + \int \underbrace{[V_H(\mathbf{r}) + V_{xc}(\mathbf{r}) + v^{loc}(\mathbf{r}) - xF]}_{V_{eff}(\mathbf{r})} \frac{\partial n(\mathbf{r})}{\partial b_A^{\mathbf{R}}} d\mathbf{r} \quad (\\
&\quad + \sum_i \left\langle \phi_i \left| v_A^{nl}(\mathbf{r}, \mathbf{r}') \right| \phi_i \right\rangle + \int v_A^{\mathbf{R}, loc}(\mathbf{r}) n(\mathbf{r}) d\mathbf{r}
\end{aligned}$$

S.5).

The electron density $n(\mathbf{r})$ is:

$$n(r) = \sum_i \left[|\phi_i|^2 + \sum_{A,R} b_A^R \sum_{nm} Q_{nm}^A \langle \phi_i | \beta_n^A \rangle \langle \beta_m^A | \phi_i \rangle \right]. \quad (\text{S.6})$$

From the above equation, $\frac{\partial n(\mathbf{r})}{\partial b_A^{\mathbf{R}}}$ is obtained:

$$\begin{aligned}
\frac{\partial n(\mathbf{r})}{\partial b_A^{\mathbf{R}}} &= \sum_i \sum_{nm} Q_{nm}^A(r) \langle \phi_i | \beta_n^A \rangle \langle \beta_m^A | \phi_i \rangle \\
&\quad + \sum_i \left[\frac{\partial \phi_i^*}{\partial b_A^{\mathbf{R}}} \phi_i + c.c. \right] + \sum_i \sum_{A,R} b_A^R \sum_{nm} Q_{nm}^A(r) \left[\left\langle \frac{\partial \phi_i}{\partial b_A^{\mathbf{R}}} \left| \beta_n^A \right\rangle + c.c. \right] \quad (\text{S.7})
\end{aligned}$$

Based on Eq. S.7, Eq. S.5 can be written:

$$\begin{aligned}
\left. \frac{\partial E(F_x)}{\partial b_A^{\mathbf{R}}} \right|_N &= \sum_i \sum_{nm} \underbrace{\left(D_{nm}^0 + \int V_{eff}(\mathbf{r}) Q_{nm}^A(r) d\mathbf{r} \right)}_{D_{nm}^{I,A}} \langle \phi_i | \beta_n^A \rangle \langle \beta_m^A | \phi_i \rangle + \int v_A^{\mathbf{R},loc}(\mathbf{r}) n(\mathbf{r}) d\mathbf{r} \\
&+ \sum_i \left(\left\langle \frac{\partial \phi_i}{\partial b_A^{\mathbf{R}}} \left| \underbrace{\left(-\frac{\hbar^2}{2m} \nabla^2 + V_{eff}(\mathbf{r}) + \sum_{\mathbf{R},A} b_A^{\mathbf{R}} \sum_{nm} D_{nm}^{I,A} |\beta_n^A\rangle \langle \beta_m^A| \right)}_{\hat{H}} \right| \phi_i \right\rangle + c.c. \right) \quad (\text{S.8}) \\
&= \sum_i \sum_{nm} D_{nm}^{I,A} \langle \phi_i | \beta_n^A \rangle \langle \beta_m^A | \phi_i \rangle + \int v_A^{\mathbf{R},loc}(\mathbf{r}) n(\mathbf{r}) d\mathbf{r} + \sum_i \left(\left\langle \frac{\partial \phi_i}{\partial b_A^{\mathbf{R}}} | \hat{H} | \phi_i \right\rangle + c.c. \right)
\end{aligned}$$

Using $\hat{H} |\phi_i\rangle = \varepsilon_i S |\phi_i\rangle$, we obtain

$$\left(\left\langle \frac{\partial \phi_i}{\partial b_A^{\mathbf{R}}} | \hat{H} | \phi_i \right\rangle + c.c. \right) = \varepsilon_i \left(\left\langle \frac{\partial \phi_i}{\partial b_A^{\mathbf{R}}} | S | \phi_i \right\rangle + c.c. \right) \quad (\text{S.9})$$

where S is a Hermitian overlap operator:

$$S = 1 + \sum_{\mathbf{R},A} \sum_{nm} b_A^{\mathbf{R}} q_{nm}^A |\beta_n^A\rangle \langle \beta_m^A| \quad (\text{S.10})$$

The generalized orthonormality condition $\langle \phi_i | S | \phi_j \rangle = \delta_{ij}$ gives

$$\left(\left\langle \frac{\partial \phi_i}{\partial b_A^{\mathbf{R}}} | S | \phi_i \right\rangle + c.c. \right) + \left\langle \phi_i \left| \frac{\partial S}{\partial b_A^{\mathbf{R}}} \right| \phi_i \right\rangle = 0 \quad (\text{S.11})$$

and Eq. S.9 can be written:

$$\left(\left\langle \frac{\partial \phi_i}{\partial b_A^{\mathbf{R}}} | \hat{H} | \phi_i \right\rangle + c.c. \right) = -\varepsilon_i \left\langle \phi_i \left| \frac{\partial S}{\partial b_A^{\mathbf{R}}} \right| \phi_i \right\rangle = -\varepsilon_i \sum_{nm} q_{nm}^A \langle \phi_i | \beta_n^A \rangle \langle \beta_m^A | \phi_i \rangle \quad (\text{S.12})$$

Substituting Eq. S.12 into Eq. S.8, the derivative of the total energy with respect to $b_A^{\mathbf{R}}$ is:

$$\left. \frac{\partial E(F_x)}{\partial b_A^{\mathbf{R}}} \right|_N = \sum_i \sum_{nm} D_{nm}^{I,A} \langle \phi_i | \beta_n^A \rangle \langle \beta_m^A | \phi_i \rangle + \int v_A^{\mathbf{R},loc}(\mathbf{r}) n(\mathbf{r}) d\mathbf{r} - \sum_i \varepsilon_i \sum_{nm} q_{nm}^A \langle \phi_i | \beta_n^A \rangle \langle \beta_m^A | \phi_i \rangle$$

(S.13)

Similarly, the derivative of the energy with respect to $b_A^{\mathbf{R}}$ for a fixed potential $v^{\text{nl}}(\mathbf{r}, \mathbf{r}')$ is:

$$\left. \frac{\partial E(F_x)}{\partial b_A^{\mathbf{R}}} \right|_{v(\mathbf{r}, \mathbf{r}')} = \frac{\partial E(F_x)}{\partial N} \frac{\partial N}{\partial b_A^{\mathbf{R}}} = \mu_{chem} Z_A^{\mathbf{R}}, \quad (\text{S.14})$$

Here, the constraint $N = \sum_{A, \mathbf{R}} Z_A^{\mathbf{R}}$ is applied to insure a neutral molecule. μ_{chem} is the chemical potential of the system, and $Z_A^{\mathbf{R}}$ is the atomic number of atom A at site $\mathbf{R}$.

Finally, the derivative of the energy with respect to the coefficients $b_A^{\mathbf{R}}$ is:

$$\frac{\partial E(F_x)}{\partial b_A^{\mathbf{R}}} = \sum_i \sum_{nm} D_{nm}^{I,A} \langle \phi_i | \beta_n^A \rangle \langle \beta_m^A | \phi_i \rangle + \int v_A^{\mathbf{R},loc}(\mathbf{r}) n(\mathbf{r}) d\mathbf{r} - \sum_i \varepsilon_i \sum_{nm} q_{nm}^A \langle \phi_i | \beta_n^A \rangle \langle \beta_m^A | \phi_i \rangle + \mu_{chem} Z_A^{\mathbf{R}}$$

(S.15)

Note that the above equations also apply to the norm-conserving pseudopotentials, as special cases, when we set Q_{nm}^A and q_{nm}^A to zero,

The polarizability α can be calculated using finite-field methods [4]:

$$\alpha_{ii} = -\frac{1}{F^2} [E_{tot}(+F_i) + E_{tot}(-F_i) - 2E(0)] \quad (\text{S.16})$$

where i is the index x, y, z. Based on Eq. S.15, the derivative of the polarizability with respect to the $b_A^{\mathbf{R}}$ coefficient is:

$$\frac{\partial \alpha_{ii}}{\partial b_A^{\mathbf{R}}} = -\frac{1}{F^2} \left[\frac{\partial E(+F_i)}{\partial b_A^{\mathbf{R}}} + \frac{\partial E(-F_i)}{\partial b_A^{\mathbf{R}}} - 2 \frac{\partial E(0)}{\partial b_A^{\mathbf{R}}} \right] \quad (\text{S.17})$$

The hyperpolarizability β_μ is defined by $\beta_\mu = \frac{\sum_i \beta_i \mu_i}{|\mu|}$ ($i = x, y, z$). The electric dipole moment μ_i^e is calculated with the finite-field method as [4]:

$$\mu_i^e = \left[-\frac{2}{3} \{E(F_i) - E(-F_i)\} + \frac{1}{12} \{E(2F_i) - E(-2F_i)\} \right] / F_i \quad (\text{S.18})$$

The molecular dipole is:

$$\mu_i = \mu_i^e + \sum_{\mathbf{R}, A} b_A^{\mathbf{R}} Z_A^{\mathbf{R}} R_i^A \quad (\text{S.19})$$

The derivative of the molecular dipole with respect to $b_A^{\mathbf{R}}$ is

$$\frac{\partial \mu_i}{\partial b_A^{\mathbf{R}}} = \frac{\partial \mu_i^e}{\partial b_A^{\mathbf{R}}} + Z_A^{\mathbf{R}} R_i^A \quad (\text{S.20})$$

The β_i components are $\beta_i = \sum_j \beta_{ij}$ ($j = x, y, z$). Each β_{ij} component is calculated using the finite-field method [4]:

$$\beta_{ii} = \left[-\frac{1}{2} \{E(2F_i) - E(-2F_i)\} + \{E(F_i) - E(-F_i)\} \right] / F_i^3 \text{ in case of } j = i, \quad (\text{S.21})$$

$$\beta_{ij} = \left[\frac{1}{2} \{E(-F_i, -F_j) - E(F_i, F_j) + E(-F_i, F_j) - E(F_i, -F_j)\} + \{E(F_i) - E(-F_i)\} \right] / F_i F_j^2 \quad \text{for } j \neq i. \quad (\text{S.22})$$

Like the calculation of $\frac{\partial \alpha_{ii}}{\partial b_A^{\mathbf{R}}}$, $\frac{\partial \mu_i^e}{\partial b_A^{\mathbf{R}}}$ and $\frac{\partial \beta_i}{\partial b_A^{\mathbf{R}}}$ are computed based on Eq. S.15. Finally, $\frac{\partial \beta_\mu}{\partial b_A^{\mathbf{R}}}$ is given by:

$$\frac{\partial \beta_{\mu}}{\partial b_A^{\mathbf{R}}} = \frac{1}{|\mu|} \sum_{i=1}^3 \left(\beta_i \frac{\partial \mu_i}{\partial b_A^{\mathbf{R}}} + \mu_i \frac{\partial \beta_i}{\partial b_A^{\mathbf{R}}} \right) - \frac{1}{|\mu|^3} \left(\sum_{i=1}^3 \beta_i \mu_i \right) \sum_{i=1}^3 \mu_i \frac{\partial \mu_i}{\partial b_A^{\mathbf{R}}} \quad (\text{S.23})$$

3. Results for a system with two designable sites

Figs. S2 and S3 show the optimization results beginning with uniform initial coefficients in the case of two functional groups (-CH₃ and -SH). The results converge at the maximum with 87% -SH at one site and 74% -SH at the other site within a few α calculations, and the gradients decrease to zero. The optimized results indicate that the -SH group dominates both designable sites. This means the linear polarizability optimized molecule is H₂S₂. Fig. S4 shows results in the case of six functional groups (-CH₃, -OH, -NH₂, -F, -Cl, and -SH), at each of two sites, beginning with uniform initial coefficients. H₂S₂ is the molecule with the maximum electronic polarizability. Table S1 gives the polarizabilities of all possible structures.

Fig. S5 shows optimized coefficients for hyperpolarizability, beginning with uniform initial coefficients in the case of six functional groups (-CH₃, -OH, -NH₂,

-F, -Cl and -SH). It shows that the system has maximum hyperpolarizability with 57% -SH at one site and 67% fluorine at the other site, i.e., SH-F. This result is in agreement with the direct enumeration and evaluation (Table S1). The calculated hyperpolarizability based on finite-field analysis with plane wave basis sets is in accordance with PW91PW91/aug-cc-pvtz Gaussian03 results [9].

Figure S2. The polarizability and $\max|d\alpha/dt_i|$ versus the number of α evaluations beginning with uniform coefficients and the two functional group $-\text{CH}_3$ and $-\text{SH}$ on left (L) and right (R) sites.

Figure S3. The coefficients of $-\text{CH}_3$ and $-\text{SH}$ versus the number of α evaluations beginning with uniform coefficients at the two designable sites.

Figure S4. The coefficients versus the number of α evaluations beginning with uniform coefficients on six functional groups (-CH₃, -OH, -NH₂, -F, -Cl and -SH).

4. Polarizabilities and hyperpolarizabilities of all possible molecules

Table S1 lists the computed electronic polarizabilities and hyperpolarizabilities of all possible species obtained by enumerating the structures and computing the values one by one.

Figure. S5. The coefficients at each point versus the number of β_μ evaluations beginning with uniform coefficients.

References

1. Baroni, S.; Dal Corso, A.; de Gironcoli, S.; Giannozzi, P.; Cavazzoni, C.; Ballabio, G.; Scandolo, S.; Chiarotti, G.; Focher, P.; Pasquarello, A.; Laasonen, K.; Trave, A.; Car, R.; Marzari, N.; Kokalj, A. <http://www.pwscf.org/>.
2. Laasonen, K.; Pasquarello, A.; Car, R.; Lee, C.; Vanderbilt, D. *Phys. Rev. B*, **1993**, *47*, 10142
3. Kleinman, L.; Bylander, D.M. *Phys. Rev. Lett.*, **1982**, *48*, 1425.
4. (a) Kurtz, H.A.; Stewart, J.J.P.; Dieter, K.M. *J. Comp. Chem.*, **1990**, *11*, 82.

- (b) Kurtz, H.A.; Douglas, S.D. *Rev. Comp. Chem.* **1998**, *12*, 241.
5. Duijnen, P.T.; Swart, M. *J. Phys. Chem. A*. **1998**, *102*, 2399.
6. Barbagli, E.; Maestro, M. *Chem. Phys. Lett.* **1974**, *24*, 567.
7. Koch, W.; Holthausen, M.C. *A Chemist's Guide to Density Functional Theory*, 2nd Ed.; Wiley-VCH: New York, 2001.
8. Schürer, G.; Gedeck, P.; Gottschalk, M.; Clark, T. *Int. J. Quan. Chem.* **1999**, *75*, 17.
9. Gaussian 03, Revision C.02, J. A. Pople, et al., Gaussian, Inc., Wallingford, CT, 2004.