Supporting Information

Development of Bifunctional Salen Catalysts: Rapid, Chemoselective Alkylations of α -Ketoesters

Erin F. DiMauro and Marisa C. Kozlowski*

Department of Chemistry, Roy and Diana Vagelos Laboratories, University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA

Table of Contents

General Considerations	S2-S3
Catalyst Synthesis and α -Ketoester Addition Procedures	S3-S8
Nonlinear Effects Study	S9-S12
References	S12-S14

General Considerations. Unless otherwise noted, all non-aqueous reactions were carried out under an atmosphere of dry N₂ in dried glassware. All manipulations involving Et₂Zn were carried out using standard Schlenk techniques. When necessary, solvents and reagents were dried prior to use. Toluene and dioxane were distilled from Na. CH₂Cl₂ was de-oxygenated by purging with N₂ and then dried by passing through activated alumina. THF was distilled from Na/benzophenone ketyl. Al(Oi-Pr)₃ and Zr(Oi-Pr)₄·i-PrOH were purchased from Acros and stored in an inert atmosphere glove box. V(O)(OⁱPr)₃ was purchased from Strem and used neat, without further purification. Ti(Oi-Pr)₄ was distilled and stored as a 1.4 M solution in hexanes. Ti(Ot-Bu)₄ was purchased from Acros and stored as a 1.4 M solution in hexanes. Et₂Zn was used as a freshly prepared 1.0 M solution in toluene. *n*-Bu₂Mg was used as a 1.0 M solution in heptane, purchased from Aldrich. (-)-(1*R*,2*R*)-1,2-Cyclohexanediamine and (+)-(1*S*,2*S*)-1,2-cyclohexanediamine were prepared as previously described.¹ Salen ligands for metal complexes **2**, **4** and **7** were prepared as previously described (see reference 1 in main text). The salen ligands for metal complexes **6**² and **9**³ were prepared according to literature procedures.

Ethyl oxo(phenyl)acetate and methyl oxo(phenyl)acetate were purchased from Aldrich and distilled prior to use. Ethyl 2-oxopropanoate and ethyl 3-methyl-2-oxobutanoate were purchased from Acros and distilled prior to use. Ethyl cyclohexyl(oxo)acetate,⁴ *tert*-butyl oxo(phenyl)acetate,⁵ ethyl (4-bromophenyl)(oxo)acetate,⁵ ethyl (4-methoxyphenyl)(oxo)acetate,⁵ and ethyl naphthyl(oxo)acetate⁶ were prepared as previously described. Ethyl 3,3-dimethyl-2-oxobutanoate⁷ was prepared from *tert*-butylmagnesium chloride and diethyl oxalate following a procedure similar to that described for ethyl cyclohexyl(oxo)acetate. Ethyl (2-methylphenyl)(oxo)acetate⁸ was prepared from *o*-tolylmagnesium bromide and diethyl oxalate following a procedure similar to that described for the preparation of ethyl naphthyl(oxo)acetate. All synthetic α -ketoesters were purified by distillation prior to use.

Analytical thin layer chromatography (TLC) was performed on EM Reagents 0.25 mm silica-gel 60-F plates. Preparative thin layer chromatography was performed on EM Reagents 1.00 mm silica-gel plates. Visualization was accomplished with UV light. Chromatography on silica gel was performed using a forced flow of the indicated solvent system on EM Reagents Silica Gel 60 (230-400 mesh).⁹ Conversions and enantiomeric excesses were measured using a Hewlett-Packard 5890 gas chromatograph (GC) with a Supelco β -DEXTM 120 column (12 m x 0.25 mm) or a Waters 600 high performance liquid chromatography (HPLC) with UV detection at 254 nm and Daicel Chiralpak AD or OD columns (0.46 cm x 25 cm). ¹H NMR spectra were recorded on Bruker AM-500 (500 MHz) or AM-250 (250 MHz) spectrometers at ambient temperature. Chemical shifts are reported in ppm from tetramethylsilane (0 ppm) or from the solvent resonance (CDCl₃ 7.26 ppm, CD₂Cl₂ 5.30 ppm, DMSO-d₆ 2.49 ppm, D₂O 4.80 ppm). Data are reported as follows: chemical shift, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, br = broad, m = multiplet), coupling constants, and number of protons. Proton decoupled ¹³C NMR spectra were recorded on a Bruker AM-500 (125 MHz) spectrometer at ambient temperature. Chemical shifts are reported in ppm from tetramethylsilane (0

ppm) or from the solvent resonance (CDCl₃ 77.36 ppm). Mass spectra were obtained on a low resonance Micromass Platform LC in electrospray mode and a high resonance VG autospec with an ionization mode of either CI or ES. IR spectra were taken on a Perkin-Elmer FT-IR spectrometer using a thin film on NaCl plates. Melting points were obtained on Thomas Scientific Unimelt apparatus and are uncorrected. Optical rotations were measured on a Perkin-Elmer Polarimeter 341 with a sodium lamp and are reported as follows $[\alpha]_D^{25}$, (c g/100 mL solvent).

11

Benzyl Oxo(phenyl)acetate (11). Benzoyl formic acid (2.5 g, 17.3 mmol) and benzyl alcohol (2.24 g, 20.76 mmol) were heated at reflux in benzene (15 mL) with a Dean Stark trap for 20 h. The reaction was cooled to room temperature and concentrated. Flash chromatography on SiO₂ (5% EtOAc/hexanes) afforded pure **11** as a yellow oil in 85% yield. Compound **11** has previously been synthesized by other methods.¹⁰

12

5-tert-Butyl-2-hydroxy-3-(pyrrolidin-1-ylmethyl)benzaldehyde (12). Pyrrolidine (0.56 g, 7.86 mmol) was added to a suspension of paraformaldehyde (0.26 g, 8.65 mmol) in glacial HOAc (3.25 mL). The suspension was stirred at room temperature for 3 h. 5-tert-Butyl-2-hydroxybenzaldehyde¹¹ (1.4 g, 7.86 mmol) was then added as a solution in absolute EtOH (13 mL) and the mixture was heated at reflux for 5 d. After cooling to room temperature, the mixture was brought to pH~8 with saturated NaHCO₃, extracted with CHCl₃, dried over Na₂SO₄, and concentrated. Preparative TLC on SiO₂ (50% EtOAc/hexanes) afforded **12** in 20% as a yellow oil: ¹H NMR (500 MHz, CDCl₃) δ = 10.31 (br s, 1H), 7.61 (d, *J* = 2.5 Hz, 1H), 7.32 (d, *J* = 2.3 Hz, 1H), 6.35 (br s, 1H), 3.84 (s, 2H), 2.67 (br s, 4H), 1.86 (m, 4H), 1.28 (s, 9H); ¹³C NMR (125 MHz, CDCl₃) δ = 191.8, 159.6, 141.3, 132.3, 124.5, 123.9, 122.0, 57.4, 53.6, 34.1, 31.3, 23.6; IR (film) 3500-2500 (br), 2961, 2871, 2842, 1679, 1651, 1606, 1478, 1395 cm⁻¹; HRMS (ES) calcd *m/z* 262.1807 [C₁₆H₂₄NO₂]⁺, found *m/z* 262.1796.

13

5-*tert*-Butyl-2-hydroxy-3-(dimethylamino-methyl)-benzaldehyde (13). Dimethylamine (189 mg of a 40% soln. in H₂O, 1.68 mmol) was added to a suspension of 5-*tert*-butyl-2-hydroxy-benzaldehyde (200 mg, 1.12 mmol) and formaldehyde (136 mg of a 37% soln. in H₂O, 1.68 mmol) in EtOH (1 mL). The reaction was heated at reflux for 24 h, then cooled to room temperature and concentrated. The residue was dissolved in CH₂Cl₂ and dried over Na₂SO₄, then concentrated and purified by preparative chromatography on SiO₂ (30% EtOAc/hexanes). **13** was isolated as a pale yellow powder in 80% yield (210 mg, 0.896 mmol): ¹H NMR (500 MHz, CDCl₃) δ = 10.3 (s, 1H), 7.63 (d, *J* = 2.6 Hz, 1H), 7.33 (br s, 1H), 3.67 (s, 2H), 2.36 (s, 6H), 1.30 (s, 9H); ¹³C NMR (125 MHz, CDCl₃) δ = 192.6, 159.9, 142.2, 133.4, 127.7, 125.5, 61.4, 45.0, 34.5, 31.7; HRMS (ES) calcd *m/z* 236.1650 [C₁₄H₂₂NO₂⁺], found *m/z* 236.1652.

14

8-(Tri-*n*-butylstannyl)quinoline (14). 8-Bromoquinoline (6.84 g, 32.9 mmol) was dissolved in THF (15 mL) and cooled to -78 °C. *n*-BuLi (15.5 mL, 36.2 mmol) was added dropwise while the reaction temperature was kept below -70 °C. To the resulting deep red solution was added *n*-Bu₃SnCl (13.4 mL, 49.3 mmol) in THF (10 mL), while the temperature was carefully maintained below -70 °C. Within 30 min the solution became orange and after 2 h, saturated NaHCO₃ solution (15 mL) was added, raising the temperature to -13 °C. After warming to room temperature overnight, the layers were separated and the aqueous portion was extracted with EtOAc. The combined organic extracts were dried over Na₂SO₄ and concentrated to an orange oil. Two flash chromatographic purifications on SiO₂ (100% hexanes) afforded **14** in 61% yield (8.40 g, 20.0 mmol) as a clear oil: ¹H NMR (250 MHz, CDCl₃) sample contaminated with ~10% *n*-Bu₄Sn, δ = 8.86 (dd, *J* = 1.8, 4.3 Hz, 1H), 8.11 (dd, *J* = 1.8, 8.2 Hz, 1H), 7.85 (dd, *J* = 1.4, 6.6 Hz, 1H), 7.74 (dd, *J* = 1.4, 8.1 Hz, 1H), 7.47 (dd, *J* = 6.5, 8.1 Hz, 1H), 7.32 (dd, *J* = 4.2, 8.2 Hz, 1H), 1.65-1.48 (m, 6H), 1.40-1.31 (m, 8H), 1.38-1.09 (m, 6H), 0.86 (t, *J* = 7.2 Hz, 12H); ¹³C NMR (125 MHz, CDCl₃) δ = 153.2, 149.1, 148.2, 137.4, 136.0, 127.8, 127.6, 126.2, 120.6, 29.3, 27.4, 13.7, 10.6.

15

5-*tert*-Butyl-2-hydroxy-3-quinolin-8-yl-benzaldehyde (15). 3-Bromo-5-*tert*-butyl-2-hydroxybenzaldehyde² (300 mg, 1.17 mmol) and 8-(tri-*n*-butylstannyl)quinoline (732 mg, 1.75 mmol) were dissolved in dry dioxane (9 mL) and added to a mixture of Pd(PPh₃)₄ (57.4 mg, 0.058 mmol) and CuBr (16.8 mg, 0.117 mmol) via cannula transfer. The suspension was then heated to reflux for 10 h, allowed to cool to room temperature, and concentrated. The resultant oil was taken up in EtOAc and washed with water. The aqueous portion was extracted with CH₂Cl₂. The combined organic fractions were dried over Na₂SO₄ and concentrated. Purification by flash chromatography on SiO₂ (10% EtOAc/hexanes) afforded **15** in 80% yield (269 mg, 0.98 mmol) as a yellow solid: ¹H NMR (500 MHz, CDCl₃) δ = 11.27 (br s, 1H), 10.11 (s, 1H), 8.93 (s, 1H), 8.29 (d, *J* = 8.2 Hz, 1H), 7.90 (d, *J* = 6.9 Hz, 1H), 7.80 (d, *J* = 5.7 Hz, 1H), 7.73 (s, 1H), 7.68 (m, 2H), 7.45 (m, 1H), 1.38 (s, 9H); ¹³C NMR (125 MHz, CDCl₃) δ = 196.1, 157.3, 150.0, 142.2, 137.3, 137.0, 136.5, 131.7, 129.2, 128.7, 128.5, 128.4, 128.2, 126.5, 121.2, 121.1, 34.3, 31.3; IR (film) 3038, 2960, 2867, 1648, 1456, 1363, 1253 cm⁻¹; HRMS (ES) calcd *m/z* 306.1494 [C₂₀H₂₀NO₂⁺], found *m/z* 306.1494.

General Procedure for the Synthesis of Salens (*R,R*)-16 - (*R,R*)-18. The aldehyde (2.0 equiv) was dissolved in EtOH and (-)-(1*R*,2*R*)-1,2-cyclohexanediamine (1.0 equiv) was added. The mixture was stirred for 1 d at room temperature and then concentrated. The salen was taken up in CH₂Cl₂ and dried over Na₂SO₄. Filtration and reconcentration, followed by drying with gentle heat in vacuo for several hours afforded the salens in 100% yield as powdery, free-flowing solids.

(-)-(*R,R*)-16. Yellow solid: m.p. 66-74 °C; [α]_D²⁰ = -240 (c 0.5 in CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ = 13.43 (br s, 2H), 8.27 (s, 2H), 7.35 (br s, 2H), 6.90 (br s, 2H), 3.81-3.60 (m, 4H), 3.29 (m, 2H), 2.57 (br s, 8H), 1.91-1.68 (m, 14H), 1.45 (br t, *J* = 9.8 Hz, 2H), 1.27 (s, 18H); ¹³C NMR (125 MHz, CDCl₃) δ = 165.2, 157.3, 140.8, 131.0, 127.0, 125.2, 118.0, 72.9, 54.4, 34.1, 33.5, 31.8, 31.7, 24.9, 23.7; IR (film) 3480-2600 (br), 2959, 2932, 2861, 2790, 1629, 1471, 1362, 1273 cm⁻¹; HRMS (ES) calcd *m/z* 601.4482 [C₃₈H₅₇N₄O₂⁺], found *m/z* 601.4488.

(-)-(R,R)-(17). Yellow powder: m.p. 70-76 °C; $[\alpha]_D^{20} = -265$ (c 0.25 in CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ = 13.51 (br s, 2H), 8.27 (s, 2H), 7.31 (br s, 2H), 7.08 (br s, 2H), 3.66-3.49 (br m, 4H), 3.32-3.27 (br m, 2H), 2.30 (s, 12H), 1.92-1.85 (m, 4H), 1.70 (m, 2H), 1.45 (m, 2H), 1.24 (s, 18H); ¹³C NMR (125 MHz, CDCl₃) δ = 165.4, 158.0, 141.1, 131.9, 127.8, 118.2, 73.0, 58.0, 45.4, 34.3, 33.6, 31.8, 24.6; IR (film) 3600-2600, 2950, 2857, 2767, 1629, 1600, 1468, 1363, 1272 cm⁻¹; HRMS (ES) calcd *m/z* 549.4168 [C₃₄H₅₃N₄O₂⁺], found *m/z* 549.4183.

(-)-(R,R)-(18). Yellow-brown solid: m.p. 148-154 °C; $[\alpha]_D^{20} = -452$ (c 0.3 in CHCl₃); ¹H NMR (500 MHz, CDCl₃) δ = 13.6 (br s, 2H), 8.89 (s, 2H), 8.32 (br s, 2H), 8.20 (br s, 2H), 7.84 (d, *J* = 8.05 Hz, 2H), 7.78 (d, *J* = 6.8 Hz, 2H), 7.62 (t, *J* = 7.6 Hz, 2H), 7.48-7.22 (m, 4H), 3.25 (br s, 2H), 1.90-1.40 (m, 8H), 1.30 (s, 18H); ¹³C NMR (125 MHz, CDCl₃) δ = 165.2, 150.2, 136.7, 136.1, 132.2, 131.6, 131.2, 129.0, 128.5, 128.4, 128.1, 127.6, 126.4, 126.1, 120.8, 118.0, 72.5, 34.5, 33.3, 31.5, 24.2; IR(film) 3380 (br), 2957, 2861, 1628, 1496, 1456, 1362, 1288, 1254 cm⁻¹; HRMS (ES) calcd *m/z* 689.3855 [C₄₆H₄₉N₄O₂⁺], found *m/z* 689.3831.

Treatment of Ethyl α -Ketoesters with EtMgBr: Formation of the α -Hydroxyester Addition/Reduction Products. Mixtures of the addition and reduction standards were prepared by reaction of the α -ketoester (1 equiv) with ethylmagnesium bromide (1 equiv) in THF at -10 °C for 1 h. (Table S1). Racemic samples of the pure α -hydroxyester reduction products were prepared by reaction of the α -ketoester (1 equiv) with the NaBH₄ (1 equiv) in EtOH at 0 °C for 45 min-1 h.

Table S1. Addition of EtMgBr to α -ethyl ketoesters (eq 1, M = MgBr, n = 1, R⁴ = Et).

Entry	R ³	Reduction	Addition
		Conv. (%) ^a	Conv. (%) ^a
1	<i>p</i> -MeO-C ₆ H ₄	16	81
2	<i>p</i> -Br-C ₆ H ₄	22	69
3	Ph	19	60
4	<i>o</i> -Me-C ₆ H ₄	19	38
5	β -naphthyl	30	37
6	Me	52	38
7	Cy	54	31
8	<i>i</i> -Pr	48	21
9	<i>t</i> -Bu	63	21

^aDetermined by ¹H NMR spectroscopy.

General Procedure for Formation of the Metal Salen Complexes. The salen (0.030 mmol) was introduced into a dry Schlenk flask, and the system purged with N₂. The salen was dissolved in dry toluene (1 mL) and Ti(O*i*-Pr)₄ (18 μ L, 1.4 M in heptanes, 0.025 mmol) was added. After stirring at

room temperature for 1 h, the toluene, heptanes and *i*-PrOH were removed in vacuo (removal of the *i*-PrOH was found to increase reactivity and selectivity). Fresh toluene (1 mL) was added to provide the catalyst solution which was used in the α -ketoester additions as described below. Care must be taken to avoid the introduction of water into this titanium catalyst solution as lower reactivity and selectivity results.

The other catalyst derivatives were prepared in the same manner using the following reagents: Ti(*Ot*-Bu)₂ catalyst from Ti(*Ot*-Bu)₄; Al(*Oi*-Pr) catalyst from Al(*Oi*-Pr)₃¹²; V(O)(*Oi*-Pr) catalyst from V(O)(*Oi*-Pr)₃¹³; Zr(*Oi*-Pr)₂ catalyst from Zr(*Oi*-Pr)₄·*i*-PrOH¹⁴. The Zn and Mg catalysts were made from Et₂Zn and *n*-Bu₂Mg, respectively. For these last two catalysts, it is possible to forgo the solvent removal step as no *i*-PrOH is generated during the catalyst formation.

General Procedure for the Addition of Diethylzinc to α -Ketoesters. The catalyst solution (see above) was cooled to -40 °C and Et₂Zn (305 μ L, 1.0 M in toluene, 0.305 mmol) was added slowly. After 5-30 min, the α -ketoester (0.25 mmol) was added dropwise. At various intervals, 100 μ L aliquots were removed from the reaction, quenched with saturated NH₄Cl, and then extracted with pentane.

The reaction progress was monitored by GC (Carrier gas: N₂; Detector: FID, 270 °C; Injector: 250 °C; Pressure: 5-7 psi). GC oven temperatures and retention times for the α -ketoesters and their corresponding α -hydroxyester addition and reduction products are listed below. Since GC and ¹H NMR data from the reaction mixtures were generally in agreement, percent conversion was calculated using unadjusted GC peak areas.

Where isolated yields are reported, the reaction was run for 2 h at -40 °C then quenched with saturated NH₄Cl and extracted with CH₂Cl₂. The extracts were dried over Na₂SO₄, and concentrated. Purification by preparative TLC on SiO₂ (30% EtOAc/hexanes) or flash chromatography on SiO₂ (10% EtOAc/hexanes) afforded methyl 2-hydroxy-2-phenylbutanoate and ethyl 2-hydroxy-2-phenylbutanoate as pale yellow oils. The ¹H and ¹³C NMR spectra of these compounds were identical to those reported.¹⁵ Absolute configurations were determined by optical rotation and comparison to the reported values.¹⁶

GC oven temperatures and retention times for the α -ketoesters, addition products and reduction products are listed below.

Methyl oxo(phenyl)acetate *t* = 19.4 min; **methyl 2-hydroxy-2-phenylbutanoate**¹⁵ *t*_R = 30.8 min, *t*_S = 34.0 min; **methyl hydroxy(phenyl)acetate**¹⁷ *t*₁ = 34.8 min, *t*₂ = 42.7 min; oven = 100 °C.

Ethyl oxo(phenyl)acetate *t* = 28.5 min; **ethyl 2-hydroxy-2-phenylbutanoate**¹⁵ *t*_R = 36.9 min, *t*_S = 39.8 min; **ethyl hydroxy(phenyl)acetate**¹⁷ *t* = 42.5 min; oven = 100 °C.

***tert*-Butyl oxo(phenyl)acetate** *t* = 59.1 min; ***t*-butyl 2-hydroxy-2-phenylbutanoate**¹⁸ *t*₁ = 65.3 min, *t*₂ = 67.8 min; ***tert*-butyl hydroxy(phenyl)acetate**¹⁹ *t*₁ = 74.1 min, *t*₂ = 86.7 min; oven = 90 °C.

Ethyl (4-methoxyphenyl)(oxo)acetate $t = 47.4$ min; **ethyl 2-hydroxy-2-(4-methoxyphenyl)-butanoate**²⁰ $t_1 = 39.7$ min, $t_2 = 41.3$ min; **ethyl hydroxy(4-methoxyphenyl)acetate**²¹ $t_1 = 41.4$ min, $t_2 = 42.9$ min; oven = 130 °C.

Ethyl (4-bromophenyl)(oxo)acetate $t = 24.2$ min; **ethyl 2-(4-bromophenyl)-2-hydroxybutanoate**²² $t_1 = 31.0$ min, $t_2 = 32.4$ min; **ethyl (4-bromophenyl)(hydroxy)acetate**²³ $t_1 = 44.7$ min, $t_2 = 48.4$ min; oven = 135 °C.

Ethyl 2-naphthyl(oxo)acetate $t = 70.2$ min; **ethyl 2-hydroxy-2-(2-naphthyl)butanoate**²⁴ $t_1 = 76.7$ min, $t_2 = 78.8$ min; **ethyl hydroxy(2-naphthyl)acetate**²¹ $t_1 = 81.4$ min, $t_2 = 82.0$ min; oven = 145 °C.

Ethyl cyclohexyl(oxo)acetate $t = 18.9$ min; **ethyl 2-cyclohexyl-2-hydroxybutanoate**²⁵ $t_1 = 28.9$ min, $t_2 = 32.7$ min; **ethyl cyclohexyl(hydroxy)acetate**⁴ $t_1 = 25.6$ min, $t_2 = 26.2$ min; oven = 100 °C.

Ethyl 3,3-dimethyl-2-oxobutanoate $t = 8.4$ min; **ethyl 2-ethyl-2-hydroxy-3,3-dimethyl-butyrate**²⁶ $t_1 = 30.9$ min, $t_2 = 42.7$ min; **ethyl 2-hydroxy-3,3-dimethylbutanoate**²⁷ $t_1 = 5.3$ min, $t_2 = 7.1$ min; oven = 60 °C.

Ethyl 3-methyl-2-oxobutanoate $t = 14.5$ min; **ethyl 2-ethyl-2-hydroxy-3-methylbutanoate**²⁸ $t_1 = 36.5$ min, $t_2 = 40.3$ min; **ethyl 2-hydroxy-3-methylbutanoate**²⁹ $t_1 = 26.7$ min, $t_2 = 27.5$ min; oven³⁰ = 80 °C..

Ethyl (2-methylphenyl)(oxo)acetate $t = 25.8$ min; **ethyl 2-hydroxy-2-(2-methylphenyl)butanoate**³¹ $t = 44.4$ min; **ethyl hydroxy(2-methylphenyl)acetate**³² $t_1 = 36.8$ min, $t_2 = 38.5$ min; oven = 110 °C. Enantiomeric excess measured by HPLC (OD hexanes, 0.5 mL/min) after conversion to the trifluoroacetylated derivative (**ethyl 2-(2-methylphenyl)-2-[trifluoroacetyl]oxybutanoate**) $t_1 = 19.2$ min, $t_2 = 23.9$ min.

Ethyl 2-oxopropanoate $t = 4.7$ min; **ethyl 2-hydroxy-2-methylbutanoate**³³ $t = 15.0$ min; **ethyl 2-hydroxypropanoate**³⁴ $t_1 = 8.2$ min, $t_2 = 9.8$ min; oven = 50 °C. The ee was measured on the acetylated derivative (**ethyl 2-acetoxybutanoate**) $t_1 = 11.7$, $t_2 = 13.4$ min; oven = 40 °C.

Benzyl oxo(phenyl)acetate $t = 8.3$ min; **benzyl 2-hydroxy-2-phenylbutanoate**³⁵ $t_1 = 8.9$ min, $t_2 = 11.7$ min; **benzyl hydroxy(phenyl)acetate**³⁶ $t_1 = 12.7$ min, $t_2 = 14.9$ min; Conversion measured by ¹H NMR. Enantiomeric excess measured by HPLC (AD 10% *i*-PrOH/hexane, 1.0 mL/min).

Enrichment of the PhCOCO₂Me/Et₂Zn Adduct by Recrystallization. A portion (600 mg, 3.14 mmol, 78% ee) of the isolated adduct from a reaction employing 5 mmol of PhCOCO₂Me and 5 mol% of **2** with 1.2 equiv. of Et₂Zn was treated with ethanolic KOH to provide the α -hydroxy acid (**10**) in quantitative yield. This material was recrystallized four times from CCl₄ to provide 72% of the α -hydroxy acid (400 mg, 2.26 mmol). A portion of this material was treated with diazomethane to provide the corresponding methyl ester (98% ee) which was analyzed by GC as described above.

Nonlinear Effects Study. These experiments were done at -78 °C due to technical difficulties at -40 °C. When the reaction from Table 3, entry 9 was performed at -78 °C, 96% conversion and 74% ee (*R*) were observed after 2 h. Two different procedures were employed for the nonlinear studies. In the first, (**A**), catalyst solutions made separately from the (*S,S*)- and (*R,R*)-salens were combined in the appropriate proportions to provide the catalyst solution of the indicated catalyst enantiomeric excess. In the second, (**B**), catalyst solutions made separately from the racemic and (*R,R*)-salens were combined in the appropriate proportions to provide the catalyst solution of the indicated catalyst enantiomeric excess. The results from both trials were similar (see Figures S1 and S2 below) indicating that the catalyst species were equilibrating to the same starting point under the reaction conditions. For example, an unreactive species was not irreversibly formed from the racemic salen upon treatment with titanium to form the catalyst.

A. (*R,R*)-Cyclohexanediamine piperidine salen, (*R,R*)-1 (85 mg, 0.136 mmol), was dissolved in toluene (1 mL) in a 10 mL Schlenk flask in an inert atmospheres glove box. Neat Ti(*Oi*-Pr)₄ was added (40.6 μ L, 0.136 mmol) and the solution was allowed to stir for 1 h. The toluene and *i*-PrOH were then removed in vacuo and the dry catalyst was redissolved in fresh toluene (0.54 mL) to make a 0.25 M solution of (*R,R*)-2. This procedure was simultaneously performed using the (*S,S*)-cyclohexanediamine piperidine salen, (*S,S*)-1, to make a 0.25 M solution of (*S,S*)-2. The subsequent ketoester alkylation reactions were performed simultaneously in flame dried, N₂ purged test-tubes. The test-tubes were each charged with a total volume of 100 μ L (0.025 mmol) of these freshly prepared catalyst solutions as outlined in Table S2.

Additional toluene (0.9 mL) was added to each test-tube and the tubes were removed from the glove box. After cooling to -78 °C, Et₂Zn (300 μ L of a 1.0 M soln. in hexanes, 0.3 mmol) was slowly added to each test tube. After stirring for 30 min, methyl oxo(phenyl)acetate (35.5 μ L, 0.25 mmol) was added dropwise to each tube. After 2 h, the reactions were quenched by injection of 1.5 mL cold water. After extraction with pentane, the crude reaction mixtures were analyzed by GC (see Table S2) as described in the General Procedure above.

Table S2. Nonlinear effect study (method A) using (*S,S*)-2 and (*R,R*)-2 catalyst solutions in the addition of Et₂Zn to PhCOCO₂Me.

Tube	μ L (<i>S,S</i>)-2 solution	μ L (<i>R,R</i>)-2 solution	catalyst %ee in (<i>S,S</i>)-2	addition product ^a % ee
1	50	50	0	1.6 (<i>R</i>) ^b
2	62.5	37.5	25	18.7 (<i>S</i>)
3	75	25	50	39.0 (<i>S</i>)
4	0	100	100	76.8 (<i>S</i>)

^a No reduction product was observed. ^bThe non-zero reading arises from a small amount of error in volumes of the catalyst solutions combined or in the GC assay.

Figure S1. Nonlinear effect study (method A) using (*S,S*)-**2** and (*R,R*)-**2** catalyst solutions in the addition of Et₂Zn to PhCOCO₂Me.

B. (*R,R*)-Cyclohexanediamine piperidine salen, (*R,R*)-**1** (100 mg, 0.16 mmol) was dissolved in toluene (1 mL) in a 10 mL Schlenk flask in an inert atmospheres glove box. Neat Ti(*Oi*-Pr)₄ (48 μL, 0.16 mmol) was then added and the solution was allowed to stir for 1 h. The toluene and *i*-PrOH were then removed in vacuo and the dry catalyst was redissolved in fresh toluene (1.94 mL) to make a 0.083 M solution of (*R,R*)-**2**. This procedure was simultaneously performed using racemic cyclohexanediamine piperidine salen, (+/-)-**1**, to make a 0.083 M solution of (+/-)-**2**. This racemic solution was slightly cloudy. The ketoester alkylation reactions were performed simultaneously in flame dried, N₂ purged test-tubes. The test-tubes were each charged with a total volume of 300 μL (0.025 mmol) of freshly prepared catalyst solutions as outlined in Table S3.

Toluene (0.7 mL) was then added to each test-tube and the tubes were removed from the glove box. After cooling to -78 °C, Et₂Zn (300 μL of a 1.0 M soln. in hexanes, 0.3 mmol) was slowly added to each test tube. After stirring for 30 min, methyl oxo(phenyl)acetate (35.5 μL, 0.25 mmol) was added dropwise to each tube. After 2 h, the reactions were quenched by injection of 1.5 mL cold water. After extraction with pentane, the crude reaction mixtures were analyzed by GC (see Table S3) as described in the General Procedure above.

Table S3. Nonlinear effect study (method B) using rac-**2** and (*R,R*)-**2** catalyst solutions in the addition of Et₂Zn to PhCOCO₂Me.

Tube	μL (+/-)- 2 solution	μL (<i>R,R</i>)- 2 solution	catalyst %ee in (<i>R,R</i>)- 2	addition product ^a % ee
1	300	0	0	1.8 (<i>S</i>) ^b
2	150	150	50	44.6 (<i>R</i>)
3	75	225	75	53.5 (<i>R</i>)
4	0	300	100	75.8 (<i>R</i>)

^a No reduction product was observed. ^bThe non-zero reading with catalyst from racemic salen arises from a small amount of error in the GC assay.

Figure S2. Nonlinear effect study (method B) using rac-**2** and (*R,R*)-**2** catalyst solutions in the addition of Et₂Zn to PhCOCO₂Me.

A plot combining the catalyst ee vs product ee results from methods A and B is shown in below in Figure S3. The reaction is characterized by the absence of a nonlinear effect, which is somewhat unusual for titanium containing catalysts.³⁷

Figure S3. Nonlinear effect study combining the results from methods A and B.

- (1) a) Galsbol, F.; Steenbol, P.; Sondergaard Sorensen, B. *Acta Chem. Scand.* **1972**, 26, 3605-3611. b) Larrow, J. F.; Jacobsen, E. N.; Gao, Y.; Hong, Y.; Nie, X.; Zepp, C. M. *J. Org. Chem.* **1994**, 59, 1939-1942. c.) Larrow, J. F.; Jacobsen, E. N. *Org. Synth.* **1998**, 75, 1-11.
- (2) Lam, F.; Xu, J. X.; Chan, K. S. *J. Org. Chem.* **1996**, 61, 8414-8418.
- (3) Deng, L.; Jacobsen, E. N. *J. Org. Chem.* **1992**, 57, 4320-4323.
- (4) Boivin, S.; Outurquin, F.; Paulmier, C. *Tetrahedron* **1997**, 53, 16767-16782.
- (5) Hu, S.; Neckers, D. C. *J. Org. Chem.* **1996**, 61, 6407-6415.
- (6) Middleton, W. J.; Bingham, E. M. *J. Org. Chem.* **1980**, 45, 2883-2887.
- (7) b.p. 62-66 °C, 10 mm Hg. Previous synthesis: Rabjohn, N.; Herbert, H. C. *J. Org. Chem.* **1970**, 35, 3726-3729.
- (8) b.p. 90-110 °C, 2 mm Hg. Previous synthesis: Hoch, J.; Choisy, J. M. *C. R. Hebd. Seances Acad. Sci.* **1959**, 248, 3314-3316.
- (9) Still, W. C.; Kahn, M.; Mitra, A. *J. Org. Chem.* **1978**, 43, 2923-2925.
- (10) Domagala, J. M. *Tetrahedron Lett.* **1980**, 21, 4997-5000.
- (11) Casiraghi, G.; Casnati, G.; Puglia, G.; Sartori, G.; Terenghi, G. *J. Chem. Soc., Perkin Trans. 1* **1980**, 1862-1865.

- (12) For examples of salen-Al complexes, see: Atwood, D. A.; Harvey, M. J. *Chem. Rev.* **2001**, *101*, 37-55, and references cited therein.
- (13) For examples of salen-V complexes, see: a) Choudhary, N.; Hughes, D. L.; Kleinkes, U.; Larkworthy, L. F.; Leigh, G. J.; Maiwald, M.; Marmion, C. J.; Sanders, J. R.; Smith, G. W.; Sudbrake, C. *Polyhedron*, **1997**, *16*, 1517-1528. b) Dutton, J. C.; Fallon, G. D.; Murray, K. S. *Inorg. Chem.* **1998**, *27*, 34-38.
- (14) For examples of salen-Zr complexes, see: a) Watanabe, A.; Uchida, T.; Ito, K.; Katsuki, T. *Tetrahedron Lett.* **2002**, *43*, 4481-4485. b) Cocazza, F.; Solari, E.; Floriani, C.; Chiesi-Villa, C.; Guastini, C. *J. Chem. Soc., Dalton Trans.* **1990**, 1335-1344.
- (15) a) **methyl 2-hydroxy-2-phenylbutanoate**: Julia, M.; Pfeuty-Saint Jalmes, V.; Ple, K.; Verpeaux, J.-N.; Hollingworth, G. *Bull. Soc. Chim. Fr.* **1996**, *133*, 15-24; b) **ethyl 2-hydroxy-2-phenylbutanoate**: Nozaki, H.; Aratani, T.; Toraya, T.; Noyori, R. *Tetrahedron* **1971**, *27*, 905-913.
- (16) a) **methyl 2-hydroxy-2-phenylbutanoate**: McKenzie, A.; Richie, A. *Chem. Ber.* **1937**, *70-B*, 23-36. b) **ethyl 2-hydroxy-2-phenylbutanoate**: Nozaki, H.; Aratani, T.; Toraya, T. *Tetrahedron Lett.* **1968**, 4097-4098.
- (17) Basavaiah, D.; Rama Krishna, P. *Tetrahedron* **1995**, *51*, 2403-2416.
- (18) Previous synthesis and partial characterization: see reference 15a.
 ^1H NMR (500 MHz, CDCl_3) δ = 7.60 (d, J = 8.6 Hz, 2H), 7.35-7.20 (m, 3H), 2.19 (dq, J = 14.6, 7.3 Hz, 1H), 1.97 (dq, J = 14.6, 7.3 Hz, 1H), 1.45 (s, 9H), 0.93 (t, J = 7.3 Hz, 3H).
- (19) Adam, W.; Metz, M.; Prechtel, F.; Renz, M. *Synthesis* **1994**, *6*, 563-566.
- (20) ^1H NMR (500 MHz, CDCl_3) δ = 7.50 (d, J = 8.8 Hz, 2H), 6.87 (d, J = 8.8 Hz, 2H), 2.19 (dq, J = 14.6, 7.3 Hz, 1H), 2.0 (dq, J = 14.6, 7.3 Hz, 1H), 1.23 (t, J = 6.4 Hz, 3H), 0.91 (t, J = 7.3 Hz, 3H).
- (21) Monenschein, H.; Drager, G.; Jung, A.; Kirschning, A. *Chem. -Eur. J.* **1999**, *5*, 2270-2280.
- (22) ^1H NMR (500 MHz, CDCl_3): δ = 7.50-7.22 (m, 4H), 4.29-4.18 (m, 2H), 2.18 (dq, J = 14.2, 7.1 Hz, 1H), 1.98 (dq, J = 14.2, 7.1 Hz, 1H), 1.28 (t, J = 6.1 Hz, 3H), 0.90 (t, J = 7.3 Hz, 3H).
- (23) Bolesov, I. G.; Gladys, S. A.; Kozmin, S. A.; Levina, R. Y. *J. Org. Chem. USSR* **1970**, *6*, 2431-2434.
 ^1H NMR (500 MHz, CDCl_3) δ = 7.49 (d, J = 9.0 Hz, 2H), 7.31 (d, J = 8.8 Hz, 2H), 5.12 (d, J = 4.7 Hz, 1H), 4.30-4.15 (m, 2H), 3.47 (d, J = 5.3 Hz, 1H), 1.23 (t, J = 7.1 Hz, 3H).
- (24) ^1H NMR (500 MHz, CDCl_3) δ = 8.10 (s, 1H), 7.83 (m, 3H), 7.48 (m, 3H), 4.31-4.15 (m, 2H), 2.33 (dq, J = 14.2, 7.1 Hz, 1H), 2.14 (dq, J = 14.2, 7.1 Hz, 1H), 1.28 (t, J = 7.1 Hz, 3H), 0.97 (t, J = 7.3 Hz, 3H).
- (25) Mallinckrodt Chem. Works US Patent 2349796, **1942**.

- ^1H NMR (500 MHz, CDCl_3) δ = 4.3 (q, J = 7.2 Hz, 2H), 3.2 (br s, 1H), 1.77 (dq, J = 14.8, 7.4 Hz, 1H), 1.66 (dq, J = 14.8, 7.4 Hz, 1H), 1.36 (t, J = 7.2 Hz, 3H), 1.34-1.10 (m, 11H), 0.82 (t, J = 7.4 Hz, 3H).
- (26) ^1H NMR (500 MHz, CDCl_3) δ = 4.2 (q, J = 7.2 Hz, 2H), 3.3 (br s, 1H), 1.9 (q, J = 7.3 Hz, 1H), 1.65 (q, J = 7.3 Hz, 1H), 1.34 (t, J = 7.2 Hz, 3H), 1.1 (s, 9H), 0.80 (t, J = 7.3 Hz, 3H).
- (27) Hartman, B. C.; Livinghouse, T.; Rickborn, B. J. *Org. Chem.* **1973**, 38, 4346-4348.
- (28) Darzens, G. C. R. *Hebd. Seances Acad. Sci.* **1911**, 152, 445-446.
- ^1H NMR (500 MHz, CDCl_3) δ = 4.25 (q, J = 7.1 Hz, 2H), 3.12 (br s, 1H), 1.94 (septet, J = 6.9 Hz, 1H), 1.75 (dq, J = 14.8, 7.4 Hz, 1H), 1.68 (dq, J = 14.8, 7.4 Hz, 2H), 1.31 (t, J = 7.1 Hz, 3H), 0.95 (d, J = 6.9 Hz, 3H), 0.88 (t, J = 6.9 Hz, 3H), 0.83 (d, J = 6.9 Hz, 3H).
- (29) Brown, H. C.; Cho, B. T.; Park, W. S. *J. Org. Chem.* **1986**, 51, 3396-3398.
- ^1H NMR (500 MHz, CDCl_3) δ = 4.29-4.19 (m, 2H), 4.02 (br s, 1H), 2.71 (br s, 1H), 2.07 (m, 1H), 1.30 (t, J = 7.2 Hz, 3H), 1.02 (d, J = 6.9 Hz, 3H), 0.87 (d, J = 6.8 Hz, 3H).
- (30) HP 6890 GC; Supelco β -DEXTM 120 (30 m $\times$ 0.25 mm); pressure = 12.7 psi; flow = 1.0 mL/min
- (31) ^1H NMR (500 MHz, CDCl_3) δ = 7.45-7.05 (m, 4H), 4.29-4.15 (m, 2H), 2.36 (s, 3H), 2.24 (m, 2H), 1.23 (t, J = 7.1 Hz, 3H), 0.98 (t, J = 7.3 Hz, 3H).
- (32) a) Roger, R.; Harper, F. C. *Recl. Trav. Chim. Pays-Bas.* **1937**, 56, 202-207. b) Dhanoa, D. S.; Bagley, S. W.; Chang, R. S. L.; Lotti, V. J.; Chen, T.-B.; Kivlighn, S. D.; Zingaro, G. J.; Siegl, P. K. S.; Chakravarty, P. K.; Patchett, A. A.; Greenlee, W. J. *J. Med. Chem.* **1993**, 36, 3738-3742.
- ^1H NMR (500 MHz, CDCl_3) δ = 7.30-7.17 (m, 4H), 5.36 (s, 1H), 4.31-4.15 (m, 2H), 2.44 (s, 3H), 1.23 (t, J = 7.2 Hz, 3H).
- (33) Higginbotham, L.; Lapworth, A. J. *Chem. Soc.* **1923**, 123, 1325-1332.
- ^1H NMR (500 MHz, CDCl_3) δ = 4.23 (q, J = 5.6 Hz, 2H), 1.85 (q, J = 6.6 Hz, 1H), 1.65 (q, J = 6.6 Hz, 1H), 1.39 (s, 3H), 1.30 (t, J = 6.5 Hz, 3H), 0.87 (t, J = 7.4 Hz, 3H).
- (34) Mahindaratne, M. P. D.; Wimalasena, K. *J. Org. Chem.* **1998**, 63, 2858-2866.
- (35) ^1H NMR (500 MHz, CDCl_3) δ = 7.59-7.20 (m, 10H), 5.24 (d, J = 12.3 Hz, 1H), 5.15 (d, J = 12.3 Hz, 1H), 2.25 (dq, J = 14.6, 7.3 Hz, 1H), 2.05 (dq, J = 14.6, 7.3 Hz, 1H), 0.89 (J = 7.3 Hz, 3H).
- (36) Carpentier, J.-F.; Mortreux, A. *Tetrahedron: Asymmetry* **1997**, 8, 1083-1099.
- (37) For a discussion of nonlinear effects with titanium catalysts see: Mikami, K. *Pure Appl. Chem.* **1996**, 68, 639-644.