Development of a Library of 6-Arylcoumarins as Candidate Fluorescent Sensors

Tomoya Hirano*, Kenichi Hiromoto, Hiroyuki Kagechika*

School of Biomedical Science, Institute of Biomaterials and Bioengineering, Tokyo Medical and Dental University, 2-3-10 Kanda-Surugadai, Chiyoda-ku, Tokyo 101-0062, Japan.

E-mail: hiraomc@tmd.ac.jp, kageomc@tmd.ac.jp

Supporting Information

Materials and Reagents

All reagents were purchased from Sigma-Aldrich Chemical Co., Tokyo Kasei Kogyo Co, Wako Pure Chemical Industries, and Kanto Kagaku Co., Inc. Silica gel for column chromatography was purchased from Kanto Kagaku Co., Inc.. Extracted solutions were dried over anhydrous Na_2SO_4 and solvents were evaporated under reduced pressure. ^1H and ^{13}C NMR spectra were recorded on a Bruker Advance 500 spectrometer. Mass spectral data was obtained on JEOL JMS-T1000LC AccuTOF or Brucker Daltonics microTOF-2focus in the positive and negative ion detection modes. Melting points were taken on a Yanagimoto micro melting point apparatus and are uncorrected. Elemental analyses were carried out by Yanaco MT-6 CHN CORDER spectrometer. UV spectra were recorded with JASCO V-550, and fluorescence spectra were recorded with JASCO FP-6600.

Ethyl 6-bromo-7-(diethylamino)-2-oxo-2*H*-chromene-3-carboxylate **1**.

1

Compound **1** was prepared from 4-(diethylamino)salicylaldehyde according to the procedure described for the reference¹³.

Mp 76-77°C (lit. 74.5-76°C); ^1H NMR (500 MHz, CDCl_3) δ 8.36 (s, 1H), 7.72 (s, 1H), 6.85 (s, 1H), 4.36 (q, J = 7.1 Hz, 2H), 3.28 (q, J = 7.0 Hz, 4H), 1.36 (t, J = 7.1 Hz, 3H), 1.11 (t, J = 7.0 Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ 163.1, 156.6, 155.4, 155.0, 147.3, 134.3, 114.6, 113.5, 112.8, 108.6, 61.6, 45.9 (2C), 14.1, 12.2 (2C); HRMS (ESI) calcd for $\text{C}_{16}\text{H}_{18}\text{BrNO}_4\text{Na}$ ($\text{M}+\text{Na}$)⁺ 390.0317. Found 390.0307.

General Procedure: Pd-Catalyzed Suzuki-Miyaura Coupling.

Ethyl 7-(diethylamino)-2-oxo-6-phenyl-2*H*-chromene-3-carboxylate **3a.**

3a

To a stirred solution of **2a** (34.3 mg, 0.168 mmol), CsF (43.5 mg, 0.286 mmol) and PdCl₂(dppf)·CH₂Cl₂ (9.4 mg, 0.012 mmol) in dry DMF (1 ml) was added **1** (20.3 mg, 0.055 mmol) at room temperature under an atmosphere of argon. The reaction mixture was heated under 60°C for 2h, cooled to room temperature. The resultant mixture was diluted with EtOAc, washed with saturated aqueous NH₄Cl, water and brine, dried, and concentrated. Purification of the residue by flash column chromatography (silica gel, 20% EtOAc/hexane) gave **3a** (14.5 mg, 72%) as yellow cubes. A small amount of this compound was recrystallized in *i*-Pr₂O to give an analytical sample.

Mp 123-124°C; ¹H NMR (500 MHz, CDCl₃) δ 8.44 (s, 1H), 7.40 (m, 4H), 7.31 (m, 1H), 7.28 (s, 1H), 6.85 (s, 1H), 4.37 (q, *J* = 7.1 Hz, 2H), 3.03 (q, *J* = 7.1 Hz, 4H), 1.37 (t, *J* = 7.1 Hz, 3H), 0.95 (t, *J* = 7.0 Hz, 6H); ¹³C NMR (125 MHz, CDCl₃) δ 163.8, 157.6, 156.4, 155.2, 148.8, 140.5, 132.7, 131.6, 128.7 (2C), 128.2 (2C), 127.3, 112.8, 111.1, 105.7, 61.4, 45.8 (2C), 14.3, 12.1 (2C); HRMS (ESI) calcd for C₂₂H₂₃NO₄Na (M+Na)⁺ 388.1525. Found 388.1502; Anal. Calcd for C₂₂H₂₃NO₄: C, 72.31; H, 6.34; N, 3.83. Found: C, 72.33; H, 6.45; N, 3.56.

Ethyl 6-(4-aminophenyl)-7-(diethylamino)-2-oxo-2*H*-chromene-3-carboxylate **3b.**

3b

Compound **3b** was prepared from **2b** according to the procedure described for **3a**. Yield 85% (a pale orange powder). A small amount of this compound was recrystallized in MeOH/*i*-Pr₂O to give an analytical sample.

Mp 170-171°C; ¹H NMR (500 MHz, CDCl₃) δ 8.42 (s, 1H), 7.23 (s, 1H), 7.21 (d, *J* = 8.4 Hz, 2H), 6.80 (s, 1H), 6.73 (d, *J* = 8.3 Hz, 2H), 4.36 (q, *J* = 7.1 Hz, 2H), 3.76 (br s, 2H), 3.05 (q, *J* = 7.1 Hz,

4H), 1.36 (t, J = 7.1 Hz, 3H), 0.95 (t, J = 7.1 Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ 163.8, 157.8, 156.1, 155.4, 148.9, 145.2, 132.3, 131.7, 130.8, 129.1 (2C), 115.5 (2C), 112.4, 111.1, 105.6, 61.4, 45.5 (2C), 14.3, 12.1 (2C); HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{24}\text{N}_2\text{O}_4\text{Na}$ ($\text{M}+\text{Na}$) $^+$ 403.1634. Found 403.1608; Anal. Calcd for $\text{C}_{22}\text{H}_{24}\text{N}_2\text{O}_4 \cdot 1/8\text{H}_2\text{O}$: C, 69.05; H, 6.39; N, 7.32. Found: C, 69.05; H, 6.36; N, 7.34.

Ethyl 6-(4-acetamidophenyl)-7-(diethylamino)-2-oxo-2*H*-chromene-3-carboxylate 3c.

3c

Compound **3c** was prepared from **2c** according to the procedure described for **3a**. Yield 63% (yellow needles). A small amount of this compound was recrystallized in EtOAc to give an analytical sample.

Mp 125-126°C; ^1H NMR (500 MHz, CDCl_3) δ 8.43 (s, 1H), 7.57 (d, J = 8.1 Hz, 2H), 7.48 (s, 1H), 7.38 (d, J = 8.2 Hz, 2H), 7.25 (s, 1H), 6.82 (s, 1H), 4.36 (q, J = 7.1 Hz, 2H), 3.03 (q, J = 7.0 Hz, 4H), 2.19 (s, 3H), 1.36 (t, J = 7.1 Hz, 3H), 0.95 (t, J = 7.0 Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ 168.4, 163.7, 157.7, 156.3, 155.3, 148.8, 137.3, 136.1, 132.6, 131.1, 128.7 (2C), 120.0 (2C), 112.8, 111.2, 105.9, 61.5, 45.7 (2C), 24.6, 14.3, 12.1 (2C); HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{26}\text{N}_2\text{O}_5\text{Na}$ ($\text{M}+\text{Na}$) $^+$ 445.1739. Found 445.1713; Anal. Calcd for $\text{C}_{24}\text{H}_{26}\text{N}_2\text{O}_5$: C, 68.23; H, 6.20; N, 6.63. Found: C, 68.03; H, 6.23; N, 6.57.

Ethyl 7-(diethylamino)-6-(4-(dimethylamino)phenyl)-2-oxo-2*H*-chromene-3-carboxylate 3d.

3d

Compound **3d** was prepared from **2d** according to the procedure described for **3a**. Yield 66% (yellow cubes). A small amount of this compound was recrystallized in $i\text{Pr}_2\text{O}$ to give an analytical sample.

Mp 166-168°C; ^1H NMR (500 MHz, CDCl_3) δ 8.42 (s, 1H), 7.29 (d, J = 8.7 Hz, 2H), 7.24 (s, 1H),

6.80 (s, 1H), 6.73 (d, J = 8.4 Hz, 2H), 4.36 (q, J = 7.1 Hz, 2H), 3.06 (q, J = 7.1 Hz, 4H), 2.97 (s, 6H), 1.36 (t, J = 7.1 Hz, 3H), 0.96 (t, J = 7.1 Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ 163.9, 157.8, 155.9, 155.5, 149.6, 148.9, 132.2, 131.9, 128.7 (2C), 128.2, 112.5 (2C), 112.3, 111.1, 105.6, 61.3, 45.5 (2C), 40.4 (2C), 14.3, 12.1 (2C); HRMS (ESI) calcd for $\text{C}_{24}\text{H}_{28}\text{N}_2\text{O}_4\text{Na}$ ($\text{M}+\text{Na}$) $^+$ 431.1947. Found 431.1930; Anal. Calcd for $\text{C}_{24}\text{H}_{28}\text{N}_2\text{O}_4$: C, 70.57; H, 6.91; N, 6.86. Found: C, 70.41; H, 6.86; N, 6.86.

Ethyl 7-(diethylamino)-6-(4-nitrophenyl)-2-oxo-2*H*-chromene-3-carboxylate 3e.

3e

Compound **3e** was prepared from **2e** according to the procedure described for **3a**. Yield 28% (a pale yellow powder). A small amount of this compound was recrystallized in $i\text{-Pr}_2\text{O}$ to give an analytical sample.

Mp 123-124°C; ^1H NMR (500 MHz, CDCl_3) δ 8.45 (s, 1H), 8.28 (d, J = 8.8 Hz, 2H), 7.64 (d, J = 8.8 Hz, 2H), 7.33 (s, 1H), 6.91 (s, 1H), 4.38 (q, J = 7.1 Hz, 2H), 3.03 (q, J = 7.1 Hz, 4H), 1.38 (t, J = 7.2 Hz, 3H), 0.98 (t, J = 7.1 Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ 163.5, 157.1, 156.9, 154.9, 148.4, 147.1, 147.0, 132.6, 129.4, 129.0 (2C), 124.2 (2C), 114.0, 111.6, 106.7, 61.7, 46.1 (2C), 14.3, 11.9 (2C); HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{22}\text{N}_2\text{O}_6\text{Na}$ ($\text{M}+\text{Na}$) $^+$ 433.1376. Found 433.1381; Anal. Calcd for $\text{C}_{22}\text{H}_{22}\text{N}_2\text{O}_6 \cdot 1/4\text{H}_2\text{O}$: C, 63.68; H, 5.47; N, 6.75. Found: C, 63.90; H, 5.45; N, 6.49.

Ethyl 7-(diethylamino)-6-(4-hydroxyphenyl)-2-oxo-2*H*-chromene-3-carboxylate 3f.

3f

Compound **3f** was prepared from **2f** according to the procedure described for **3a**. Yield 74% (yellow needles). A small amount of this compound was recrystallized in EtOAc to give an analytical sample.

Mp 112-113°C; ^1H NMR (500 MHz, CDCl_3) δ 8.44 (s, 1H), 7.29 (d, J = 8.6 Hz, 2H), 7.25 (s, 1H), 6.88 (d, J = 8.6 Hz, 2H), 6.84 (s, 1H), 5.15 (br s, 1H), 4.37 (q, J = 7.1 Hz, 2H), 3.04 (q, J = 7.1 Hz,

4H), 1.37 (t, $J = 7.1$ Hz, 3H), 0.96 (t, $J = 7.1$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ 163.9, 158.1, 156.2, 155.5, 155.4, 149.1, 132.6, 132.5, 131.5, 129.4 (2C), 115.8 (2C), 112.3, 111.1, 105.6, 61.5, 45.6 (2C), 14.3, 12.1 (2C); HRMS (ESI) calcd for $\text{C}_{22}\text{H}_{22}\text{NO}_5$ ($\text{M}-\text{H}$) $^-$ 380.1498. Found 380.1491; Anal. Calcd for $\text{C}_{22}\text{H}_{23}\text{NO}_5$: C, 69.28; H, 6.08; N, 3.67. Found: C, 69.02; H, 6.20; N, 3.65.

Ethyl 7-(diethylamino)-6-(4-methoxyphenyl)-2-oxo-2*H*-chromene-3-carboxylate 3g.

3g

Compound **3g** was prepared from **2g** according to the procedure described for **3a**. Yield 63% (a pale yellow needles). A small amount of this compound was recrystallized in $i\text{-Pr}_2\text{O}$ to give an analytical sample.

Mp 70-71°C; ^1H NMR (500 MHz, CDCl_3) δ 8.43 (s, 1H), 7.35 (d, $J = 8.8$ Hz, 2H), 7.26 (s, 1H), 6.93 (d, $J = 8.8$ Hz, 2H), 6.85 (s, 1H), 4.37 (q, $J = 7.1$ Hz, 2H), 3.84 (s, 3H), 3.04 (q, $J = 7.1$ Hz, 4H), 1.37 (t, $J = 7.1$ Hz, 3H), 0.96 (t, $J = 7.1$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ 163.8, 159.0, 157.6, 156.2, 155.1, 148.8, 132.7, 132.5, 131.4, 129.3 (2C), 114.2 (2C), 112.9, 111.3, 106.0, 61.5, 55.3, 45.8 (2C), 14.3, 12.1 (2C); HRMS (ESI) calcd for $\text{C}_{23}\text{H}_{25}\text{NO}_5\text{Na}$ ($\text{M}+\text{Na}$) $^+$ 418.1630. Found 418.1608rt; Anal. Calcd for $\text{C}_{23}\text{H}_{25}\text{NO}_5$: C, 69.86; H, 6.37; N, 3.54. Found: C, 69.62; H, 6.40; N, 3.54.

Ethyl 6-(4-acetylphenyl)-7-(diethylamino)-2-oxo-2*H*-chromene-3-carboxylate 3h.

3h

Compound **3h** was prepared from **2h** according to the procedure described for **3a**. Yield 58% (a pale yellow cubes). A small amount of this compound was recrystallized in $i\text{-Pr}_2\text{O}$ to give an analytical sample.

Mp 79-80°C; ^1H NMR (500 MHz, CDCl_3) δ 8.44 (s, 1H), 8.00 (d, $J = 8.4$ Hz, 2H), 7.55 (d, $J = 8.4$ Hz, 2H), 7.30 (s, 1H), 6.87 (s, 1H), 4.37 (q, $J = 7.1$ Hz, 2H), 3.02 (q, $J = 7.1$ Hz, 4H), 2.62 (s, 3H), 1.37 (t, $J = 7.1$ Hz, 3H), 0.96 (t, $J = 7.1$ Hz, 6H); ^{13}C NMR (125 MHz, CDCl_3) δ 197.4, 163.6, 157.3, 156.7,

155.0, 148.6, 145.4, 136.0, 132.6, 130.4, 128.9 (2C), 128.4 (2C), 113.4, 111.3, 106.2, 61.6, 45.9 (2C), 26.6, 14.3, 12.0 (2C); HRMS (ESI) calcd for $C_{24}H_{25}NO_5Na$ ($M+Na$)⁺ 430.1630. Found 430.1606; Anal. Calcd for $C_{24}H_{25}NO_5$: C, 70.74; H, 6.18; N, 3.44. Found: C, 70.55; H, 6.29; N, 3.42.

Ethyl 6-(4-cyanophenyl)-7-(diethylamino)-2-oxo-2*H*-chromene-3-carboxylate 3i.

3i

Compound **3i** was prepared from **2i** according to the procedure described for **3a**. Yield 45% (pale yellow needles). A small amount of this compound was recrystallized in EtOAc to give an analytical sample.

Mp 174-176°C; ¹H NMR (500 MHz, CDCl₃) δ 8.44 (s, 1H), 7.70 (d, *J* = 8.3 Hz, 2H), 7.58 (d, *J* = 8.3 Hz, 2H), 7.29 (s, 1H), 6.89 (s, 1H), 4.37 (q, *J* = 7.1 Hz, 2H), 3.01 (q, *J* = 7.1 Hz, 4H), 1.37 (t, *J* = 7.1 Hz, 3H), 0.97 (t, *J* = 7.0 Hz, 6H); ¹³C NMR (125 MHz, CDCl₃) δ 163.5, 157.2, 156.8, 154.9, 148.5, 145.2, 132.6 (2C), 132.6, 129.7, 128.9 (2C), 118.6, 113.7, 111.5, 111.2, 106.5, 61.6, 45.9 (2C), 14.3, 11.9 (2C); HRMS (ESI) calcd for $C_{23}H_{22}N_2O_4Na$ ($M+Na$)⁺ 413.1477. Found 413.1462; Anal. Calcd for $C_{23}H_{22}N_2O_4$: C, 70.75; H, 5.68; N, 7.17. Found: C, 70.61; H, 5.75; N, 6.99.

Ethyl 7-(diethylamino)-6-(4-(methoxycarbonyl)phenyl)-2-oxo-2*H*-chromene-3-carboxylate 3j.

3j

Compound **3j** was prepared from **2j** according to the procedure described for **3a**. Yield 46% (yellow cubes). A small amount of this compound was recrystallized in *i*-Pr₂O to give an analytical sample.

Mp 115-117°C; ¹H NMR (500 MHz, CDCl₃) δ 8.44 (s, 1H), 8.07 (d, *J* = 8.2 Hz, 2H), 7.52 (d, *J* = 8.3 Hz, 2H), 7.31 (s, 1H), 6.88 (s, 1H), 4.37 (q, *J* = 7.1 Hz, 2H), 3.93 (s, 3H), 3.02 (q, *J* = 7.1 Hz, 4H), 1.37 (t, *J* = 7.1 Hz, 3H), 0.96 (t, *J* = 7.0 Hz, 6H); ¹³C NMR (125 MHz, CDCl₃) δ 166.7, 163.6, 157.3, 156.6, 155.0, 148.6, 145.2, 132.6, 130.4, 130.1 (2C), 129.1, 128.2 (2C), 113.2, 111.2, 106.0, 61.5, 52.1, 45.8 (2C), 14.2, 12.0 (2C); HRMS (ESI) calcd for $C_{24}H_{25}NO_6Na$ ($M+Na$)⁺ 446.1580. Found

446.1583; Anal. Calcd for C₂₄H₂₅NO₆: C, 68.07; H, 5.95; N, 3.31. Found: C, 67.78; H, 5.94; N, 3.30.

Ethyl 6-(4-(benzyloxycarbonyl)phenyl)-7-(diethylamino)-2-oxo-2*H*-chromene-3-carboxylate 3k.

3k

Compound **3k** was prepared from **2k** according to the procedure described for **3a**. Yield 63% (pale yellow cubes). A small amount of this compound was recrystallized in *i*-Pr₂O to give an analytical sample.

Mp 121-123°C; ¹H NMR (500 MHz, CDCl₃) δ 8.44 (s, 1H), 8.11 (d, *J* = 8.4 Hz, 2H), 7.52 (d, *J* = 8.5 Hz, 2H), 7.46 (d, *J* = 7.0 Hz, 2H), 7.39 (m, 2H), 7.34 (m, 1H), 7.30 (s, 1H), 6.86 (s, 1H), 5.37 (s, 2H), 4.37 (q, *J* = 7.1 Hz, 2H), 3.01 (q, *J* = 7.1 Hz, 4H), 1.37 (t, *J* = 7.1 Hz, 3H), 0.95 (t, *J* = 7.1 Hz, 6H); ¹³C NMR (125 MHz, CDCl₃) δ 166.1, 163.6, 157.4, 156.7, 155.0, 148.6, 145.3, 136.0, 132.6, 130.4, 130.2 (2C), 129.1, 128.6 (2C), 128.33, 128.27 (2C), 128.2 (2C), 113.3, 111.3, 106.1, 66.8, 61.5, 45.9 (2C), 14.3, 12.0 (2C); HRMS (ESI) calcd for C₃₀H₂₉NO₆Na (M+Na)⁺ 522.1893. Found 522.1888; Anal. Calcd for C₃₀H₂₉NO₆ · 1/8H₂O: C, 71.80; H, 5.88; N, 2.79. Found: C, 71.70; H, 6.01; N, 2.81.

Ethyl 6-(4-amino-3-nitrophenyl)-7-(diethylamino)-2-oxo-2*H*-chromene-3-carboxylate 3l.

3l

Compound **3l** was prepared from **2l** according to the procedure described for **3a**. Yield 62% (a pale orange powder). A small amount of this compound was recrystallized in EtOAc/hexane to give an analytical sample.

Mp 187-188°C; ¹H NMR (500 MHz, CDCl₃) δ 8.44 (s, 1H), 8.23 (d, *J* = 2.1 Hz, 1H), 7.52 (dd, *J* = 8.6, 2.1 Hz, 1H), 7.31 (s, 1H), 6.89 (s, 1H), 6.86 (d, *J* = 8.6 Hz, 1H), 6.21 (br s, 2H), 4.37 (q, *J* = 7.2 Hz, 2H), 3.06 (q, *J* = 7.1 Hz, 4H), 1.37 (t, *J* = 7.1 Hz, 3H), 0.99 (t, *J* = 7.1 Hz, 6H); ¹³C NMR (125 MHz, CDCl₃) δ 163.5, 157.4, 156.2, 155.2, 148.6, 144.0, 135.8, 132.1, 132.1, 129.8, 128.9, 125.0, 119.2, 113.3, 111.7, 106.7, 61.5, 45.6 (2C), 14.2, 11.9 (2C); HRMS (ESI) calcd for C₂₂H₂₂N₃O₆ (M-H)⁻

424.1509. Found 424.1513; Anal. Calcd for $C_{22}H_{23}N_3O_6$: C, 62.11; H, 5.45; N, 9.88. Found: C, 62.02; H, 5.65; N, 9.76.

4-(7-(Diethylamino)-3-(ethoxycarbonyl)-2-oxo-2*H*-chromene-6-yl)benzoic acid **3m.**

3m

To a stirred solution of **3k** (110.4 mg, 0.221 mmol) in MeOH/THF (2:1, v/v, 3 ml) was added 10% Pd/C (17.0 mg) at room temperature under an atmosphere of argon. H_2 gas was introduced to the reaction mixture. After being stirred for 40 min, the resultant mixture was filtered through Hyflo Super-Cel®. The filtrate was concentrated. Purification of the residue by flash column chromatography (silica gel, 50% EtOAc/hexane) gave **3m** (66.0 mg, 73%) as a pale yellow powder. A small amount of this compound was recrystallized in EtOAc to give an analytical sample.

Mp 249-250°C; 1H NMR (500 MHz, $CDCl_3$) δ 11.06 (br s, 1H), 8.46 (s, 1H), 8.15 (d, J = 8.3 Hz, 2H), 7.56 (d, J = 8.3 Hz, 2H), 7.33 (s, 1H), 6.86 (s, 1H), 4.36 (q, J = 7.1 Hz, 2H), 3.03 (q, J = 7.1 Hz, 4H), 1.36 (t, J = 7.1 Hz, 3H), 0.96 (t, J = 7.1 Hz, 6H); ^{13}C NMR (125 MHz, $CDCl_3$) δ 170.9, 163.6, 157.4, 156.7, 154.9, 148.6, 146.1, 132.7, 130.8 (2C), 130.3, 128.4 (2C), 128.1, 113.4, 111.4, 106.2, 61.6, 46.0 (2C), 14.3, 12.0 (2C); HRMS (ESI) calcd for $C_{23}H_{22}NO_6$ ($M-H$) $^-$ 408.1447. Found 408.1450; Anal. Calcd for $C_{23}H_{23}NO_6 \cdot 1/4H_2O$: C, 66.74; H, 5.72; N, 3.38. Found: C, 66.98; H, 5.76; N, 3.35.

Ethyl 6-(3,4-diaminophenyl)-7-(diethylamino)-2-oxo-2*H*-chromene-3-carboxylate **3n.**

3n

To a stirred solution of **3l** (120.9 mg, 0.284 mmol) in MeOH (2.5 ml) was added 10% Pd/C (39.1 mg) at room temperature under an atmosphere of argon. H_2 gas was introduced to the reaction mixture. After being stirred for 2 h, the resultant mixture was filtered through Hyflo Super-Cel®. The filtrate was concentrated. Purification of the residue by flash column chromatography (silica gel, 2.5% MeOH/ $CHCl_3$) gave **3n** (72.5 mg, 65%) as an orange powder. A small amount of this

compound was recrystallized in MeOH/*i*-Pr₂O to give an analytical sample.

Mp 185-186°C; ¹H NMR (500 MHz, CDCl₃) δ 8.41 (s, 1H), 7.23 (s, 1H), 6.82-6.75 (m, 4H), 4.36 (q, *J* = 7.1 Hz, 2H), 3.68 (br s, 4H), 3.07 (q, *J* = 7.1 Hz, 4H), 1.37 (t, *J* = 7.1 Hz, 3H), 0.96 (t, *J* = 7.1 Hz, 6H); ¹³C NMR (125 MHz, CDCl₃) δ 163.9, 157.9, 156.1, 155.3, 149.0, 134.7, 134.0, 132.5, 132.4, 131.5, 120.0, 116.9, 116.4, 112.0, 110.8, 105.2, 61.3, 45.5 (2C), 14.2, 12.1 (2C); HRMS (ESI) calcd for C₂₂H₂₄N₃O₄ (M-H)⁻ 394.1767. Found 394.1760; Anal. Calcd for C₂₂H₂₅N₃O₄ · 1/4H₂O: C, 66.07; H, 6.43; N, 10.51. Found: C, 66.01; H, 6.27; N, 10.44.

Ethyl 6-(1*H*-benzo[*d*]1,2,3-triazol-5-yl)-7-(diethylamino)-2-oxo-2*H*-chromene-3- carboxylate 3o.

3o

To a stirred solution of **3n** (47.8 mg, 0.131 mmol) in MeOH/H₂O (1:4, v/v, 2 ml) was added AcOH (0.2 ml) and NaNO₂ (21.4 mg 0.310 mmol) at room temperature. After being stirred for 5 min, the reaction mixture was diluted with EtOAc. The organic layer was washed with water and brine, dried, and concentrated. Purification of the residue by flash column chromatography (silica gel, 50% EtOAc/hexane) gave **3o** (50.2 mg, 94%) as a yellow powder. A small amount of this compound was recrystallized in MeOH/*i*-Pr₂O to give an analytical sample.

Mp 232-233°C; ¹H NMR (500 MHz, CDCl₃) δ 8.51 (s, 1H), 8.02 (s, 1H), 7.98 (d, *J* = 8.6 Hz, 1H), 7.55 (dd, *J* = 8.6, 1.0 Hz, 1H), 7.41 (s, 1H), 6.91 (s, 1H), 4.38 (q, *J* = 7.1 Hz, 2H), 3.05 (q, *J* = 7.1 Hz, 4H), 1.36 (t, *J* = 7.1 Hz, 3H), 0.96 (t, *J* = 7.0 Hz, 6H); ¹³C NMR (125 MHz, CDCl₃) δ 163.7, 158.1, 156.5, 155.2, 149.0, 139.3, 138.4, 138.3, 133.4, 131.1, 127.0, 115.5, 114.1, 112.5, 111.4, 106.0, 61.7, 52.2, 46.0 (2C), 14.3, 12.1 (2C); HRMS (ESI) calcd for C₂₂H₂₁N₄O₄ (M-H)⁻ 405.1563. Found 405.1564; Anal. Calcd for C₂₂H₂₂N₄O₄: C, 65.01; H, 5.46; N, 13.78. Found: C, 64.72; H, 5.60; N, 13.68.

Table S1. Absorbance and Fluorescence maximum wavelengths of 6-arylcoumarin compounds.

Solvent	CH ₃ CN			CH ₃ OH			Water		
	Abs. Max.	Ex. Max.	Em. Max.	Abs. Max.	Ex. Max.	Em. Max.	Abs. Max.	Ex. Max.	Em. Max.
3a	410	407	462	415	407	462	431	401	465
3b	419	n.d.	n.d.	423	n.d.	n.d.	436	n.d.	n.d.
3c	411	403	471	416	410	477	431	410	480
3d	420	n.d.	n.d.	425	n.d.	n.d.	435	n.d.	n.d.
3e	401	392	444	404	397	449	425	408	457
3f	414	408	465	421	417	475	433	414	482
3g	412	406	466	418	414	472	433	418	484
3h	406	404	458	411	411	467	427	409	474
3i	403	399	454	406	404	458	427	400	464
3j	406	402	455	410	409	463	427	415	467
3l	409	n.d.	n.d.	412	n.d.	n.d.	428	n.d.	n.d.
3m	407	400	454	415	410	465	430	411	470
3n	420	n.d.	n.d.	425	n.d.	n.d.	437	n.d.	n.d.
3o	408	407	462	412	410	466	430	404	477

Absorbance maximum wavelength and excitation, emission maximum wavelength of each compound (3 μ M) was measured in acetonitrile, methanol and water (1 mM sodium phosphate buffer (pH 7.4)). All solutions contained 0.3% DMSO as a cosolvent. n.d.: no fluorescent peak was detected.

1

1

3a

3a

3b

3b

3c

3c

3d

3d

3e

3e

3f

3f

3g

3g

3h

3h

3i

3i

3j

3j

3k

3k

31

31

3m

3m

3n

3n

30

30