Supporting Information

Development of a New Class of Chiral Phosphorus Ligands: P-Chirogenic Diaminophosphine Oxides. A Unique Source of Enantioselection in Pd-Catalyzed Asymmetric Construction of Quaternary Carbons

Tetsuhiro Nemoto, Takamasa Masuda, Takayoshi Matsumoto, and Yasumasa Hamada*

Contents

General (S-1)

1. Experimental and Analytical Data for the Mechanistic Studies (S-2 ~ S-10)

- A. Nonlinear Effect
- B. Initial Rate Kinetics
- C. BSA-Induced P(V) to P(III) Transformation of Diaminophosphine Oxides **12**
- D. FAB MS and ^{31}P NMR Analyses
- E. Discussions

2. General Procedures, Characterization of the New Compounds, and Discussion (S-11 ~ S-14)

- A. Asymmetric Construction of Quaternary Carbons Using Cyclic Nucleophiles
- B. Asymmetric Construction of Quaternary Carbons Using Acyclic Nucleophiles
- C. Asymmetric Allylic Alkylation of **10** with Dimethyl Malonate

3. Experimental Procedures for the Synthesis of P-Chirogenic Diaminophosphine Oxides (S-15 ~ S-22)

- A. Synthesis of the Aspartic Acid-Derived P-Chirogenic Diaminophosphine Oxides (DIAPHOXs)
- B. Synthesis of the Glutamic Acid-Derived P-Chirogenic Diaminophosphine Oxide **24**
- C. Synthesis of the Homophenylalanine-Derived P-Chirogenic Diaminophosphine Oxide **25** and **26**

4. Analytical Data (S-23 ~ S-51)

- A. FAB MS and ^{31}P NMR Spectrum for the Structural Analysis
- B. ^{13}C NMR Spectrum for the Purity Documentation (all new compounds in the asymmetric reaction and the ligand synthesis, and known compounds synthesized by a new method)

General: Infrared (IR) spectra were recorded on FT/IR 230 Fourier transform infrared spectrophotometer (JASCO). NMR spectra were recorded on ecp 400 spectrometer (JEOL), operating at 400 MHz for ^1H NMR, 100 MHz for ^{13}C NMR, and 160 MHz for ^{31}P NMR. Chemical shifts in CDCl_3 , and $\text{DMSO}-d_6$ were reported downfield from TMS (= 0 ppm) for ^1H NMR. For ^{13}C NMR, chemical shifts were reported downfield from TMS (= 0 ppm) or in the scale relative to the solvent signal [CHCl_3 (77.0 ppm) and DMSO (39.5 ppm)] as an internal reference. For ^{31}P NMR, chemical shifts were reported in the scale relative to H_3PO_4 signal (0 ppm in CDCl_3) as an internal reference. Optical rotations were measured on P-1020 polarimeter (JASCO). FAB mass spectra were measured on AX500 (JMX). EI mass spectra were measured on Gcmate (JEOL). The enantiomeric excess (ee) was determined by HPLC analysis. HPLC was performed on HPLC

systems consisting of the following: pump, PU-980 (JASCO); detector, UV-970 (JASCO), measured at 254 nm; column, DAICEL CHIRALPAK AD, DAICEL CHIRALCEL OD-H, DAICEL CHIRALCEL OJ; mobile phase, hexane–2-propanol; flow rate, 0.3–0.8 mL/min. Reactions were carried out in dry solvent under argon atmosphere. Other reagents were purified by the usual methods.

1. Experimental and Analytical Data for the Mechanistic Studies

A. Nonlinear Effect

General Procedure: Reaction Type A (**10** to **11**, 60% ee of **6**)

To a stirred mixture of 1,3-diphenyl-2-propenyl acetate (**10**) (102.5 mg, 0.406 mmol), dimethyl malonate (140 μ L, 1.22 mmol), Pd cat (1.48 mg, 0.002 mmol), **6** (5.08 mg, 0.013 mmol), enantiomer of **6** (1.27 mg, 0.00325 mmol) in CH_2Cl_2 (1.6 mL) was added BSA (300 μ L, 1.22 mmol) at room temperature. After being stirred for 24 h, the reaction was concentrated under reduced pressure, and the obtained residue was purified by flash column chromatography (SiO_2 , hexane/ethyl acetate 30/1) to give (1*S*)-1,3-diphenyl-2-propenyl-propanedioic acid dimethyl ester (**11**) as colorless oil (129 mg, 99%, 80% ee).

General Procedure: Reaction Type B (**7a** to **9a**: 60% ee of **6**)

To a stirred mixture of cinnamyl acetate (**7a**) (33.3 μ L, 0.2 mmol), keto ester **8a** (48 μ L, 0.3 mmol), Pd cat (1.83 mg, 0.005 mmol), **6** (6.3 mg, 0.016 mmol), enantiomer of **6** (1.6 mg, 0.004 mmol) and $\text{Zn}(\text{OAc})_2$ (3.7 mg, 0.02 mmol) in toluene (1.6 mL) was added BSA (198 μ L, 0.8 mmol) at room temperature. After being stirred for 8 h, the reaction was concentrated under reduced pressure, and the obtained residue was purified by flash column chromatography (SiO_2 , hexane/ethyl acetate 30/1 to 15/1) to give (1*S*)-2-oxo-1-(3-phenyl-allyl)-cyclohexanecarboxylic acid ethyl ester (**7a**) as colorless oil (55.6 mg, 97%, 66% ee).

Table S-1. and **Table S-2.** Asymmetric Amplification

Table S-1.

$(\eta^3\text{-C}_3\text{H}_5\text{PdCl})_2$ (1 mol %) 6 (X % ee, 4 mol %) BSA (3.0 eq), dimethyl malonate (3.0 eq) 10 $\xrightarrow{\text{CH}_2\text{Cl}_2$ (0.25 M), rt, 24 h} 11		
Run	Ligand ee	Product ee
1	0% ee	0%
2	20% ee	31% ee
3	40% ee	54% ee
4	60% ee	80% ee
5	80% ee	95% ee
6	100% ee	98% ee

Table S-2.

$(\eta^3\text{-C}_3\text{H}_5\text{PdCl})_2$ (2.5 mol %) 6 (X % ee, 10 mol %) $\text{Zn}(\text{OAc})_2$ (10 mol %) BSA (4.0 eq), 8a (1.5 eq) 7a $\xrightarrow{\text{toluene}$ (0.125 M), rt, 8 h} 9a		
Run	Ligand ee	Product ee
1	0% ee	0%
2	20% ee	26% ee
3	40% ee	55% ee
4	60% ee	66% ee
5	80% ee	82% ee
6	100% ee	92% ee

Figure S-1. Nonlinear Effect

B. Initial Rate Kinetics

General Procedure for Kinetic Experiments: Reaction Type A

[Pd cat.] = 0.004 M, [Acetate **10**] = 0.197 M, [dimethyl malonate] = 0.595 M, [BSA] = 0.6 M (2 mol % catalyst).

To a stirred mixture of **10** (240 μ L, 1 mmol), Pd cat (3.7 mg, 0.01 mmol), **6** (15.7 mg, 0.04 mmol), in CH_2Cl_2 (5 mL) was added BSA (740 μ L, 3 mmol) at 25 $^\circ\text{C}$. After being stirred for 30 min, dimethyl malonate (340 μ L, 3 mmol) was added to the reaction, and the resulting mixture was stirred at the same temperature. Aliquots were taken at recorded times according to the following procedure. Approximately 0.4 mL of the reaction mixture was withdrawn with a syringe, and was immediately introduced into a test tube containing approximately 1.0 mL of water and 2.0 mL of ethyl acetate. The obtained mixture was thoroughly agitated, and the organic layer was withdrawn. After the solvent was evaporated in vacuo, the obtained crude residue was analyzed by ^1H NMR. Yields of the products were measured by comparison of integrated area of the olefinic protons of **10** and **11**.

General Procedure for Kinetic Experiments : Reaction Type B

[Pd cat.] = 0.003 M, [Acetate **7a**] = 0.15 M, [Keto Ester **8a**] = 0.225 M, [$\text{Zn}(\text{OAc})_2$] = 0.015 M, [BSA] = 0.6 M (2 mol % catalyst).

To a stirred mixture of cinnamyl acetate (**7a**) (100 μ L, 0.6 mmol), Pd cat (2.2 mg, 0.006 mmol), **6** (9.4 mg, 0.024 mmol), and $\text{Zn}(\text{OAc})_2$ (11.0 mg, 0.06 mmol) in toluene (4 mL) was added BSA (593 μ L, 2.4 mmol) at 30 $^\circ\text{C}$. After being stirred for 30 min, keto ester **8a** was added to the reaction, and the resulting mixture was stirred at the same temperature. Aliquots were taken at recorded times according to the following procedure. Approximately 0.4 mL of the reaction mixture was withdrawn with a syringe, and was immediately introduced into a test tube containing approximately 1.0 mL of water and 2.0 mL of ethyl acetate. The obtained mixture was thoroughly agitated, and the organic layer was withdrawn. After the solvent was evaporated in vacuo, the obtained crude residue was analyzed by ^1H NMR. Yields of the products were measured by comparison of integrated area of the olefinic protons of **7a** and **9a**.

Note: The procedure for kinetics is slightly different from the one of asymmetric reaction. First, we carried out kinetic studies using the asymmetric reaction procedure. However, the obtained reaction profile revealed that no reaction occurred in the first stage of the reaction (~ 15 min). We expected that this delay might be due to the gradual ligand tautomerization induced by BSA. Thus, in the kinetics procedure, reactions were stirred for 30 min in the presence of BSA, before the addition of dimethyl malonate or keto ester **8a**.

Kinetics Data (Reaction Type A)

Kinetics on the Catalyst

Initial Rate

0.004 M: $1.3241 \times 10^{-4} \text{ Ms}^{-1}$
 0.002 M: $0.6641 \times 10^{-4} \text{ Ms}^{-1}$
 0.0014 M: $0.4913 \times 10^{-4} \text{ Ms}^{-1}$

First Order Dependency on the Catalyst.

[cat.] = 0.0014 ~ 0.004 M, [10] = 0.197 M, [dimethyl malonate] = 0.595 M

Kinetics on the Acetate 10

Initial Rate

0.1313 M: $0.8825 \times 10^{-4} \text{ Ms}^{-1}$
 0.197 M: $0.9073 \times 10^{-4} \text{ Ms}^{-1}$
 0.2626 M: $0.8498 \times 10^{-4} \text{ Ms}^{-1}$

Zero Order Dependency on the Acetate **10**.

[10] = 0.1313~ 0.2626 M, [cat.] = 0.004 M, [dimethyl malonate] = 0.595 M

Kinetics on Dimethyl Malonate

Initial Rate

0.298 M: $0.6361 \times 10^{-4} \text{ Ms}^{-1}$
 0.437 M: $0.9073 \times 10^{-4} \text{ Ms}^{-1}$
 0.595 M: $1.3241 \times 10^{-4} \text{ Ms}^{-1}$

First Order Dependency on dimethyl malonate.

[dimethyl malonate] = 0.298 ~ 0.595 M, [cat.] = 0.004 M, [10] = 0.197 M

Reaction Profile: Initial Rate $V = d/dt [11] = K_{rds} \cdot [\text{cat.}]^1 \cdot [\text{acetate } 10]^0 \cdot [\text{dimethyl malonate}]^1$

Rate Determining Step: Nucleophilic addition of dimethyl malonate to the π -allylpalladium (II) intermediate.

Kinetics Data (Reaction Type B)

Kinetics on the Catalyst

[cat.] = 0.0075 ~ 0.003 M, [7a] = 0.15 M,
[8a] = 0.225 M, [Zn(OAc)₂] = 0.015 M

Initial Rate

0.003 M: $y = 3.3633x - 12.14$ ($R^2 = 0.9997$); $8.4083 \times 10^{-5} \text{ Ms}^{-1}$
0.00225 M: $y = 2.31x - 12.16$ ($R^2 = 0.9982$); $5.7750 \times 10^{-5} \text{ Ms}^{-1}$
0.0015 M: $y = 1.8225x - 22.67$ ($R^2 = 0.9924$); $4.5563 \times 10^{-5} \text{ Ms}^{-1}$
0.00113 M: $y = 1.206x - 8.730$ ($R^2 = 0.9963$); $3.0150 \times 10^{-5} \text{ Ms}^{-1}$
0.00075 M: $y = 0.7062x - 8.367$ ($R^2 = 0.9987$); $1.7655 \times 10^{-5} \text{ Ms}^{-1}$

First Order Dependency
on the Catalyst.

Kinetics on Cinnamyl Acetate

[7a] = 0.05 M ~ 0.15 M, [cat.] = 0.0015 M, [8a] = 0.225 M,
[Zn(OAc)₂] = 0.015 M

Initial Rate

0.150 M: $y = 0.0018x - 0.0182$ ($R^2 = 0.9918$); $2.9688 \times 10^{-5} \text{ Ms}^{-1}$
0.120 M: $y = 0.0018x - 0.0131$ ($R^2 = 0.9916$); $2.9200 \times 10^{-5} \text{ Ms}^{-1}$
0.100 M: $y = 0.0018x - 0.0073$ ($R^2 = 0.9999$); $2.9458 \times 10^{-5} \text{ Ms}^{-1}$
0.075 M: $y = 0.0018x - 0.0081$ ($R^2 = 0.9737$); $2.9666 \times 10^{-5} \text{ Ms}^{-1}$
0.050 M: $y = 0.0019x - 0.011$ ($R^2 = 0.9905$); $3.1875 \times 10^{-5} \text{ Ms}^{-1}$

Zero Order Dependency
on cinnamyl acetate.

Kinetics on the Keto Ester 8a

[8a] = 0.15 M ~ 0.3 M, [cat.] = 0.0015 M, [7a] = 0.1 M,
[Zn(OAc)₂] = 0.015 M

Initial Rate

0.3 M: $y = 0.0021x - 0.0153$ ($R^2 = 0.9589$); $3.4667 \times 10^{-5} \text{ Ms}^{-1}$
0.225 M: $y = 0.0018x - 0.0073$ ($R^2 = 0.9999$); $2.9458 \times 10^{-5} \text{ Ms}^{-1}$
0.18 M: $y = 0.0012x - 0.0065$ ($R^2 = 0.9999$); $1.9732 \times 10^{-5} \text{ Ms}^{-1}$
0.15 M: $y = 0.0011x - 0.0087$ ($R^2 = 0.9949$); $1.7708 \times 10^{-5} \text{ Ms}^{-1}$

First Order Dependency
on the Keto Ester 8a.

Kinetics on Zn(OAc)₂

[Zn(OAc)₂] = 0.00225 M ~ 0.015 M, [cat.] = 0.0015 M,
[**7a**] = 0.15 M, [**8a**] = 0.225 M

Initial Rate

10 mol % (0.015 M): $y = 1.1875x - 12.15$ ($R^2 = 0.9918$); $2.9657 \times 10^{-5} \text{ Ms}^{-1}$

6 mol % (0.009 M): $y = 0.9696x - 2.9357$ ($R^2 = 0.9946$); $2.4240 \times 10^{-5} \text{ Ms}^{-1}$

3 mol % (0.0045 M): $y = 0.70x - 3.9$ ($R^2 = 0.9865$); $1.7500 \times 10^{-5} \text{ Ms}^{-1}$

1.5 mol % (0.00225 M): $y = 0.285x - 4.3$ ($R^2 = 0.9567$); $0.7125 \times 10^{-5} \text{ Ms}^{-1}$

First Order Dependency
on Zn(OAc)₂.

Reaction Profile: Initial Rate $V = d/dt [\mathbf{9a}] = K_{\text{rds}} \cdot [\text{cat.}]^1 \cdot [\text{acetate } \mathbf{7a}]^0 \cdot [\text{keto ester } \mathbf{8a}]^1 \cdot [\text{Zn(OAc)}_2]^1$

Rate Determining Step: Nucleophilic addition of the nucleophile to the π -allylpalladium (II) intermediate.

C. BSA-Induced P(V) to P(III) Transformation of Diaminophosphine Oxide **12**

General Procedure for NMR Experiments: To a stirred solution of 1,3-diphenyl-[1,3,2]diazaphospholidine 2-oxide (**12**) (5.2 mg, 0.02 mmol) in CDCl₃ (1.0 mL) was added BSA (100 μL , 0.4 mmol) at room temperature, and the solution was stirred for 15 min. After transferring to a NMR tube by a syringe, ³¹P NMR analysis was performed.

Compound 12: white brown solid; Mp 171–172 °C; IR (neat) ν 3403, 1601, 1491, 1475, 1285, 1156, 1134, 1036, 982, 750, 691 cm⁻¹; ¹H NMR (CDCl₃): δ 3.81–3.85 (m, 2H), 3.94–3.98 (m, 2H), 7.06 (tt, $J = 1.2 \text{ Hz}$, 7.2 Hz, 2H), 7.22 (dd, $J = 1.2 \text{ Hz}$, 8.0 Hz, 4H), 7.36 (dd, $J = 7.2 \text{ Hz}$, 8.0 Hz, 4H), 7.95 (d, $J = 628 \text{ Hz}$, 1H); ¹³C NMR (CDCl₃): δ 43.4 (d, $J = 8.8 \text{ Hz}$), 116.2 (d, $J = 5.3 \text{ Hz}$), 122.3 (x 2), 129.6 (x 2), 141.2; ³¹P NMR (CDCl₃): δ 6.7; FAB-LRMS m/z 259 (MH⁺), 258 (M⁺); FAB-HRMS. Calcd for C₁₄H₁₅N₂OP: 258.0922 (M⁺). Found: 258.0905 (M⁺).

12 was reacted with BSA to provide a trivalent phosphorus compound **14**. The structure was confirmed by comparing the ³¹P NMR data of an authentic sample, which was prepared according to the reported method using *N,N*-diethylsilylamine. In contrast, a predominant peak was observed at –12.5 ppm in the presence of TMSCl. Considering the Lewis basic property of phosphoramides, it would be reasonable to propose **13** as the corresponding structure (Scheme S-1). BSA-induced P(V) to P(III) transformation is a general property for secondary phosphine oxides. In fact, diphenylphosphine oxide (**SI-3**) was transformed into the corresponding trivalent phosphorus compound **SI-4** in the presence of BSA. Diphenylphosphine oxide is known to tautomerized in only the presence of transition metals, such as allylpalladium chloride dimer. However, no tautomerization occurred in the case of diaminophosphine oxide **12** under the same conditions, indicating that BSA is indispensable reagent for P(V) to P(III) transformation of diaminophosphine oxides (Scheme S-2).

Scheme S-1. BSA-Induced P(V) to P(III) Transformation

Scheme S-2. Transition Metal-Induced P(V) to P(III) Transformation

D. FAB MS and ^{31}P NMR Analyses

FAB MS Data

Page S-23: FAB MS of **6**.

Page S-24: FAB MS of **6** and BSA (1:30).

Page S-25: FAB MS of $(\eta^3\text{-C}_3\text{H}_5\text{PdCl})_2$, **6**, and BSA (Pd:**6**:BSA = 1:1:30).

Page S-26: FAB MS of $(\eta^3\text{-C}_3\text{H}_5\text{PdCl})_2$, **6**, and BSA (Pd:**6**:BSA = 1:2:60).

Page S-27: FAB MS of $(\eta^3\text{-C}_3\text{H}_5\text{PdCl})_2$, **25**, and BSA (Pd:**6**:BSA = 1:1:30).

^{31}P NMR Data

Page S-28: ^{31}P NMR of **6** in CDCl_3 .

Page S-29: ^{31}P NMR of **6** and BSA (1:30) in CDCl_3 .

Page S-30: ^{31}P NMR of $(\eta^3\text{-C}_3\text{H}_5\text{PdCl})_2$, **6**, and BSA (Pd:**6**:BSA = 1:1:30) in CDCl_3 .

Page S-31: ^{31}P NMR of $(\eta^3\text{-C}_3\text{H}_5\text{PdCl})_2$, **6**, and BSA (Pd:**6**:BSA = 1:2:60) in CDCl_3 .

Page S-32: ^{31}P NMR of $(\eta^3\text{-C}_3\text{H}_5\text{PdCl})_2$, **6**, and BSA (Pd:**6**:BSA = 1:3:90) in CDCl_3 .

Page S-33: ^{31}P NMR of $(\eta^3\text{-C}_3\text{H}_5\text{PdCl})_2$, **25**, and BSA (Pd:**6**:BSA = 1:1:30) in CDCl_3 .

General Procedure for FAB MS Analysis: To a stirred solution of **6** (3.9 mg, 0.01 mmol) and $(\eta^3\text{-C}_3\text{H}_5\text{PdCl})_2$ (1.8 mg, 0.005 mmol) in CH_2Cl_2 (0.8 mL) was added excess BSA (50 μL , 0.3 mmol) at room temperature. After 15 min, the solution was evaporated in vacuo, and the obtained residue was utilized for FAB MS (+) analysis (matrix: NBA).

FAB MS analysis of the residue obtained from a CH_2Cl_2 solution of **6** and BSA (**6**:BSA = 1:30) gave us useful information to determine the structure of the actual ligand **17** (Page S-24). Experimental results described in Scheme S-3 support our conclusion. Cationic complex **16** was detected in the FAB MS analysis of the residue obtained from a CH_2Cl_2 solution of $(\eta^3\text{-C}_3\text{H}_5\text{PdCl})_2$, **6**, and BSA (Pd:**6**:BSA = 1:1:30) (Page S-25). Moreover, two singlet signals were detected at 100.0 ppm and 101.7 ppm in ^{31}P NMR analysis of a CDCl_3 solution of $(\eta^3\text{-C}_3\text{H}_5\text{PdCl})_2$, **6**, and BSA (Pd:**6**:BSA = 1:1:30) (Page S-30). We first expected that coordination of the nitrogen atom on the sidearm might contribute to the stabilization of the complex, resulting in the formation of **SI-8** and **SI-9**. However, similar tendency was also observed with use of **25**, a ligand without nitrogen functionality on the sidearm (Page S-27 and S-33). Although it is difficult to make conclusion about the exact structures of the observed complexes, these results appear to indicate that the nitrogen atom is not necessarily important to form some Pd complexes with one molecule of the ligand (Scheme S-4). Increase in the ratio of **6** to the Pd catalyst resulted in decreased detection of the two singlet peaks, and a new singlet signal was observed at 98.0 ppm (Page S-31 and S-32). A π -allylpalladium (II) complex with two molecules of **17** (**SI-11**) is expected to be the corresponding structure (Scheme S-5). No useful information, however, was obtained in the FAB MS analysis of the residue obtained from a CH_2Cl_2 solution of $(\eta^3\text{-C}_3\text{H}_5\text{PdCl})_2$, **6**, and BSA (Pd:**6**:BSA = 1:2:60) (Page S-26).

Note: (1) We used CDCl_3 as the solvent for NMR studies. Asymmetric allylic alkylation of **10** with dimethyl malonate in CHCl_3 gave **11** in 98% yield and 97% ee. This result suggests that the same species generated in the solution, and functions as the active species. (2) In ^{31}P NMR experiments, excess BSA (30 eq to **6**) was used for the P(V) to P(III) transformation. The reason why we used excess BSA is to reproduce the similar situation to the reaction mixture. Actually, P(V) to P(III) transformation was observed when 1~2 equiv of BSA was used. The efficiency, however, was much lower than the case of excess BSA. This might be one of the reasons why the amount of BSA affects the reactivity of quaternary carbon center formation.

Scheme S-3.

Scheme S-4.

Scheme S-5.

E. Discussions

Clarification of the Active Catalyst

Clues for Clarification

- (1) *rds* = Nucleophilic attack to a π -allylpalladium complex (from Kinetic Studies) = The stage of enantioselection.
- (2) Single catalyst component is participating in the rate-determining step (from Kinetic Studies).
- (3) The optimal Pd-ligand ratio is 1:2, and no reaction occurs when Pd-ligand ratio is 1:1 (Table 3).
- (4) Nonlinear effect was observed in both types of reaction.

It is known that there are several origins of nonlinear effect. Taking the clue (1) and clue (2) into account, however, our attention can be focused on the following two possibilities in the case of Reaction Type A.

- (I) Pd complex with two molecules of **17** functions as the active catalyst in the stage of enantioselection.
- (II) Reservoir effect (Scheme S-6).

Scheme S-6. Reservoir Effect

If reservoir effect is the origin of the observed nonlinear effect, a Pd complex with one molecule of **17** (**SI-12**) is proposed as the active catalyst, and there is equilibrium between **SI-12** and a homochiral complex **SI-13** (or other species), which functions as a resting complex in the reaction mixture. No reaction occurred, however, when the catalyst was prepared from Pd catalyst and **6** in a ratio of 1:1 (Table 3, run 3). If the observed nonlinear effect is explained by reservoir effect, **SI-12**, the active catalyst, must be formed in this reaction condition by the aforementioned equilibrium, and should promote the asymmetric catalysis. This clearly indicates that the observed nonlinear effect originates from the formation of a Pd complex with two molecules of **17**, which is participating in the stage of enantioselection (= nucleophilic attack to a π -allylpalladium complex) as the active species. The same discussion is possible in the case of Reaction Type B.

In the case of Reaction Type B, we can postulate an alternative possibility for the reason of the observed nonlinear effect, other than two candidates described above (Scheme S-7). This interpretation would be excluded by the fact that no interaction between the phosphorus atom and the Zn atom was observed in a ^{31}P NMR analysis of a CDCl_3 solution of **6**, $\text{Zn}(\text{OAc})_2$, and BSA (**6**: $\text{Zn}(\text{OAc})_2$:BSA = 2:5:60).

Discussions for the Result Using Enol Silyl Ether **20** as the Substrate

In the reaction mechanism of Reaction Type B (Scheme 6), enol silyl ether **20** is proposed as the actual nucleophile. When **20** itself was used as the substrate, however, remarkable decrease in the reactivity was observe, despite the ee value remaining unchanged [Scheme S-8 (a)]. On the other hand, the same reaction proceeded to completion with the use of **20**, which was prepared from **8a** and BSA in a separate reaction vessel, and then added to the reaction mixture [Scheme S-8 (b)]. This result implies that *N*-trimethylsilylacetamide, generated in situ by the reaction of BSA with a β -keto ester, might be related to this asymmetric catalysis in some way. Considering the unchanged ee value, *N*-trimethylsilylacetamide might not be related directly to the enantiofacial recognition.

2. General Procedures, Characterization of the New Compounds, and Discussion

A. Asymmetric Construction of Quaternary Carbons Using Cyclic Nucleophiles.

All of the acetates were prepared from the corresponding allylic alcohol using usual procedure (Ac_2O , NEt_3 , THF, 0 °C). Compounds **7a** and **7h** are commercially available. For the known compounds **7b** and **7d**, see: Matsuzaka *et al.* *J. Org. Chem.* **1988**, 53, 3832. For the known compound **7c**, see: Lehmann *et al.* *Tetrahedron* **1995**, 51, 8863. For the known compound **7e**, see: Hayashi *et al.* *Synth. Commun.* **1989**, 19, 2109. For the known compound **7f**, see: Meuzelaar *et al.* *Tetrahedron* **2000**, 56, 2895. For the known compound **7g**, see: Goux *et al.* *Tetrahedron* **1996**, 52, 9521. β -Keto esters **8a** and **8d** are commercially available. β -Keto esters **8b–c**, and **8e–k** were prepared according to the reported procedure. For the known compound **8b**, see: Plieninger *et al.* *Chem. Ber.* **1954**, 87, 1760. For the known compounds **8c**, **8f**, and **8h**, see: Mander *et al.* *Tetrahedron Lett.* **1983**, 24, 5425. For the known compound **8e**, see: Christoffers *et al.* *Eur. J. Org. Chem.* **2000**, 1633. For the known compound **8g**, see: Cossy *et al.* *Tetrahedron* **1995**, 51, 11751. All of the allylated products were prepared according to the general procedure. Spectral and analytical data of compounds **9a–d**, **9f–m**, and **9o** have already reported in the precedent communication, see: ref 9.

General Procedure

To a stirred mixture of cinnamyl acetate (**7a**) (83.3 μL , 0.5 mmol), keto ester (**8a**) (120 μL , 0.75 mmol), $(\eta^3\text{-C}_3\text{H}_5\text{PdCl})_2$ (1.8 mg, 0.0025 mmol), **6** (7.8 mg, 0.02 mmol), and $\text{Zn}(\text{OAc})_2$ (9.2 mg, 0.05 mmol) in toluene (3.33 mL) was added BSA (0.494 mL, 2.0 mmol) at room temperature. After being stirred for 16 h, the reaction was concentrated under reduced pressure, and the obtained residue was purified by flash column chromatography (SiO_2 , hexane/ethyl acetate 30/1 to 15/1) to give (1*S*)-2-oxo-1-(3-phenyl-allyl)-cyclohexanecarboxylic acid ethyl ester (**9a**) as colorless oil (143 mg, 99%, 93% ee). $[\alpha]_{\text{D}}^{18} -80.9$ (*c* 0.85, CHCl_3 , 93% ee). The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALCEL OD-H, 2-propanol/hexane 5/95, flow rate 0.4 mL/min, t_{R} 14.5 min [(*R*)-isomer] and 16.8 min [(*S*)-isomer], detection at 254 nm).

Compound 9b: colorless oil; $[\alpha]_{\text{D}}^{25} -64.2$ (*c* 0.97, CHCl_3 , 91% ee). The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALCEL OD-H, 2-propanol/hexane 5/95, flow rate 0.5 mL/min, t_{R} 17.7 min [(*R*)-isomer] and 20.4 min [(*S*)-isomer], detection at 254 nm).

Compound 9c: colorless oil; $[\alpha]_{\text{D}}^{18} -65.2$ (*c* 1.01, CHCl_3 , 94% ee). The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALCEL OD-H, 2-propanol/hexane 5/95, flow rate 0.4 mL/min, t_{R} 18.2 min (minor isomer) and 20.1 min (major isomer), detection at 254 nm).

Compound 9d: colorless oil; $[\alpha]_{\text{D}}^{26} +51.8$ (*c* 0.88, CHCl_3 , 85% ee). The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALCEL OD-H, 2-propanol/hexane 5/95, flow rate 0.4 mL/min, t_{R} 21.5 min [(*R*)-isomer] and 23.5 min [(*S*)-isomer], detection at 254 nm).

Compound 9e: colorless oil; IR (neat) ν 3030, 2960, 2918, 1748, 1728, 1454, 1264, 1219, 1157, 745, 695 cm^{-1} ; ^1H NMR (CDCl_3): δ 1.87–2.08 (m, 3H), 2.22–2.31 (m, 1H), 2.39–2.52 (m, 2H), 2.54 (ddd, $J = 1.2$ Hz, 7.6 Hz, 14.0 Hz, 1H), 2.85 (ddd, $J = 1.2$ Hz, 7.6 Hz, 14.0 Hz, 1H), 5.14 (d, $J = 13.2$ Hz, 1H), 5.18 (d, $J = 13.2$ Hz, 1H), 6.04 (dt, $J = 15.6$ Hz, 7.6 Hz, 1H), 6.41 (dd, $J = 1.2$ Hz, 15.6 Hz, 1H), 7.19–7.38 (m, 10H); ^{13}C NMR (CDCl_3): δ 19.5, 32.2, 37.0, 38.1, 60.3, 67.1, 124.3, 126.2 (x 2), 127.4, 128.0 (x 2), 128.3, 128.5 (x 2), 128.6 (x 2), 134.1, 135.5, 136.9, 170.7, 214.3; EI-LRMS m/z 334 (M^+); $[\alpha]_{\text{D}}^{10} +59.6$ (c 0.55, CHCl_3 , 76% ee); EI-HRMS. Calcd for $\text{C}_{22}\text{H}_{22}\text{O}_3$ (M^+): 334.1569. Found: 334.1574. The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALPAK AD, 2-propanol/hexane 1/99, flow rate 0.7 mL/min, t_{R} 35.8 min (minor isomer) and 40.6 min (major isomer), detection at 254 nm).

Compound 9f: colorless oil; $[\alpha]_{\text{D}}^{25} -23.9$ (c 0.54, CHCl_3 , 78% ee). The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALCEL OD-H, 2-propanol/hexane 5/95, flow rate 0.4 mL/min, t_{R} 16.4 min (minor isomer) and 18.0 min (major isomer), detection at 254 nm).

Compound 9g: colorless oil; $[\alpha]_{\text{D}}^{24} -54.7$ (c 2.00, CHCl_3 , 72% ee). The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALPAK AD, 2-propanol/hexane 5/95, flow rate 0.5 mL/min, t_{R} 12.4 min (minor isomer) and 15.1 min (major isomer), detection at 254 nm).

Compound 9h: colorless oil; $[\alpha]_{\text{D}}^{24} -19.0$ (c 1.23, CHCl_3 , 93% ee). The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALCEL OD-H, 2-propanol/hexane 5/95, flow rate 0.4 mL/min, t_{R} 22.9 min [(*R*)-isomer] and 24.4 min [(*S*)-isomer], detection at 254 nm).

Compound 9i: colorless oil; $[\alpha]_{\text{D}}^{18} -14.1$ (c 0.56, CHCl_3 , 92% ee); The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALCEL OD-H, 2-propanol/hexane 5/95, flow rate 0.4 mL/min, t_{R} 20.0 min (minor isomer) and 22.1 min (major isomer), detection at 254 nm).

Compound 9j: colorless oil; $[\alpha]_{\text{D}}^{18} -22.1$ (c 1.17, CHCl_3 , 91% ee); The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALPAK AD, 2-propanol/hexane 5/95, flow rate 0.3 mL/min, t_{R} 31.3 min (minor isomer) and 35.3 min (major isomer), detection at 254 nm).

Compound 9k: colorless oil; $[\alpha]_{\text{D}}^{21} -12.9$ (c 0.97, CHCl_3 , 91% ee); The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALPAK AD, 2-propanol/hexane 5/95, flow rate 0.4 mL/min, t_{R} 35.8 min (minor isomer) and 38.8 min (major isomer), detection at 254 nm).

Compound 9l: colorless oil; $[\alpha]_{\text{D}}^{18} -20.7$ (c 1.4, CHCl_3 , 82% ee). The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALCEL OD-H, 2-propanol/hexane 5/95, flow rate 0.4 mL/min, t_{R} 22.7 min (minor isomer) and 25.4 min (major isomer), detection at 254 nm).

Compound 9m: colorless oil; $[\alpha]_{\text{D}}^{20} -19.0$ (c 0.79, CHCl_3 , 80% ee). The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALCEL OD-H, 2-propanol/hexane 5/95, flow rate 0.4 mL/min, t_{R} 12.4 min (minor isomer) and 13.8 min (major isomer), detection at 254 nm).

Compound 9n: colorless oil; IR (neat) ν 3441, 1731, 1681, 1637, 1602, 1237, 741, 698 cm^{-1} ; ^1H NMR (CDCl_3): δ 2.14 (ddd, $J = 4.8$ Hz, 10.4 Hz, 13.6 Hz, 1H), 2.53 (ddd, $J = 4.8$ Hz, 4.8 Hz, 13.6 Hz, 1H), 2.69–2.77 (m, 2H), 2.91 (ddd, $J = 4.8$ Hz, 4.8 Hz, 17.2 Hz, 1H), 3.06 (ddd, $J = 4.8$ Hz, 10.4 Hz, 17.2 Hz, 1H), 3.66 (s, 3H), 3.97 (d, $J = 4.4$ Hz, 2H), 5.67–5.78 (m, 2H), 7.20–7.34 (m, 7H), 7.46–7.49 (m, 1H), 8.04–8.06 (m, 1H); ^{13}C NMR (CDCl_3): δ 25.8, 30.6, 37.3, 52.5, 57.5, 70.5, 71.9, 126.8, 127.6, 127.8 (x 2), 128.1, 128.3 (x 2), 128.7, 128.7, 131.0, 131.8, 133.6, 138.2, 143.2, 172.0, 194.9;

FAB-LRMS m/z 365 (MH^+); $[\alpha]_D^{22}$ -22.5 (c 0.755, $CHCl_3$, 88% ee). FAB-HRMS. Calcd for $C_{23}H_{25}O_4$ (MH^+): 365.1753. Found: 365.1745. The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALCEL OD-H, 2-propanol/hexane 5/95, flow rate 0.4 mL/min, t_R 46.3 min (major isomer) and 54.8 min (minor isomer), detection at 254 nm).

Compound 9o: colorless oil; $[\alpha]_D^{25}$ -29.7 (c 0.87, $CHCl_3$, 63% ee). The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALCEL OD-H, 2-propanol/hexane 5/95, flow rate 0.4 mL/min, t_R 15.2 min [(*R*)-isomer] and 17.1 min [(*S*)-isomer], detection at 254 nm).

Table S-3. Screening of Acetate Salt in the Asymmetric Substitution of **7a** with **8a**

$(\eta^3-C_3H_5PdCl)_2$ (2.5 mol %)
6 (10 mol %)
 additive (10 mol %)
 BSA (3 eq), **8a** (1.25 eq)
 toluene, rt, 24 h

run	additive	yield	ee
1	—	10%	53% ee
2	<i>n</i> -Bu ₄ NOAc	21%	18% ee
3	LiOAc	53%	8% ee
4	KOAc	No Reaction	—
5	CsOAc	No Reaction	—
6	Mg(OAc) ₂ ·4H ₂ O	99%	66% ee
7	Ba(OAc) ₂	No Reaction	—
8	Cu(OAc) ₂	No Reaction	—
9	Yb(OAc) ₃ ·4H ₂ O	No Reaction	—
10	In(OAc) ₃	17%	71% ee
11	Zn(OAc) ₂ ·2H ₂ O	81%	89% ee
12	Zn(OAc) ₂	80%	91% ee

B. Asymmetric Construction of Quaternary Carbons Using Acyclic Nucleophiles.

Compound **9p–r** were prepared according to the general procedure. For the known compound **8j–k**, see: Kuwano et al. *J. Org. Chem.* **1998**, 63, 3499. For the known compounds **9q–r**, see: ref 12b.

General Procedure

To a stirred mixture of cinnamyl acetate (**7a**) (14.7 mg, 0.083 mmol), keto ester (**8i**) (30.2 mg, 0.125 mmol), $(\eta^3-C_3H_5PdCl)_2$ (0.76 mg, 0.002 mmol), **6** (3.26 mg, 0.008 mmol), in toluene (0.55 mL) was added BSA (82 μ L, 0.332 mmol) at 0 °C. After being stirred for 48 h at the same temperature, the reaction was concentrated under reduced pressure, and the obtained residue was purified by flash column chromatography (SiO_2 , hexane/ethyl acetate 4/1) to give 2-acetylcyclohexanecarboxylate-5-phenylpent-4-enoic acid methyl ester (**9p**) as a white solid (29.4 mg, 99%, 80% ee). Mp: 82–85 °C; IR (KBr) ν 2929, 1723, 1644, 1507, 1296, 1204, 974, 742, 691, 638 cm^{-1} ; ^1H NMR ($CDCl_3$): δ 1.19–1.78 (m, 10H), 2.04 (s, 3H), 2.81 (tt, J = 3.6 Hz, 11.6 Hz, 1H), 3.19 (ddd, J = 1.2 Hz, 7.6 Hz, 15.2 Hz, 1H), 3.40 (ddd, J = 1.2 Hz, 7.6 Hz, 15.2 Hz, 1H), 3.77 (s, 3H), 5.76 (ddd, J = 7.6 Hz, 7.6 Hz, 15.6 Hz, 1H), 6.42 (d, J = 15.6 Hz, 1H), 6.99 (s, 1H), 7.21–7.31

(m, 5H); ^{13}C NMR (CDCl_3): δ 23.0, 25.3, 25.4, 25.4, 29.4, 30.0, 35.3, 45.0, 53.1, 73.1, 122.4, 126.2 (x 2), 127.6, 128.5 (x 2), 134.7, 136.8, 168.7, 169.1, 205.0; EI-LRMS m/z 357 (M^+); $[\alpha]_{\text{D}}^{20} +18.4$ (c 0.99, CHCl_3 , 80% ee); EI-HRMS. Calcd for $\text{C}_{21}\text{H}_{27}\text{NO}_4$ (M^+): 357.1940. Found: 357.1947. The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALCEL OD-H, 2-propanol/hexane 1/9, flow rate 0.8 mL/min, t_{R} 10.3 min (minor isomer) and 12.6 min (major isomer), detection at 254 nm).

Compound 8i: Mp: 151–152 °C; IR (neat) ν 3299, 2926, 2856, 1742, 1718, 1655, 1549, 1451, 1375, 1268, 1206, 1015, 687 cm^{-1} ; ^1H NMR (CDCl_3): δ 1.18–1.50 (m, 5H), 1.65–1.84 (m, 4H), 1.97–2.04 (m, 1H), 2.07 (s, 3H), 2.79 (tt, J = 3.6 Hz, 11.2 Hz, 1H), 3.80 (s, 3H), 5.41 (d, J = 6.4 Hz, 1H), 6.69 (d, J = 6.4 Hz, 1H); ^{13}C NMR (CDCl_3): δ 22.8, 25.1, 25.6, 25.7, 27.7, 29.1, 48.6, 53.2, 60.7, 166.9, 169.7, 204.1; FAB-LRMS m/z 242 (MH^+); FAB-HRMS. Calcd for $\text{C}_{12}\text{H}_{20}\text{NO}_4$ (MH^+): 242.1392. Found: 242.1391.

Compound 9q: white solid; $[\alpha]_{\text{D}}^{25} -5.0$ (c 0.83, CHCl_3 , 84% ee). The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALCEL OJ, 2-propanol/hexane 1/9, flow rate 0.5 mL/min, t_{R} 43.0 min (major isomer) and 55.7 min (minor isomer), detection at 254 nm).

Compound 9r: white solid; $[\alpha]_{\text{D}}^{25} -9.6$ (c 0.48, CHCl_3 , 78% ee). The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALCEL OD-H, 2-propanol/hexane 5/95, flow rate 0.4 mL/min, t_{R} 15.2 min [(*S*)-isomer] and 17.1 min [(*R*)-isomer], detection at 254 nm).

Table S-4. Optimization of the reaction of **7a** with **8i**

run	additive	solvent	temp.	time	yield	ee
1	$\text{Zn}(\text{OAc})_2$	toluene (0.125 M)	rt	24 h	73%	71% ee
2	LiOAc	toluene (0.125 M)	rt	24 h	94%	72% ee
3	$\text{Mg}(\text{OAc})_2 \cdot 4\text{H}_2\text{O}$	toluene (0.125 M)	rt	24 h	23%	69% ee
4	$\text{In}(\text{OAc})_3$	toluene (0.125 M)	rt	24 h	21%	74% ee
5	<i>n</i> -Bu ₄ NOAc	toluene (0.125 M)	rt	24 h	82%	72% ee
6	————	toluene (0.125 M)	rt	24 h	95%	73% ee
7	————	toluene (0.125 M)	4 °C	24 h	83%	78% ee
8	————	CH_2Cl_2 (0.125 M)	4 °C	24 h	84%	66% ee
9	————	toluene (0.15 M)	0 °C	48 h	99%	80% ee

C. Asymmetric Allylic Alkylation of **10** with Dimethyl Malonate

Compounds **10** was prepared according to the general procedure. For the known compounds **11**, see: Mino et al. *Tetrahedron:Asymmetry* **2001**, *12*, 2435.

General Procedure

To a stirred solution of **10** (126 mg, 0.5 mmol), (η^3 -C₃H₅PdCl)₂ (1.8 mg, 0.005 mmol), **6** (7.8 mg, 0.02 mmol), and dimethyl malonate (0.171 mL, 1.5 mmol) in CH₂Cl₂ (2 mL) was added BSA (0.370 mL, 1.5 mmol) at room temperature. After being stirred for 24 h, the reaction was concentrated under reduced pressure, and the obtained residue was purified by flash column chromatography (SiO₂, hexane/ethyl acetate 30/1) to give (1*S*)-1,3-diphenyl-2-propenyl-propanedioic acid dimethyl ester (**11**) as a colorless oil (152.2 mg, 94%, 99% ee). $[\alpha]_D^{22}$ -12.8 (*c* 0.38, CHCl₃, 99% ee). The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALCEL OD-H, 2-propanol/hexane 1/99, flow rate 0.3 mL/min, *t_R* 41.0 min [(*R*)-isomer] and 43.6 min [(*S*)-isomer], detection at 254 nm).

3. Experimental Procedures for the Synthesis of P-Chirogenic Diaminophosphine Oxides

A. Synthesis of the Aspartic Acid-Derived P-Chirogenic Diaminophosphine Oxides (DIAPHOXs)

Benzyloxycarbonyl-L-aspartic anhydride (**1**) was prepared by using the known procedure (*J. Chem. Soc.* **1954**, 2870).

Compound 2: To a stirred mixture of acid anhydride **1** (6.11 g, 24.5 mmol) in DMF (50 mL) was added aniline (2.46 mL, 27.0 mmol) at room temperature, and then the mixture was kept stirring for 3 h at the same temperature. Aniline (2.91 mL, 31.9 mmol) and WSCI (5.18 g, 27.0 mmol) were added to the reaction, and the mixture was stirred for 24 h at room temperature. After evaporation of the half of DMF, the reaction was diluted with ethyl acetate, and washed with 1N HCl and water (x 3). Evaporation of the organic solvent under reduced pressure, followed by ether wash of the residue gave pure **2** as a white solid (8.98 g, 87%). Mp 222–225 °C; IR (KBr) ν 3442, 3289, 1698, 1654, 1534, 1444, 1264, 692 cm⁻¹; ¹H NMR (DMSO-*d*₆): δ 2.71 (dd, *J* = 8.4 Hz, 15.2 Hz, 1H), 2.83 (dd, *J* = 5.2 Hz, 15.2 Hz, 1H), 4.62 (ddd, *J* = 3.6 Hz, 5.2 Hz, 8.4 Hz, 1H), 5.04 (s, 2H), 7.02–7.07 (m, 2H), 7.27–7.35 (m, 9H), 7.57–7.63 (m, 4H), 7.70 (d, *J* = 3.6 Hz, 1H), 9.95 (s, 1H), 10.11 (s, 1H); ¹³C NMR (DMSO-*d*₆): δ 38.5, 52.2, 65.5, 119.0 (x 2), 119.4 (x 2), 123.1, 123.3, 127.7 (x 2), 127.8, 128.3 (x 2), 128.6 (x 2), 128.7 (x 2), 136.8, 139.0, 139.1, 155.8, 167.9, 170.0; FAB-LRMS *m/z* 418 (MH⁺); $[\alpha]_D^{25}$ +3.46 (*c* 0.52, DMF); Anal. Calcd for C₂₄H₂₄N₃O₄: C, 69.05, H, 5.55, N, 10.07. Found: C, 68.82, H, 5.49, N, 10.08.

Compound 3: A suspension of **2** (7.955 g, 19 mmol) and 5% Pd-C (808 mg, 0.38 mmol) in DMF (150 mL) and 2-propanol (50 mL) was stirred under H₂ atmosphere at room temperature. After 2 h, the mixture was filtered through celite, and the filtrate was evaporated under reduced pressure. After dilution with ethyl acetate, the organic layer was washed with NaHCO₃, water (x 1), and then dried over Na₂SO₄. After concentration in vacuo, the crude residue was washed with ether to afford the corresponding amine as a white solid. To a stirred solution of the obtained solid (2.46 g, 8.68 mmol) and triethylamine (1.81 mL, 13.0 mmol) in THF (150 mL) was added benzoyl chloride (1.26 mL, 10.85 mmol) at 0 °C. After being stirred for 1 h, the reaction was quenched with saturated aqueous NaHCO₃. The resulting mixture was extracted

with ethyl acetate (x 2), and the combined organic layers were washed with 1N HCl, water, and brine. Evaporation of organic solvent under reduced pressure, followed by ether wash of the residue gave pure **3** as a white solid (3.29 g, 97%). Mp 222–223 °C; IR (KBr) ν 3461, 3296, 1660, 1637, 1600, 1528, 1499, 1444, 1321, 753, 692 cm^{-1} ; ^1H NMR ($\text{DMSO-}d_6$): δ 2.88 (dd, J = 8.4 Hz, 15.6 Hz, 1H), 2.94 (dd, J = 6.0 Hz, 15.6 Hz, 1H), 5.00 (ddd, J = 6.0 Hz, 7.2 Hz, 8.4 Hz, 1H), 7.00–7.06 (m, 2H), 7.26–7.31 (m, 4H), 7.44–7.64 (m, 7H), 7.87–7.89 (m, 2H), 8.77 (d, J = 7.2 Hz, 1H), 10.00 (s, 1H), 10.17 (s, 1H); ^{13}C NMR ($\text{DMSO-}d_6$): δ 38.1, 51.4, 119.1 (x 2), 119.4 (x 2), 123.1, 123.3, 127.5 (x 2), 128.2 (x 2), 128.7 (x 2), 128.7 (x 2), 131.4, 133.9, 139.0, 139.1, 166.4, 168.2, 169.9; EI-LRMS m/z 387 (M^+); $[\alpha]_D^{25} +13.6$ (c 0.50, DMF); EI-HRMS. Calcd for $\text{C}_{23}\text{H}_{27}\text{N}_3$ (M^+): 387.1583. Found: 387.1588.

Compound 4: To a stirred suspension of LAH (2.03 g, 53.5 mmol) in THF (178 mL) was added **3** (3.46 g, 8.92 mmol) at 0 °C. The resulting suspension was heated on oil bath and refluxed for 13 h. After cooling down to room temperature, the reaction was quenched by the addition of water and 1N NaOH. The obtained white solid was filtered, and refluxed in THF (60 mL) for 3 h. After filtration, the combined organic layers were washed with brine, dried over Na_2SO_4 , and then evaporated in vacuo. The residue was purified by flash column chromatography (SiO_2 , hexane/ethyl acetate 3/1 to 1/2) to give triamine **4** as pale yellow oil (2.21 g, 72%). IR (neat) ν 3404, 3023, 2923, 2828, 1603, 1505, 1319, 1257, 749, 693 cm^{-1} ; ^1H NMR (CDCl_3): δ 1.74–1.85 (m, 2H), 2.93–2.98 (m, 1H), 3.06 (dd, J = 6.0 Hz, 12.4 Hz, 1H), 3.15–3.20 (m, 3H), 3.77 (s, 1H), 6.52–6.71 (m, 6H), 7.13–7.30 (m, 9H); ^{13}C NMR (CDCl_3): δ 32.0, 41.2, 46.4, 50.9, 54.6, 112.7 (x 2), 112.9 (x 2), 117.1, 117.4, 127.0, 128.0 (x 2), 128.4 (x 2), 129.1 (x 2), 129.2 (x 2), 140.2, 148.2, 148.4; EI-LRMS m/z 345 (M^+); $[\alpha]_D^{19} +1.01$ (c 1.39, CHCl_3); EI-HRMS. Calcd for $\text{C}_{23}\text{H}_{27}\text{N}_3$ (M^+): 345.2205. Found: 345.2210.

P-Chirogenic Diaminophosphine Oxide **6**:

Method A: To a stirred solution of **4** (1.99 g, 5.77 mmol) and triethylamine (2.90 mL, 20.8 mmol) in toluene (29 mL) at –78 °C was added dropwise phosphorous trichloride (0.553 mL, 6.35 mmol) over 3 min. The reaction temperature was gradually warmed up to room temperature, and then the mixture was kept stirring for 12 h. The reaction mixture was diluted with CHCl_3 , and the organic layer was washed with H_2O , and then dried over Na_2SO_4 . After concentration in vacuo, the obtained residue was dissolved into ethyl acetate, and then SiO_2 (50 mg/mmol substrate) and H_2O (0.13 mL, 7.21 mmol) was added. After being stirred for 20 h at room temperature under air, the reaction was diluted with CHCl_3 , and then filtered through a short pad of silica gel, and the silica residue was washed with CHCl_3 and ethyl acetate intensively. After concentration of the organic washings in vacuo, the obtained yellow solid was washed with ethyl acetate, and then filtered to get **6** as a white solid. Additionally, purification of the yellow residue by flash column chromatography (SiO_2 , hexane/ethyl acetate 5/1 to 1/1) afforded **6** (Total yield: 1.36 g, 60%).

Method B: To a stirred solution of **4** (2.20 g, 6.35 mmol) and triethylamine (3.44 mL, 24.7 mmol) in toluene (42 mL) at –78 °C was added dropwise phosphorous trichloride (0.610 mL, 6.98 mmol) over 5 min. The reaction temperature was gradually warmed up to room temperature, and then the mixture was kept stirring for 12 h. After filtration of the generated ammonium salt, the organic phase was concentrated in vacuo. The obtained residue was purified by flash column chromatography (SiO_2 , hexane/ethyl acetate 5/1 to 1/1) to afford **6** (1.26 g, 51%). Mp 131–132 °C; IR (KBr) ν 3370, 2927, 1601, 1524, 1502, 1327, 1303, 1219, 1170, 1127, 1034, 950, 751, 730, 700 cm^{-1} ; ^1H NMR (CDCl_3): δ 1.67–1.76 (m, 1H), 2.00–2.08 (m, 1H), 3.04–3.18 (m, 2H), 3.31–3.37 (m, 1H), 3.45 (s, 1H), 3.65 (m, 1H), 3.76–3.82 (m, 1H), 4.20 (dd, J = 13.2 Hz (PNCH), 15.2 Hz (HCH), 1H), 4.44 (dd, J = 10.4 Hz (PNCH), 15.2 Hz (HCH), 1H), 6.50–6.53 (m, 2H),

6.71–6.75 (m, 1H), 6.99–7.04 (m, 1H), 7.15–7.19 (m, 4H), 7.29–7.47 (m, 7H) 7.70 (d, $J = 628$ Hz (O=PH), 1H); ^{13}C NMR (CDCl_3): δ 32.7 (d, $J = 3.9$ Hz), 39.6, 46.5 (d, $J = 6.1$ Hz), 48.9 (d, $J = 10.4$ Hz), 52.1 (d, $J = 7.2$ Hz), 112.8 (x 2), 115.9 (d, $J = 5.3$ Hz) (x 2), 117.9, 121.9, 127.8, 128.6 (x 2), 128.7 (x 2), 129.3 (x 2), 129.5 (x 2), 136.7, 141.4 (d, $J = 7.7$ Hz), 147.6; ^{31}P NMR (CDCl_3): δ 12.5; EI-LRMS m/z 391 (M^+); FAB-LRMS m/z 392 (MH^+); $[\alpha]_{\text{D}}^{18} -38.6$ (c 0.5, CHCl_3 , >99% ee); FAB-HRMS. Calcd for $\text{C}_{23}\text{H}_{27}\text{N}_3\text{OP}$: 392.1845 (MH^+). Found: 392.1854; Anal. Calcd for $\text{C}_{23}\text{H}_{26}\text{N}_3\text{OP}$: C, 70.57; H, 6.69; N, 10.73. Found, C, 70.27; H, 6.64; N, 10.66. The enantiomeric excess was determined by HPLC analysis (DAICEL CHIRALCEL OD-H, 2-propanol/hexane 1/1, flow rate 0.4 mL/min, t_{R} 49.5 min (ligand prepared from D-Asp) and 54.2 min (ligand prepared from L-Asp), detection at 254 nm).

N-Acetylated ligand (22): To a stirred solution of **6** (39.1 mg, 0.1 mmol) and triethylamine (70 μL , 0.5 mmol) in CH_2Cl_2 (1 mL) was added acetyl chloride (24 μL , 0.36 mmol) at 0 $^\circ\text{C}$, and the reaction was gradually warmed up to room temperature. After 12 h, the reaction was diluted with ethyl acetate, washed with brine, and dried over Na_2SO_4 . After concentration in vacuo, the obtained residue was purified by flash column chromatography (SiO_2 , hexane/ethyl acetate 1/3) to afford **22** (29.7 mg, 69%) as colorless sticky oil. IR (neat) ν 3463, 2927, 1651, 1596, 1496, 1455, 1401, 1301, 1230, 1163, 1116, 951, 755, 702 cm^{-1} ; ^1H NMR (CDCl_3): δ 1.56–1.67 (m, 1H), 1.77 (s, 3H), 1.88–1.98 (m, 1H), 3.34–3.40 (m, 1H), 3.55–3.61 (m, 2H), 3.77–3.85 (m, 2H), 4.14 (dd, $J = 8.8$ Hz (PNCH), 15.2 Hz (HCH), 1H), 4.38 (dd, $J = 10.4$ Hz (PNCH), 15.2 Hz (HCH), 1H), 6.94–7.05 (m, 3H), 7.17–7.42 (m, 12H), 7.62 (d, $J = 627$ Hz (O=PH), 1H); ^{13}C NMR (CDCl_3): δ 22.6, 30.6, 44.6, 46.2 (d, $J = 6.5$ Hz), 48.5 (d, $J = 9.5$ Hz), 51.7 (d, $J = 7.6$ Hz), 115.9 (d, $J = 5.8$ Hz) (x 2), 121.8, 127.7, 127.8 (x 2), 128.1, 128.5 (x 2), 128.6 (x 2), 129.4 (x 2), 129.9 (x 2), 136.6, 141.4, 142.5, 170.4; ^{31}P NMR (CDCl_3): δ 12.5; EI-LRMS m/z 433 (M^+); $[\alpha]_{\text{D}}^{26} -52.9$ (c 0.56, CHCl_3); EI-HRMS. Calcd for $\text{C}_{25}\text{H}_{28}\text{N}_3\text{O}_2\text{P}$: 433.1919 (M^+). Found: 433.1919.

N-Methylated Ligand (23): To a stirred solution of **6** (39.1 mg, 0.1 mmol) in CH_3CN (8 mL) was added 37% aqueous formaldehyde (24 μL , 0.3 mmol) at room temperature. After 30 min, NaBH_3CN (7.5 mg, 0.15 mmol) and one drop of acetic acid were added, and the reaction was stirred for 30 min. After dilution with ethyl acetate, the solution was washed with saturated aqueous NaHCO_3 , water, and brine, and then dried over Na_2SO_4 . After concentration in vacuo, the obtained residue was purified by flash column chromatography (SiO_2 , hexane/ethyl acetate 2.5/1 to 1/1) to afford **23** (34.7 mg, 86%) as colorless oil. IR (neat) ν 3057, 3028, 2925, 2867, 1600, 1504, 1304, 1229, 1161, 749, 691 cm^{-1} ; ^1H NMR (CDCl_3): δ 1.61–1.70 (m, 1H), 1.97–2.05 (m, 1H), 2.81 (s, 3H), 3.22–3.35 (m, 3H), 3.52–3.60 (m, 1H), 3.74–3.80 (m, 1H), 4.17 (dd, $J = 12.4$ Hz (PNCH), 15.2 Hz (HCH), 1H), 4.42 (dd, $J = 10.4$ Hz (PNCH), 15.2 Hz (HCH), 1H), 6.62 (dd, $J = 0.8$ Hz, 8.8 Hz, 2H), 6.72–6.75 (m, 1H), 7.00–7.04 (m, 1H), 7.15–7.45 (m, 11H), 7.67 (d, $J = 628$ Hz (O=PH), 1H); ^{13}C NMR (CDCl_3): δ 29.7 ($J = 4.1$ Hz), 38.5, 46.5 ($J = 6.6$ Hz), 48.5, 48.9 ($J = 10.7$ Hz), 52.5 ($J = 7.4$ Hz), 112.7 (x 2), 115.9 ($J = 4.9$ Hz) (x 2), 117.0, 121.9, 127.8, 128.7 (x 2), 128.7 (x 2), 129.3 (x 2), 129.5 (x 2), 136.5, 141.4 ($J = 7.4$ Hz), 148.8; ^{31}P NMR (CDCl_3): δ 12.4; FAB-LRMS m/z 405 (M^+); $[\alpha]_{\text{D}}^{21} -36.0$ (c 0.71, CHCl_3); FAB-HRMS. Calcd for $\text{C}_{24}\text{H}_{28}\text{N}_3\text{OP}$: 405.1970 (M^+). Found: 405.1979.

B. Synthesis of the Glutamic Acid-Derived P-Chirogenic Diaminophosphine Oxide 24 (Scheme S-9)
Scheme S-9.

Reagents and Conditions: ^a aniline derivative, DMF, rt, then aniline derivative, WSCI, rt. ^b (1) Pd-C (cat.), H₂, DMF-*i*-PrOH, rt. (2) Benzoyl chloride, NEt₃, THF, rt. ^c LiAlH₄, THF, reflux. ^d BH₃·THF, THF, reflux. ^e PCl₃, NEt₃, toluene, -78 °C to rt, then purification with silica gel column chromatography.

Benzyloxycarbonyl-L-glutamic anhydride (**SI-16**) was prepared by using the known procedure (*Biochemistry* **2001**, *40*, 8335.)

Compound SI-17: To a stirred mixture of acid anhydride **SI-16** (5.42g, 20.6 mmol) in DMF (100 mL) was added aniline (2.25 mL, 24.7 mmol) at room temperature, and then the mixture was kept stirring for 1 h at the same temperature. Aniline (2.07 mL, 22.7 mmol) and WSCI (4.95 g, 25.8 mmol) were added to the reaction, and the mixture was stirred for 24 h at room temperature. After evaporation of the half of DMF, the reaction was diluted with ethyl acetate, and washed with 1N HCl and water (x 3). Evaporation of the organic solvent under reduced pressure, followed by ether wash of the residue gave pure **SI-17** as a white solid (7.21 g, 82%). Mp 228–230 °C; IR (KBr) ν 3294, 1698, 1658, 1600, 1539, 1443, 1251, 1055, 763, 702 cm⁻¹; ¹H NMR (DMSO-*d*₆): δ 1.88–2.18 (m, 2H), 2.36–2.51 (m, 2H), 4.20 (m, 1H), 5.04 (s, 2H), 7.00–7.08 (m, 2H), 7.26–7.37 (m, 9H), 7.57–7.64 (m, 5H), 9.93 (s, 1H), 10.10 (s, 1H); ¹³C NMR (DMSO-*d*₆): δ 26.3, 31.6, 53.9, 64.4, 117.9 (x 2), 118.2 (x 2), 121.9, 122.3, 126.7 (x 2), 126.7, 127.2 (x 2), 127.5 (x 2), 127.6 (x 2), 135.8, 137.8, 138.1, 154.9, 169.2, 169.5; FAB-LRMS m/z 433 (MH⁺); [α]_D²² +5.1 (c 0.90, DMF); Anal. Calcd for C₂₅H₂₅N₃O₄: C, 69.59, H, 5.84, N, 9.74. Found: C, 69.25, H, 5.77, N, 9.67.

Compound SI-18: A suspension of **SI-17** (4.31 g, 10 mmol) and 5% Pd-C (680 mg, 0.32 mmol) in DMF (70 mL) and 2-propanol (70 mL) was stirred under H₂ atmosphere at room temperature. After 12 h, the mixture was filtered through celite, and the filtrate was evaporated under reduced pressure. After dilution with ethyl acetate, the organic layer was washed with NaHCO₃, water (x 1), and then dried over Na₂SO₄. After concentration in vacuo, the crude residue was washed with ether to afford the corresponding amine as a brown solid. To a stirred solution of the crude compound and triethylamine (1.73 mL, 12.4 mmol) in THF (80 mL) and DMF (20 mL) was added benzoyl chloride (1.16 mL, 10.3 mmol) at 0 °C. The reaction was stirred for 1.5 h, and then the solvent was under reduced pressure. After dilution with ethyl acetate, the organic layer was washed with NaHCO₃, water (x 1), and then dried over Na₂SO₄. Evaporation of organic solvent under reduced pressure, followed by ether wash of the residue gave pure **SI-18** as a white solid (3.02 g,

76%: 2 steps). Mp 240–241 °C; IR (KBr) ν 3279, 1660, 1635, 1601, 1534, 1499, 1444, 1325, 755, 692 cm^{-1} ; ^1H NMR ($\text{DMSO}-d_6$): δ 2.12–2.20 (m, 2H), 2.42–2.58 (m, 2H), 4.58–4.63 (m, 1H), 6.98–7.06 (m, 2H), 7.24–7.33 (m, 4H), 7.44–7.64 (m, 7H), 7.92–7.94 (m, 2H), 8.64 (d, $J = 7.6$ Hz, 1H), 9.94 (s, 1H), 10.14 (s, 1H); ^{13}C NMR ($\text{DMSO}-d_6$): δ 27.1, 33.0, 54.1, 119.0 (x 2), 119.3 (x 2), 123.0, 123.3, 127.6 (x 2), 128.2 (x 2), 128.6 (x 2), 128.7 (x 2), 131.4, 133.9, 139.0, 139.2, 166.5, 170.4, 170.5; FAB-LRMS m/z 402 (MH^+); $[\alpha]_D^{25} +29.1$ (c 1.0, DMF); Anal. Calcd for $\text{C}_{24}\text{H}_{23}\text{N}_3\text{O}_3$: C, 71.80, H, 5.77, N, 10.47. Found: C, 71.66, H, 5.82, N, 10.30.

Compound SI-20: To a stirred suspension of LAH (1.37 g, 53.5 mmol) in THF (100 mL) was added **SI-18** (2.44 g, 6 mmol) at 0 °C. The resulting suspension was heated on oil bath and refluxed for 8 h. After cooling down to room temperature, the reaction was quenched by the addition of water and 1N NaOH. The obtained white solid was filtered, and refluxed in THF (50 mL) for 3 h. After filtration, the combined organic layers were washed with brine, dried over Na_2SO_4 , and then evaporated in vacuo. The residue was purified by flash column chromatography (SiO_2 , hexane/ethyl acetate 3/1 to 1/2) to give a mixture of diamine **SI-19** (1.13 g, 51%) and the desired triamine **SI-20** (636 mg, 31%). [Compound **SI-19**: ^1H NMR (CDCl_3): δ 1.68–1.84 (m, 4H), 3.13–3.35 (m, 4H), 3.70 (br-s, 1H), 4.05 (br-s, 1H), 4.35–4.48 (m, 1H), 6.26 (d, $J = 8.0$ Hz, 1H), 6.58–6.72 (m, 6H), 7.14–7.18 (m, 4H), 7.34–7.39 (m, 2H), 7.45–7.49 (m, 1H), 7.60–7.69 (m, 2H); ^{13}C NMR (CDCl_3): δ 26.0, 30.6, 43.6, 48.6, 49.8, 112.8 (x 2), 112.8 (x 2), 117.4, 117.6, 126.9 (x 2), 128.6 (x 2), 129.2 (x 2), 129.3 (x 2), 131.6, 134.2, 148.1, 168.0; FAB-LRMS m/z 374 (MH^+)] The partially reduced diamine **SI-19** was converted into **SI-20** by treatment with $\text{BH}_3\cdot\text{THF}$ complex. To a stirred solution of **SI-19** (373 mg, 1 mmol) in THF (5 mL) was added THF solution of $\text{BH}_3\cdot\text{THF}$ complex (2.5 mL, 2.5 mmol), and the reaction was refluxed for 15 h. After cooling down to room temperature, the reaction was quenched by the addition of 1N NaOH. The mixture was extracted with ethyl acetate (10 mL x 3), and the combined organic layers were washed with brine, dried over Na_2SO_4 . After concentration in vacuo, the obtained residue was purified by flash column chromatography (SiO_2 , hexane/ethyl acetate 2/1 to 1/2) to give triamine **SI-20** (301 mg, 84%) as pale yellow oil. IR (neat) ν 3322, 3031, 2926, 2849, 1602, 1504, 1323, 1264, 1153, 1074, 748, 693 cm^{-1} ; ^1H NMR (CDCl_3): δ 1.74–1.85 (m, 2H), 2.93–2.98 (m, 1H), 3.06 (dd, $J = 6.0$ Hz, 12.4 Hz, 1H), 3.15–3.20 (m, 3H), 3.77 (s, 1H), 6.52–6.71 (m, 6H), 7.13–7.30 (m, 9H); ^{13}C NMR (CDCl_3): δ 25.2, 26.9, 43.2, 43.9, 48.6, 53.9, 112.5 (x 2), 112.9 (x 2), 117.2, 117.3, 128.7, 128.8 (x 2), 129.2 (x 2), 129.3 (x 2), 129.7 (x 2), 131.5, 147.2, 148.0; FAB-LRMS m/z 360 (MH^+); $[\alpha]_D^{25} +7.6$ (c 0.58, CHCl_3); FAB-HRMS. Calcd for $\text{C}_{24}\text{H}_{30}\text{N}_3$ (MH^+): 360.2440. Found: 360.2424.

Diaminophosphine Oxide 24: To a stirred solution of **SI-20** (612 mg, 1.70 mmol) and triethylamine (0.924 mL, 6.63 mmol) in toluene (17 mL) at -78 °C was added dropwise phosphorous trichloride (0.163 mL, 1.87 mmol) over 1 min. The reaction temperature was gradually warmed up to room temperature, and then the mixture was kept stirring for 13 h. After filtration of the generated ammonium salt, the organic phase was concentrated in vacuo. The obtained residue was allowed to purify by flash column chromatography (SiO_2 , hexane/ethyl acetate 1/1 to 1/2). Successive purification by flash column chromatography afforded pure diaminophosphine oxide **24** (243 mg, 35%) as a colorless oil. IR (neat) ν 3335, 3029, 2930, 2864, 1733, 1602, 1503, 1303, 1225, 1164, 751, 693 cm^{-1} ; ^1H NMR (CDCl_3): δ 1.40–1.80 (m, 4H), 2.95–3.10 (m, 2H), 3.20–3.26 (m, 1H), 3.50–3.60 (m, 1H), 3.55 (broad peak (NH), 1H), 3.68–3.74 (m, 1H), 4.18 (dd, $J = 12.8$ Hz (PNCH), 15.2 Hz (HCH), 1H), 4.38 (d, $J = 10.8$ Hz (PNCH), 15.2 Hz (HCH), 1H), 6.54–6.56 (m, 2H), 6.68–6.72 (m, 1H), 6.98–7.00 (m, 1H), 7.13–7.18 (m, 4H), 7.27–7.36 (m, 5H), 7.44–7.46 (m, 2H), 7.65 (d, $J = 627$ Hz (O=PH), 1H); ^{13}C

NMR (CDCl₃): δ 24.1, 30.2 (d, J = 4.2 Hz), 43.5, 46.3 (d, J = 6.2 Hz), 48.5 (d, J = 9.9 Hz), 53.7 (d, J = 7.2 Hz), 112.7 (x 2), 115.7 (d, J = 5.7 Hz) (x 2), 117.4, 121.8, 127.7, 128.5 (x 2), 128.6 (x 2), 129.2 (x 2), 129.4 (x 2), 136.8, 141.4 (d, J = 7.6 Hz), 147.9; ³¹P NMR (CDCl₃) δ 12.6; FAB-LRMS m/z 405 (M⁺); [α]_D²¹ -35.8 (c 0.69, CHCl₃); FAB-HRMS Calcd for C₂₄H₂₈N₃OP (MH⁺): 405.1970. Found: 405.1987.

C. Synthesis of the Homophenylalanine-Derived P-Chirogenic Diaminophosphine Oxides **25** and **26** (Scheme S-10)

Scheme S-10.

Reagents and Conditions: ^a (1) aniline, WSCI, DMF, rt. (2) Pd-C (cat.), H₂, DMF-*i*-PrOH, rt. (3) benzoyl chloride, NEt₃, THF, rt. ^b LiAlH₄, THF, reflux. ^c BH₃-THF, THF, reflux. ^d PCl₃, NEt₃, toluene, -78 °C to rt, then H₂O.

Compound SI-22: To a stirred solution of **SI-21** (515 mg, 1.65 mmol) and WSCI (379 mg, 1.98 mmol) in DMF (6.6 mL) was added aniline (188 μ L, 2.06 mmol) at 0 °C. The reaction temperature was gradually warmed up to room temperature, and then the reaction mixture was kept stirring for 12 h. After dilution with ethyl acetate, the mixture was washed with 1N HCl (x 2) and brine (x 1), and the organic layer was dried over Na₂SO₄. After concentration in vacuo, the crude residue was washed with ether to afford the crude anilide (440 mg) as a white solid. A suspension of the crude anilide (388 mg, 1 mmol) and 5% Pd-C (53 mg, 0.025 mmol) in DMF (6 mL) and 2-propanol (4 mL) was stirred under H₂ atmosphere at room temperature. After 4 h, the mixture was filtered through celite, and the filtrate was evaporated under reduced pressure. After dilution with ethyl acetate, the organic layer was washed with water (x 2), and then dried over Na₂SO₄. After concentration in vacuo, the crude residue was obtained as pale yellow oil. To a stirred solution of the crude compound and triethylamine (0.21 mL, 1.5 mmol) in THF (10 mL) was added benzoyl chloride (0.14 mL, 1.2 mmol) at 0 °C. The reaction was stirred for 1 h, and then the solvent was under reduced pressure. After dilution with ethyl acetate, the organic layer was washed with NaHCO₃, water, and then dried over Na₂SO₄. Evaporation of organic solvent under reduced pressure, followed by ether wash of the residue gave pure **SI-22** as a white solid (310 mg, 61%: 3 steps). Mp: 213–215 °C; IR (KBr) ν 3277, 1659, 1632, 1602, 1533, 1446, 1304, 697 cm⁻¹; ¹H NMR (CDCl₃): δ 2.23–2.31 (m, 1H), 2.38–2.46 (m, 1H), 2.79–2.88 (m, 2H), 4.87–4.93 (m, 1H), 6.91 (d, J = 7.6 Hz, 1H), 7.08–7.12 (m, 1H), 7.11–7.31 (m, 7H), 7.42 (dd, J = 7.6 Hz, 7.6 Hz, 2H), 7.51–7.54 (m, 3H), 7.73 (dd, J = 1.6 Hz, 7.6 Hz, 2H), 8.64 (s, 1H); ¹³C NMR

(CDCl₃): δ 39.1, 33.4, 54.1, 120.0 (x 2), 124.5, 126.3, 127.1 (x 2), 128.4 (x 2), 128.7 (x 2), 128.7 (x 2), 129.0 (x 2), 132.1, 133.4, 137.6, 140.7, 167.9, 169.6; EI-LRMS m/z 358 (M⁺); $[\alpha]_D^{25} +39.2$ (c 0.59, DMF); EI-HRMS. Calcd for C₂₃H₂₂N₂O₂ (M⁺): 358.1681. Found: 358.1675.

Compound SI-23: To a stirred suspension of LAH (76 mg, 2.0 mmol) in THF (5 mL) was added **SI-22** (179.2 mg, 0.5 mmol) at 0 °C, and the mixture was refluxed for 12 h. After cooling down to room temperature, the reaction was quenched by the addition of water and 1N NaOH. The obtained white solid was filtered, and refluxed in THF (50 mL) for 3 h. After filtration, the combined organic layers were washed with brine, dried over Na₂SO₄, and then evaporated in vacuo. The residue was purified by flash column chromatography (SiO₂, hexane/ethyl acetate 3/1) to give a mixture of partially reduced monoamides (54.3 mg, 32%) and the desired diamine **SI-23** (108.4 mg, 66%). The obtained monoamides are subsequently reduced by 3 equivalents of BH₃·THF to afford the desired diamine 70 % yield (Total yield of **SI-23**: 88%; yellow oil). IR (neat) ν 3374, 3025, 2924, 2854, 1603, 1505, 1454, 1318, 748, 695 cm⁻¹; ¹H NMR (CDCl₃): δ 1.81–1.98 (m, 2H), 2.64–2.76 (m, 2H), 2.87–2.94 (m, 1H), 3.05 (dd, J = 6.8 Hz, 12.0 Hz, 1H), 3.24 (dd, J = 4.0 Hz, 12.0 Hz, 1H), 3.75 (d, J = 12.8 Hz, 1H), 3.79 (d, J = 12.8 Hz, 1H), 6.60–6.72 (m, 3H), 7.15–7.34 (m, 12H); ¹³C NMR (CDCl₃): δ 32.4, 34.5, 46.4, 50.8, 55.7, 113.0 (x 2), 117.2, 125.9, 127.0, 128.1 (x 2), 128.3 (x 2), 128.5 (x 2), 128.5 (x 2), 129.2 (x 2), 140.4, 141.9, 148.7; EI-LRMS m/z 330 (M⁺); $[\alpha]_D^{22} +13.1$ (c 1.26, CHCl₃); EI-HRMS. Calcd for C₂₃H₂₆N₂ (M⁺): 330.2096. Found: 330.2080.

Diaminophosphine Oxides 25 and 26: To a stirred solution of **SI-23** (100 mg, 0.30 mmol) and triethylamine (0.167 mL, 1.2 mmol) in toluene (3 mL) at –78 °C was added dropwise phosphorous trichloride (29 μ L, 0.33 mmol) over 1 min. The reaction temperature was gradually warmed up to room temperature. After stirring for 12 h, H₂O was added to the reaction, and then the mixture was stirred for 6 h. After filtration of the generated ammonium salt, the organic phase was concentrated in vacuo. The obtained residue was allowed to purify by flash column chromatography (SiO₂, hexane/ethyl acetate/triethylamine 300/100/4). Pure diaminophosphine oxide **25** (40.5 mg, 36%) and the diastereomer **26** (39.5 mg, 35%) were obtained as colorless oil.

Compound 25: IR (neat) ν 3446, 3060, 3027, 2925, 2862, 1601, 1503, 1455, 1303, 1232, 1159, 753, 700 cm⁻¹; ¹H NMR (CDCl₃): δ 1.70–1.80 (m, 1H), 1.97–2.07 (m, 1H), 2.49 (ddd, J = 6.4 Hz, 10.0 Hz, 14.0 Hz, 1H), 2.62 (ddd, J = 5.2 Hz, 10.4 Hz, 17.0 Hz, 1H), 3.29–3.35 (m, 1H), 3.54–3.62 (m, 1H), 3.73–3.79 (m, 1H), 4.20 (dd, J = 12.4 Hz (PNCH), 13.2 Hz (HCH), 1H), 4.40 (dd, J = 10.8 Hz (PNCH), 13.2 Hz (HCH), 1H), 7.00–7.45 (m, 15H), 7.68 (d, J = 627 Hz (O=PH), 1H); ¹³C NMR (CDCl₃): δ 30.6, 34.5 (d, J = 3.8 Hz), 46.4 (d, J = 6.5 Hz), 48.6 (d, J = 10.3 Hz), 53.6 (d, J = 7.3 Hz), 115.8 (d, J = 5.3 Hz) (x 2), 121.8, 126.2, 127.7, 128.2 (x 2), 128.6 (x 2), 128.6 (x 2), 128.7 (x 2), 129.5 (x 2), 136.8, 140.6, 141.6 (d, J = 7.6 Hz); ³¹P NMR (CDCl₃) δ 12.7; EI-LRMS m/z 376 (M⁺); $[\alpha]_D^{21} -27.7$ (c 0.45, CHCl₃); EI-HRMS. Calcd for C₂₃H₂₅N₂OP (M⁺): 376.1705. Found: 376.1720.

Compound 26: IR (neat) ν 3443, 2925, 1601, 1503, 1455, 1303, 1232, 1159, 753, 700 cm⁻¹; ¹H NMR (CDCl₃): δ 1.92–2.17 (m, 2H), 2.55 (ddd, J = 7.2 Hz, 10.0 Hz, 13.6 Hz, 1H), 2.79 (ddd, J = 5.2 Hz, 10.0 Hz, 13.6 Hz, 1H), 3.40–3.50 (m, 1H), 3.50–3.56 (m, 1H), 3.72–3.78 (m, 1H), 4.15 (dd, J = 11.2 Hz (PNCH), 15.6 Hz (HCH), 1H), 4.46 (dd, J = 8.8 Hz (PNCH), 15.6 Hz (HCH), 1H), 7.00–7.36 (m, 15H), 7.77 (d, J = 613 Hz (O=PH), 1H); ¹³C NMR (CDCl₃): δ 31.2, 33.4, 45.7 (d, J = 8.0 Hz), 49.3 (d, J = 10.3 Hz), 53.1 (d, J = 8.1 Hz), 115.8 (d, J = 5.4 Hz) (x 2), 121.8, 126.1, 127.8, 128.1 (x 2), 128.2 (x 2), 128.5 (x 2), 128.8 (x 2), 129.4 (x 2), 136.6 (d, J = 3.9 Hz), 140.7, 141.7 (d, J = 7.6 Hz); ³¹P NMR

(CDCl₃) δ 12.2; FAB-LRMS m/z 377 (MH⁺); [α]_D²⁵ +21.1 (c 0.53, CHCl₃).

The similarity of ¹H NMR profile corresponding to the 5-membered ring moiety of **25** to those of other ligands indicates that **25** has the same stereochemical arrangement on the phosphorus atom (Table S-4).

Table S-5.

Proton	6	22	23	24	25	26
H(a)	3.34 ppm	3.37 ppm	3.33 ppm	3.23 ppm	3.32 ppm	3.53 ppm
H(b)	3.79 ppm	3.81 ppm	3.78 ppm	3.71 ppm	3.76 ppm	3.75 ppm
H(c)	3.65 ppm	3.58 ppm	3.57 ppm	3.55 ppm	3.58 ppm	3.45 ppm
H(d)	7.70 ppm (J = 628 Hz)	7.62 ppm (J = 627 Hz)	7.67 ppm (J = 628 Hz)	7.65 ppm (J = 627 Hz)	7.68 ppm (J = 628 Hz)	7.77 ppm (J = 613 Hz)

4. Analytical Data

A. FAB MS and ³¹P NMR Spectrum for the Structural Analysis (S-23 ~ S-33)

B. ¹³C NMR Spectrum for the Purity Documentation (all new compounds in the asymmetric catalysis and the ligand synthesis, and known compounds synthesized by a new method) (S-34 ~ S-51)

[Mass Spectrum]

Date : 05-Jan-2005 16:41

Sample: yktn-170105-1

Note : -

Inlet : Direct

Ion Mode : FAB+

Spectrum Type : Normal Ion [MF-Linear]

RT : 0.16 min Scan# : 2

BP : m/z 91.0000 Int. : 201.39

Output m/z range : 0.0000 to 600.8902

Cut Level : 0.00 %

S-23

[Mass Spectrum]

Data : yk-tn-041203-ligand-2002 Date : 03-Dec-2004 16:58

Sample: yk-tn-041203-ligand

Note : -

Inlet : Direct

Ion Mode : FAB+

Spectrum Type : Normal Ion [MF-Linear]

RT : 0.16 min Scan# : (2,3)

BP : m/z 73.0000 Int. : 133.54

Output m/z range : 0.0000 to 605.3412 Cut Level : 0.00 %

S-24

[Mass Spectrum]

Data : yktn-041125-nba-dry001

Date : 25-Nov-2004 15:23

Sample: yktn-041126-nba-dry

Note : -

Inlet : Direct

Ion Mode : FAB+

Spectrum Type : Normal Ion [MF-Linear]

RT : 0.16 min Scan# : 2

BP : m/z 91.0000 Int. : 7.32

Output m/z range : 0.0000 to 1500.0000

Cut Level : 0.00 %

S-25

[Mass Spectrum]

Data : Yk-tn-041203-Pd-ligand-1-2-0 Date : 03-Dec-2004 17:07

Sample: Yk-tn-041203-Pd-ligand-1-2-

Note : -

Inlet : Direct Ion Mode : FAB+

Spectrum Type : Normal Ion [MF-Linear]

RT : 0.16 min Scan# : 2

BP : m/z 91.0000 Int. : 22.65

Output m/z range : 0.0000 to 1500.0000 Cut Level : 0.00 %

[Mass Spectrum]

Date : 05-Jan-2005 16:54
 Data : yktn-170105-1002
 Sample: yktn-170105-2-

Note : -
 Inlet : Direct Ion Mode : FAB+
 Spectrum Type : Normal Ion [MF-Linear]
 RT : 0.16 min Scan# : (2,3)
 BP : m/z 73.0000 Int. : 144.18
 Output m/z range : 0.0000 to 1500.0000 Cut Level : 0.00 %

S-27

DFILE Z:\YK-TN-05-13-31P_1_convert
 COMNT YK-TN-05-13-31P
 DATIM 25-SEP-2003 12:00:14
 OBNUC 31P
 EXMOD single_pulse_bil
 OBFRQ 161.00 MHz
 OBSET 834.00 KHz
 OBFIN 693.1 Hz
 POINT 32768
 FREQU 113636.4 Hz
 SCANS 166
 ACQTM 0.288 sec
 PD 1.000 sec
 PW1 4.0 us
 IRNUC 1H
 CTEMP 23.3 C
 SLVNT D2O
 EXREF 12.53 ppm
 BF 1.20 Hz
 RGAIN 15

DFILE Z:\No.8\YK-TN-08-70-31P-BSA
COMNT YK-TN-08-70-31P-BSA
DATIM 23-AUG-2004 16:51:13
OBNUC 31P
EXMOD single_pulse_bil
OBFRQ 161.00 MHz
OBSET 834.00 KHz
OBFIN 693.1 Hz
POINT 32768
FREQU 113636.4 Hz
SCANS 51
ACQTM 0.288 sec
PD 1.000 sec
PW1 4.0 us
IRNUC 1H
CTEMP 21.8 C
SLVNT CDCL3
EXREF 12.45 ppm
BF 1.20 Hz
RGAIN 15

S-29

Z:\No.9\YK-TN-09-37-1-1_1_convert.als
YK-TN-09-37-1-1

DFILE
COMNT
DATIM
OBNUC
EXMOD
OBFRQ
OBSET
OBFIN
POINT
FREQU
SCANS
ACQTM
PD
PWL
IRNUC
CTEMP
SLVNT
EXREF
BF
RGAIN

Z:\No.9\YK-TN-09-37-1-1_1_1_c
YK-TN-09-37-1-1
28-OCT-2004 16:39:36
31P
single_pulse_bil
161.00 MHz
834.00 KHz
693.1 Hz
32768
113636.4 Hz
154
0.288 sec
1.000 sec
4.0 us
1H
22.3 c
CDCL3
0.00 ppm
1.20 Hz
15

DFILE
COMNT
DATIM
OBNUC
EXMOD
OBFRQ
OBSET
OBFIN
POINT
FREQU
SCANS
ACQTM
PD
PW1
IRNUC
CTEMP
SLVNT
EXREF
BF
RGAIN

Z:\No.9\YK-TN-09-49-31P_1_c
YK-TN-09-49-31P
9-NOV-2004 16:04:07
31P
single_pulse bil
161.00 MHz
834.00 KHz
693.1 Hz
32768
113636.4 Hz
503
0.288 sec
1.000 sec
4.0 us
1H
22.6 c
CDCL3
0.00 ppm
1.20 Hz
15

101.722
99.965
98.015

12.686

Z:\No.9\YK-TN-09-50-31P_1_convert.als
YK-TN-09-50-31P

DFILE
COMNT
DATIM
OBNUC
EXMOD
OBFRO
OBSET
OBFIN
POINT
FREQU
SCANS
ACQTM
PD
PW1
IRNUC
CTEMP
SLVNT
EXREF
BF
RGAIN

31P
single_pulse_bil
161.00 MHz
834.00 KHz
693.1 Hz
32768
113636.4 Hz
201
0.288 sec
1.000 sec
4.0 us
1H
22.4 C
CDCL3
0.00 ppm
1.20 Hz
15

Z:\No10\YK-TN-10-13-31P_1_convert.als
YK-TN-10-13-31P

DFILE
COMNT
DATIM
OBNUC
EXMOD
OBFRO
OBSET
OBFIN
POINT
FREQU
SCANS
ACQTM
PD
PW1
IRNUC
CTEMP
SLVNT
EXREF
BF
RGAIN

Z:\No10\YK-TN-10-13-31P_1_c
YK-TN-10-13-31P
6-JAN-2005 15:15:50
31P
single_pulse_bil
161.00 MHz
834.00 KHz
693.1 Hz
32768
113636.4 Hz
97
0.288 sec
1.000 sec
4.0 us
1H
22.7 C
CDCL3
12.77 ppm
1.20 Hz
15

12.770

101.485
99.664

D:\nemoto\YK-TN-1-25-13C-DMSO-2_1.als
YK-TN-1-25-13C-DMSO-2

DFILE D:\nemoto\YK-TN-1-25-13C-DMSO-2_1.a
COMNT YK-TN-1-25-13C-DMSO-2
DATIM 29-JAN-2003 19:20:12
OBNUC 13C
EXMOD single pulse dec
OBFRQ 100.00 MHz
OBSET 536.00 KHz
OBFIN 362.3 Hz
POINT 65536
FREQU 25188.9 Hz
SCANS 429
ACQTM 2.602 sec
PD 1.500 sec
PWL 3.5 us
IRNUC 1H
CTEMP 21.6 c
SLVNT DMSO
EXREF 39.50 ppm
BF 1.20 Hz
RGAIN 24

2

D:\nemoto\YK-TN-1-37-13C-DMSO_1.als
YK-TN-1-37-1H-DMSO

DFILE D:\nemoto\YK-TN-1-37-13C-DMSO_1.als
COMNT YK-TN-1-37-1H-DMSO
DATIM 4-FEB-2003 19:05:21
OBNUC 13C
EXMOD single pulse dec
OBFRQ 100.00 MHz
OBSET 536.00 KHz
OBFIN 362.3 Hz
POINT 65536
FREQU 25188.9 Hz
SCANS 323
ACQTM 2.602 sec
PD 1.500 sec
PWL 3.5 us
IRNUC 1H
CTEMP 21.1 c
SLVNT DMSO
EXREF 39.50 ppm
BF 1.20 Hz
RGAIN 25

3

S-34

D:\nemoto\YK-TN-1-43-product-13C_1.als
YK-TN-1-43-product-13C

DFILE D:\nemoto\YK-TN-1-43-product-13C_1.
COMMT YK-TN-1-43-product-13C
DATIM 11-FEB-2003 20:18:43
OBNUC 13C
EXMOD single_pulse_dec
OBFRQ 100.00 MHz
OBSET 536.00 KHz
OBFIN 362.3 Hz
POINT 65536
FREQU 25188.9 Hz
SCANS 210
ACQTM 2.602 sec
PD 1.500 sec
PWL 3.5 us
IRNUC 1H
CTEMP 24.7 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 25

4.

Z:\YK-TN-02-19-CDCL3-13C_1.als
YK-TN-2-19-CDCL3-13C

DFILE Z:\YK-TN-02-19-CDCL3-13C_1.
COMMT YK-TN-2-19-CDCL3-13C
DATIM 2-APR-2003 11:53:33
OBNUC 13C
EXMOD single_pulse_dec
OBFRQ 100.00 MHz
OBSET 536.00 KHz
OBFIN 362.3 Hz
POINT 65536
FREQU 25188.9 Hz
SCANS 181
ACQTM 2.602 sec
PD 1.500 sec
PWL 3.5 us
IRNUC 1H
CTEMP 25.3 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 26

6

S-35

D:\13C NMR-7a.als
YK-TN-030823-cinnamyl.ethyl.6.13C

DFILE D:\13C NMR-7a.als
COMNT YK-TN-030823-cinnamyl.ethyl.6.13C
DATIM 23-AUG-2003 19:02:24
OBNUC 13C
EXMOD single pulse dec
OBFRQ 100.00 MHz
OBSET 536.00 KHz
OBFIN 362.3 Hz
POINT 65536
FREQU 25188.9 Hz
SCANS 43
ACQTM 2.602 sec
PD 1.500 sec
PW1 3.5 us
IRNUC 1H
CTEMP 23.2 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 25

9a

D:\13C NMR-7b.als
YK-TN-04-71-Bnester-13C

DFILE D:\13C NMR-7b.als
COMNT YK-TN-04-71-Bnester-13C
DATIM 13-SEP-2003 20:45:49
OBNUC 13C
EXMOD single pulse dec
OBFRQ 100.00 MHz
OBSET 536.00 KHz
OBFIN 362.3 Hz
POINT 65536
FREQU 25188.9 Hz
SCANS 151
ACQTM 2.602 sec
PD 1.500 sec
PW1 3.5 us
IRNUC 1H
CTEMP 23.3 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 25

9b

S-36

Z:\13C NMR-7c.als
YKMT-0273-BCM

DFILE Z:\13C NMR-7c.als
COMNT YKMT-0273-BCM
DATIM
OBNUC 13C
EXMOD SINGL-BCM
OBFRQ 100.40 MHz
OBSET 130.00 KHz
OBFIN 5500.0 Hz
POINT 32768
FREQU 27100.3 Hz
SCANS 320
ACQTM 1.209 sec
PD 3.700 sec
PW1 4.9 us
IRNUC 1H
CTEMP 23.7 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 24

9c

Z:\13C NMR-7d.als
YKMT-0273-5ME-BCM

DFILE Z:\13C NMR-7d.als
COMNT YKMT-0273-5ME-BCM
DATIM
OBNUC 13C
EXMOD SINGL-BCM
OBFRQ 100.40 MHz
OBSET 130.00 KHz
OBFIN 5500.0 Hz
POINT 32768
FREQU 27100.3 Hz
SCANS 320
ACQTM 1.209 sec
PD 3.700 sec
PW1 4.9 us
IRNUC 1H
CTEMP 24.5 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 24

9d

S-37

DFILE Z:\2004\YKMT-040110-5-BN-13
COMNT 13C-COM
DATIM
OBNUC 13C
EXMOD SINGL-BCM
OBFRQ 100.40 MHz
OBSET 130.00 KHz
OBFIN 5500.0 Hz
POINT 32768
FREQU 27100.3 Hz
SCANS 120
ACQTM 1.209 sec
PD 3.700 sec
PWL 4.9 us
IRNUC 1H
CTEMP 22.6 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 24

9e

DFILE D:\13C NMR-7e.als
COMNT YK-TN-00-7-Fr36-40-13C
DATIM 1-DEC-2003 19:50:14
OBNUC 13C
EXMOD single_pulse_dec
OBFRQ 100.00 MHz
OBSET 536.00 KHz
OBFIN 362.3 Hz
POINT 65536
FREQU 25188.9 Hz
SCANS 40
ACQTM 2.602 sec
PD 1.500 sec
PWL 3.5 us
IRNUC 1H
CTEMP 23.5 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 25

9f

S-38

DFILE D:\13C NMR-7f.als
COMNT YK-TN-00-8-cinnamyl-13C
DATIM 24-NOV-2003 20:50:59
OBNUC 13C
EXMOD single pulse dec
OBFRQ 100.00 MHz
OBSET 536.00 KHz
OBFIN 362.3 Hz
POINT 65536
FREQU 25188.9 Hz
SCANS 64
ACQTM 2.602 sec
PD 1.500 sec
PWL 3.5 us
IRNUC 1H
CTEMP 23.3 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 25

9g

DFILE D:\13C NMR-7g.als
COMNT YK-TN-04-72-tetralone-13C
DATIM 13-SEP-2003 21:03:23
OBNUC 13C
EXMOD single pulse dec
OBFRQ 100.00 MHz
OBSET 536.00 KHz
OBFIN 362.3 Hz
POINT 65536
FREQU 25188.9 Hz
SCANS 100
ACQTM 2.602 sec
PD 1.500 sec
PWL 3.5 us
IRNUC 1H
CTEMP 23.3 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 25

9h

S-39

D:\YK-TN-06-27-Me-13C_1.als
YK-TN-06-27-Me-13C

DFILE D:\YK-TN-06-27-Me-13C_1.als
COMNT YK-TN-06-27-Me-13C
DATIM 17-DEC-2003 22:45:50
OBNUC 13C
EXMOD single_pulse_dec
OBFRQ 100.00 MHz
OBSET 536.00 KHz
OBFIN 362.3 Hz
POINT 65536
FREQU 25188.9 Hz
SCANS 44
ACQTM 2.602 sec
PD 1.500 sec
PWL 3.5 us
IRNUC 1H
CTEMP 25.0 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 25

9i

D:\13C NMR-7i.als
YK-TN-00-cl-13C-data

DFILE D:\13C NMR-7i.als
COMNT YK-TN-00-cl-13C-data
DATIM 17-DEC-2003 16:55:14
OBNUC 13C
EXMOD single_pulse_dec
OBFRQ 100.00 MHz
OBSET 536.00 KHz
OBFIN 362.3 Hz
POINT 65536
FREQU 25188.9 Hz
SCANS 16
ACQTM 2.602 sec
PD 1.500 sec
PWL 3.5 us
IRNUC 1H
CTEMP 22.7 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 26

9j

S-40

DFILE D:\YK-TM-05-03-13C3_1.als
COMNT YK-TM-05-03-13C3
DATIM 16-DEC-2003 18:44:08
OBNUC 13C
EXMOD single_pulse_dec
OBFRQ 100.00 MHz
OBSET 536.00 KHz
OBFIN 362.3 Hz
POINT 65536
FREQU 25188.9 Hz
SCANS 25
ACQTM 2.602 sec
PD 1.500 sec
PWL 3.5 us
IRNUC 1H
CTEMP 22.5 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 25

9k

DFILE D:\13C NMR-7k.als
COMNT YK-TN-05-43-13C-data
DATIM 11-NOV-2003 21:04:47
OBNUC 13C
EXMOD single_pulse_dec
OBFRQ 100.00 MHz
OBSET 536.00 KHz
OBFIN 362.3 Hz
POINT 65536
FREQU 25188.9 Hz
SCANS 43
ACQTM 2.602 sec
PD 1.500 sec
PWL 3.5 us
IRNUC 1H
CTEMP 23.3 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 25

9l

5-41

D:\13C NMR-71.als
YK-TN-06-00-cyclo-prepara2nd-13C

DFILE D:\13C NMR-71.als
COMNT YK-TN-06-00-cyclo-prepara2nd-13C
DATIM 1-DEC-2003 23:10:24
OBNUC 13C
EXMOD single pulse dec
OBFRQ 100.00 MHz
OBSET 536.00 KHz
OBFIN 362.3 Hz
POINT 65536
FREQU 25188.9 Hz
SCANS 76
ACQTM 2.602 sec
PD 1.500 sec
PW1 3.5 us
IRNUC 1H
CTEMP 23.4 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 25

9m

Z:\2004\YKMT-OBACETATE-13C.NMF;1
YKMT-OBACETATE-13C

DFILE Z:\2004\YKMT-OBACETATE-13C
COMNT YKMT-OBACETATE-13C
DATIM
OBNUC 13C
EXMOD SINGL-BCM
OBFRQ 100.40 MHz
OBSET 130.00 KHz
OBFIN 5500.0 Hz
POINT 32768
FREQU 27100.3 Hz
SCANS 92
ACQTM 1.209 sec
PD 3.700 sec
PW1 4.9 us
IRNUC 1H
CTEMP 23.6 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 24

9m

5-42

DFILE E:\No.11\YK-TN-11-050418
COMNT YK-TN-11-050418-13C
DATIM 18-APR-2005 18:59:15
OBNUC 13C
EXMOD single_pulse_dec
OBFRQ 100.00 MHz
OBSET 536.00 KHz
OBFIN 362.3 Hz
POINT 65536
FREQU 25188.9 Hz
SCANS 43
ACQTM 2.602 sec
PD 1.500 sec
PWL 3.5 us
IRNUC 1H
CTEMP 22.9 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 28

9o

Z:\YKMT-6\YKMT-CHEX-13C.NMF;1
13C-COM

DFILE Z:\YKMT-6\YKMT-CHEX-13C.NMF
COMNT 13C-COM
DATIM
OBNUC 13C
EXMOD SINGL-BCM
OBFRQ 100.40 MHz
OBSET 130.00 KHz
OBFIN 5500.0 Hz
POINT 32768
FREQU 27100.3 Hz
SCANS 38
ACQTM 1.209 sec
PD 3.700 sec
PWL 4.9 us
IRNUC 1H
CTEMP 22.4 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 24

9l

S-43

Z:\YKMT-IPR-13C.NMF;1
13C-COM

DFILE Z:\YKMT-IPR-13C.NMF;1
COMNT 13C-COM
DATIM
OBNUC 13C
EXMOD SINGL-BCM
OBFRQ 100.40 MHz
OBSET 130.00 KHz
OBFIN 5500.0 Hz
POINT 32768
FREQU 27100.3 Hz
SCANS 37
ACQTM 1.209 sec
PD 3.700 sec
PW1 4.9 us
IRNUC 1H
CTEMP 22.6 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 24

9q

Z:\YKMT-ME-13C.NMF;1
13C-COM

DFILE Z:\YKMT-ME-13C.NMF;1
COMNT 13C-COM
DATIM
OBNUC 13C
EXMOD SINGL-BCM
OBFRQ 100.40 MHz
OBSET 130.00 KHz
OBFIN 5500.0 Hz
POINT 32768
FREQU 27100.3 Hz
SCANS 137
ACQTM 1.209 sec
PD 3.700 sec
PW1 4.9 us
IRNUC 1H
CTEMP 22.6 c
SLVNT CDCL3
EXREF 77.00 ppm
BF 1.20 Hz
RGAIN 24

9r

S-44

DFILE Z:\2004\YKMT-CHEX-SUBS
 COMNT 13C-COM
 DATIM
 OBNUC 13C
 EXMOD SINGL-BCM
 OBFRQ 100.40 MHz
 OBSET 130.00 KHz
 OBFIN 5500.0 Hz
 POINT 32768
 FREQU 27100.3 Hz
 SCANS 120
 ACQTM 1.209 sec
 PD 3.700 sec
 PW1 4.9 us
 IRNUC 1H
 CTEMP 23.5 c
 SLVNT CDCL3
 EXREF 77.00 ppm
 BF 1.20 Hz
 RGAIN 24

8i

Z:\YKBT-1-6.NMF;1
 13C-COM

DFILE Z:\YKBT-1-6.NMF;1
 COMNT 13C-COM
 DATIM
 OBNUC 13C
 EXMOD SINGL-BCM
 OBFRQ 100.40 MHz
 OBSET 130.00 KHz
 OBFIN 5500.0 Hz
 POINT 32768
 FREQU 27100.3 Hz
 SCANS 20
 ACQTM 1.209 sec
 PD 3.700 sec
 PW1 4.9 us
 IRNUC 1H
 CTEMP 23.1 c
 SLVNT CDCL3
 EXREF 77.00 ppm
 BF 1.20 Hz
 RGAIN 24

11

(S-45)

DFILE Z:\No.9\YK-TN-09-1026-1.
 COMNT YK-TN-09-1026-13C
 DATIM 26-OCT-2004 19:11:35
 OBNUC 13C
 EXMOD single_pulse_dec
 OBFRQ 100.00 MHz
 OBSET 536.00 KHz
 OBFIN 362.3 Hz
 POINT 65536
 FREQU 25188.9 Hz
 SCANS 120
 ACQTM 2.602 sec
 PD 1.500 sec
 PW1 3.5 us
 IRNUC 1H
 CTEMP 23.4 c
 SLVNT CDCL3
 EXREF 77.00 ppm
 BF 1.20 Hz
 RGAIN 26

12

:\\YK-TN-05-65-ligand-Ac-13C_1.als
 K-TN-05-65-ligand-Ac-13C

DFILE Z:\\YK-TN-05-65-ligand-Ac
 COMNT YK-TN-05-65-ligand-Ac-13
 DATIM 10-NOV-2003 15:19:59
 OBNUC 13C
 EXMOD single_pulse_dec
 OBFRQ 100.00 MHz
 OBSET 536.00 KHz
 OBFIN 362.3 Hz
 POINT 65536
 FREQU 25188.9 Hz
 SCANS 161
 ACQTM 2.602 sec
 PD 1.500 sec
 PW1 3.5 us
 IRNUC 1H
 CTEMP 23.6 c
 SLVNT CDCL3
 EXREF 77.00 ppm
 BF 1.20 Hz
 RGAIN 25

22

(S-46)

DFIL 4: \2004\ykmt-n-me-13c_1
 COMNT ykmt-n-me-13c
 DATIM 8-NOV-2004 22:15:17
 OBNUC 13C
 EXMOD single_pulse_dec
 OBFRQ 100.00 MHz
 OBSET 536.00 KHz
 OBFIN 362.3 Hz
 POINT 65536
 FREQU 25188.9 Hz
 SCANS 503
 ACQTM 2.602 sec
 PD 1.500 sec
 PW1 3.5 us
 IRNUC 1H
 CTEMP 23.6 c
 SLVNT CDCL3
 EXREF 77.00 ppm
 BF 1.20 Hz
 RGAIN 26

23

Z:\YK-TN-07-40-GluL-13C_1.als
 YK-TN-07-40-GluL-13C

DFIL Z:\YK-TN-07-40-GluL-13C
 COMNT YK-TN-07-40-GluL-13C
 DATIM 13-MAR-2004 15:43:19
 OBNUC 13C
 EXMOD single_pulse_dec
 OBFRQ 100.00 MHz
 OBSET 536.00 KHz
 OBFIN 362.3 Hz
 POINT 65536
 FREQU 25188.9 Hz
 SCANS 77
 ACQTM 2.602 sec
 PD 1.500 sec
 PW1 3.5 us
 IRNUC 1H
 CTEMP 23.5 c
 SLVNT CDCL3
 EXREF 77.00 ppm
 BF 1.20 Hz
 RGAIN 26

24

(C-47)

DFILE Z:\No10\YK-TN-10-04-pr
 COMNT YK-TN-10-04-preparaLP-
 DATIM 20-DEC-2004 19:11:15
 OBNUC 13C
 EXMOD single_pulse_dec
 OBFRQ 100.00 MHz
 OBSET 536.00 KHz
 OBFIN 362.3 Hz
 POINT 65536
 FREQU 25188.9 Hz
 SCANS 255
 ACQTM 2.602 sec
 PD 1.500 sec
 PW1 3.5 us
 IRNUC 1H
 CTEMP 23.0 c
 SLVNT CDCL3
 EXREF 77.00 ppm
 BF 1.20 Hz
 RGAIN 26

25

Z:\No.8\YK-TN-08-GluZ-13C_1.als
 YK-TN-08-GluZ-13C

DFILE Z:\No.8\YK-TN-08-GluZ-13C
 COMNT YK-TN-08-GluZ-13C
 DATIM 2-JUL-2004 17:11:48
 OBNUC 13C
 EXMOD single_pulse_dec
 OBFRQ 100.00 MHz
 OBSET 536.00 KHz
 OBFIN 362.3 Hz
 POINT 65536
 FREQU 25188.9 Hz
 SCANS 482
 ACQTM 2.602 sec
 PD 1.500 sec
 PW1 3.5 us
 IRNUC 1H
 CTEMP 23.0 c
 SLVNT DMSO
 EXREF 39.50 ppm
 BF 1.20 Hz
 RGAIN 25

51-17

(S-48)

DFILE E:\No.9\YK-TN-041114-Gluamide-1:
 COMNT YK-TN-041114-Gluamide-13C
 DATIM 14-NOV-2004 17:28:05
 OBNUC 13C
 EXMOD single_pulse_dec
 OBFRQ 100.00 MHz
 OBSET 536.00 KHz
 OBFIN 362.3 Hz
 POINT 65536
 FREQU 25188.9 Hz
 SCANS 1264
 ACQTM 2.602 sec
 PD 1.500 sec
 PW1 3.5 us
 IRNUC 1H
 CTEMP 23.3 c
 SLVNT DMSO
 EXREF 39.50 ppm
 BF 1.20 Hz
 RGAIN 25

SI-18

::\No.11\YK-TN-11-050416-13C_1.als
 YK-TN-11-050416-13C

DFILE Z:\No.11\YK-TN-11-050416-13C
 COMNT YK-TN-11-050416-13C
 DATIM 16-APR-2005 18:40:33
 OBNUC 13C
 EXMOD single_pulse_dec
 OBFRQ 100.00 MHz
 OBSET 536.00 KHz
 OBFIN 362.3 Hz
 POINT 65536
 FREQU 25188.9 Hz
 SCANS 223
 ACQTM 2.602 sec
 PD 1.500 sec
 PW1 3.5 us
 IRNUC 1H
 CTEMP 22.9 c
 SLVNT CDCL3
 EXREF 77.00 ppm
 BF 1.20 Hz
 RGAIN 29

SI-20

(S-49)

DFILE 2:\NO10\YK-TN-10-02-SM
 COMNT YK-TN-10-02-SM-13C
 DATIM 26-DEC-2004 15:03:50
 OBNUC 13C
 EXMOD single_pulse_dec
 OBFRQ 100.00 MHz
 OBSET 536.00 KHz
 OBFIN 362.3 Hz
 POINT 65536
 FREQU 25188.9 Hz
 SCANS 1000
 ACQTM 2.602 sec
 PD 1.500 sec
 PW1 3.5 us
 IRNUC 1H
 CTEMP 22.7 c
 SLVNT CDCL3
 EXREF 77.00 ppm
 BF 1.20 Hz
 RGAIN 25

SI-22

E:\No10\YK-TN-10-02-mp-13C_1.als
 YK-TN-10-02-mp-13C

DFILE E:\No10\YK-TN-10-02-mp-13
 COMNT YK-TN-10-02-mp-13C
 DATIM 16-DEC-2004 21:49:05
 OBNUC 13C
 EXMOD single_pulse_dec
 OBFRQ 100.00 MHz
 OBSET 536.00 KHz
 OBFIN 362.3 Hz
 POINT 65536
 FREQU 25188.9 Hz
 SCANS 201
 ACQTM 2.602 sec
 PD 1.500 sec
 PW1 3.5 us
 IRNUC 1H
 CTEMP 22.7 c
 SLVNT CDCL3
 EXREF 77.00 ppm
 BF 1.20 Hz
 RGAIN 26

SI-23

(S-50)

DFILE Z:\No.11\YK-TN-11-45-di
 COMNT YK-TN-11-45-diastereome
 DATIM 20-JUN-2005 17:09:22
 OBNUC 13C
 EXMOD single_pulse_dec
 OBFRQ 100.00 MHz
 OBSET 536.00 KHz
 OBFIN 361.1 Hz
 POINT 65536
 FREQU 25188.9 Hz
 SCANS 370
 ACQTM 2.602 sec
 PD 1.500 sec
 PW1 3.5 us
 IRNUC 1H
 CTEMP 22.8 c
 SLVNT CDCL3
 EXREF 77.00 ppm
 BF 1.20 Hz
 RGAIN 28

26

S-51