Di(azacrown) Conjugates of 2'-*O*-Methyl Oligoribonucleotides as Sequence-selective Artificial Ribonucleases

*Teija Niittymäki, Pasi Virta, Kaisa Ketomäki and Harri Lönnberg**

Department of Chemistry, University of Turku, FIN-20014 Turku, Finland.

harlon@utu.fi

SUPPORTING INFORMATION

Table of Contents

General procedures	S2
Figure S1. ¹ H NMR (CDCl ₃ , 500 MHz) spectrum of 14	S3
Figure S2. ³¹ P NMR (CDCl ₃ , 200 MHz) spectrum of 14	S3
Figure S3. ¹ H NMR (CDCl ₃ , 500 MHz) spectrum of 23a	S4
Figure S4. ¹ H NMR (CDCl ₃ , 500 MHz) spectrum of 23a	S5
Figure S5. ¹ H NMR (CDCl ₃ , 500 MHz) spectrum of 24	S6
Figure S6. ³¹ P NMR (CD ₃ CN, 200 MHz) spectrum of 24	S6
Figure S7. ¹ H NMR (CDCl ₃ , 500 MHz) spectrum of 23b	S7
Figure S8. ¹ H NMR (CDCl ₃ , 500 MHz) spectrum of 23b	S8
Figure S9. ¹ H NMR (CD ₃ CN, 500 MHz) spectrum of 25	S9
Figure S10. ³¹ P NMR (CD ₃ CN, 200 MHz) spectrum of 25	S9
Figure S11. ¹ H NMR (CD ₃ CN, 500 MHz) spectrum of 28	S10

Figure S12. ^{31}P NMR (CD_3CN , 200 MHz) spectrum of 28	S10
Figure S13. RP HPLC chromatogram of purified 1a	S11
Figure S14. RP HPLC chromatogram of crude and purified 2a	S11
Figure S15. RP HPLC chromatogram of purified 3a	S12
Figure S16. RP HPLC chromatogram of purified 1b	S12
Figure S17. RP HPLC chromatogram of purified 2b	S13
Figure S18. RP HPLC chromatogram of purified 3b	S13

General procedures

The NMR spectra were recorded in deuteriochloroform and deuterioacetonitrile at 25°C on a 500 MHz spectrometer. For examples of the NMR spectra, see **S1-S12**. The mass spectra were recorded using EI (**11** and **12**), ESI (**1a-3a**, **1b**, **2b**, **14**, **18a,b-23a,b**, **24-28**) and MALDI (**3b**) ionization.

A Thermo Hypersil C18 (250 × 10 mm, 5μm) RP column was used for the purification of the conjugates (**1a,b-3a,b**) (flow rate of 3.0 mL/min). A linear gradient from 10% B in A to 30% B in A in 30 min was applied. A: 50 mmol L⁻¹ NH₄OAc in H₂O and B: 50 mmol L⁻¹ NH₄OAc in H₂O-acetonitrile (1:1, v/v). The wavelength of UV detection was 260nm. For examples of RP HPLC profiles, see **S13-S18**.

Figure S1. ^1H NMR (CDCl_3 , 500 MHz) spectrum of **14**

Figure S2. ^{31}P NMR (CDCl₃, 200 MHz) spectrum of **14**

Figure S3. ¹H NMR (CDCl₃, 500 MHz) spectrum of **23a**

Figure S4. ^1H NMR (CDCl_3 , 500 MHz) spectrum of **23a**

Figure S5. ^1H NMR (CDCl_3 , 500 MHz) spectrum of **24**

Figure S6. ³¹P NMR (CD₃CN, 200 MHz) spectrum of **24**

Figure S7. ¹H NMR (CDCl₃, 500 MHz) spectrum of **23b**

Figure S8. ^1H NMR (CDCl_3 , 500 MHz) spectrum of **23b**

Figure S9. ^1H NMR (CD_3CN , 500 MHz) spectrum of **25**

Figure S10. ^{31}P NMR (CD_3CN , 200 MHz) spectrum of **25**

Figure S11. ^1H NMR (CD_3CN , 500 MHz) spectrum of **28**

Figure S12. ^{31}P NMR (CD_3CN , 200 MHz) spectrum of **28**

Figure S13. RP HPLC chromatogram of purified **1a**

2a

Figure S15. RP HPLC chromatogram of purified **3a**

Figure S16. RP HPLC chromatogram of purified **1b**

Figure S17. RP HPLC chromatogram of purified **2b**

Figure S18. RP HPLC chromatogram of purified **3b**