SUPPORTING INFORMATION

Diastereoselective, Titanium-Mediated Cyclization of ω -Vinyl Tethered Imides

Soumava Santra, Nikolai Masalov, Oleg L. Epstein, and Jin Kun Cha*
Department of Chemistry, Wayne State University, 5101 Cass Ave,
Detroit, MI 48202

Typical procedure: reductive coupling of *N*-3-butenylsuccinimide (1**).** To a solution of titanium(IV) isopropoxide (0.45 mL, 1.5 mmol) in 4 mL of anhydrous THF, which had been cooled to $-78\text{ }^{\circ}\text{C}$, was added dropwise a 1.6 M solution of *n*-BuLi in hexane (1.9 mL, 3 mmol) under a gentle stream of argon. The flask was placed between an argon inlet and a bubbler. After the resulting clear orange solution had been stirred at the same temperature for 15–20 min, the bath was removed and the reaction mixture was allowed to warm to room temperature during 10–15 min. The mixture became dark green and was stirred at room temperature for 15 min. The dark homogeneous solution was cooled to $0\text{ }^{\circ}\text{C}$ (ice bath), and a solution of **1** (77 mg, 0.5 mmol) in anhydrous THF (1 mL) was then added dropwise. The resulting deep red reaction mixture was stirred at $0\text{ }^{\circ}\text{C}$ for 20 min, quenched by slow addition of 0.5 mL of water (the suspension became dark blue), exposed to air, and stirred until it became white (15–30 min). Ethyl acetate (5 mL) and anhydrous MgSO_4 (1–2 g) were added to the reaction mixture and the resulting slurry was stirred for an additional 15 min. After filtration, solids were washed thoroughly with EtOAc–MeOH (9:1) and the combined organic filtrates were concentrated under reduced pressure. The residue was purified by column chromatography on silica gel, using EtOAc as eluent to afford 70 mg (90%) of **2**.

4a. IR (neat) 3377, 1679 cm^{-1} ; ^1H NMR (500 MHz, benzene- d_6) δ 7.45–7.53 (m, 2 H), 7.17–7.23 (m, 2 H), 7.10 (m, 1 H), 4.64 (t, $J = 8.6$ Hz, 1 H), 4.03 (br s, 1 H), 2.62 (dt, $J = 7.1, 17.1$ Hz, 1 H), 2.10 (dt, $J = 7.1, 12.7$ Hz, 1 H), 2.02 (ddd, $J = 2.6, 10.1, 17.1$ Hz, 1 H), 1.96 (m, 1 H), 1.66 (m, 1 H), 1.53 (dt, $J = 9.3, 13.7$ Hz, 1 H), 1.28 (m, 1 H), 0.88 (d, $J = 6.6$ Hz, 3 H); ^{13}C NMR (125 MHz, benzene- d_6) δ 175.1, 143.6, 128.5, 127.1, 127.0, 97.4, 58.2, 44.4, 43.7, 33.2, 32.7, 11.7; HRMS (M^+) calcd for $\text{C}_{14}\text{H}_{17}\text{NO}_2$ 231.1259, found 231.1257.

5a. IR (neat) 3251, 1668 cm^{-1} ; ^1H NMR (500 MHz, acetone- d_6) δ 7.12–7.29 (m, 5 H), 4.72 (br s, 1 H), 4.64 (d, $J = 9.1$ Hz, 1 H), 2.90 (m, 1 H), 2.62 (dt, $J = 9.1, 12.7$ Hz, 1 H), 2.20–2.42 (m, 3 H), 2.17 (dd, $J = 8.1, 12.7$ Hz, 1 H), 1.88 (dd, $J = 6.6, 12.2$ Hz, 1 H), 1.03 (d, $J = 6.6$ Hz, 3 H); ^{13}C NMR (125 MHz, acetone- d_6) δ 171.5, 143.3, 129.0, 127.4, 126.9, 98.9, 56.5, 45.5, 40.4, 35.3, 33.6, 12.4; HRMS (M^+) calcd for $\text{C}_{14}\text{H}_{17}\text{NO}_2$ 231.1259, found 231.1256.

5b. IR (neat) 3411, 1682 cm^{-1} ; ^1H NMR (400 MHz, acetone- d_6) δ 7.08–7.29 (m, 5 H), 4.63 (br s, 1 H), 4.50 (m, 1 H), 2.40 (m, 1 H), 2.12–2.24 (m, 2 H), 2.06 (m, 2 H), 1.87 (m, 1 H), 1.65 (m, 2 H), 1.51 (m, 1 H), 1.05 (d, $J = 6.5$ Hz, 3 H); ^{13}C NMR (100 MHz, acetone- d_6) δ 175.3, 144.1, 128.5, 127.2, 126.7, 91.3, 55.2, 40.6, 34.5, 33.6, 29.2, 27.0, 15.8; HRMS ($\text{M}^+ - \text{H}_2\text{O}$) calcd for $\text{C}_{15}\text{H}_{17}\text{NO}$ 227.1310, found 227.1311.

5c. IR (neat) 3354, 1633 cm^{-1} ; ^1H NMR (500 MHz, acetone- d_6) δ 7.07–7.27 (m, 5 H), 4.98 (d, $J = 9.1$ Hz, 1 H), 4.57 (br s, 1 H), 2.27 (m, 4 H), 2.16 (m, 2 H), 1.75–1.92 (m, 2 H), 1.61 (dd, $J = 6.1, 12.2$ Hz, 1 H), 1.00 (d, $J = 6.6$ Hz, 3 H); ^{13}C NMR (125 MHz, acetone- d_6) δ 169.9, 145.3, 128.9, 126.9, 126.1, 90.6, 59.4, 42.3, 39.0, 32.9, 32.0, 18.0, 12.6; HRMS (M^+) calcd for $\text{C}_{15}\text{H}_{19}\text{NO}_2$ 245.1416, found 245.1414.

5d. IR (neat) 3232, 1611 cm^{-1} ; ^1H NMR (500 MHz, acetone- d_6) δ (easily epimerized via the ring opened form) 7.10–7.30 (m, 5 H), 5.43 (t, $J = 4.4$ Hz, 1 H), 4.53 (br s, 1 H), 2.55 (m, 1 H), 2.30–2.47 (m, 2 H), 2.24 (m, 1 H), 2.05–2.20 (m, 2 H), 1.75–1.90 (m, 3 H), 1.44 (m, 1 H), 1.21 (1.38) (m, 1 H), 0.92 (d, $J = 7.1$ Hz, 3 H); ^{13}C NMR (125 MHz, acetone- d_6) δ 170.6, 144.2, 128.7, 126.5, 86.0, 52.9, 36.8, 34.8, 34.7, 32.4, 26.3, 24.3, 16.7, 16.5; HRMS ($\text{M}+\text{H}^+$) calcd for $\text{C}_{16}\text{H}_{22}\text{NO}_2$ 260.1651, observed 260.1655.

7a. IR (neat) 3336, 1673 cm^{-1} ; ^1H NMR (500 MHz, acetone- d_6) δ 7.20–7.40 (m, 5 H), 4.95 (br s, 1 H), 3.30–3.60 (m, 3 H), 2.93 (m, 1 H), 2.20–2.40 (m, 2 H), 2.00–2.10 (m, 2 H), 0.92 (d, $J = 6.7$ Hz, 3 H); ^{13}C NMR (125 MHz, acetone- d_6) δ 174.5, 142.2, 129.4, 128.7, 127.6, 98.2, 52.9, 51.6, 49.0, 33.4, 33.3, 10.6.

7b. IR (neat) 3314, 1669 cm^{-1} ; ^1H NMR (500 MHz, acetone- d_6) δ 7.18–7.35 (m, 5 H), 4.63 (br s, 1 H), 3.86 (ddd, $J = 2.0, 5.1, 13.1$ Hz, 1 H), 2.99 (ddt, $J = 1.5, 3.8, 13.1$ Hz, 1 H), 2.87 (dt, $J = 3.6, 11.8$ Hz, 1 H), 2.44 (m, 1 H), 2.23 (m, 1 H), 2.12 (m, 2 H), 1.83 (m, 1 H), 1.74 (m, 1 H), 1.63 (dt, $J = 5.0, 12.7$ Hz, 1 H), 0.73 (d, $J = 7.1$ Hz, 3 H); ^{13}C NMR

(125 MHz, acetone- d_6) δ 172.8, 145.7, 129.3, 128.4, 127.1, 90.3, 47.2, 45.6, 36.9, 34.3, 33.6, 30.4, 13.3.

7c. IR (neat) 3357, 1624 cm^{-1} ; ^1H NMR (500 MHz, acetone- d_6) δ 7.20–7.40 (m, 5 H), 4.7 (br s, 1 H), 3.75 (dd, $J = 9.1, 11.6$ Hz, 1 H), 3.44 (dd, $J = 9.1, 11.6$ Hz, 1 H), 3.22 (dt, $J = 9.1, 11.6$ Hz, 1 H), 2.32 (m, 1 H), 2.22 (m, 2 H), 2.09 (m, 2 H), 1.79 (m, 1 H), 1.62 (m, 1 H), 0.90 (d, $J = 6.6$ Hz, 3 H); ^{13}C NMR (125 MHz, acetone- d_6) δ 169.1, 142.2, 129.4, 128.7, 127.6, 90.3, 53.1, 52.3, 47.5, 33.8, 31.9, 17.5, 11.3.

7d. IR (neat) 3357, 1672 cm^{-1} ; ^1H NMR (500 MHz, acetone- d_6) δ 7.14–7.33 (m, 5 H), 4.62 (br s, 1 H), 4.55 (ddd, $J = 2.5, 4.6, 13.2$ Hz, 1 H), 3.02 (dt, $J = 3.1, 13.2$ Hz, 1 H), 2.88 (dt, $J = 3.6, 11.9$ Hz, 1 H), 2.20–2.35 (m, 2 H), 2.09–2.18 (m, 2 H), 1.86–1.95 (m, 2 H), 1.71 (m, 2 H), 1.59 (m, 1 H), 0.75 (d, $J = 6.6$ Hz, 3 H); ^{13}C NMR (125 MHz, acetone- d_6) δ 169.3, 146.5, 129.3, 128.4, 126.9, 86.2, 47.1, 45.9, 36.9, 36.3, 35.0, 33.7, 16.9, 13.8; HRMS (MH^+) calcd for $\text{C}_{16}\text{H}_{22}\text{NO}_2$ 260.1651, observed 260.1660.

Dalt 500 spectrometer

Unity 500 spectrometer

Mercury 400 spectrometer

5b

5b

Mercury 400 spectrometer

Unity 500 spectrometer

5c

5d

Unity 500 spectrometer

NMR 500 spectrometer

7a

7a

Unity 500 spectrometer

Unity 500 spectrometer

7b

Dulky 500 spectrometer

7c

Daley 500 spectrometer

7d