DMSO Mediated Elimination Reactions in Aryl 1,2-Dihalopropanoates: Scope and Mechanistic Insights

Wei, Li, ^{*1} Jianchang Li, ¹ Melissa Lin, ² Sumrit Wacharasindhu, ² Keiko Tabei, ² and Tarek S. Mansour²

¹Chemical and Screening Sciences, Wyeth Research, 200 Cambridge Park Drive, Cambridge, MA 02140, USA and ²Chemical and Screening Sciences, Wyeth Research, 401 North Middletown Road, Pearl River, NY 10965, USA

weili@wyeth.com

LIST OF CONTENTS

General experimental	S2
¹H, ¹³C NMR spectra	S4-42
Dibenzosuberone	S4
Methyl Cinnamate	S6
(E)-Methyl 3-(4-chlorophenyl)acrylate	S8
Cinnamic Acid	S10
(E)-3-(4-chlorophenyl)acrylic acid	S11
Cinnamamide	S12
(E)-chalcone	S14
2H-chromen-2-one	S16
β-Nitrostyrene	S18
(E)-ethyl 3-(2-chloropyridin-3-yl)acrylate	S20
(E)-ethyl 3-(6-chloropyridin-3-yl)acrylate	S21
(E)-ethyl 3-(2-chloropyridin-4-yl)acrylate	S23
(E)-ethyl 3-(pyridin-3-yl)acrylate	S25
(E)-1,2-di(pyridin-3-yl)ethane	S27
(E)-N,N-dimethyl-4-(2-(pyridin-4-yl)vinyl)aniline	S29
(Z)-2-Bromo-3-(3-methyl-thiophen-2-yl)-acrylic acid methyl ester	S31
(Z)-Ethyl 2-bromo-3-(4-bromo-5-phenylthiophen-2-yl)acrylate	S33
(E)-Ethyl 3-(4-bromothiophen-2-yl)acrylate	S35
(Z)-Ethyl 2-bromo-3-(4-bromothiophen-2-yl)acrylate	S37
(E)-Ethyl 3-(4-phenylthiophen-2-yl)acrylate	S39
(Z)-Ethyl 2-Bromo-3-(4-phenyl-thiophen-2-yl)-acrylate	S41

General experimental: All reagents and solvents were purchased and used without further purification. All the starting dihalides are commercially available and were used as received after purity check. All products were characterized by ^1H and ^{13}C NMR, MS, or elemental analysis. Mass spectrometers used were Hewlett Packard 5972 series and 5890 series II Plus GC for very small molecules, Micromass LCT (ESI or APCI) with Agilent 1100 LC Pump (for LCMS), and Micromass Q-Tof (ESI or APCI) with Agilent 1100 LC Pump (for LCMSMS). Although the reactions in Table 1 were carried out in neat DMSO, using solvents such as THF or ethyl acetate gave similar results. NMR experiments were performed on either a Bruker AVANCE 400 MHz spectrometer equipped with a 5 mm broadband probe or a AVANCE 500 MHz spectrometer equipped with a 5mm TXI CryoProbe using DMSO-*d*₆ as the solvent for both proton and carbon NMR unless otherwise indicated. Fluorine-carbon coupling constants are also included in the ^{13}C NMR data for compounds containing fluorine substituents. The following abbreviations are used to designate the multiplicity: s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet.

General DMSO reductive debromination procedures: The 1,2-dibromo compounds used in these studies were prepared by addition of bromine in either carbon tetrachloride or carbon disulfide according to literature procedures.¹ A typical procedure for the reductive debromination is illustrated by the preparation of dibenzosuberone: a solution of *trans*-10,11-dibromodibenzosuberone (3.66 g, 0.01 M) in 30 ml of DMSO is heated at 75 °C for 2 hour. The reaction mixture is poured into water (30 ml), and the resulting mixture

1 a) Cossy, J.; Pete J. P. *Bulletin de la Societe Chimique de France* **1979**, 559-67. b) Sanchez B. J.; Hernandez M. D.; McPhee D. J. *Boletin de la Sociedad Quimica del Peru* **1982**, 48, 195-205.

is extracted with ethyl acetate (60 ml x 2). The extract is washed with water and brine, dried over anhydrous sodium sulfate, and concentrated under reduced pressure to give 2.01 g (yield 98%) of the dibenzosuberone as a white solid. ^1H NMR (CDCl_3) δ 8.21 (dd, $J = 5.9, 1.3$ Hz, 2 H), 7.61-7.42 (m, 6H), 7.06 (s, 2 H). ^{13}C NMR (400 MHz, $\text{DMSO}-d_6$) δ 192.4, 137.7 (2C), 134.4 (2C), 132.5 (2C), 131.4 (2C), 130.9 (2C), 129.5 (2C), 129.1 (2C).

Dibenzosuberone (entry 1 in Table 1)

Dibenzosuberone (entry 1 in Table 1)

Methyl Cinnamate (entry 2 in Table 1)

Methyl Cinnamate (entry 2 in Table 1)

(E)-Methyl 3-(4-chlorophenyl)acrylate (entry 3 in Table 1)

(E)-Methyl 3-(4-chlorophenyl)acrylate (entry 3 in Table 1)

Cinnamic Acid (entry 4 in Table 1)

(E)-3-(4-chlorophenyl)acrylic acid (entry 5 in Table 1)

Cinnamamide (entry 6 in Table 1)

Cinnamamide (entry 6 in Table 1)

(E)-chalcone (entry 7 in Table 1)

(E)-chalcone (entry 7 in Table 1)

2H-chromen-2-one: (entry 8 in Table 1)

2H-chromen-2-one: (entry 8 in Table 1)

β -Nitrostyrene (entry 9 in Table 1)

β -Nitrostyrene (entry 9 in Table 1)

(E)-ethyl 3-(2-chloropyridin-3-yl)acrylate (entry 1 in Table 2)

(E)-ethyl 3-(6-chloropyridin-3-yl)acrylate (entry 2 in Table 2)

05-297396LNC/L33659-29-Al/Sumrit Wacharasindhu/CDCl3/L05-9467

Current Data Parameters
NAME 297396
EXTNO 11
PROCNO 1

F2 - Acquisition Parameters
Date_ 20051003
Time 20.34

INSTRUM spect
PULPROG 5 mm QNP 1H/13
TD 65536
SOLVENT CDCl3
NS 3072
DS 2
SWH 29239.766 Hz
FIDRES 0.46163 Hz
AQ 1.1207156 sec
RG 2048
DM 17.100 usec
DE 4.50 usec
TE 300.0 K
D1 0.20000000 sec
d11 0.03000000 sec
d12 0.00002000 sec

===== CHANNEL f1 =====
NUC1 13C
P1 5.00 usec
PL1 0.00 dB
SFO1 100.6757321 MHz

===== CHANNEL f2 =====
CPDPRG2 waltz16
NUC2 1H
PCPD2 91.50 usec
PL2 1.00 dB
PL12 15.00 dB
PL13 16.00 dB
SFO2 400.3316013 MHz

F2 - Processing parameters
SI 32768
SF 100.6630190 MHz
WDW EM
SSB 0
LB 1.00 Hz
GB 0
PC 1.40

1D NMR plot parameters
CX 20.00 cm
CY 14.00 cm
FIP 213.000 ppm
F1 21642.55 Hz
F2P -5.000 ppm
F2 -503.31 Hz
PPMCM 11.00000 ppm/cm
HZCM 1107.29321 Hz/cm

Chemical structure: CCOC(=O)/C=C/c1cc(Cl)nc1

Peak list (ppm, integration):

Chemical Shift (ppm)	Integration
1.4	3.00
4.1	2.00
4.3	1.00
6.8	1.00
7.2	1.00
7.5	1.00

(E)-ethyl 3-(2-chloropyridin-4-yl)acrylate (entry 3 in Table 2)

(E)-ethyl 3-(2-chloropyridin-4-yl)acrylate (entry 3 in Table 2)

(E)-ethyl 3-(pyridin-3-yl)acrylate (entry 4 in Table 2)

(E)-ethyl 3-(pyridin-3-yl)acrylate (entry 4 in Table 2)

05-297386LNH/L33659-77-A1/Sumrit Wacharasindhu/CDC13/L05-9464

(E)-1,2-di(pyridin-3-yl)ethene (entry 5 in Table 2)

(E)-N,N-dimethyl-4-(2-(pyridin-4-yl)vinyl)aniline (entry 6 in Table 2)

(E)-N,N-dimethyl-4-(2-(pyridin-4-yl)vinyl)aniline (entry 6 in Table 2)

(Z)-2-Bromo-3-(3-methyl-thiophen-2-yl)-acrylic acid methyl ester (entry 2 in Table 3)

(Z)-2-Bromo-3-(3-methyl-thiophen-2-yl)-acrylic acid methyl ester (entry 2 in Table 3)

(Z)-Ethyl 2-bromo-3-(4-bromo-5-phenylthiophen-2-yl)acrylate (entry 3 in Table 3)

(Z)-Ethyl 2-bromo-3-(4-bromo-5-phenylthiophen-2-yl)acrylate (entry 3 in Table 3):

(E)-Ethyl 3-(4-bromothiophen-2-yl)acrylate (entry 4A in Table 3)

(E)-Ethyl 3-(4-bromothiophen-2-yl)acrylate (entry 4A in Table 3):

(Z)-Ethyl 2-bromo-3-(4-bromothiophen-2-yl)acrylate (entry 4B in Table 3)

(Z)-Ethyl 2-bromo-3-(4-bromothiophen-2-yl)acrylate (entry 4B in Table 3)

(E)-Ethyl 3-(4-phenylthiophen-2-yl)acrylate (entry 5A in Table 3)

(E)-Ethyl 3-(4-phenylthiophen-2-yl)acrylate (entry 5A in Table 3)

(Z)-Ethyl 2-Bromo-3-(4-phenyl-thiophen-2-yl)-acrylate (entry 5B in Table 3)

(Z)-Ethyl 2-Bromo-3-(4-phenyl-thiophen-2-yl)-acrylate (entry 5B in Table 3)