Supporting Information for:

Dimethylzinc-Mediated Alkynylation of Imines

Lorenzo Zani,[†] Silvia Alesi,[‡] Pier Giorgio Cozzi,[‡] and Carsten Bolm^{†,}*

Institute of Organic Chemistry, RWTH Aachen University, Landoltweg 1, D-52056, Aachen, Germany,
and Dipartimento di Chimica “G. Ciamician”, Alma Mater Studiorum, Università di Bologna, Via Selmi
2, I-40126, Bologna, Italy

Table of Contents

1. General information	S1
2. Alkynylation of N-substituted aromatic imines 1	S2
2.1. Typical procedure for the alkynylation of imines 1	S2
2.2. Analytical data of compounds 3aa-3oa	S2
3. Three component synthesis of propargylic imines 3	S8
3.1. Typical procedure for the three component synthesis of propargylic amines 3	S8
3.2. Optimization of the reaction conditions	S8
3.3. Analytical data of compounds 3pa-3Ba	S9
4. Copies of the NMR spectra of compounds 3aa-3na and 3pa-3Ba	S13

1. General Information

All reactions were carried out under an inert atmosphere of argon or nitrogen, using standard Schlenk techniques. Toluene was purified by distillation over sodium-benzophenone ketyl radical, and CH₂Cl₂ was distilled from calcium hydride prior to use. Ethyl acetate, diethyl ether, petroleum ether (PE) and pentane for flash column chromatography were distilled before use. ¹H-NMR spectra were recorded either on a 300 MHz or 400 MHz spectrometer using TMS as internal standard and CDCl₃ as solvent,

unless otherwise stated. ^{13}C -NMR spectra were recorded either on a 75 MHz or 100 MHz spectrometer using CDCl_3 as solvent, unless otherwise stated. Coupling constants (J) are given in Hertz (Hz). IR absorptions are given in wavenumbers (cm^{-1}). Mass peaks are identified by the corresponding m/z values. Melting points are reported as intervals and are uncorrected.

2. Alkynylation of N-substituted aromatic imines **1**

2.1. Typical procedure for the alkynylation of imines **1**.

In an oven-dried Schlenk flask under an inert atmosphere of argon alkyne **2a-d** (0.75–1.25 mmol, 1.5–2.5 equiv) was dissolved in anhydrous toluene (4.5 mL). A 2.0 M solution of dimethylzinc in toluene (0.38–0.63 mL, 0.75–1.25 mmol, 1.5–2.5 equiv) was then carefully added, and the resulting mixture was stirred at room temperature for 30 min. The appropriate imine **1** (0.5 mmol) was then added in one portion, and the temperature was increased to the desired value (50–70 °C). The resulting solution was stirred for 24 h, after which a white precipitate appeared in some cases. The reaction was quenched with water (10 mL). The aqueous phase was extracted with CH_2Cl_2 (3x15 mL) and the organic phase was washed with brine (25 mL) and dried over MgSO_4 . Evaporation of the solvent under reduced pressure furnished the crude product typically as a solid, which was purified by flash column chromatography and/or by recrystallization.

2.2. Analytical data of compounds **3aa-3md**.

N-(*p*-toluensulfonyl)-3-amino 1,3-diphenylprop-1-yne (3aa). Prepared starting from imine **1a** (0.130 g, 0.5 mmol) and phenylacetylene (**2a**, 0.077 g, 0.75 mmol). Purification by flash column chromatography (ethyl acetate/pentane 1:4) afforded pure **3aa**. Yield 0.145 g (0.4 mmol, 80%); white solid; mp 188–189 °C. ^1H -NMR (CDCl_3 , 400 MHz) δ 2.24 (s, 3H), 4.91 (d, J = 9.2 Hz, 1H), 5.48 (d, J = 9.2 Hz, 1H), 7.01–7.08 (m, 2H), 7.11–7.34 (m, 8H), 7.44–7.52 (m, 2H), 7.70–7.79 (m, 2H). ^{13}C -NMR (CDCl_3 , 100 MHz) δ 21.4, 49.8, 85.4, 86.7, 121.9, 127.3, 127.5, 128.1, 128.4, 128.6, 128.7, 129.5, 129.7, 131.5, 137.4, 143.5. IR (KBr): 3265, 2223, 1595, 1431, 1327, 1290, 1154, 1047 cm^{-1} . MS (EI, 70 eV): m/z = 360 $[\text{M}-1]^+$, 222, 206 $[\text{M}-\text{Ts}]^+$, 191, 105, 91, 77. Anal. Calcd. for $\text{C}_{22}\text{H}_{19}\text{NO}_2\text{S}$: C, 73.31; H, 5.03; N, 3.88. Found: C, 73.09; H, 5.17; N, 3.75.

N-(methansulfonyl)-3-amino 1,3-diphenylprop-1-yne (3ba). Prepared starting from imine **1b** (0.092 g, 0.5 mmol) and phenylacetylene (**2a**, 0.077 g, 0.75 mmol). Purification by flash column

chromatography (ethyl acetate/pentane 1:4) afforded pure **3ba**. Yield 0.112 g (0.39 mmol, 78%); white solid; mp 100-101 °C. ¹H-NMR (CDCl₃, 400 MHz) δ 3.00 (s, 3H), 4.92 (d, *J* = 8.2 Hz, 1H), 5.56 (d, *J* = 8.2 Hz, 1H), 7.23-7.36 (m, 6H), 7.37-7.42 (m, 2H), 7.52-7.57 (m, 2H). ¹³C-NMR (CDCl₃, 100 MHz) δ 42.0, 49.9, 86.1, 87.1, 121.8, 127.4, 128.5, 128.8, 128.9, 129.0, 131.6, 137.2. IR (KBr): 3198, 2199, 1596, 1490, 1445, 1320, 1146, 1054 cm⁻¹. MS (EI, 70 eV): *m/z* = 285 [M]⁺, 284 [M-1]⁺, 222, 206 [M-Ms]⁺, 191, 178, 105, 77. Anal. Calcd. for C₁₆H₁₅NO₂S: C, 67.34; H, 5.30; N, 4.91. Found: C, 67.02; H, 5.10; N, 4.83.

N-(2,4,6-trimethylbenzensulfonyl)-3-amino 1,3-diphenylprop-1-yne (3ca). Prepared starting from imine **1c** (0.144 g, 0.5 mmol) and phenylacetylene (**2a**, 0.077 g, 0.75 mmol). Purification by flash column chromatography (ethyl acetate/pentane 1:4) afforded pure **3ca**. Yield 0.160 g (0.41 mmol, 82%); yellow oil that solidified upon standing. ¹H-NMR (CDCl₃, 400 MHz) δ 2.11 (s, 3H), 2.60 (s, 6H), 4.94 (d, *J* = 8.5 Hz, 1H), 5.43 (d, *J* = 8.8 Hz, 1H), 6.80 (s, 2H), 7.02-7.07 (m, 2H), 7.14-7.31 (m, 8H), 7.45-7.51 (m, 2H). ¹³C-NMR (CDCl₃, 100 MHz) δ 21.0, 23.2, 49.5, 84.9, 86.4, 122.0, 127.3, 128.1, 128.49, 128.53, 128.7, 131.6, 131.9, 134.4, 137.3, 139.0, 142.2.

N-(*p*-nitrobenzensulfonyl)-3-amino 1,3-diphenylprop-1-yne (3da). Prepared starting from imine **1d** (0.145 g, 0.5 mmol) and phenylacetylene (**2a**, 0.077 g, 0.75 mmol). Purification by flash column chromatography (ethyl acetate/pentane 1:4) afforded pure **3da**. Yield 0.175 g (0.45 mmol, 91%); white solid; mp 167-168 °C. ¹H-NMR (CDCl₃, 400 MHz) δ 5.20 (d, *J* = 8.7 Hz, 1H), 5.64 (d, *J* = 8.7 Hz, 1H), 7.04-7.13 (m, 2H); 7.19-7.41 (m, 6H), 7.49-7.57 (m, 2H), 8.02-8.09 (m, 2H), 8.16-8.23 (m, 2H). ¹³C-NMR (CDCl₃, 100 MHz) δ 50.2, 84.8, 87.6, 121.3, 124.1, 127.4, 128.4, 128.8, 128.9, 128.9, 129.2, 131.3, 136.5, 146.2, 149.9. IR (KBr): 3269, 1693, 1605, 1528, 1343, 1310, 1162, 1040 cm⁻¹. MS (EI, 70 eV): *m/z* = 391 [M-1]⁺, 222, 206 [M-Ts]⁺, 191, 128, 105, 77. Anal. Calcd. for C₂₁H₁₆N₂O₄S: C, 64.27; H, 4.11; N, 7.14. Found: C, 64.18; H, 4.11; N, 6.95.

N-(diphenylphosphoryl)-3-amino 1,3-diphenylprop-1-yne (3ea). Prepared starting from imine **1e** (0.122 g, 0.4 mmol) and phenylacetylene (**2a**, 0.061 g, 0.6 mmol). Purification by flash column chromatography (ethyl acetate/pentane 1:1) afforded pure **3ea**. Yield 0.113 g (0.27 mmol, 69%); white solid; mp 160-161 °C. ¹H-NMR (CDCl₃, 400 MHz) δ 3.56 (t, *J* = 9.2 Hz, 1H), 5.40 (t, *J* = 9.9 Hz, 1H), 7.27-7.59 (m, 14H); 7.66-7.74 (m, 2H), 7.81-7.91 (m, 2H), 8.05-8.13 (m, 2H). ¹³C-NMR (CDCl₃, 100 MHz) δ 47.2, 85.5, 88.8, 122.6, 127.2, 127.9, 128.1, 128.3, 128.5, 128.6, 131.6, 131.7, 131.8, 131.9, 132.6, 132.7, 133.1, 140.2, 140.2. IR (KBr): 3162, 1961, 1901, 1435, 1188, 1114, 1056 cm⁻¹. MS (EI, 70

eV): m/z = 407 $[M]^+$, 330 $[M-C_6H_5]^+$, 206 $[M-P(O)Ph_2]^+$, 191, 155, 77. Anal. Calcd. for $C_{27}H_{22}NOP$: C, 79.59; H, 5.44; N, 3.44. Found: C, 79.75; H, 5.18; N, 3.35.

N-(2-methoxyphenyl)-3-amino 1,3-diphenylprop-1-yne (3ha).

Prepared starting from imine **1h** (0.106 g, 0.5 mmol) and phenylacetylene (**2a**, 0.128 g, 1.25 mmol). Purification by flash column chromatography (diethyl ether / PE 1:20) afforded pure **3ha**. Yield 0.119 g (0.38 mmol, 76%); yellow oil. 1H -NMR ($CDCl_3$, 400 MHz) δ 3.83 (s, 3H), 4.78 (s, 1H), 5.50 (s, 1H), 6.71-6.91 (m, 4H), 7.23-7.44 (m, 8H), 7.64-7.70 (m, 2H). ^{13}C -NMR ($CDCl_3$, 100 MHz) δ 50.4, 55.5, 84.9, 88.7, 109.6, 111.7, 117.7, 121.1, 122.9, 127.4, 128.0, 128.16, 128.20, 128.7, 131.8, 136.4, 139.9, 147.1. IR (neat): 3418, 3062, 2934, 2834, 1600, 1510, 1243, 1223 cm^{-1} . MS (EI, 70 eV): m/z = 313 $[M]^+$, 282 $[M-CH_3O]^+$, 236, 225, 191 $[M-C_7H_8NO]^+$. Anal. Calcd. for $C_{22}H_{19}NO$: C, 84.31; H, 6.11; N, 4.47. Found: C, 84.43; H, 6.22; N, 4.20.

N-(p-toluenesulfonyl)-1-amino-1-phenylnon-2-yne (3ab) Prepared starting from imine **1a** (0.130 g, 0.5 mmol) and 1-octyne (**2b**, 0.083 g, 0.75 mmol). Purification by flash column chromatography (ethyl acetate / PE 1:3) afforded pure **3ab**. Yield 0.125 g (0.35 mmol, 69%); white solid; mp 82-83 °C. 1H -NMR ($CDCl_3$, 400 MHz) δ 0.81 (t, J = 7.0 Hz, 3H), 1.11-1.28 (m, 8H), 1.89 (dt, J = 6.9 Hz, J = 2.0 Hz, 2H), 2.35 (s, 3H), 4.73 (dd, J = 8.8 Hz, J = 2.0 Hz, 1H), 5.23 (d, J = 8.8 Hz, 1H), 7.17-7.27 (m, 5H); 7.37-7.43 (m, 2H), 7.67-7.73 (m, 2H). ^{13}C -NMR ($CDCl_3$, 100 MHz) δ 14.1, 18.6, 21.6, 22.6, 28.3, 28.6, 31.3, 49.5, 76.6, 87.6, 127.2, 127.4, 128.1, 128.4, 129.3, 137.5, 138.0, 143.1. IR (KBr): 3288, 2928, 2857, 2227, 1597, 1330, 1157, 1031 cm^{-1} . MS (EI, 70 eV): m/z = 299, 214 $[M-Ts]^+$, 155, 143, 91, 77. Anal. Calcd. for $C_{22}H_{27}NO_2S$: C, 71.51; H, 7.36; N, 3.79. Found: C, 71.65; H, 7.70; N, 3.66.

N-(p-toluenesulfonyl)-3-amino-3-phenyl-1-trimethylsilylprop-1-yne (3ac). Prepared starting from imine **1a** (0.130 g, 0.5 mmol) and trimethylsilylethyne (**2c**, 0.075 g, 0.75 mmol). Purification by flash column chromatography (ethyl acetate / PE 1:5) afforded pure **3ac**. Yield 0.091 g (0.26 mmol, 51%); white solid; mp 139-140 °C. 1H -NMR ($CDCl_3$, 400 MHz) δ 0.04 (s, 9H), 2.43 (s, 3H), 4.81 (d, J = 8.5 Hz, 1H), 5.34 (d, J = 9.2 Hz, 1H), 7.25-7.36 (m, 5H), 7.46-7.52 (m, 2H), 7.76-7.82 (m, 2H). ^{13}C -NMR ($CDCl_3$, 100 MHz) δ -0.3, 21.6, 49.7, 91.6, 101.4, 127.2, 127.4, 128.3, 128.5, 129.5, 137.2, 137.3, 143.3. IR (KBr): 3256, 2175, 1952, 1911, 1425, 1328, 1249, 1162, 1056 cm^{-1} . MS (EI, 70 eV): m/z = 356 $[M-1]^+$, 260, 234, 218, 202 $[M-Ts]^+$, 159, 91. Anal. Calcd. for $C_{19}H_{23}NO_2SSi$: C, 63.83; H, 6.48; N, 3.92. Found: C, 63.75; H, 6.35; N, 3.78.

N-(*p*-toluensulfonyl)-3-amino-3-phenyl-1-(4-trifluoromethyl)phenylprop-1-yne (3ad). Prepared starting from imine **1a** (0.130 g, 0.5 mmol) and 4-(trifluoromethyl)phenylacetylene (**2d**, 0.128 g, 0.75 mmol). Purification by flash column chromatography (ethyl acetate / PE 1:4) afforded pure **3ad**. Yield 0.183 g (0.43 mmol, 86%); white solid; mp 140-141 °C. ¹H-NMR (CDCl₃, 300 MHz) δ 2.24 (s, 3H), 4.96 (d, *J* = 8.9 Hz, 1H), 5.50 (d, *J* = 9.1 Hz, 1H), 7.12-7.34 (m, 7H), 7.40-7.49 (m, 4H), 7.70-7.77 (m, 2H). ¹³C-NMR (CDCl₃, 75 MHz) δ 21.4, 49.7, 85.2, 88.1, 125.0, 125.1, 125.8, 127.3, 127.5, 128.6, 128.7, 128.8, 129.6, 131.8, 136.9, 137.4, 143.6. IR (KBr): 3278, 1924, 1807, 1736, 1606, 1328, 1163 cm⁻¹. MS (EI, 70 eV): *m/z* = 410 [M-F]⁺, 274 [M-Ts]⁺, 259, 173, 91, 77. Anal. Calcd. for C₂₃H₁₈NO₂SF₃: C, 64.33; H, 4.22; N, 3.26. Found: C, 63.99; H, 4.69; N, 3.25. HRMS (EI): *m/z* calcd. For C₂₃H₁₈NO₂SF₃-SO₂H: 364.1313. Found: 364.1313.

N-(*p*-nitrobenzensulfonyl)-1-amino-1-phenylnon-2-yne (3db). Prepared starting from imine **1d** (0.145 g, 0.5 mmol) and 1-octyne (**2b**, 0.083 g, 0.75 mmol). Purification by flash column chromatography (ethyl acetate/PE 1:4), followed by recrystallization from MTBE afforded pure **3db**. Yield 0.159 g (0.39 mmol, 79%); yellow solid; mp 103-104 °C. ¹H-NMR (CDCl₃, 300 MHz) δ 0.87 (t, *J* = 6.9 Hz, 3H), 1.14-1.36 (m, 8H), 1.96 (dt, *J* = 6.9 Hz, *J* = 2.1 Hz, 2H), 5.04 (d, *J* = 7.8 Hz, 1H), 5.39 (d, *J* = 8.2 Hz, 1H), 7.27-7.36 (m, 3H), 7.39-7.48 (m, 2H), 7.97-8.07 (m, 2H), 8.25-8.34 (m, 2H). ¹³C-NMR (CDCl₃, 75 MHz) δ 14.0, 18.5, 22.5, 28.2, 28.3, 28.5, 31.2, 49.9, 76.1, 88.3, 123.9, 127.3, 128.6; 128.7, 128.7, 137.3, 146.4, 149.9. IR (KBr): 3268, 2927, 2230, 1602, 1525, 1345, 1309, 1166, 1044 cm⁻¹. MS (EI, 70 eV): *m/z* = 330, 291 [M-C₈H₁₅]⁺, 214 [M-Ns]⁺, 143, 91, 77. Anal. Calcd. for C₂₁H₂₄N₂SO₄: C, 62.98; H, 6.04; N, 6.99. Found: C, 63.38; H, 6.06; N, 6.94.

N-(*p*-toluensulfonyl)-3-amino-3-(4-methylphenyl)-1-phenylprop-1-yne (3ia). Prepared starting from imine **1i** (0.109 g, 0.4 mmol) and phenylacetylene (**2a**, 0.061 g, 0.75 mmol). Purification by flash column chromatography (ethyl acetate / PE 1:4) afforded pure **3ia**. Yield 0.116 g (0.31 mmol, 77%); white solid; mp 190-191 °C. ¹H-NMR (CDCl₃, 400 MHz) δ 2.32 (s, 3H), 2.34 (s, 3H), 4.89 (d, *J* = 9.1 Hz, 1H), 5.52 (d, *J* = 9.1 Hz, 1H), 7.07-7.17 (m, 4H), 7.19-7.33 (m, 5H), 7.38-7.46 (m, 2H), 7.77-7.85 (m, 2H). ¹³C-NMR (CDCl₃, 100 MHz) δ 21.2, 21.5, 49.6, 85.7, 86.5, 122.0, 127.1, 127.4, 128.0, 128.4, 129.3, 129.4, 131.4, 134.4, 137.3, 138.2, 143.4. IR (KBr): 3269, 2221, 1430, 1329, 1155, 1047 cm⁻¹. MS (EI, 70 eV): *m/z* = 310, 236, 220 [M-Ts]⁺, 205, 128, 105, 91, 77. HRMS (EI): *m/z* calcd. For C₂₃H₂₁NO₂S-SO₂H: 310.1596. Found: 310.1595.

N-(*p*-toluensulfonyl)-1-amino-1-(4-methylphenyl)non-2-yne (3ib). Prepared starting from imine **1i** (0.109 g, 0.4 mmol) and 1-octyne (**2**, 0.066 g, 0.6 mmol). Purification by flash column chromatography (ethyl acetate / PE 1:3) afforded pure **3ib**. Yield 0.092 g (0.24 mmol, 60%); white solid; mp 107-108 °C. ¹H-NMR (CDCl₃, 300 MHz) δ 0.88 (t, *J* = 6.9 Hz, 3H), 1.16-1.36 (m, 8H), 1.95 (dt, *J* = 6.9 Hz, *J* = 2.1 Hz, 2H), 2.32 (s, 3H), 2.43 (s, 3H), 4.72 (d, *J* = 8.7 Hz, 1H), 5.26 (d, *J* = 8.7 Hz, 1H), 7.06-7.16 (m, 2H), 7.24-7.37 (m, 4H), 7.73-7.81 (m, 2H). ¹³C-NMR (CDCl₃, 75 MHz) δ 14.0, 18.5, 21.1, 21.5, 22.5, 28.3, 28.5; 31.3, 49.2, 76.8, 87.3, 127.2, 127.5, 129.2, 129.3, 135.2, 137.6, 138.0, 143.2. IR (KBr): 3271, 2930, 2224, 1916, 1334, 1160, 1028 cm⁻¹. MS (EI, 70 eV): *m/z* = 313, 274 [M-C₈H₁₅]⁺, 228 [M-Ts]⁺, 157, 105, 91, 77. Anal. Calcd. for C₂₃H₂₉NO₂S: C, 72.03; H, 7.62; N, 3.65. Found: C, 71.64; H, 7.76; N, 3.73.

N-(*p*-toluensulfonyl)-3-amino-3-(4-methoxyphenyl)-1-phenylprop-1-yne (3ja). Prepared starting from imine **1j** (0.145 g, 0.5 mmol) and phenylacetylene (**2a**, 0.077 g, 0.75 mmol). Purification by flash column chromatography (ethyl acetate / PE 1:3) afforded pure **3ja**. Yield 0.171 g (0.44 mmol, 87%); white solid; mp 186-187 °C. ¹H-NMR (CDCl₃, 400 MHz) δ 2.32 (s, 3H), 3.80 (s, 3H), 4.90 (d, *J* = 8.9 Hz, 1H), 5.51 (d, *J* = 9.0 Hz, 1H), 6.84-6.90 (m, 2H), 7.08-7.15 (m, 2H), 7.20-7.33 (m, 5H), 7.43-7.50 (m, 2H), 7.78-7.84 (m, 2H). ¹³C-NMR (CDCl₃, 100 MHz) δ 21.5, 49.3, 55.4, 85.7, 86.5, 114.0, 121.9, 127.4, 128.0, 128.4, 128.5, 129.4, 129.4, 131.4, 137.3, 143.4, 159.5. IR (KBr): 3263, 2221, 1509, 1329, 1250, 1157, 1034 cm⁻¹. MS (EI, 70 eV): *m/z* = 391 [M]⁺, 326, 236 [M-Ts]⁺, 235, 221, 178, 91, 77. Anal. Calcd. for C₂₃H₂₁NO₃S: C, 70.56; H, 5.41; N, 3.58. Found: C, 70.42; H, 5.36; N, 3.57.

N-(*p*-toluensulfonyl)-3-amino-3-(2-naphthyl)-1-phenylprop-1-yne (3ka). Prepared starting from imine **1k** (0.143 g, 0.5 mmol) and phenylacetylene (**2a**, 0.077 g, 0.75 mmol). Purification by recrystallization from ethyl acetate afforded pure **3ka**. Yield 0.166 g (0.43 mmol, 85%); white solid; mp 206-207 °C. ¹H-NMR (CD₂Cl₂, 400 MHz) δ 2.21 (s, 3H), 5.06 (d, *J* = 9.2 Hz, 1H), 5.60 (d, *J* = 9.2 Hz, 1H), 7.07-7.30 (m, 7H), 7.39-7.47 (m, 2H), 7.51-7.57 (m, 1H), 7.68-7.81 (m, 5H), 7.90 (s, 1H). ¹³C-NMR (CD₂Cl₂, 100 MHz) δ 21.2, 50.0, 85.4, 86.7, 125.0, 126.0, 126.1, 126.4, 126.5, 127.4, 127.5, 128.0, 128.1, 128.6, 128.6, 129.5, 131.5, 133.0, 133.1, 134.8, 137.2, 143.8. IR (KBr): 3259, 2220, 1427, 1329, 1156, 1039 cm⁻¹. MS (EI, 70 eV): *m/z* = 411 [M]⁺, 346, 272, 256 [M-Ts]⁺, 241, 228, 155, 128, 91, 77. Anal. Calcd. for C₂₆H₂₁NO₂S: C, 75.89; H, 5.14; N, 3.40. Found: C, 75.95; H, 5.10; N, 3.36.

N-(*p*-toluensulfonyl)-3-amino-3-(2-bromophenyl)-1-phenylprop-1-yne (3la). Prepared starting from imine **1l** (0.169 g, 0.5 mmol) and phenylacetylene (**2a**, 0.077 g, 0.75 mmol). Purification by flash

column chromatography (ethyl acetate / PE 1:4) afforded pure **3la**. Yield 0.205 g (0.47 mmol, 93%); white solid; mp 169-170 °C. ¹H-NMR (CDCl₃, 300 MHz) δ 2.24 (s, 3H), 5.10 (d, *J* = 8.3 Hz, 1H), 5.75 (d, *J* = 8.3 Hz, 1H), 7.04-7.26 (m, 9H), 7.43-7.52 (m, 2H), 7.67-7.74 (m, 2H). ¹³C-NMR (CD₂Cl₂, 75 MHz) δ 21.4, 50.1, 85.0, 86.5, 121.9, 123.0, 127.5, 127.8, 128.1, 128.6, 129.4, 129.6, 130.0, 131.6, 133.5, 136.7, 137.2, 143.5. IR (KBr): 3268, 1923, 1594, 1329, 1156, 1054 cm⁻¹. MS (EI, 70 eV): *m/z* = 441 [M]⁺, 439, 378, 376, 302, 300, 286 [M-Ts]⁺, 284, 204, 105, 91, 77. Anal. Calcd. for C₂₂H₁₈NO₂BrS: C, 60.01; H, 4.12; N, 3.18. Found: C, 60.05; H, 4.48; N, 3.05.

N-(*p*-toluensulfonyl)-3-amino-3-(2-furyl)-1-phenylprop-1-yne (3ma). Prepared starting from imine **1m** (0.125 g, 0.5 mmol) and phenylacetylene (**2a**, 0.077 g, 0.75 mmol). Purification by flash column chromatography (ethyl acetate / PE 2:7) afforded pure **3ma**. Yield 0.140 g (0.40 mmol, 80%); pale brown solid; mp 151-152 °C. ¹H-NMR (CDCl₃, 400 MHz) δ 2.25 (s, 3H), 4.98 (d, *J* = 9.0 Hz, 1H), 5.53 (d, *J* = 9.0 Hz, 1H), 6.22 (dd, *J* = 3.3 Hz, *J* = 1.9 Hz, 1H), 6.31 (d, *J* = 3.3 Hz, 1H), 7.05-7.29 (m, 8H), 7.69-7.76 (m, 2H). ¹³C-NMR (CD₂Cl₂, 100 MHz) δ 21.5, 44.2, 83.4, 85.3, 108.3, 110.4, 121.6, 127.3, 128.0, 128.6, 129.4, 131.5, 137.2, 143.1, 143.5, 149.4. IR (KBr): 3263, 2225, 1433, 1333, 1157, 1039 cm⁻¹. MS (EI, 70 eV): *m/z* = 303, 250 [M-C₈H₅]⁺, 196 [M-Ts]⁺, 181 [M-C₇H₈NO₂S]⁺, 168, 152, 105, 91, 77. Anal. Calcd. for C₂₀H₁₇NO₃S: C, 68.35; H, 4.88; N, 3.89. Found: C, 68.00; H, 5.20; N, 3.89.

N-(*p*-toluensulfonyl)-3-amino-3-(2-furyl)-1-(4-trifluoromethylphenyl) prop-1-yne (3md). Prepared starting from imine **1m** (0.125 g, 0.5 mmol) and 4-(trifluoromethyl)phenylacetylene (**2d**, 0.128 g, 0.75 mmol). Purification by recrystallization from MTBE afforded pure **3md**. Yield 0.199 g (0.47 mmol, 95%); white solid; mp 117-118 °C. ¹H-NMR (CDCl₃, 400 MHz) δ 2.25 (s, 3H), 5.06 (d, *J* = 8.9 Hz, 1H), 5.55 (d, *J* = 8.9 Hz, 1H), 6.24 (dd, *J* = 3.2 Hz, *J* = 1.9 Hz, 1H), 6.30 (d, *J* = 3.2 Hz, 1H), 7.12-7.31 (m, 5H), 7.41-7.48 (m, 2H), 7.69-7.76 (m, 2H). ¹³C-NMR (CD₂Cl₂, 100 MHz) δ 21.5, 44.0, 83.9, 86.0, 108.5, 110.5, 125.0, 125.0, 125.4, 127.4, 129.5, 131.8, 137.2, 143.2, 143.6, 148.8. IR (KBr): 3270, 2923, 1926, 1616, 1329, 1162, 1132 cm⁻¹. MS (EI, 70 eV): *m/z* = 400, 354, 280, 264 [M-Ts]⁺, 249, 139, 91, 77. Anal. Calcd. for C₂₁H₁₆NO₃SF₃: C, 60.14; H, 3.85; N, 3.34. Found: C, 60.03; H, 4.15; N, 3.32.

N-(*p*-toluensulfonyl)-3-amino-3-cyclohexyl-1-phenylprop-1-yne (3na). Prepared starting from imine **1n** (0.133 g, 0.5 mmol) and phenylacetylene (**2a**, 0.077 g, 0.75 mmol). Purification by recrystallization from EtOAc afforded pure **3na**. Yield 0.155 g (0.42 mmol, 84%); white solid; mp 202-203 °C. ¹H-NMR (CDCl₃, 400 MHz) δ 0.95-1.23 (m, 5H), 1.66-1.87 (m, 6H), 2.25 (s, 3H), 4.04 (dd, *J* = 9.9 Hz, *J* = 6.0 Hz, 1H), 4.59 (d, *J* = 9.9 Hz, 1H), 6.93-7.00 (m, 2H), 7.11-7.22 (m, 5H), 7.70-7.76 (m, 2H). ¹³C-NMR

(CD₂Cl₂, 100 MHz) δ 21.6, 25.86, 25.94, 26.3, 28.6, 29.3, 43.3, 51.6, 85.2, 86.2, 122.2, 127.5, 128.0, 128.3, 129.5, 131.5, 137.4, 143.4. IR (KBr): 3853, 3744, 3452, 3279, 2925, 2360, 2337, 1650, 1331, 1156 cm⁻¹. MS (EI, 70 eV): m/z = 367 [M⁺], 284 [M-C₆H₁₁]⁺, 212 [M-Ts]⁺, 155 [Ts]⁺, 139, 91. Anal. Calcd. for C₂₂H₂₅NO₂S: C, 71.90; H, 6.86; N, 3.81. Found: C, 71.54; H, 7.14; N, 3.75.

N-(*p*-toluensulfonyl)-3-amino-5-methyl-1-phenylhex-1-yne (3oa). Prepared starting from imine **1o** (0.120 g, 0.5 mmol) and phenylacetylene (**2a**, 0.077 g, 0.75 mmol). Purification by flash column chromatography (ethyl acetate / PE 1:7) afforded pure **3na**. Yield 0.025 g (0.08 mmol, 15%); white solid. ¹H-NMR (CDCl₃, 400 MHz) δ 0.90 (t, J = 6.5 Hz, 6H), 1.16-1.29 (m, 1H), 1.50-1.64 (m, 2H), 2.27 (s, 3H), 4.28 (dt, J = 9.6 Hz, J = 7.7 Hz, 1H), 4.51 (d, J = 9.6 Hz, 1H), 6.95-7.00 (m, 2H), 7.13-7.28 (m, 5H), 7.74-7.80 (m, 2H). ¹³C-NMR (CD₂Cl₂, 100 MHz) δ 21.5, 22.3, 22.4, 24.9, 44.9, 45.8, 84.3, 87.4, 127.5, 128.0, 128.3, 129.5, 131.4, 135.8, 137.4, 143.4.

3. Three-component synthesis of propargylic imines **3**

3.1. Typical procedure for the three-component synthesis of propargylic imines **3**

METHOD A. In an oven-dried Schlenk flask under an inert atmosphere of argon were placed the appropriate aldehyde **7** (0.2 mmol) and 2-methoxyaniline (**6**, 0.2 mmol, 1.0 equiv), followed by anhydrous toluene (2.0 mL). After 30 min stirring, a 2.0 M solution of dimethylzinc in toluene (0.35 mL, 0.7 mmol, 3.5 equiv) was added. The reaction mixture was then stirred for another 30 min, before adding phenylacetylene (**2a**, 0.5 mmol, 2.5 equiv). The resulting solution was stirred at room temperature for 48-60 h. The reaction mixture was diluted with diethyl ether (5 mL), and quenched with water (10 mL). The resulting heterogeneous mixture was filtered over Celite[®]. The aqueous phase was separated and washed with diethylether (2x5 mL). The combined organic layers were dried over Na₂SO₄ and evaporated under reduced pressure to give the crude product, which was then purified by flash column chromatography (see Supporting Information for details).

METHOD B (concentrated conditions). In an oven-dried Schlenk flask under an inert atmosphere of argon were placed the appropriate aldehyde **7** (0.4 mmol) and 2-methoxyaniline (**6**, 0.4 mmol, 1.0 equiv). A 2.0 M solution of dimethylzinc in toluene (0.7 mL, 1.4 mmol, 3.5 equiv) was immediately added. The reaction mixture was then stirred for 15 min, before the addition of phenylacetylene (**1a**, 1.0 mmol, 2.5 equiv). The resulting solution was stirred at room temperature for 48-96 h. Subsequently, the protocol followed the procedure reported above for Method A.

3.2. Optimization of the reaction conditions (Method A)

entry	solvent	Me ₂ Zn (equiv)	phenylacetylene (2a , equiv)	yield (%)
1	CH ₂ Cl ₂	2.5	1.5	32
2	CH ₂ Cl ₂	3.5	2.5	72
3	CH ₂ Cl ₂	4.5	3.0	53
4	Toluene	3.5	2.5	76
5	THF	3.5	2.5	0

3.3. Analytical data of compounds 3pa-3Ba.

N-(2-methoxyphenyl)-3-amino-3-(4-chlorophenyl)-1-phenylprop-1-yne (3pa). Prepared starting from aldehyde **7b** (0.030 g, 0.2 mmol) and phenylacetylene (**2a**, 0.051 g, 0.5 mmol) according to Method A. Purification by flash column chromatography (CH₂Cl₂ / cyclohexane 2:3) afforded pure **3pa**. Yield 0.053 g (0.15 mmol, 76%); yellow oil. ¹H-NMR (CDCl₃, 400 MHz) δ 3.83 (s, 3H), 4.81 (brs, 1H), 5.43(s, 1H), 6.78-6.85 (m, 4H), 7.26-7.33 (m, 3H), 7.36 (d, *J* = 8.1 Hz, 2H), 7.42-7.45 (m, 2H), 7.62 (d, *J* = 8.1 Hz, 2H). ¹³C-NMR (CDCl₃, 100 MHz) δ 49.8, 55.4, 85.1, 88.1, 109.5, 111.6, 117.9, 121.0, 122.6, 128.2, 128.4, 128.6, 128.8, 131.7, 133.7, 136.1, 138.5, 147.1. IR (neat): 3409, 3058, 2945, 2939, 1606, 1507, 1480, 1023 cm⁻¹. Anal. Calcd. for C₂₂H₁₈ClNO: C, 75.97; H, 5.22; N, 4.03. Found: C, 75.80; H, 5.29; N, 4.02.

N-(2-methoxyphenyl)-3-amino-3-(3-bromophenyl)-1-phenylprop-1-yne (3qa). Prepared starting from aldehyde **7c** (0.037 g, 0.2 mmol) and phenylacetylene (**2a**, 0.051 g, 0.5 mmol) according to Method A. Purification by flash column chromatography (diethyl ether / cyclohexane 1:9) afforded pure

3qa. Yield 0.057 g (0.15 mmol, 73%); yellow oil. $^1\text{H-NMR}$ (CDCl_3 , 300 MHz) δ 3.83 (s, 3H), 4.83 (brs, 1H), 5.43 (s, 1H), 6.68-6.80 (m, 4H), 7.24-7.30 (m, 3H), 7.41-7.50 (m, 3H), 7.58-7.63 (m, 2H), 7.82 (s, 1H). $^{13}\text{C-NMR}$ (CDCl_3 , 75 MHz) δ 49.9, 55.4, 85.3, 87.9, 109.6, 111.7, 118.0, 121.1, 122.6, 122.8, 125.9, 128.2, 128.4, 130.3, 130.3, 131.1, 131.8, 136.1, 142.3, 147.2. IR (neat): 3436, 3177, 2965, 2939, 2893, 1600, 1513, 1454, 1374 cm^{-1} . Anal. Calcd. for $\text{C}_{22}\text{H}_{18}\text{BrNO}$: C, 67.36; H, 4.62; N, 3.57. Found: C, 67.39; H, 4.70; N, 3.53.

N-(2-methoxyphenyl)-3-amino-3-(4-bromophenyl)-1-phenylprop-1-yne (3ra). Prepared starting from aldehyde **7d** (0.037 g, 0.2 mmol) and phenylacetylene (**2a**, 0.051 g, 0.5 mmol) according to Method A. Purification by flash column chromatography (CH_2Cl_2 / cyclohexane 2:3) afforded pure **3ra**. Yield 0.057 g (0.15 mmol, 73%); yellow oil. $^1\text{H-NMR}$ (CDCl_3 , 400 MHz) δ 3.83 (s, 3H), 4.90 (brs, 1H), 5.43 (s, 1H), 6.78-6.85 (m, 4H), 7.26-7.33 (m, 3H), 7.42-7.45 (m, 2H), 7.57-7.63 (m, 4H). $^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz) δ 49.8, 55.4, 85.1, 88.1, 109.8, 111.9, 118.0, 121.3, 122.1, 122.9, 128.5, 128.7, 129.3, 132.0, 132.1, 136.3, 139.3, 147.4. IR (neat): 3430, 3170, 2961, 2928, 2893, 1604, 1510, 1450, 1371 cm^{-1} . Anal. Calcd. for $\text{C}_{22}\text{H}_{18}\text{BrNO}$: C, 67.36; H, 4.62; N, 3.57. Found: C, 67.38; H, 4.65; N, 3.56.

N-(2-methoxyphenyl)-3-amino-3-(2-naphthyl)-1-phenylprop-1-yne (3sa). Prepared starting from aldehyde **7e** (0.031 g, 0.2 mmol) and phenylacetylene (**2a**, 0.051 g, 0.5 mmol) according to Method A. Purification by flash column chromatography (diethyl ether / cyclohexane 1:9) afforded pure **3sa**. Yield 0.067 g (0.18 mmol, 92%); yellow oil. $^1\text{H-NMR}$ (CDCl_3 , 400 MHz) δ 3.84 (s, 3H); 4.92 (brs, 1H), 5.62 (s, 1H), 6.78-6.85 (m, 4H), 7.26-7.33 (m, 3H), 7.42-7.45 (m, 4H), 7.78 (d, $J = 7.4$ Hz, 2H), 7.80-7.90 (m, 2H), 8.15 (s, 1H). $^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz) δ 50.6, 55.4, 85.1, 88.7, 109.6, 111.7, 117.8, 121.1, 122.9, 125.4, 126.1, 126.2, 126.3, 127.7, 128.1, 128.2, 128.2, 128.6, 131.8, 133.1, 133.4, 136.5, 137.3, 147.2. IR (neat): 3423, 3058, 2919, 2833, 1594, 1500, 1241, 1208 cm^{-1} . Anal. Calcd. for $\text{C}_{26}\text{H}_{21}\text{NO}$: C, 85.92; H, 5.82; N, 3.85. Found: C, 86.03; H, 5.85; N, 3.81.

N-(2-methoxyphenyl)-3-amino-3-biphenyl-1-phenylprop-1-yne (3ta). Prepared starting from aldehyde **7f** (0.037 g, 0.2 mmol) and phenylacetylene (**2a**, 0.051 g, 0.5 mmol) according to Method A. Purification by flash column chromatography (CH_2Cl_2 / cyclohexane 3:7) afforded pure **3ta**. Yield 0.23 g (0.06 mmol, 30%); yellow oil. $^1\text{H-NMR}$ (CDCl_3 , 400 MHz) δ 3.84 (s, 3H); 4.81 (brs, 1H); 5.58 (s, 1H); 6.73-6.77 (m, 1H); 6.82 (d, $J = 7.2$ Hz, 1H), 6.90-6.92 (m, 2H), 7.28-7.31 (m, 3H), 7.36-7.48 (m, 5H), 7.59-7.62 (m, 2H), 7.64 (t, $J = 8.7$ Hz, 2H), 7.45 (d, $J = 8.7$ Hz, 2H). $^{13}\text{C-NMR}$ (CDCl_3 , 100 MHz)

δ 50.0, 55.4, 84.9, 88.6, 105.5, 109.5, 111.6, 117.6, 121.1, 122.9, 127.1, 127.5, 127.8, 128.1, 128.18, 128.24, 128.8, 131.8, 136.4, 138.9, 140.9, 147.2. IR (neat): 3423, 3051, 2925, 2826, 1593, 1500, 1235, 1020 cm^{-1} . Anal. Calcd. for $\text{C}_{28}\text{H}_{23}\text{NO}$: C, 86.34; H, 5.95; N, 3.60. Found: C, 86.40; H, 5.98; N, 3.56.

N-(2-methoxyphenyl)-3-amino-3-(3-nitrophenyl)-1-phenylprop-1-yne (3ua). Prepared starting from aldehyde **7g** (0.030 g, 0.2 mmol) and phenylacetylene (**2a**, 0.051 g, 0.5 mmol) according to Method A. Purification by flash column chromatography (CH_2Cl_2 / cyclohexane 3:7) afforded pure **3ua**. Yield 0.047 g (0.13 mmol, 66%); yellow oil. ^1H -NMR (CDCl_3 , 300 MHz) δ 3.85 (s, 3H); 5.10 (brs, 1H), 5.60 (s, 1H), 6.67-6.86 (m, 4H), 7.26-7.33 (m, 3H), 7.42-7.45 (m, 2H), 7.57 (t, $J = 8.1\text{ Hz}$, 1H), 8.01 (d, $J = 7.8\text{ Hz}$, 1H), 8.20 (dd, $J = 8.1\text{ Hz}$, $J = 0.9\text{ Hz}$, 1H), 8.48 (s, 1H). ^{13}C -NMR (CDCl_3 , 75 MHz) δ 49.9, 55.4, 85.9, 87.2, 109.8, 111.7, 118.5, 121.0, 122.3, 123.0, 128.3, 128.6, 129.6, 131.8, 133.3, 135.7, 142.4, 147.3, 146.6. IR (neat): 3423, 3051, 2925, 2826, 1593, 1500, 1235, 1020 cm^{-1} . Anal. Calcd. for $\text{C}_{22}\text{H}_{18}\text{N}_2\text{O}_3$: C, 73.73; H, 5.06; N, 7.82. Found: C, 73.80; H, 5.13; N, 7.75.

N-(2-methoxyphenyl)-3-amino-3-(4-trifluoromethylphenyl)-1-phenylprop-1-yne (3va). Prepared starting from aldehyde **7h** (0.035 g, 0.2 mmol) and phenylacetylene (**2a**, 0.051 g, 0.5 mmol) according to Method A. Purification by flash column chromatography (diethyl ether / cyclohexane 1:9) afforded pure **3va**. Yield 0.060 g (0.16 mmol, 79%); yellow oil. ^1H -NMR (CDCl_3 , 300 MHz) δ 3.84 (s, 3H), 5.58 (s, 1H), 6.68-6.85 (m, 4H); 7.27-7.35 (m, 3H), 7.38-7.44 (m, 2H), 7.65 (d, $J = 8.4\text{ Hz}$, 2H), 7.79 (d, $J = 8.4\text{ Hz}$, 2H). ^{13}C -NMR (CDCl_3 , 75 MHz) δ 50.0, 55.4, 85.4, 87.7, 109.7, 111.7, 118.2, 120.1, 121.1, 122.5, 125.67, 125.72, 127.6, 128.3, 128.5, 131.8, 136.0, 144.0, 147.2. IR (neat): 3390, 3270, 2919, 2853, 1475, 1380, cm^{-1} . Anal. Calcd. for $\text{C}_{23}\text{H}_{18}\text{F}_3\text{NO}$: C, 72.43; H, 4.76; N, 3.67. Found: C, 72.48; H, 4.79; N, 3.62.

4-[1-(2-Methoxy-3-phenylamino)-3-phenylpropyn-2-yl]-benzoic acid methyl ester (3wa). Prepared starting from aldehyde **7i** (0.033 g, 0.2 mmol) and phenylacetylene (**2a**, 0.050 g, 0.5 mmol) according to Method A. Purification by flash column chromatography (diethyl ether / cyclohexane 1:4) afforded pure **3wa**. Yield 0.048 g (0.13 mmol, 65%); yellow oil. ^1H -NMR (CDCl_3 , 300 MHz) δ 3.81 (s, 3H), 3.88 (s, 3H), 5.50 (s, 1H), 6.69-6.84 (m, 4H), 7.22-7.28 (m, 3H), 7.36-7.40 (m, 2H), 7.70 (d, $J = 8.4\text{ Hz}$, 2H), 8.04 (d, $J = 8.4\text{ Hz}$, 2H). ^{13}C -NMR (CDCl_3 , 75 MHz) δ 50.1, 52.1, 55.4, 85.3, 87.9, 109.6, 111.7, 118.0, 121.0, 122.5, 127.2, 128.2, 128.4, 129.8, 130.0, 131.7, 136.0, 145.0, 147.2, 166.7. IR (neat): 3390, 3270,

2919, 2853, 1475, 1380 cm^{-1} . Anal. Calcd. for $\text{C}_{24}\text{H}_{21}\text{NO}_3$: C, 77.71; H, 5.70; N, 3.77. Found: C, 77.76; H, 5.78; N, 3.71.

N-(2-methoxyphenyl)-3-amino-3-(pentafluorophenyl)-1-phenylprop-1-yne (3xa). Prepared starting from aldehyde **7j** (0.055 g, 0.2 mmol) and phenylacetylene (**2a**, 0.051 g, 0.5 mmol) according to Method A. Purification by flash column chromatography (CH_2Cl_2 / cyclohexane 35:65) afforded pure **3xa**. Yield 0.051 g (0.13 mmol, 63%); yellow oil. ^1H -NMR (CDCl_3 , 400 MHz) δ 3.84 (s, 3H), 5.95 (s, 1H), 6.72-6.82 (m, 3H), 6.87 (dt, $J = 7.6$ Hz, $J = 1.6$ Hz, 1H), 7.27-7.34 (m, 3H), 7.43-7.46 (m, 2H). ^{13}C -NMR (CDCl_3 , 100 MHz) δ 40.4, 55.6, 84.4, 85.1, 110.2, 111.2, 114.5 (t, $J = 56$ Hz), 119.0, 121.2, 122.0, 128.3, 128.8, 131.9, 134.8, 136.5 (dt, $J = 73$ Hz, $J = 24$ Hz), 139.9 (dt, $J = 66$ Hz, $J = 20$ Hz), 139.0 (m), 143.7 (m), 146.1 (m), 147.7. IR (neat): 3390, 3270, 2919, 2853, 1475, 1380 cm^{-1} . Anal. Calcd. for $\text{C}_{22}\text{H}_{14}\text{F}_5\text{NO}$: C, 65.51; H, 3.50; N, 3.47. Found: C, 65.48; H, 3.41; N, 3.49.

N-(2-methoxyphenyl)-3-amino-1-phenyl-3-(3-pyridyl)-prop-1-yne (3ya). Prepared starting from aldehyde **7k** (0.021 g, 0.2 mmol) and phenylacetylene (**2a**, 0.051 g, 0.5 mmol) according to Method A. Purification by flash column chromatography (ethyl acetate / cyclohexane 55:45) afforded pure **3ya**. Yield 0.041 g (0.13 mmol, 65%); yellow oil. ^1H -NMR (CDCl_3 , 300 MHz) δ 3.84 (s, 3H), 5.60 (s, 1H), 6.78-6.93 (m, 4H), 7.30-7.34 (m, 3H), 7.42-7.46 (m, 3H), 8.09 (d, $J = 7.5$ Hz, 1H), 8.64 (d, $J = 5.1$ Hz, 1H), 8.97 (s, 1H). ^{13}C -NMR (CDCl_3 , 75 MHz) δ 48.2, 55.4, 85.6, 87.3, 109.7, 111.8, 118.3, 121.0, 122.4, 123.6, 128.2, 128.5, 131.7, 135.0, 135.7, 135.9, 147.3, 148.9, 149.1. IR (neat): 3390, 3270, 2919, 2853, 1475, 1380, cm^{-1} . Anal. Calcd. for $\text{C}_{21}\text{H}_{18}\text{N}_2\text{O}$: C, 80.23; H, 5.77; N, 8.91. Found: C, 80.20; H, 5.79; N, 8.93.

N-(2-methoxyphenyl)-3-amino-3-cyclohexyl-1-phenylprop-1-yne (3za). Prepared starting from aldehyde **7l** (0.044 g, 0.4 mmol) and phenylacetylene (**2a**, 0.102 g, 1.0 mmol) according to Method B. Purification by flash column chromatography (CH_2Cl_2 / cyclohexane 15:85) afforded pure **3za**. Yield 0.109 g (0.34 mmol, 85%); yellow oil. ^1H -NMR (CDCl_3 , 300 MHz) δ 1.35-1.47 (m, 6H), 1.74-2.08 (m, 5H), 3.90 (s, 3H), 4.22 (d, $J = 5.7$ Hz, 1H), 4.40 (brs, 1H), 6.62-6.97 (m, 4H), 7.29-7.32 (m, 3H), 7.39-7.44 (m, 2H). ^{13}C -NMR (CDCl_3 , 75 MHz) δ 26.0, 26.1, 26.4, 42.3, 51.3, 55.3, 83.6, 89.1, 109.4, 110.1, 116.9, 121.1, 123.2, 127.8, 128.1, 131.6, 136.8, 147.0. Anal. Calcd. for $\text{C}_{22}\text{H}_{25}\text{NO}$: C, 82.72; H, 7.89; N, 4.38. Found: C, 82.70; H, 7.84; N, 4.40.

N-(2-methoxyphenyl)-3-amino-4,4-dimethyl-1-phenylpent-1-yne (3Aa). Prepared starting from aldehyde **7m** (0.034 g, 0.4 mmol) and phenylacetylene (**2a**, 0.102 g, 1.0 mmol) according to Method B. Purification by flash column chromatography (CH₂Cl₂ / cyclohexane 1:4) afforded pure **3Aa**. Yield 0.079 g (0.27 mmol, 67%); yellow oil. ¹H-NMR (CDCl₃, 300 MHz) δ 1.22 (s, 9H), 3.91 (s, 3H), 4.08 (s, 1H), 4.55 (brs, 1H), 6.72-6.78 (m, 1H), 6.83-6.90 (m, 2H), 6.89-6.95 (m, 1H), 7.29-7.38 (m, 3H), 7.38-7.43 (m, 2H). ¹³C-NMR (CDCl₃, 75 MHz) δ 26.5, 35.7, 55.4, 55.9, 83.3, 89.4, 109.6, 111.0, 116.9, 121.1, 123.3, 127.8, 128.1, 131.6, 137.3, 137.1. Anal. Calcd. for C₂₀H₂₃NO: C, 81.87; H, 7.90; N, 4.77. Found: C, 81.84; H, 7.88; N, 4.80.

N-(2-methoxyphenyl)-3-amino-1-phenyldec-1-yne (3Ba). Prepared starting from aldehyde **7n** (0.052 g, 0.4 mmol) and phenylacetylene (**2a**, 0.102 g, 1.0 mmol) according to Method B. Purification by flash column chromatography (CH₂Cl₂ / cyclohexane 1:4) afforded pure **3Ba**. Yield 0.060 g (0.18 mmol, 45%); yellow oil. ¹H-NMR (CDCl₃, 300 MHz) δ 0.82-1.00 (m, 3H), 1.20-1.42 (m, 9H), 1.60-1.80 (m, 2H), 1.89-2.02 (m, 1H), 3.90 (s, 3H), 4.34 (t, *J* = 6.3 Hz, 1H), 4.46 (brs, 1H), 6.73-6.98 (m, 4H), 7.29-7.31 (m, 3H), 7.39-7.43 (m, 2H). ¹³C-NMR (CDCl₃, 75 MHz) δ 14.1, 22.6, 26.1, 29.2, 29.2, 31.8, 35.9, 45.9, 55.3, 82.6, 90.3, 109.4, 111.3, 117.2, 121.1, 123.2, 127.9, 128.1, 131.7, 136.6, 147.0. Anal. Calcd. for C₂₃H₂₉NO: C, 82.34; H, 8.71; N, 4.18. Found: C, 82.31; H, 8.69; N, 4.20.

4. Copies of the NMR spectra of compounds 3aa-3na and 3pa-3Ba

S2