Supporting Information

Direct Asymmetric Hydroxyamination Reaction Catalyzed by an Axially Chiral Secondary Amine Catalyst

Taichi Kano, Mitsuhiro Ueda, Jun Takai, and Keiji Maruoka*

*Department of Chemistry, Graduate School of Science, Kyoto University
Sakyo, Kyoto 606-8502, Japan*

General Information. Infrared (IR) spectra were recorded on a Shimadzu IR Prestige-21 spectrometer. ¹H NMR spectra were measured on a JEOL JNM-FX400 (400 MHz) spectrometer. Data were reported as follows: chemical shifts in ppm from tetramethylsilane (in the case of CDCl₃) or CD₂HOD (δ = 3.30, in the case of CD₃OD) as an internal standard, integration, multiplicity (s = singlet, d = doublet, t = triplet, q = quartet, dd = double-doublet, m = multiplet, br = broad, app = apparent), coupling constants (Hz), and assignment. ¹³C NMR spectra were measured on a JEOL JNM-FX400 (100 MHz) spectrometer with complete proton decoupling. Chemical shifts were reported in ppm from the residual solvent as an internal standard. High performance liquid chromatography (HPLC) was performed on Shimadzu 10A instruments at 254 nm using 4.6 nm x 25 cm Daicel Chiralpak and Chiralcel. High-resolution mass spectra (HRMS) were performed on Applied Biosystems Mariner 8295 API-TOF workstation and BRUKER microTOF. Optical rotations were measured on a JASCO DIP-1000 digital polarimeter. For thin layer chromatography (TLC) analysis throughout this work, Merck precoated TLC plates (silica gel 60 GF₂₅₄, 0.25 mm) were used. The products were purified by flash column chromatography on silica gel 60 (Merck 1.09386.9025, 230-400 mesh).

Tetrahydrofuran (THF) was purchased from Kanto Chemical Co. Inc. as “Dehydrated”. The commercially available aldehydes were distilled and stored under argon atmosphere at -17 °C. (S)-2-Pyrrolidinemethanol (*S*)-**2a**, nitrosobenzene, (S)-2-amino-3-methylbutanol, (S)-2-amino-3-phenylpropanol and (S)-2-amino-3-cyclohexylpropanol were purchased and used without further purification. 3-Cyclohexylpropanal,¹ 5-benzyloxypentanal,² 4-methoxynitrosobenzene,³ (S)-3,5-dihydro-4*H*-dinaphth[2,1-*c*:1',2'-*e*]azepine (*S*)-**1a**,⁴ (S)-2,2'-bis(neopentyloxycarbonyl)-1,1'-binaphthyl (*S*)-**7**⁵ and (S)- α,α -diphenyl-2-pyrrolidinemethanol (*S*)-**2b**⁶ were synthesized according to literature procedures and used after column chromatography on silica gel.

Synthesis and Characterization of Chiral Secondary Amine Catalysts 1b-1e

a) $\text{Mg}(\text{TMP})_2$, THF; then Br_2 ; b) DIBAL-H, THF; c) BBr_3 , CH_2Cl_2 ; d) allylamine, CH_3CN ; e) 10 mol% $\text{Pd}(\text{OAc})_2$, dppp, $i\text{-Pr}_2\text{NEt}$, CO, DMSO, MeOH; f) LiAlH_4 , THF; g) MeLi, THF; h) PhLi, Et_2O ; i) $\text{Pd}(\text{OAc})_2$, PPh_3 , NDMBA, CH_2Cl_2 ; j) $\text{BF}_3 \cdot \text{OEt}_2$, Et_3SiH , CH_2Cl_2 .

Diol (S)-9.⁷ To a solution of magnesium bis(2,2,6,6-tetramethylpiperidide) (10.0 mmol) in THF was added (S)-7 (1.0 mmol, 482 mg) in THF (4.0 mL) dropwise at 0 °C. After stirring for 12 h at room temperature, the reaction mixture was treated with Br_2 (410 μL , 8.0 mmol) at -78 °C and then stirred for 3 h at room temperature. The mixture was poured into cooled 1N HCl and extracted with ethyl acetate. The organic layer was washed with saturated Na_2SO_3 , dried over Na_2SO_4 and then concentrated. The residue was purified by flash column chromatography on silica gel (eluting with hexane/ethyl acetate = 10:1) to give a mixture including the desired (S)-8. This mixture (483 mg), which consisted of three components (3,3'-dibromo, 3-bromo and non-substituted forms), was used for the next step without further purification.

To the mixture obtained above in CH_2Cl_2 (20 mL) was added a 0.99 M toluene solution of diisobutyl aluminum hydride (6.0 mmol, 6.1 mL) dropwise at 0 °C. The mixture was kept stirring at room temperature until the starting material was completely consumed (check by TLC). The reaction mixture was carefully poured into cooled 1N HCl and extracted with CH_2Cl_2 . The organic layer was washed with brine, dried over Na_2SO_4 and concentrated. The residue was purified by flash column chromatography on silica gel (eluting with CH_2Cl_2 only) to give (S)-9 (209 mg, 0.44 mmol, 44% yield). $[\alpha]_D^{28}$ -57.7 ($c = 0.92$, CHCl_3); $^1\text{H-NMR}$ (400 MHz, CDCl_3) δ 8.31 (2H, s, Ar-H), 7.85 (2H, d, $J = 8.3$ Hz, Ar-H), 7.50 (2H, app t, $J = 7.3$ Hz, Ar-H), 7.28 (2H, app t, $J = 5.4$ Hz, Ar-H), 6.92 (2H, d,

$J = 8.5$ Hz, Ar-H), 4.76 (2H, d, $J = 9.8$ Hz, -CHH-), 4.07 (2H, d, $J = 12.0$ Hz, -CHH-), 3.47 (2H, br s, -OH); ^{13}C NMR (100 MHz, CDCl_3) δ 136.8, 135.9, 134.6, 133.7, 132.4, 127.4, 127.2, 127.1, 126.6, 123.0, 62.1; IR (neat) 3273, 2949, 2868, 2359, 2342, 1028, 976, 906, 800, 754, 733 cm^{-1} ; HRMS (ESI-TOF) Calcd. for $\text{C}_{22}\text{H}_{16}\text{Br}_2\text{O}_2\text{Na}$: 494.9390 ($[\text{M} + \text{Na}]^+$), Found: 494.9390 ($[\text{M} + \text{Na}]^+$).

Tetrabromide (S)-10. A mixture of (S)-**9** (209 mg, 0.44 mmol) and boron tribromide (84 μL , 0.89 mmol) in CH_2Cl_2 (30 mL) was stirred for 1 h at 0 $^\circ\text{C}$, and water was then carefully added dropwise. After extraction with CH_2Cl_2 , the organic layer was washed with brine, dried over Na_2SO_4 and concentrated. The resulting residue was purified by flash column chromatography on silica gel (eluting with hexane only) to give (S)-**10** (188 mg, 0.31 mmol, 71% yield). $[\alpha]_{\text{D}}^{33}$ -109.8 ($c = 1.0$, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 8.34 (2H, s, Ar-H), 7.85 (2H, d, $J = 8.3$ Hz, Ar-H), 7.52 (2H, t, $J = 7.3$ Hz, Ar-H), 7.31-7.25 (2H, m, Ar-H), 6.99 (2H, d, $J = 8.5$ Hz, Ar-H), 4.40 (2H, d, $J = 10.5$ Hz, -CHHBr), 4.33 (2H, d, $J = 10.2$ Hz, -CHHBr); ^{13}C NMR (100 MHz, CDCl_3) δ 136.8, 134.1, 133.3, 132.7, 131.2, 128.1, 127.3, 127.2, 127.1, 122.6, 32.9; IR (neat) 3057, 1579, 1555, 1493, 1429, 1312, 1213, 1024, 993, 907, 885, 860, 748, 731, 575 cm^{-1} ; HRMS (APCI-TOF) Calcd. for $\text{C}_{22}\text{H}_{14}\text{Br}_4$: 593.7824 ($[\text{M}]^+$), Found: 593.7850 ($[\text{M}]^+$).

Allylamine (S)-11. To a solution of (S)-**10** (188 mg, 0.31 mmol) in acetonitrile (1.6 mL) was added allylamine (71 μL , 0.94 mmol) at room temperature. The mixture was stirred for 10 h, and then poured into water. After extraction with ethyl acetate, the organic layer was washed with brine and dried over Na_2SO_4 . Evaporation of solvents and purification of the residue by flash column chromatography on silica gel (eluting with hexane/ethyl acetate = 3:1) afforded (S)-**11** (136 mg, 0.28 mmol, 90% yield). $[\alpha]_{\text{D}}^{34}$ 314.8 ($c = 1.0$, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 8.27 (2H, s, Ar-H), 7.84 (2H, d, $J = 8.0$ Hz, Ar-H), 7.47 (2H, app d, $J = 3.9$ Hz, Ar-H), 7.26 (4H, app s, Ar-H), 6.04-5.94 (1H, m, -CH=CH₂), 5.27 (1H, app d, $J = 17.4$ Hz, -CH=CHH), 5.18 (1H, app d, $J = 10.3$ Hz, -CH=CHH), 4.39 (2H, d, $J = 13.0$ Hz, -CHHN-), 3.62 (1H, dd, $J = 13.4, 4.4$ Hz, -NCHHCH-), 3.04-3.01 (3H, m, -CHHN-, -NCHHCH-); ^{13}C NMR (100 MHz, CDCl_3) δ 136.8, 135.9, 133.7, 132.8, 132.0, 130.3, 127.4, 127.3, 126.6, 126.4, 122.8, 117.6, 59.0, 53.6; IR (neat) 3067, 2924, 2805, 2359, 1580, 1558, 1497, 1138, 991, 970, 922, 908, 882, 775, 746, 733, 584 cm^{-1} ; HRMS (ESI-TOF) Calcd. for $\text{C}_{25}\text{H}_{20}\text{Br}_2\text{N}$: 493.9938 ($[\text{M} + \text{H}]^+$), Found: 493.9935 ($[\text{M} + \text{H}]^+$).

Methyl ester (S)-12. Allylamine (S)-**11** (75 mg, 0.15 mmol), $\text{Pd}(\text{OAc})_2$ (3.4 mg, 0.015 mmol), bis(diphenylphosphino)propane (dppp) (6.3 mg, 0.015 mmol) and $i\text{Pr}_2\text{NEt}$ (117 μL , 0.67 mmol) in DMSO (5.0 mL) and MeOH (5.0 mL) were charged into autoclave under argon atmosphere. After pressurized with CO (10 atm), the mixture was heated to 90 $^\circ\text{C}$ with stirring for 36 h. After cooling to room temperature, the reaction mixture was poured into water and extracted with ethyl acetate.

The organic layer was washed with brine, dried over Na₂SO₄ and then concentrated. The residue was purified by flash column chromatography on silica gel (eluting with hexane/ethyl acetate = 3:1) to give (*S*)-**12** (41 mg, 0.092 mmol, 61% yield). [α]_D²⁷ 529.9 (*c* = 1.1, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 8.60 (2H, s, Ar-H), 8.01 (2H, d, *J* = 8.3 Hz, Ar-H), 7.50 (2H, app t, *J* = 7.8 Hz, Ar-H), 7.32 (2H, app t, *J* = 8.5 Hz, Ar-H), 7.23 (2H, d, *J* = 8.5 Hz, Ar-H), 6.03-5.93 (1H, m, -CH=CH₂), 5.14-5.10 (2H, m, -CH=CH₂), 4.79 (2H, d, *J* = 13.2 Hz, -CHHN-), 4.00 (6H, s, -COOCH₃), 3.31 (1H, dd, *J* = 14.6, 5.6 Hz, -NCHHCH-), 2.97 (2H, d, *J* = 13.2 Hz, -CHHN-), 2.94 (1H, dd, *J* = 13.0, 7.3 Hz, -NCHHCH-); ¹³C NMR (100 MHz, CDCl₃) δ 168.1, 136.7, 136.2, 132.6, 132.0, 131.8, 131.5, 129.2, 128.3, 127.9, 127.1, 126.2, 117.0, 58.8, 52.5, 49.7; IR (neat) 2949, 1717, 1643, 1622, 1442, 1418, 1331, 1304, 1275, 1242, 1206, 1192, 1169, 1152, 1138, 1057, 1024, 1003, 908, 791, 752, 731 cm⁻¹; HRMS (ESI-TOF) Calcd. for C₂₉H₂₆NO₄: 452.1856 ([M + H]⁺), Found: 452.1856 ([M + H]⁺).

Alcohol (*S*)-13. To a solution of (*S*)-**12** (135 mg, 0.30 mmol) in THF (1.5 mL) was added lithium aluminum hydride (LiAlH₄) (23 mg, 0.60 mmol) at 0 °C. The mixture was stirred for 2 h at 0 °C, and saturated NH₄Cl was then carefully added dropwise. After extraction with ethyl acetate, the organic layer was washed with brine, dried over Na₂SO₄ and concentrated. The resulting residue was purified by flash column chromatography on silica gel (eluting with CH₂Cl₂/MeOH = 20:1~10:1) to give (*S*)-**13** (105 mg, 0.27 mmol, 89% yield). [α]_D²⁹ 326.3 (*c* = 1.0, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.99 (2H, s, Ar-H), 7.90 (2H, d, *J* = 8.3 Hz, Ar-H), 7.45 (2H, app t, *J* = 8.0 Hz, Ar-H), 7.38 (2H, d, *J* = 8.5 Hz, Ar-H), 7.25-7.21 (2H, m, Ar-H), 5.98-5.88 (1H, m, -CH=CH₂), 5.22-5.20 (2H, m, -CH=CH₂), 5.05 (2H, d, *J* = 12.5 Hz, -CHHOH), 4.88 (2H, d, *J* = 12.7 Hz, -CHHOH), 4.04 (2H, d, *J* = 13.0 Hz, -CHHN-), 3.37 (1H, dd, *J* = 13.7, 5.8 Hz, -NCHHCH-), 3.08 (1H, dd, *J* = 13.4, 6.6 Hz, -NCHHCH-), 3.02 (2H, d, *J* = 13.2 Hz, -CHHN-); ¹³C NMR (100 MHz, CDCl₃) δ 136.9, 136.4, 135.1, 132.9, 131.3, 130.9, 128.2, 127.9, 127.3, 125.9 (two peaks overlap), 118.2, 64.6, 58.4, 49.8; IR (neat) 3333, 3059, 2880, 1447, 1096, 1024, 908, 750, 731 cm⁻¹; HRMS (ESI-TOF) Calcd. for C₂₇H₂₆NO₂: 396.1958 ([M + H]⁺), Found: 396.1951 ([M + H]⁺).

Alcohol (*S*)-14. To a solution of (*S*)-**12** (500 mg, 1.1 mmol) in THF (16 mL) was added a 0.98 M ether solution of methyl lithium (5.7 mL, 5.6 mmol) at -78 °C. The mixture was stirred for 2 h at -78 °C, and saturated NH₄Cl was then carefully added dropwise. After extraction with ethyl acetate, the organic layer was washed with brine, dried over Na₂SO₄ and concentrated. The resulting residue was purified by flash column chromatography on silica gel (eluting with CH₂Cl₂/MeOH = 10:1) to give (*S*)-**14** (326 mg, 0.72 mmol, 65% yield). [α]_D²⁹ 93.2 (*c* = 0.97, CHCl₃); ¹H NMR (400 MHz, CD₃OD) δ 8.10 (2H, s, Ar-H), 7.93 (2H, d, *J* = 8.3 Hz, Ar-H), 7.40 (2H, app t, *J* = 7.1 Hz, Ar-H), 7.12 (2H, app t, *J* = 8.5 Hz, Ar-H), 6.92 (2H, d, *J* = 8.6 Hz, Ar-H), 5.97-5.89 (1H, m, -CH=CH₂), 5.18-5.14 (2H, m, -CH=CH₂), 5.04 (2H, d, *J* = 11.4 Hz, -CHHN-), 3.80 (1H, app d, *J* = 13.2 Hz,

-NCH $\overline{H}$ CH-), 3.16 (2H, d, J = 12.4 Hz, -CH $\overline{H}$ N-), 3.01-2.97 (1H, m, -NCH $\overline{H}$ CH-), 1.90 (6H, s, -CH $\overline{3}$), 1.69 (6H, s, -CH $\overline{3}$); ^{13}C NMR (100 MHz, CDCl_3) δ 143.6, 138.1, 132.2, 130.6, 128.4 (two peaks overlap), 127.2 (two peaks overlap), 125.8 (two peaks overlap), 125.0, 117.8, 73.4, 58.9, 50.6, 32.8, 32.1; IR (neat) 2974, 2974, 2926, 2359, 2342, 1780, 1699, 1261, 1161, 1138, 750 cm^{-1} ; HRMS (ESI-TOF) Calcd. for $\text{C}_{31}\text{H}_{34}\text{NO}_2$: 452.2584 ($[\text{M} + \text{H}]^+$), Found: 452.2584 ($[\text{M} + \text{H}]^+$).

Alcohol (S)-15. To a solution of (S)-**12** (300 mg, 0.66 mmol) in Et_2O (6.6 mL) was added a 2.1 M dibutyl ether solution of phenyl lithium (1.9 mL, 4.0 mmol) at -78°C . The mixture was stirred for 2 h at -78°C , and saturated NH_4Cl was then carefully added dropwise. After extraction with ethyl acetate, the organic layer was washed with brine, dried over Na_2SO_4 and concentrated. The resulting residue was recrystallized from CH_2Cl_2 /hexane to give (S)-**15** (375 mg, 0.54 mmol, 81% yield). $[\alpha]_{\text{D}}^{30}$ -88.3 (c = 1.0, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.63 (2H, d, J = 8.3 Hz, Ar- $\overline{H}$), 7.41-7.28 (19H, m, Ar- $\overline{H}$), 7.22-7.16 (9H, m, Ar- $\overline{H}$), 5.82-5.69 (1H, m, -CH=CH $\overline{2}$), 4.86 (1H, app d, J = 10.2 Hz, -CH=CH $\overline{H}$), 4.76 (1H, app d, J = 17.1 Hz, -CH=CH $\overline{H}$), 4.00 (2H, d, J = 13.6 Hz, -CH $\overline{H}$ N-), 3.24 (1H, dd, J = 13.4, 4.4 Hz, -NCH $\overline{H}$ CH-), 2.78 (2H, d, J = 13.4 Hz, -CH $\overline{H}$ N-), 2.52 (1H, dd, J = 13.2, 8.6 Hz, -NCH $\overline{H}$ CH-); ^{13}C NMR (100 MHz, CDCl_3) δ 148.1, 146.5, 143.3, 137.9, 136.0, 132.7, 131.4, 130.7, 129.7, 128.6, 128.3 (two peaks overlap), 127.8 (two peaks overlap), 127.2 (two peaks overlap), 126.8, 126.0, 125.6, 117.3, 83.1, 59.0, 51.4; IR (neat) 3289, 3057, 2359, 1597, 1491, 1447, 1373, 1321, 1240, 1172, 1018, 908, 893, 752, 733, 700 cm^{-1} ; HRMS (ESI-TOF) Calcd. for $\text{C}_{51}\text{H}_{42}\text{NO}_2$: 700.3210 ($[\text{M} + \text{H}]^+$), Found: 700.3209 ($[\text{M} + \text{H}]^+$).

Amino alcohol (S)-1b. A mixture of (S)-**13** (415 mg, 1.05 mmol), *N,N*-dimethylbarbituric acid (NDMBA) (540 mg, 3.5 mmol), $\text{Pd}(\text{OAc})_2$ (26 mg, 0.12 mmol), and triphenylphosphine (121 mg, 0.46 mmol) in CH_2Cl_2 (8.5 mL) was stirred at room temperature for 2 h under argon atmosphere. After addition of CH_2Cl_2 , the organic layer was washed with saturated NaHCO_3 , dried over Na_2SO_4 and concentrated. The resulting residue was purified by flash column chromatography on silica gel (eluting with CH_2Cl_2 /MeOH/ NH_3 = 20:1:0.1) to give (S)-**1b** (237 mg, 0.67 mmol, 64% yield). $[\alpha]_{\text{D}}^{26}$ 470.5 (c = 1.0, CHCl_3); ^1H NMR (400 MHz, CD_3OD) δ 8.05 (2H, s, Ar- $\overline{H}$), 7.94 (2H, d, J = 8.3 Hz, Ar- $\overline{H}$), 7.43 (2H, app t, J = 7.8 Hz, Ar- $\overline{H}$), 7.20 (4H, app t, J = 8.0 Hz, Ar- $\overline{H}$), 5.02 (2H, d, J = 13.0 Hz, -CH $\overline{H}$ OH), 4.96 (2H, d, J = 12.9 Hz, -CH $\overline{H}$ OH), 4.24 (2H, d, J = 12.9 Hz, -CH $\overline{H}$ N-), 3.19 (2H, d, J = 13.0 Hz, -CH $\overline{H}$ N-); ^{13}C NMR (100 MHz, CD_3OD) δ 137.6, 137.3, 134.3, 133.5, 132.1, 129.3 (two peaks overlap), 128.0, 126.9, 126.8, 64.2, 43.4; IR (neat) 3296, 2868, 1666, 1595, 1447, 1360, 1072, 1022, 891, 750 cm^{-1} ; HRMS (ESI-TOF) Calcd. for $\text{C}_{24}\text{H}_{22}\text{NO}_2$: 356.1645 ($[\text{M} + \text{H}]^+$), Found: 356.1651 ($[\text{M} + \text{H}]^+$).

Amino alcohol (S)-1c. (S)-**1c** was prepared in a similar manner as described above using (S)-**14**

instead of (*S*)-**13** (74% yield). $[\alpha]_{\text{D}}^{25}$ 406.6 ($c = 1.0$, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.99 (2H, s, Ar-H), 7.91 (2H, d, $J = 8.3$ Hz, Ar-H), 7.47 (2H, app t, $J = 6.3$ Hz, Ar-H), 7.23 (4H, app t, $J = 7.3$ Hz, Ar-H), 5.22 (2H, d, $J = 11.5$ Hz, -CHHN-), 3.38 (2H, d, $J = 12.4$ Hz, -CHHN-), 1.92 (6H, s, -CH₃), 1.70 (6H, s, -CH₃); ^{13}C NMR (100 MHz, CDCl_3) δ 143.5, 137.8, 132.4, 130.4, 128.4 (two peaks overlap), 127.2, 125.9, 125.8, 125.4, 73.4, 44.3, 32.5, 32.2; IR (neat) 3377, 3053, 2976, 1667, 1587, 1447, 1379, 1361, 1150, 941, 909, 891, 748, 731, 598 cm^{-1} ; HRMS (ESI-TOF) Calcd. for $\text{C}_{28}\text{H}_{30}\text{NO}_2$: 412.2271 ($[\text{M} + \text{H}]^+$), Found: 412.2268 ($[\text{M} + \text{H}]^+$).

Amino alcohol (*S*)-1d.⁸ (*S*)-**1d** was prepared in a similar manner as described above using (*S*)-**15** instead of (*S*)-**13** (95% yield). $[\alpha]_{\text{D}}^{28}$ -18.2 ($c = 1.0$, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.65 (2H, d, $J = 8.0$ Hz, Ar-H), 7.43-7.35 (20H, m, Ar-H), 7.25 (8H, app t, $J = 6.4$ Hz, Ar-H), 3.97 (2H, d, $J = 13.0$ Hz, -CHHN-), 2.80 (2H, d, $J = 12.9$ Hz, -CHHN-); ^{13}C NMR (100 MHz, CDCl_3) δ 147.8, 146.7, 142.8, 137.5, 131.6, 130.7, 129.8, 128.7, 128.1, 127.9, 127.8 (two peaks overlap), 127.5, 127.2 (two peaks overlap), 126.9, 126.0, 125.6, 83.2, 44.9; IR (neat) 3495, 3057, 1597, 1491, 1447, 1323, 1180, 1159, 1018, 909, 891, 752, 733, 700, 584 cm^{-1} ; HRMS (ESI-TOF) Calcd. for $\text{C}_{48}\text{H}_{38}\text{NO}_2$: 660.2897 ($[\text{M} + \text{H}]^+$), Found: 660.2893 ($[\text{M} + \text{H}]^+$).

Amine (*S*)-1e. A mixture of (*S*)-**1d** (100 mg, 0.15 mmol), $\text{BF}_3 \cdot \text{OEt}_2$ (96 μL , 0.76 mmol) and triethylsilane (0.24 mL, 1.5 mmol) in CH_2Cl_2 (1.5 mL) was stirred at room temperature for 10 h under argon atmosphere. After addition of triethylamine (0.22 mL, 1.5 mmol), the mixture was filtered through a pad of Celite, and the filtrate was concentrated. The resulting residue was purified by flash column chromatography on silica gel (eluting with hexane/ethyl acetate/triethylamine = 3:1:0.04) to give (*S*)-**1e** (68 mg, 0.11 mmol, 71% yield). $[\alpha]_{\text{D}}^{27}$ 59.0 ($c = 0.78$, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.68 (2H, d, $J = 8.0$ Hz, Ar-H), 7.34 (8H, app dd, $J = 14.2, 7.1$ Hz, Ar-H), 7.28 (8H, app t, $J = 5.7$ Hz, Ar-H), 7.20 (3H, app t, $J = 7.1$ Hz, Ar-H), 7.14 (9H, app t, $J = 6.6$ Hz, Ar-H), 6.01 (2H, s, -CHPh₂), 3.91 (2H, d, $J = 13.0$ Hz, -CHHN-), 2.97 (2H, d, $J = 13.0$ Hz, -CHHN-); ^{13}C NMR (100 MHz, CDCl_3) δ 143.6, 143.2, 139.6, 136.3, 133.7, 132.3, 130.2, 129.7, 129.6, 128.4, 128.3, 128.1, 127.3, 126.6, 126.4, 125.3, 54.1, 43.6; IR (neat) 3059, 3024, 1593, 1492, 1449, 1078, 1030, 907, 750, 733, 700, 567 cm^{-1} ; HRMS (ESI-TOF) Calcd. for $\text{C}_{48}\text{H}_{38}\text{N}$: 628.2999 ($[\text{M} + \text{H}]^+$), Found: 628.2999 ($[\text{M} + \text{H}]^+$).

General Procedure for the Organocatalytic α -Hydroxyamination Reaction between Nitrosobenzene and Aldehydes. A mixture of (*S*)-**1d** (13.2 mg, 0.020 mmol, 10 mol%) and nitrosobenzene (21.4 mg, 0.20 mmol) in THF (1.0 mL) was stirred at 0 $^\circ\text{C}$ for 3 minutes. To the mixture was then added an aldehyde (0.60 mmol) dropwise. After stirring for 1 h at 0 $^\circ\text{C}$, the reaction mixture was then transferred to a methanol suspension of NaBH_4 (23 mg, 0.60 mmol) at

0 °C. After 15 minutes, the reaction mixture was treated with saturated NaCl, extracted with CH₂Cl₂ (5 mL x 3) and dried over Na₂SO₄. The organic layer was concentrated and the resulting residue was purified by flash column chromatography on silica gel to give the corresponding 2-(*N*-phenylhydroxyamino)alcohol. The enantiomeric purity was determined by chiral HPLC analysis.

2-(*N*-Phenylhydroxyamino)propanol:

Prepared according to the general procedure with propanal (43 µL, 0.60 mmol). The crude material was purified by column chromatography on silica gel (eluting with hexane/ethyl acetate = 3:1) to give the title compound as a pale yellow oil [90% yield (30.2 mg)].

The enantiomeric purity was determined by HPLC analysis (Daicel Chiralcel AD-H, hexane/ethanol = 10:1, flow rate = 1.0 mL/min, retention time; 10.7 min (minor) and 15.0 min (major)).

$[\alpha]_D^{24}$ 8.0 ($c = 1.0$, CHCl₃; 99% ee); ¹H NMR (400 MHz, CD₃OD) δ 7.22 (2H, app t, $J = 7.3$ Hz, Ar-H), 7.11 (2H, dd, $J = 8.8, 1.2$ Hz, Ar-H), 6.86 (1H, app tt, $J = 7.3, 1.2$ Hz, Ar-H), 3.80 (1H, dd, $J = 10.7, 7.1$ Hz, -CHHOH), 3.74-3.65 (1H, m, -NCH-), 3.55 (1H, dd, $J = 10.8, 5.9$ Hz, -CHHOH), 1.00 (3H, d, $J = 6.6$ Hz, -CH₃); ¹³C NMR (100 MHz, CD₃OD) δ 153.8, 129.4, 121.8, 117.3, 64.8, 63.7, 11.0; IR (neat) 3300, 2976, 2936, 2882, 1597, 1489, 1452, 1034, 760, 737, 694 cm⁻¹; HRMS (ESI-TOF) Calcd. for C₉H₁₄NO₂: 168.1019 ([M + H]⁺), Found: 168.1018 ([M + H]⁺).

2-(*N*-Phenylhydroxyamino)-3-methylbutanol:

Prepared according to the general procedure with 3-methylbutanal (64 µL, 0.60 mmol). The crude material was purified by column chromatography on silica gel (eluting with CH₂Cl₂/MeOH = 30:1) to give the title compound as a pale yellow oil [70% yield (27.5 mg)].

The enantiomeric purity was determined by HPLC analysis (Daicel Chiralcel OJ-H, hexane/ethanol = 20:1, flow rate = 1.0 mL/min, retention time; 12.8 min (minor) and 18.3 min (major)).

$[\alpha]_D^{29}$ 11.6 ($c = 1.0$, CHCl₃; 97% ee); ¹H NMR (400 MHz, CD₃OD) δ 7.18 (2H, app t, $J = 7.3$ Hz, Ar-H), 7.05 (2H, d, $J = 7.8$ Hz, Ar-H), 6.77 (1H, app t, $J = 7.3$ Hz, Ar-H), 3.82-3.74 (2H, m, -CH₂OH), 3.25-3.20 (1H, m, -NCH-), 2.20-2.09 (1H, m, -CH₃CH-), 1.02 (6H, app t, $J = 7.1$ Hz, -CH(CH₃)₂); ¹³C NMR (100 MHz, CD₃OD) δ 154.9, 129.4, 120.6, 115.9, 73.7, 61.7, 29.2, 21.5, 20.5; IR (neat) 3298, 2959, 2872, 1597, 1489, 1458, 1244, 1057, 1017, 752, 694 cm⁻¹; HRMS (ESI-TOF) Calcd. for C₁₁H₁₈NO₂: 196.1332 ([M + H]⁺), Found: 196.1334 ([M + H]⁺).

2-(*N*-Phenylhydroxyamino)hexanol:

Prepared according to the general procedure with hexanal (72 µL, 0.60 mmol). The crude material was purified by column chromatography on silica gel (eluting with hexane/ethyl acetate = 3:1) to give the title compound as a pale yellow oil [76% yield (31.7 mg)].

The enantiomeric purity was determined by HPLC analysis (Daicel Chiralcel AD-H, hexane/ethanol = 10:1, flow rate = 0.5 mL/min, retention time; 13.8 min (minor) and 15.8 min (major)).

$[\alpha]_D^{24}$ 20.2 ($c = 0.5$, CHCl_3 ; 96% ee); ^1H NMR (400 MHz, CD_3OD) δ 7.20 (2H, app t, $J = 7.3$ Hz, Ar-H), 7.08 (2H, dd, $J = 9.0, 1.2$ Hz, Ar-H), 6.81 (1H, app t, $J = 7.3$ Hz, Ar-H), 3.72 (1H, dd, $J = 11.0, 6.6$ Hz, -CHHOH), 3.62-3.49 (2H, m, -CHHOH, -NCH-), 1.70-1.62 (1H, m, -CHCHH-), 1.57-1.46 (1H, m, -CHCHH-), 1.38-1.23 (4H, m, $-(\text{CH}_2)_2\text{CH}_3$), 0.85 (3H, t, $J = 6.8$ Hz, $-\text{CH}_3$); ^{13}C NMR (100 MHz, CD_3OD) δ 154.3, 129.4, 121.2, 116.6, 68.3, 63.1, 30.2, 28.1, 23.9, 14.4; IR (neat) 3312, 2955, 2931, 2870, 1597, 1489, 1454, 1053, 1029, 758, 694 cm^{-1} ; HRMS (ESI-TOF) Calcd. for $\text{C}_{12}\text{H}_{20}\text{NO}_2$: 210.1489 ($[\text{M} + \text{H}]^+$), Found: 210.1491 ($[\text{M} + \text{H}]^+$).

2-(*N*-Phenylhydroxyamino)-3-phenylpropanol:

Prepared according to the general procedure with 3-phenylpropanal (79 μL , 0.60 mmol). The crude material was purified by column chromatography on silica gel (eluting with hexane/ethyl acetate = 4:1) to give the title compound as a pale yellow oil [80% yield (38.8 mg)].

The enantiomeric purity was determined by HPLC analysis (Daicel Chiralcel AD-H, hexane/2-propanol = 10:1, flow rate = 1.0 mL/min, retention time; 12.1 min (minor) and 14.9 min (major)).

$[\alpha]_D^{24}$ -99.3 ($c = 0.8$, CHCl_3 ; 98% ee); ^1H NMR (400 MHz, CD_3OD) δ 7.24-7.19 (4H, m, Ar-H), 7.13-7.10 (5H, m, Ar-H), 6.85 (1H, app t, $J = 7.3$ Hz, Ar-H), 3.84-3.74 (2H, m, -CHHOH, -NCH-), 3.57 (1H, dd, $J = 10.2, 3.6$ Hz, -CHHOH), 2.86 (1H, dd, $J = 13.7, 4.9$ Hz, -CHHPh), 2.74 (1H, dd, $J = 13.7, 7.8$ Hz, -CHHPh); ^{13}C NMR (100 MHz, CD_3OD) δ 153.9, 141.0, 130.0, 129.4, 129.1, 126.8, 121.7, 117.0, 71.1, 62.1, 33.8; IR (neat) 3298, 3061, 3026, 2928, 2884, 1597, 1487, 1452, 1215, 1028, 754, 723, 696 cm^{-1} ; HRMS (ESI-TOF) Calcd. for $\text{C}_{15}\text{H}_{18}\text{NO}_2$: 244.1332 ($[\text{M} + \text{H}]^+$), Found: 244.1333 ($[\text{M} + \text{H}]^+$).

2-(*N*-Phenylhydroxyamino)-3-cyclohexylpropanol:

Prepared according to the general procedure with 3-cyclohexylpropanal (84 mg, 0.60 mmol). The crude material was purified by column chromatography on silica gel (eluting with hexane/ethyl acetate = 3:1) to give the title compound as a pale yellow oil [86% yield (42.7 mg)].

The enantiomeric purity was determined by HPLC analysis (Daicel Chiralcel AD-H, hexane/ethanol = 20:1, flow rate = 1.0 mL/min, retention time; 11.8 min (minor) and 13.3 min (major)).

$[\alpha]_D^{21}$ 14.8 ($c = 0.87$, CHCl_3 ; 99% ee); ^1H NMR (400 MHz, CD_3OD) δ 7.20 (2H, app t, $J = 7.3$ Hz, Ar-H), 7.08 (2H, d, $J = 7.8$ Hz, Ar-H), 6.81 (1H, app t, $J = 7.3$ Hz, Ar-H), 3.71 (1H, dd, $J = 10.5, 6.8$ Hz, -CHHOH), 3.68-3.62 (1H, m, -NCH-), 3.55 (1H, dd, $J = 10.7, 5.4$ Hz, -CHHOH), 1.76-1.53 (6H, m, $-\text{CH}_2-$), 1.39-1.27 (2H, m, $-\text{CH}_2-$), 1.23-1.08 (3H, m, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$), 0.91-0.78 (2H, m, $-\text{CH}_2-$); ^{13}C NMR (100 MHz, CD_3OD) δ 154.3, 129.3, 121.2, 116.6, 65.6, 63.3, 36.0, 35.6, 34.9, 34.6, 27.8,

27.5, 27.4; IR (neat) 3316, 2920, 2851, 1597, 1489, 1449, 1028, 758, 694 cm^{-1} ; HRMS (ESI-TOF) Calcd. for $\text{C}_{15}\text{H}_{24}\text{NO}_2$: 250.1802 ($[\text{M} + \text{H}]^+$), Found: 250.1804 ($[\text{M} + \text{H}]^+$).

2-(*N*-Phenylhydroxyamino)-10-undecenol:

Prepared according to the general procedure with 10-undecenal (120 μL , 0.60 mmol). The crude material was purified by column chromatography on silica gel (eluting with hexane/ethyl acetate = 4:1) to give the title compound as a pale yellow oil [77% yield (42.9 mg)].

The enantiomeric purity was determined by HPLC analysis (Daicel Chiralcel AD-H, hexane/ethanol = 20:1, flow rate = 1.0 mL/min, retention time; 9.2 min (minor) and 12.8 min (major)).

$[\alpha]_{\text{D}}^{30}$ 9.4 ($c = 1.0$, CHCl_3 ; 99% ee); ^1H NMR (400 MHz, CD_3OD) δ 7.19 (2H, app t, $J = 7.3$ Hz, Ar-H), 7.08 (2H, d, $J = 7.6$ Hz, Ar-H), 6.81 (1H, app t, $J = 7.3$ Hz, Ar-H), 5.83-5.72 (1H, m, $-\text{CH}=\text{CH}_2$), 4.98-4.87 (2H, m, $-\text{CH}=\text{CH}_2$), 3.72 (1H, dd, $J = 11.0, 6.8$ Hz, $-\text{CHHOH}$), 3.60-3.51 (2H, m, $-\text{CHHOH}$, $-\text{NCH}_2-$), 2.00 (2H, app q, $J = 7.6$ Hz, $-\text{CH}_2\text{CH}=\text{CH}_2$), 1.71-1.59 (1H, m, $-\text{NCHCH}_2-$), 1.54-1.44 (1H, m, $-\text{NCHCH}_2-$), 1.35-1.20 (10H, m, $-(\text{CH}_2)_5\text{CH}_2\text{CH}=\text{CH}_2$); ^{13}C NMR (100 MHz, CD_3OD) δ 154.3, 140.0, 129.3, 121.2, 116.6, 114.5, 68.3, 63.1, 34.9, 30.7, 30.4, 30.14, 30.10, 28.3, 27.9; IR (neat) 3354, 2924, 2854, 2363, 1489, 1462 cm^{-1} ; HRMS (ESI-TOF) Calcd. for $\text{C}_{17}\text{H}_{28}\text{NO}_2$: 278.2115 ($[\text{M} + \text{H}]^+$), Found: 278.2118 ($[\text{M} + \text{H}]^+$).

2-(*N*-Phenylhydroxyamino)-5-benzyloxypentanol:

Prepared according to the general procedure with 5-benzyloxypentanal (115 mg, 0.60 mmol). The crude material was purified by column chromatography on silica gel (eluting with hexane/ethyl acetate = 1:1) to give the title compound as a pale yellow oil [86% yield (51.7 mg)].

The enantiomeric purity was determined by HPLC analysis (Daicel Chiralcel AD-H, hexane/ethanol = 5:1, flow rate = 0.5 mL/min, retention time; 18.7 min (minor) and 20.8 min (major)).

$[\alpha]_{\text{D}}^{29}$ 6.0 ($c = 1.0$, CHCl_3 ; 97% ee); ^1H NMR (400 MHz, CD_3OD) δ 7.31-7.25 (5H, m, Ar-H), 7.18 (2H, app t, $J = 7.4$ Hz, Ar-H), 7.06 (2H, dd, $J = 9.0, 1.2$ Hz, Ar-H), 6.80 (1H, app t, $J = 7.3$ Hz, Ar-H), 4.43 (2H, s, $-\text{CH}_2\text{Ph}$), 3.73-3.68 (1H, m, $-\text{CHHOH}$), 3.61-3.55 (2H, m, $-\text{CHHOH}$, $-\text{NCH}_2-$), 3.46 (2H, t, $J = 5.6$ Hz, $-\text{CH}_2\text{CH}_2\text{OBn}$), 1.78-1.60 (4H, m, $-\text{NCH}(\text{CH}_2)_2-$); ^{13}C NMR (100 MHz, CD_3OD) δ 154.2, 139.5, 129.4, 129.2, 128.8, 128.5, 121.2, 116.5, 73.8, 71.4, 67.8, 63.0, 27.9, 25.4; IR (neat) 3331, 2924, 2857, 1595, 1489, 1452, 1362, 1207, 1094, 1072, 1028, 752, 696, 586 cm^{-1} ; HRMS (ESI-TOF) Calcd. for $\text{C}_{18}\text{H}_{24}\text{NO}_3$: 302.1751 ($[\text{M} + \text{H}]^+$), Found: 302.1751 ($[\text{M} + \text{H}]^+$).

Determination of the Absolute Configuration of 2-(*N*-Phenylhydroxyamino)propanol. The absolute configuration was determined to be *S* by converting of the title compound to the known 2-(phenylamino)propanol⁹ and by comparing the sign of optical rotation: A mixture of (*S*)-**1d** (13.2 mg, 0.020 mmol, 10 mol%) and nitrosobenzene (21.4 mg, 0.20 mmol) in THF (1.0 mL) was stirred

at 0 °C for 3 minutes. To the mixture was added propanal (43.2 μ L, 0.60 mmol) dropwise. After stirring for 1 h at 0 °C, the reaction mixture was treated with LiAlH₄ (76 mg, 2.0 mmol) at 0 °C. After 10 minutes, the reaction mixture was heated to 73 °C with stirring for 4 h. After cooling to 0 °C, the reaction mixture was treated with saturated NH₄Cl and extracted with ethyl acetate (5 mL x 3). The organic layer was washed with H₂O and brine; dried over Na₂SO₄ and then concentrated. The resulting residue was purified by flash column chromatography on silica gel (eluting with hexane/ethyl acetate = 3:1) to give 2-(phenylamino)propanol as a pale yellow oil [77% yield (23.2 mg)].

The enantiomeric purity was determined by HPLC analysis (Daicel Chiralcel AD-H, hexane/2-propanol = 20:1, flow rate = 1.0 mL/min, retention time; 13.8 min (minor) and 14.8 min (major)).

$[\alpha]_{\text{D}}^{28}$ 19.8 (c = 1.0, CHCl₃; 96% ee); Other spectrum data have been reported in the previous work.⁷

Determination of the Absolute Configuration of 2-(*N*-Phenylhydroxyamino)-3-methylbutanol (Entry 2 in Table 2). The absolute configuration was determined to be *S* by converting of the title compound to 2-(phenylamino)-3-methylbutanol in a similar manner as described above and then by comparing the sign of optical rotation with (*S*)-2-(phenylamino)-3-methylbutanol, which was prepared by the coupling reaction of (*S*)-2-amino-3-methylbutanol with iodobenzene according to the literature procedure.¹⁰

(*S*)-2-(phenylamino)-3-methylbutanol:

$[\alpha]_{\text{D}}^{27}$ -26.1 (c = 1.0, CHCl₃); ¹H NMR (400 MHz, CDCl₃) δ 7.18 (2H, app t, J = 7.3 Hz, Ar-H), 6.73-6.67 (3H, m, Ar-H), 3.77 (1H, dd, J = 11.0, 4.2 Hz, -CHHOH), 3.53 (1H, dd, J = 11.0, 7.1 Hz, -CHHOH), 3.33-3.29 (1H, m, -NCH-), 1.95-1.83 (1H, m, -CH(CH₃)₂), 0.97 (6H, dd, J = 10.8, 6.8 Hz, -CH(CH₃)₂); ¹³C NMR (100 MHz, CDCl₃) δ 148.4, 129.4, 117.7, 113.7, 62.7, 61.0, 30.2, 19.2, 19.1; IR (neat) 3392, 2959, 1601, 1499, 1319, 1063, 748, 692 cm⁻¹; HRMS (ESI-TOF) Calcd. for C₁₁H₁₈NO: 180.1383 ([M + H]⁺), Found: 180.1387 ([M + H]⁺).

Determination of the Absolute Configuration of 2-(*N*-Phenylhydroxyamino)-3-phenylpropanol (Entry 4 in Table 2). The absolute configuration was determined to be *S* by converting of the title compound to 2-(phenylamino)-3-phenylpropanol in a similar manner as described above and then by comparing the sign of optical rotation with (*S*)-2-(phenylamino)-3-phenylpropanol, which was prepared by the coupling reaction of (*S*)-2-amino-3-phenylpropanol with iodobenzene according to the literature procedure.¹⁰

(S)-2-(N-Phenylhydroxyamino)-3-phenylpropanol:

$[\alpha]_D^{29}$ 2.8 ($c = 1.0$, CHCl_3); ^1H NMR (400 MHz, CDCl_3) δ 7.29 (2H, app t, $J = 7.6$ Hz, Ar-H), 7.24-7.17 (5H, m, Ar-H), 6.75 (1H, app t, $J = 7.4$ Hz, Ar-H), 6.68 (2H, d, $J = 7.8$ Hz, Ar-H), 3.78-3.69 (2H, m, -CHHOH, -NCH-), 3.50 (1H, dd, $J = 10.8, 5.1$ Hz, -CHHOH), 2.93 (1H, dd, $J = 13.9, 5.1$ Hz, -CHHPh), 2.84 (1H, dd, $J = 13.7, 7.3$ Hz, -CHHPh), 1.91 (1H, s, -CH₂OH); ^{13}C NMR (100 MHz, CDCl_3) δ 147.1, 137.9, 129.4, 129.3, 128.5, 126.5, 118.1, 113.9, 63.1, 55.8, 37.3; IR (neat) 3385, 2928, 1599, 1495, 1317, 1030, 746, 692 cm^{-1} ; HRMS (ESI-TOF) Calcd. for $\text{C}_{15}\text{H}_{18}\text{NO}$: 228.1383 ($[\text{M} + \text{H}]^+$), Found: 228.1381 ($[\text{M} + \text{H}]^+$).

2-(4-Methoxyphenylamino)-3-cyclohexylpropanol (5):

A mixture of (S)-**1d** (13.2 mg, 0.020 mmol, 10 mol%), 4-methoxynitrosobenzene (27.4 mg, 0.20 mmol) in THF (1.0 mL) was stirred at 0 °C for 3 minutes. To the mixture was added 3-cyclohexylpropanal (84 mg, 0.60 mmol) dropwise. After stirring for 1 h, the reaction mixture was treated with LiAlH_4 (76 mg, 2.0 mmol) at 0 °C. After 10 minutes, the mixture was heated to 73 °C with stirring for 4 h. After cooling to 0 °C, the mixture was treated with saturated NH_4Cl and extracted with ethyl acetate (5 mL x 3). The organic layer was washed with H_2O and brine; dried over Na_2SO_4 and then concentrated. The resulting residue was purified by flash column chromatography on silica gel (eluting with hexane/ethyl acetate = 5:1) to give the title compound as a pale yellow oil [74% yield (39.2 mg)].

The enantiomeric purity was determined by HPLC analysis (Daicel Chiralcel AD-H, hexane/2-propanol = 50:1, flow rate = 1.0 mL/min, retention time; 63.1 min (minor) and 66.1 min (major)).

$[\alpha]_D^{29}$ -19.2 ($c = 0.9$, CH_3OH ; 95% ee); ^1H NMR (400 MHz, CDCl_3) δ 6.77 (2H, d, $J = 9.0$ Hz, Ar-H), 6.63 (2H, d, $J = 9.0$ Hz, Ar-H), 3.74 (3H, s, -OCH₃), 3.73 (1H, dd, $J = 10.2, 3.7$ Hz, -CHHOH), 3.50-3.45 (1H, m, -NCH-), 3.42 (1 H, dd, $J = 10.5, 6.1$ Hz, -CHHOH), 1.76-1.60 (5H, m, -CH₂-, -NCHCHH-), 1.44-1.30 (3H, m, -CH₂-, -NCHCHH-), 1.27-1.08 (3H, m, -CH₂-, -CH₂CH-), 0.98-0.81 (2H, m, -CH₂-); ^{13}C NMR (100 MHz, CDCl_3) δ 152.3, 141.7, 115.2, 114.8, 64.8, 55.8, 53.9, 40.2, 34.4, 33.63, 33.57, 26.5, 26.3, 26.2; IR (neat) 3383, 2920, 2849, 1510, 1447, 1234, 1179, 1038, 818, 754 cm^{-1} ; HRMS (ESI-TOF) Calcd. for $\text{C}_{16}\text{H}_{26}\text{NO}_2$: 264.1958 ($[\text{M} + \text{H}]^+$), Found: 264.1954 ($[\text{M} + \text{H}]^+$).

Determination of the Absolute Configuration of 2-(4-Methoxyphenylamino)-3-cyclohexylpropanol (5). The absolute configuration was determined to be *S* by comparing the sign of optical rotation with (S)-2-(4-methoxyphenylamino)-3-cyclohexylpropanol, which was prepared by the coupling reaction of (S)-2-amino-3-cyclohexylpropanol with 4-iodoanisole according to the literature procedure.¹⁰

***t*-Butyl 3-cyclohexyl-2-(4-methoxyphenylamino)propylcarbamate (6):**

A mixture of (*S*)-**1d** (16.5 mg, 0.025 mmol, 10 mol%) and 4-methoxynitrosobenzene (33.9 mg, 0.25 mmol) in THF (1.24 mL) was stirred at 0 °C for 3 minutes. To the mixture was added 3-cyclohexylpropanal (104 mg, 0.74 mmol) dropwise. After stirring for 1 h, the reaction mixture was treated with *O*-benzylhydroxylamine (122 mg, 0.99 mmol) and magnesium sulfate at 0 °C. After further 1 h, the reaction mixture was treated with LiAlH₄ (282 mg, 7.4 mmol) at same temperature and stirred there for 10 minutes. The reaction mixture was then heated to 73 °C with stirring for 4 h. After cooling to 0 °C, the mixture was treated with saturated Rochelle salt and stirred at room temperature for 10 h. To the resulting mixture was added di-*t*-butyl dicarbonate (0.35 mL, 1.5 mmol) at room temperature. After 5 h, the mixture was extracted with ether (5 mL x 3). The organic layer was washed with saturated NaHCO₃, H₂O and brine; dried over Na₂SO₄ and then concentrated. The resulting residue was purified by flash column chromatography on silica gel (eluting with hexane/ethyl acetate = 5:1) to give the title compound as a pale yellow oil [56% yield (49.9 mg)]. The enantiomeric purity was determined by HPLC analysis (Daicel Chiralcel AS-H, hexane/2-propanol = 20:1, flow rate = 0.5 mL/min, retention time; 14.7 min (major) and 16.6 min (minor)).

$[\alpha]_D^{26}$ -61.8 (*c* = 1.0, CHCl₃; 95% ee); ¹H NMR (400 MHz, CDCl₃) δ 6.77 (2H, d, *J* = 9.0 Hz, Ar-H), 6.58 (2H, d, *J* = 8.8 Hz, Ar-H), 4.82 (1H, s, ArNH-), 3.74 (3H, s, -OCH₃), 3.50-3.43 (1H, m, -NCH-), 3.36-3.28 (1H, m, -CHCHHNH-), 3.09-3.03 (1H, m, -CHCHHNH-), 1.76-1.61 (5H, m, -CH₂-, -NCHCHH-), 1.44 (9H, s, -C(CH₃)₃), 1.37-1.09 (6H, m, -CH₂-, -CH₂CH-, -NCHCHH-), 0.92-0.85 (2H, m, -CH₂-); ¹³C NMR (100 MHz, CDCl₃) δ 152.1, 128.8, 128.1, 115.0, 114.5, 79.2, 55.8, 52.0, 44.7, 41.5, 34.3, 33.6, 33.5, 28.4, 26.5, 26.2, 26.1; IR (neat) 3368, 2976, 2922, 2851, 1697, 11510, 1449, 1366, 1234, 1169, 1040, 820 cm⁻¹; HRMS (ESI-TOF) Calcd. for C₂₁H₃₅N₂O₃: 363.2642 ([M + H]⁺), Found: 363.2641 ([M + H]⁺).

References

- (1) Aggarwal, V. K.; Bae, I.; Lee, H.-Y. *Tetrahedron* **2004**, *60*, 9725.
- (2) Pihko, P. M.; Aho, J. E. *Org. Lett.* **2004**, *6*, 3849.
- (3) Albert, D. *Synthesis* **2004**, *5*, 706.
- (4) Ooi, T.; Kameda, M.; Maruoka, K. *J. Am. Chem. Soc.* **2003**, *125*, 5139.
- (5) Kano, T.; Takai, J.; Tokuda, O.; Maruoka, K. *Angew. Chem., Int. Ed.* **2005**, *44*, 3055.
- (6) Zuo, G.; Zhang, Q.; Xu, J. *Heteroatom Chemistry* **2003**, *14*, 42.
- (7) Ooi, T.; Uematsu, Y.; Maruoka, K. *J. Org. Chem.* **2003**, *68*, 4576.
- (8) Ooi, T.; Ohara, D.; Tamura, M.; Maruoka, K. *J. Am. Chem. Soc.* **2004**, *126*, 6844.
- (9) Ozawa, F.; Okamoto, H.; Kawagishi, S.; Yamamoto, S.; Minami, T.; Yoshifuji, M. *J. Am. Chem.*

Soc. **2002**, *124*, 10968.

(10) Job. G. E.; Buchwald, S. L. *Org. Lett.* **2002**, *4*, 3703.