Supporting Information

Direct Boronation of Allyl Alcohols with Diboronic Acid using Palladium Pincer-Complex Catalysis. A Remarkably Facile Allylic Displacement of the Hydroxy Group Under Mild Reaction Conditions

Vilhelm J. Olsson, Sara Sebelius, Nicklas Selander and Kálmán J. Szabó*

Stockholm University, Arrhenius Laboratory, Department of Organic Chemistry
SE-106 91 Stockholm, Sweden. E-mail: kalman@organ.su.se. Fax: +46-8-15 49 08

Contents:

- Detailed experimental procedures and characterization of the products
- ^1H , ^{13}C and ^{19}F NMR spectra of **5a-j**

Experimental Procedures and Characterization of the Prepared Compounds

The reported reactions were conducted under argon atmosphere in order to standardize the reaction conditions, however it was found that using ambient atmosphere does not significantly influence the yields and the purity of the products. Depending on the solubility of the products the ^1H -NMR and ^{13}C -NMR spectra were recorded in acetone- d_6 (internal standard: 2.05 ppm, ^1H ; 30.60 ppm, ^{13}C), DMSO- d_6 (internal standard: 2.54 ppm, ^1H ; 40.45 ppm, ^{13}C), CD_3OD (internal standard: 3.31 ppm, ^1H ; 49.0 ppm, ^{13}C) and CDCl_3 (internal standard: 7.26 ppm, ^1H ; 77.36 ppm, ^{13}C) solutions at room temperature by a Varian 400 spectrometer. In mixed solvents (e.g. reaction mixtures) DMSO- d_6 was used as internal standard (2.54 ppm). Due to quadrupolar relaxation, the carbon attached to the boron atom was not always detected in ^{13}C -NMR. The ^{19}F -NMR spectra were recorded at either 282.2 or 376.3 MHz, using α,α,α -trifluorotoluene as standard.

Palladium pincer-complex **1** was prepared according to the procedure published by Yao and co-workers¹ except that the final product was purified by CH_2Cl_2 :ether 20:1 as eluent. The allylic alcohols were purchased from Aldrich or prepared according to literature procedures.^{2a-c} Diboronic acid (**2**) was purchased from Boron Molecular Ltd and was used as received.

General Procedure A. The corresponding allyl alcohol **3a-i** (0.15 mmol) was dissolved in a DMSO/MeOH mixture (0.3/0.3 ml) followed by addition of diboronic acid **2** (0.18 mmol) and pincer complex **1** (0.0075 mmol, 5 mol%). The reaction mixture was stirred for the allotted temperatures and times listed in Table 1. The ^1H -NMR spectrum of the allylboronic acid products **4a-h** was determined from the reaction mixture. Thereafter, KHF_2 (0.9 mmol, 70 mg) in water (0.7 mL) was added to the reaction mixture and the resulting suspension was stirred for 45 minutes at room temperature. The precipitate was separated and the filtrate containing the crude product was evaporated. The solid remaining after evaporation was extracted with acetone and filtered, subsequently the acetone solution was evaporated and the remaining solid was recrystallized from acetone/ether, methanol/ether or dichloromethane.

General Procedure B. The corresponding allylic alcohol **3j-k** (0.15 mmol) was dissolved in a DMSO/MeOH mixture (0.2/0.2 ml) followed by addition of diboronic acid **2** (0.18 mmol), pincer complex **1** (0.0075 mmol, 5 mol%) and a catalytic amount of p-toluenesulfonic acid (0.0075 mmol, 5 mol%). The reaction mixture was stirred for the allotted temperatures and times listed in Table 1 under Ar atmosphere. The work up procedure was the same as for procedure A.

Potassium Trifluoro(3-Phenyl-2-Propenyl)borate (5a) from Cinnamyl Alcohol (3a). This compound was prepared according to the above general procedure A from **3a** except that a different mixture of DMSO/MeOH (0.35/0.1 ml) was used as solvent. The ^1H NMR spectrum of allylboronic acid product **4a** was determined from the crude mixture. ^1H NMR (DMSO- d_6 /CD $_3$ OD): 7.29 (d, 7.4 Hz, 2H), 7.23 (t, 7.4 Hz, 2H), 7.11 (t, 7.4 Hz, 1H), 6.31 (dt, 7.0 Hz, 15.6 Hz, 1H), 6.23 (d, 15.6 Hz, 1H), 1.76 (d, 7.0 Hz, 2H). The NMR data for the isolated compound **5a**: ^1H NMR (acetone- d_6): 7.26 (d, 7.9 Hz, 2H), 7.19 (t, 7.9 Hz, 2H), 7.04 (t, 7.9 Hz, 1H), 6.50 (dt, 7.8 Hz, 15.8 Hz, 1H), 6.12 (d, 15.8 Hz, 1H), 1.29 (br, 2H). ^{13}C NMR (acetone- d_6): 141.7, 137.5, 129.8, 127.0, 126.7, 126.6. ^{19}F NMR (acetone- d_6): -140.0.

Potassium Trifluoro(3-Phenyl-2-Propenyl)Borate (5a) from Vinylbenzylalcohol (3b). This compound was prepared according to the above general procedure A from **3b** except that pure MeOH (0.35ml) was used as solvent. The spectral data for the allylboronic acid product **4a** and the isolated compound **5a** are given above.

Potassium Trifluoro(2-Octenyl)Borate (5b). This compound was prepared according to the above general procedure A. The ^1H NMR spectrum of allylboronic acid product **4b** was determined from the crude mixture. ^1H NMR (DMSO- d_6 /CD $_3$ OD): 5.39 (dt, 7.0 Hz, 15.1 Hz, 1H), 5.24 (dt, 6.6 Hz, 15.1 Hz, 1H), 1.90 (q, 7.0 Hz, 2H), 1.50 (d, 6.6 Hz, 2H), 1.25 (m, 6H), 0.84 (t, 7.0 Hz, 3H). The NMR data for the isolated compound **5b**: ^1H NMR (acetone- d_6): 5.54 (dt, 7.6 Hz, 15.0 Hz, 1H), 5.08 (dt, 6.6 Hz, 15.0 Hz, 1H), 1.90 (q, 6.7 Hz, 2H), 1.27 (m, 6H), 1.02 (br, 2H), 0.87 (t, 6.7 Hz, 3H). ^{13}C NMR (acetone- d_6): 134.8, 126.6, 34.6, 33.1, 31.6, 24.0, 15.1. ^{19}F NMR (acetone- d_6): -140.5.

Potassium Trifluoro(2-Hexenyl)Borate (5c). This compound was prepared according to the above general procedure A. The ^1H NMR spectrum of allylboronic acid product **4c** was determined from the crude mixture. ^1H NMR ($\text{DMSO-}d_6/\text{CD}_3\text{OD}$): 5.66 (q, 10.7 Hz, 2H), 1.90 (br, 2H), 1.69 (m, 5H), 1.46 (d, 10.7 Hz, 2H). The NMR data for the isolated compound **5b**: ^1H NMR (acetone- d_6): 5.54 (dt, 7.4 Hz, 15.0 Hz, 1H), 5.08 (dt, 6.8 Hz, 15.0 Hz, 1H), 1.88 (q, 7.4 Hz, 2H), 1.30 (h, 7.4 Hz, 2H), 1.02 (br, 2H), 0.85 (t, 7.4 Hz, 3H). ^{13}C NMR (acetone- d_6): 134.9, 126.5, 36.8, 24.8, 14.8. ^{19}F NMR (acetone- d_6): -141.0.

Potassium Trifluoro(3-Methyl-2-Butenyl)Borate (5d). This compound was prepared according to the above general procedure A, except that DMSO/MeOH (0.35/0.1 ml) was used as solvent. The ^1H NMR spectrum of allylboronic acid product **4d** was determined from the crude mixture. ^1H NMR ($\text{DMSO-}d_6/\text{CD}_3\text{OD}$): 5.12 (t, 7.5 Hz, 1H), 1.59 (s, 3H), 1.50 (s, 3H), 1.46 (d, 7.5 Hz, 2H). The NMR data for the isolated compound **5d**: ^1H NMR (acetone- d_6): 5.29 (t, 7.9 Hz, 1H), 1.59 (s, 3H), 1.51 (s, 3H), 0.96 (br, 2H). ^{13}C NMR (acetone- d_6): 128.9, 125.6, 26.9, 21.7, 18.4. ^{19}F NMR (acetone- d_6): -140.5.

Potassium Trifluoro(2-Cyclohexenyl)Borate (5e). This compound was prepared according to the above general procedure A. The ^1H NMR spectrum of allylboronic acid product **4e** was determined from the crude mixture. ^1H NMR ($\text{DMSO-}d_6/\text{CD}_3\text{OD}$): 5.66 (d, 10.3 Hz, 1H), 5.52 (d, 10.3 Hz, 1H), 1.90 (br, 2H), 1.62 (br, 1H), 1.60 (m, 4H). The NMR data for the isolated compound **5e**: ^1H NMR (acetone- d_6): 5.80 (d, 10.1 Hz, 1H), 5.34 (dq, 3.4 Hz, 10.1 Hz, 1H), 1.87 (m, 2H), 1.72 (m, 1H), 1.61 (m, 1H), 1.41 (m, 2H), 1.09 (br, 1H). ^{13}C NMR (acetone- d_6): 136.5, 122.9, 27.2, 26.4, 24.9. ^{19}F NMR (acetone- d_6): -146.5.

Potassium Trifluoro(4-Benzyloxy-2-Butenyl)Borate (5f). This compound was prepared according to the above general procedure A. The ^1H NMR spectrum of allylboronic acid product **4f** was determined from the crude mixture. ^1H NMR ($\text{DMSO-}d_6/\text{CD}_3\text{OD}$): 7.30 (m, 5H), 5.74 (dt, 7.5 Hz, 15.3 Hz, 1H), 5.44 (dt, 6.2 Hz, 15.3 Hz, 1H), 4.44 (s, 2H), 3.90 (d, 6.2 Hz, 2H), 1.61 (d, 7.5 Hz, 2H). The NMR data for the isolated compound **5f**: ^1H NMR (acetone- d_6): 7.22-7.35 (m, 5H), 5.88 (dt, 8.0 Hz, 15.1 Hz, 1H), 5.27 (dt, 6.7 Hz, 15.1 Hz, 1H), 4.45 (s, 2H), 3.87 (d, 6.7 Hz, 2H),

1.10 (br, 2H). ^{13}C NMR (acetone- d_6): 141.9, 141.2, 129.7, 129.2, 128.7, 122.6, 73.4, 72.2, 26.9. ^{19}F NMR (acetone- d_6): -140.9.

Potassium Trifluoro(4-Hydroxy-2-Butenyl)Borate (5g). This compound was prepared according to the above general procedure A, except that pure MeOH (0.35 ml) was used as solvent. The ^1H NMR spectrum of allylboronic acid product **4g** was determined from the crude mixture. ^1H NMR (CD_3OD): 5.71 (dt, 7.4 Hz, 15.2 Hz, 1H), 5.50 (dt, 5.9 Hz, 15.2 Hz, 1H), 3.97 (d, 5.9 Hz, 2H), 1.63 (d, 7.4 Hz, 2H). The NMR data for the isolated compound **5g**: ^1H NMR (CD_3OD): 5.81 (dt, 7.9 Hz, 15.0 Hz, 1H), 5.37 (dt, 6.7 Hz, 15.0 Hz, 1H), 3.93 (d, 6.7 Hz, 2H), 1.13 (br, 2H). ^{13}C NMR (CD_3OD): 137.7, 126.3, 64.6. ^{19}F NMR (CD_3OD): -143.0.

Potassium Trifluoro(6-Hydroxy-2-Cyclohexenyl)Borate (5h). This compound was prepared according to the above general procedure A, except that a DMSO/water mixture (0.5/0.1 ml) was used as solvent. The ^1H NMR spectrum of allylboronic acid product **4h** was determined from the crude mixture. ^1H NMR ($\text{DMSO-}d_6$): 5.60 (d, 9.6 Hz, 1H), 5.47 (d, 9.6 Hz, 1H), 3.94 (ddd, 3.1 Hz, 5.5 Hz, 9.0 Hz, 1H), 2.01 (m, 2H), 1.91 (br, 1H), 1.62 (m, 2H). The NMR data for the isolated compound **5h**: ^1H NMR ($\text{DMSO-}d_6$, assignments of the protons, in bold, are based on COSY experiment): 5.60 (d, 10.0 Hz, 1H, **H3**), 5.31 (d, 10.0 Hz, 1H, **H2**), 3.87 (br, 1H, **H6**), 3.14 (dq, 3.2 Hz, 8.4 Hz, 1H, OH), 2.06 (m, 1H, **H4eq**), 1.84 (d, 16.4 Hz, 1H, **H5eq**), 1.74 (dtd, 5.9 Hz, 5.9 Hz, 12.5 Hz, 1H, **H4ax**), 1.40 (dt, 5.9 Hz, 12.5 Hz, 1H, **H5ax**), 1.23 (br, 1H, **H1**). ^{13}C NMR ($\text{DMSO-}d_6$): 132.9, 121.7, 68.6, 30.5, 23.1. ^{19}F NMR ($\text{DMSO-}d_6$): -136.5.

The substitution pattern of **5h** was determined on the basis of the COSY spectrum. The COSY spectrum of **5h** clearly shows a crosspeak between **H1** (1.23 ppm) and **H6** (3.87 ppm) indicating that these protons are attached to neighbouring carbons bearing the

BF₃K (**C1**) and OH (**C6**) groups. The stereochemistry of **4h** and **5h** was determined by analysis of the J-coupling pattern of **H6** in the ¹H-NMR spectrum of **4h**. This peak appears as a ddd (3.1 Hz, 5.5 Hz, 9.0 Hz) at 3.94 ppm in the ¹H-NMR of **4h**. Decoupling of **H1** at 1.9 ppm leads to a simplification of the coupling pattern of **H6** at 3.94 ppm to a dd (3.1 Hz, 9.0 Hz). This indicates that the J-coupling constant between **H1** and **H6** is 5.5 Hz. This relatively small coupling constant is indicative to a gauche coupling between **H1** and **H6**; and therefore we assign the stereochemical relationship of **H1** and **H6** as syn. As a consequence, the relative stereochemistry of the BF₃K and OH groups is also syn to each other.

Potassium Trifluoro(5-Methoxycarbonyl-2-Cyclohexenyl)Borate (5i). This compound was prepared according to the above general procedure B, except that 2 equivalents (0.30 mmol) of diboronic acid **2** was used. The workup procedure was also slightly modified as the crude product obtained after evaporation of the solvent was first washed with chloroform and then extracted with acetone. The acetone solution was filtered and subsequently evaporated. Recrystallization from acetone/ether gave **5i** as a white solid. The ¹H NMR spectrum of allylboronic acid product **4i** was determined from the crude mixture. ¹H NMR (DMSO-d₆): 5.68 (m, 1H), 5.49 (m, 1H), 3.58 (s, 3H), 2.62 (m, 1H), 2.10 (m, 2H), 1.92 (dt, 12.1 Hz, 3.0 Hz, 1H), 1.75 (br, 1H) 1.63 (m, 1H). The NMR data for the isolated compound **5i**: ¹H NMR (CD₃OD, assignments of the protons, in bold, are based on COSY experiment): 5.80 (ddt, 10.1 Hz, 4.5 Hz, 2.1 Hz, 1H, **H2**), 5.39 (dq, 10.1 Hz, 3.2 Hz, 1H, **H3**), 3.64 (s, 3H, **Me**), 2.87 (ddt, 11.7, 8.0 Hz, 3.2 Hz, 1H, **H5**), 2.13 (m, 2H, **H4**), 1.94 (dt, 11.7, 3.2 Hz, 1H, **H6eq**), 1.66 (ddd, 11.7 Hz, 11.7 Hz, 6.4 Hz, 1H **H6ax**), 1.26 (br, 1H, **H1**). ¹³C NMR (CD₃OD): 180.0, 134.7, 120.5, 51.8, 39.2, 28.7, 28.2. ¹⁹F NMR (CD₃OD): -146.8.

The stereochemistry of the product was determined on the basis of the J-coupling constants of **H6ax** in **5i**. Proton **H6ax** (assigned by the COSY spectrum) appears as ddd (11.7 Hz, 11.7 Hz, 6.4 Hz) at 1.66 ppm in the ^1H NMR spectrum of **5i**. One of the two large coupling constants (11.7 Hz) is assigned to the geminal coupling to **H6eq**. This involves that there is at least one trans coupling (11.7 Hz) and a gauche coupling (6.4 Hz) to the vicinal protons H1 and H5, which indicates that these protons are mutually anti to each other. Since these protons are attached to **C1** and **C5** bearing the COOMe and BF_3K functionalities their relative stereochemistry is assigned to be anti.

Potassium Trifluoro(4-Ethoxycarbonyl-2-Butenyl)Borate (5j). This compound was prepared according to the above general procedure B, except that only 2.5 mol% of p-toluenesulfonic acid was employed. The crude product obtained after evaporation of the solvent was extracted with chloroform, and then the chloroform solution was filtrated and evaporated. Recrystallization from acetone/ether gave **5j** as a white solid. The ^1H NMR spectrum of allylboronic acid product **4j** was determined from the crude mixture. ^1H NMR ($\text{DMSO-d}_6/\text{CD}_3\text{OD}$): 5.56 (dt, 15.0 Hz, 7.1 Hz, 1H), 5.33 (dt, 15.0 Hz, 7.1 Hz, 1H), 4.04 (q, 7.1 Hz, 2H), 2.95 (d, 7.1 Hz, 2H), 1.55 (d, 7.1 Hz, 2H), 1.17 (t, 7.1 Hz, 3H). The NMR data for the isolated compound **5j**: ^1H NMR (CDCl_3): 5.71 (dt, 15.2 Hz, 7.1 Hz, 1H), 5.22 (dt, 15.2 Hz, 7.1 Hz, 1H), 4.08 (q, 7.1 Hz, 2H), 2.96 (d, 7.1 Hz, 2H), 1.21 (m, 5H). ^{13}C NMR (CDCl_3): 175.0, 138.1, 118.1, 61.0, 38.0, 24.0 (br), 14.4. ^{19}F NMR (CDCl_3): -140.9.

Competitive Boronation Experiment of 4a and Cinnamyl Acetate (Figure 1). In an NMR tube cinnamyl alcohol (**3a**) or cinnamyl acetate (0.15 mmol) was dissolved in a mixture of DMSO-d_6 and CD_3OD (0.3/0.1 ml) followed by addition of diboronic acid **2** (0.18 mmol) and catalyst **1** (0.0075 mmol, 5 mol%). The reaction vessel was placed in an NMR spectrometer probe for 11 hours at 55°C ; and the progress of the reaction was monitored by ^1H -NMR spectroscopy. Using the above reaction conditions the catalytic effects of strong acids were also studied. In this experiment catalytic amount of p-toluenesulfonic acid (0.0075 mmol, 5 mol%) was also added to the reaction mixture after addition of **1** and **2**.

References:

- (1) Yao, Q.; Kinney, E. P.; Zheng, C. *Org. Lett.* **2004**, *6*, 2997.
- (2) (a) Bäckvall, J. E.; Styrbjörn, B. E.; Nordberg, R. E.; *J. Org. Chem.* **1984**, *49*, 4619. (b) Marshall, J.A.; Xie, S.; *J. Org. Chem.* **1995**, *60*, 7230. (c) Ling, T.; Chowdhury, C.; Kramer, B.A.; Vong, B.G.; Palladino, M.A.; Theodorakis, E.A.; *J. Org. Chem.* **2001**, *66*, 8843.