S-1
SUPPORTING INFORMATION PART 1
for

Direct electrophilic α -fluorination of imines: efficient
synthesis of mono- and difluoroimines

Guido Verniest, Eva Van Hende, Riccardo Surmont and Norbert De Kimpe^{*}

*Department of Organic Chemistry, Faculty of Bioscience Engineering,
Ghent University, Coupure Links 653, 9000 Ghent, Belgium*

TABLE OF CONTENTS:

I. General methods	(S2)
II. Experimental procedures and spectroscopic data (¹ H NMR, ¹³ C NMR, ¹⁹ F NMR, IR, MS, HRMS and elemental analysis) of all new compounds 2e , 2g , 2i , 3a , 3c-g , 5a-c , 6b , 9a-e and 10a-e	(S2)
III. Copies of ¹ H NMR and ¹³ C NMR spectra of 2e , 2g , 2i , 3a-c , 3f , 3g , 4a-e , 5a-c , 6b , 8a-c , 9a-e , 10a-e	(S14)

^{*} Corresponding author. Tel.: +32 (0)9 264 59 51; fax: +32 (0)9 264 62 43; e-mail: norbert.dekimpe@UGent.be.

I. General methods

^1H NMR (300 MHz), ^{13}C NMR (75 MHz) and ^{19}F NMR (282MHz) spectra were run with a Jeol JNM-EX 270 NMR spectrometer or with a Jeol Eclipse FT 300 NMR spectrometer. The compounds were diluted in CDCl_3 with tetramethylsilane (TMS, $\delta=0$ ppm) as a reference. ^{19}F NMR spectra were recorded using CDCl_3 as a lock solvent. IR spectra were obtained from a Perkin Elmer Spectrum One infrared spectrometer. Mass spectra were recorded on a Varian MAT 112 spectrometer (70eV), using either GC-MS coupling or a direct inlet system. Some volatile compounds were recorded on a HP 6890 GC coupled with a HP 5973 MSD (Mass selective detector). LC-MS was performed with an Agilent 1100 Series VS (ES, 4000V) mass spectrometer. HRMS analyses were performed using a Finnigan MAT 95 XP-API-GC Trap tandem mass spectrometer system. Melting points of crystalline compounds were measured with a Büchi 540 apparatus. The elemental analysis of several new compounds was performed on a Perkin Elmer 2400 Elemental Analyser.

II. Synthetic procedures and spectral data

1. Synthesis of imines **2** and **6**

N*-(1-(4-Chlorophenyl)ethylidene)isopropylamine **2e*. The synthesis of imine **2e** is given as a representative example for the synthesis of both imines **2** and **6** which are used as starting material for further fluorination reactions. Spectral data of known imines **2a-d**, **2f** and **6a,b** were in accordance with literature data.¹¹ To a cooled (0°C) solution of 10.00 g of 4'-chloroacetophenone (64.72 mmol) and 15.27 g of isopropylamine (0.26 mol, 4 equiv.) in 250 ml of dry diethyl ether is added 7.36 g of titanium(IV) chloride (38.83 mmol, 0.6 equiv.) in 10 ml of dry pentane. The mixture is stirred for 15 h allowing the temperature to rise to room temperature. After completion of the reaction, the mixture is poured in a biphasic mixture of 400 ml of aqueous 0.5M NaOH solution and 100 ml of diethyl ether. After separation of the organic phase, the aqueous phase is again extracted with 200 ml diethyl ether (2 times). The combined organic phases are dried over a mixture of MgSO_4 and K_2CO_3 (10:1). Evaporation of the solvent, the crude imine **2e** was distilled. Bp 85-90°C (1 mmHg), yield 80%. *E/Z* ratio 92:8. ^1H NMR (CDCl_3): $\delta_{\text{E-isomer}}$ 1.21 (6H, d, $J = 6.3\text{Hz}$, 2 x CH_3), 2.22 (3H, s, CH_3), 3.83 (1H, sept, $J = 6.3\text{Hz}$, CHN), 7.30-7.72 (4H, m, 4 x CH). $\delta_{\text{Z-isomer}}$ 1.07 (6H, d, $J = 6.3\text{Hz}$, 2 x CH_3), 2.24 (3H, s, CH_3), 3.38 (1H, sept, $J = 6.3\text{Hz}$, CHN), 7.02-7.40 (4H, m, 4 x CH). ^{13}C

NMR (CDCl₃): $\delta_{E\text{-isomer}}$ 14.3 (CH₃), 23.2 (2 x CH₃), 50.8 (CHN), 127.6 (2 x CH_{ar}), 127.8 (2 x CH_{ar}), 134.7 (C_{quat}), 139.6 (CCl), 160.3 (C=N). $\delta_{Z\text{-isomer}}$ 1.07 (6H, d, J = 6.3Hz, 2 x CH₃), 2.24 (3H, s, CH₃), 3.38 (1H, sept, J = CHN), 7.02-7.40 (4H, m, 4 x CH). ¹³C NMR (CDCl₃): $\delta_{Z\text{-isomer}}$ 13.9 (CH₃), 23.5 (2 x CH₃), 51.9 (CHN), 126.8 (2 x CH_{ar}), 128.4 (2 x CH_{ar}), 133.4 (C_{quat}), 137.4 (CCl), 164.2 (C=N). IR (NaCl): ν_{max} 1633 cm⁻¹. MS (ES+) *m/z* (%): 196 (M+H⁺, 100). Copies of ¹H NMR and ¹³C NMR spectra are included in chapter III of the supporting information file.

N-((1E)-3,4-Dihydro-1(2H)-naphthalenyldene)isopropylamine 2g. Bp 133-138°C (11 mmHg), yield 79%. *E/Z* ratio 100:0. ¹H NMR (CDCl₃): δ 1.20 (6H, d, J = 6.2Hz, 2 x CH₃), 1.93 (2H, quint, J = 6.3Hz, CH₂), 2.60 (2H, t, J = 6.3Hz, CH₂), 2.81 (2H, t, J = 6.3Hz, CH₂), 3.89 (1H, sept, J = 6.2Hz, CHN), 7.09-7.12 (1H, m, CH), 7.18-7.30 (2H, m, 2 x CH), 8.19 (1H, dd, J = 7.7Hz, 1.7Hz, CH). ¹³C NMR (CDCl₃): δ 23.1 (CH₂), 23.9 (2 x CH₃), 27.2 (CH₂), 30.0 (CH₂), 49.9 (CHN), 126.0 (CH_{ar}), 126.4 (CH_{ar}), 128.4 (CH_{ar}), 129.5 (CH_{ar}), 135.4 (C_{quat}), 140.3 (C_{quat}), 161.4 (C=N). IR (NaCl): ν_{max} 1628, 1597, 1453, 1297 cm⁻¹. MS *m/z* (%): 187 (M⁺, 41), 172 (M⁺-CH₃, 100), 117 (43). Copies of ¹H NMR and ¹³C NMR spectra are included in chapter III of the supporting information file.

N-(1-Phenylpentylidene)isopropylamine 2i. Bp 80°C (1 mmHg), yield 94%. *E:Z* ratio 59:41 (determined by ¹H NMR). ¹H NMR (300MHz, CDCl₃): $\delta_{E\text{-isomer}}$ 0.90 (3H, t, J=7.2Hz, Me), 1.22 (6H, d, J=6.1Hz, 2xMe), 1.25-1.50 (4H, m, 2xCH₂), 2.64-2.71 (2H, m, CH₂C=N), 3.89 (1H, sept, J=6.1Hz, CHMe₂), 7.32-7.42 (3H, m, 3xCH_{ar}), 7.66-7.71 (2H, m, 2xCH_{ar,ortho}). $\delta_{Z\text{-isomer}}$ 0.87 (3H, t, J=7.2Hz, Me), 1.06 (6H, d, J=6.1Hz, 2xMe), 1.25-1.50 (4H, m, 2xCH₂), 2.46-2.53 (2H, m, CH₂C=N), 3.37 (1H, sept, J=6.1Hz, CHMe₂), 7.04-7.09 (2H, m, 2xCH_{ar,ortho}), 7.32-7.42 (3H, m, 3xCH_{ar}). ¹³C NMR (75MHz, CDCl₃): $\delta_{E\text{-isomer}}$ 13.8 (Me), 23.0 (CH₂Me), 24.0 (2xMe), 28.6 (CH₂C=N), 29.9 (CH₂CH₂Me), 50.7 (CHMe₂), 127.0 (2xCH_{ar,ortho}), 128.2 (2xCH_{ar,meta}), 128.9 (CH_{ar,para}), 141.0 (C_{ar,quat}), 166.5 (C=N). $\delta_{Z\text{-isomer}}$ 13.9 (Me), 22.6 (CH₂Me), 23.9 (2xMe), 28.7 (CH₂CH₂Me), 42.3 (CH₂C=N), 52.0 (CHMe₂), 126.2 (2xCH_{ar,ortho}), 127.3 (CH_{ar,para}), 128.3 (2xCH_{ar,meta}), 139.1 (C_{ar,quat}), 169.6 (C=N). IR (NaCl): ν_{max} 1627 cm⁻¹. MS (ES+): *m/z*(%): 204 (M+H⁺, 100). Copies of ¹H NMR and ¹³C NMR spectra are included in chapter III of the supporting information file.

N-(1-Butylpentylidene)isopropylamine 6b. Bp 45-47°C (0.4 mmHg), yield (91%). ¹H NMR (CDCl₃): δ 0.91 (3H, t, J = 6.9Hz, CH₃), 0.93 (3H, t, J = 6.9Hz, CH₃), 1.09 (6H, d, J = 6.3Hz, 2 x CH₃), 1.27-1.53 (8H, m, 4 x CH₂), 2.81 (2H, t, J = 7.8Hz, CH_{2a}C=N), 2.82 (2H, t, J = 7.8Hz, CH_{2b}C=N), 3.68 (1H, sept, J = 6.3Hz, CHN). ¹³C NMR (CDCl₃): δ 13.8 (CH₃),

14.0 (CH₃), 22.7 (CH₂), 23.1 (CH₂), 23.9 (2 x CH₃), 29.4 (CH₂), 29.6 (CH₂), 30.2 (CH₂), 40.3 (CH₂), 49.6 (CHN), 170.4 (C=N). IR (NaCl): $\nu_{\max}$ 1655, 1466 cm⁻¹. MS m/z (%): 183 (M⁺, 2), 168 (2), 154 (18), 99 (86), 84 (100). Copies of ¹H NMR and ¹³C NMR spectra are included in chapter III of the supporting information file.

2. α -Monofluorination of imines **2**

***N*-(2-Fluoro-1-phenylethylidene)isopropylamine **3a**.** In a flame dried 100 ml flask, a heterogeneous mixture of 2.35 g (7.45 mmol, 1.2 equiv.) of NFSI (*N*-fluorobenzenesulfonimide) and 1.72 g (12.42 mmol, 2 equiv.) of K₂CO₃ and 8 g of 4Å molecular sieves (dried at 120°C) in 50 ml of acetonitrile and 10 ml of DMF was vigorously stirred under N₂ atmosphere for 15 minutes at 0°C. At the same temperature, 1.00 g (6.21 mmol) of freshly distilled *N*-(1-phenylethylidene)isopropylamine **2a** was added dropwise via a syringe. The mixture was stirred for 3 hours at 0°C. Subsequently, an excess of triethylamine (5 ml) was added at 0°C and stirring was continued for 2 minutes. The reaction mixture was filtered over Celite and the solids were washed 3 times with 20 ml of diethyl ether. Subsequently, the filtrate was poured in 100 ml of aqueous 0.5M NaOH. After separation of the organic layer, the aqueous phase was extracted with 3 x 25 ml of diethyl ether. To remove the DMF, the combined organic phases were washed with a saturated aqueous NaCl-solution (2 x 100 ml) and subsequently dried with MgSO₄/K₂CO₃ (10:1). Filtration of the drying agents and evaporation of the solvents in vacuo yielded virtually pure monofluorinated imine **3a** (containing only traces of difluorinated imine **9a**). Although the crude product is pure enough to be used in further reactions, a further purification could be established by column chromatography (short column) using dry solvents and silica that was dried at high vacuum (0.01 mmHg) at 100°C for 5h and treated with Et₃N directly after cooling to room temperature. Flash chromatography (hexane/EtOAc 9:1, R_f = 0.37), yield 62%. *E/Z* ratio 62:38. ¹H NMR (CDCl₃): $\delta_{E\text{-isomer}}$ 1.12 (6H, d, J = 6.2Hz, 2 x CH₃), 3.57 (1H, sept, J = 6.2Hz, CHN), 5.06 (2H, d, J = 47.3Hz, CH₂F), 7.09-7.90 (5H, m, C₆H₅). $\delta_{Z\text{-isomer}}$ 1.25 (6H, d, J = 6.2Hz, 2 x CH₃), 4.01 (1H, sept x d, J_{HH} = 6.2Hz, J_{HF} = 1.4Hz, CHN), 5.35 (2H, d, J = 47.3Hz, CH₂F), 7.15-7.75 (5H, m, C₆H₅). ¹³C NMR (CDCl₃): $\delta_{E\text{-isomer}}$ 23.7 (2 x CH₃), 52.1 (CHN), 85.7 (d, J = 176.5Hz, CH₂F), 126.8 (2 x CH_{ar}), 128.6 (2 x CH_{ar}), 128.9 (CH_{ar}), 134.69 (C_{quat}), 164.0 (d, J = 18.5Hz, C=N). $\delta_{Z\text{-isomer}}$ 24.2 (2 x CH₃), 51.6 (CHN), 75.7 (d, J = 167.3Hz, CH₂F), 127.3 (2 x CH_{ar}), 128.4 (2 x CH_{ar}), 129.8 (CH_{ar}), 138.5 (C_{quat}), 160.3 (d, J = 13.9Hz, C=N). ¹⁹F NMR (CDCl₃): $\delta_{E\text{-isomer}}$ -220.4 (t, J = 47.3Hz, CH₂F). $\delta_{Z\text{-isomer}}$ -219.0 (t, J = 47.3Hz, CH₂F). IR (NaCl): $\nu_{\max}$ 1646 cm⁻¹. MS (ES⁺) m/z (%): 180 (M+H⁺, 100). HRMS calcd for

$C_{11}H_{15}NF$ ($M+H^+$) 180.11830, found 180.11815. Copies of 1H NMR and ^{13}C NMR spectra are included in chapter III of the supporting information file.

***N*-(2-Fluoro-1-phenylpropylidene)isopropylamine 3b.**¹² Bp 67-74°C (2.0 mmHg), yield 74%. *E/Z* ratio 97:3. Spectroscopic data were in according to literature data.¹² HRMS calcd for $C_{12}H_{17}NF$ ($M+H^+$) 194.13395, found 194.13329. Copies of 1H NMR and ^{13}C NMR spectra are included in chapter III of the supporting information file.

***N*-(2-Fluoro-1-phenylethylidene)butylamine 3c.** Not purified, yield 92% (purity 84%). *E/Z* ratio 67:33. 1H NMR ($CDCl_3$): $\delta_{E-isomer}$ 0.85 (3H, t, $J = 7.4$ Hz, CH_3), 1.28 (2H, sext, $J = 7.4$ Hz, CH_2), 1.60 (2H, quint, $J = 7.4$ Hz, CH_2), 3.33 (2H, tt, $J = 7.4$ Hz, 1.1Hz, NCH), 5.10 (2H, dt, $J = 47.2$ Hz, 1.1Hz, CH_2F), 7.17-7.59 (5H, m, 5 x CH_2). $\delta_{Z-isomer}$ 0.97 (3H, t, $J = 7.4$ Hz, CH_3), 1.45 (2H, sext, $J = 7.4$ Hz, CH_2), 1.74 (2H, quint, $J = 7.4$ Hz, CH_2), 3.64 (2H, td, $J_{HH} = 7.4$ Hz, $J_{HF} = 2.7$ Hz, NCH), 5.37 (2H, d, $J = 47.2$ Hz, CH_2F), 7.17-7.59 (5H, m, 5 x CH_2). ^{13}C NMR ($CDCl_3$): $\delta_{E-isomer}$ 24.4 (2 x CH_2), 25.5 (CH_2), 33.70 (2 x CH_2), 60.6 (NCH), 85.8 (d, $J = 175.4$ Hz, CH_2F), 126.9 (2 x CH_{ar}), 128.7 (2 x CH_{ar}), 128.9 (CH_{ar}) 134.7 (C_{quat}), 164.3 (d, $J = 18.5$ Hz, $C=N$). $\delta_{Z-isomer}$ 24.6 (2 x CH_2), 25.6 (CH_2), 34.1 (2 x CH_2), 60.1 (CHN), 75.6 (d, $J = 175.4$ Hz, CH_2F), 127.3 (2 x CH_{ar}), 128.3 (2 x CH_{ar}), 129.7 (CH_{ar}), 132.4 (C_{quat}), 162.3 (d, $J = 18.5$ Hz, $C=N$). ^{19}F NMR ($CDCl_3$): $\delta_{E-isomer}$ -220.4 (t, $J = 47.2$ Hz, CH_2F). $\delta_{Z-isomer}$ -220.7 (t, $J = 47.2$ Hz, CH_2F). IR (NaCl): ν_{max} 1671 cm^{-1} . MS (ES+) m/z (%): 220 ($M+H^+$, 100). Copies of 1H NMR and ^{13}C NMR spectra are included in chapter III of the supporting information file.

***N*-(2-Fluoro-1-phenylethylidene)cyclohexylamine 3d.** Recrystallization (diethyl ether-petroleum ether), mp 56°C, yield 37%. *E/Z* ratio 74:26. 1H NMR ($CDCl_3$): $\delta_{E-isomer}$ 1.02-1.91 (10H, m, 5 x CH_2), 3.22 (1H, quint, $J = 7.4$ Hz, CHN), 5.07 (2H, d, $J = 47.3$ Hz, CH_2F), 7.15-7.20 (2H, m, 2 x CH_{ar}), 7.38-7.44 (3H, m, 3 x CH_{ar}). $\delta_{Z-isomer}$ 1.02-1.91 (10H, m, 5 x CH_2), 3.59-3.70 (1H, m, CHF), 5.36 (2H, d, $J = 47.3$ Hz, CH_2F), 7.38-7.44 (3H, m, 3 x CH_{ar}), 7.72-7.77 (2H, m, 2 x CH_{ar}). ^{13}C NMR ($CDCl_3$): $\delta_{E-isomer}$ 24.4 (2 x CH_2), 25.5 (CH_2), 33.70 (2 x CH_2), 60.6 (NCH), 85.8 (d, $J = 175.4$ Hz, CH_2F), 126.9 (2 x CH_{ar}), 128.7 (2 x CH_{ar}), 128.9 (CH_{ar}) 134.7 (C_{quat}), 164.3 (d, $J = 18.5$ Hz, $C=N$). $\delta_{Z-isomer}$ 24.6 (2 x CH_2), 25.6 (CH_2), 34.1 (2 x CH_2), 60.1 (CHN), 75.6 (d, $J = 175.4$ Hz, CH_2F), 127.3 (2 x CH_{ar}), 128.3 (2 x CH_{ar}), 129.7 (CH_{ar}), 132.4 (C_{quat}), 162.3 (d, $J = 18.5$ Hz, $C=N$). ^{19}F NMR ($CDCl_3$): $\delta_{E-isomer}$ -220.1 (t, $J = 47.3$ Hz, CH_2F). $\delta_{Z-isomer}$ -218.6 (t, $J = 47.3$ Hz, CH_2F). IR (NaCl): ν_{max} 1671 cm^{-1} . MS (ES+) m/z (%): 220 ($M+H^+$, 100). Anal. Calcd. for $C_{14}H_{18}NF$: C, 76.68; H, 8.27; N, 6.39. Found: C, 76.89; H, 8.41; N, 6.26.

***N*-(1-(4-Chlorophenyl)-2-fluoroethylidene)isopropylamine 3e.** Flash chromatography (hexane/EtOAc 9:1, $R_f = 0.29$), yield 68%. *E/Z* ratio 56:44. 1H NMR

(CDCl₃): $\delta_{E\text{-isomer}}$ 1.11 (6H, d, $J = 6.1\text{Hz}$, 2 x CH₃), 3.54 (1H, sept, $J = 6.1\text{Hz}$, CHN), 5.05 (2H, d, $J = 47.3\text{Hz}$, CH₂F), 7.12 (2H, d, $J = 8.5\text{Hz}$, 2 x CH_{ar}), 7.42 (2H, d, $J = 8.5\text{Hz}$, 2 x CH_{ar}). $\delta_{Z\text{-isomer}}$ 1.25 (6H, d, $J = 6.1\text{Hz}$, 2 x CH₃), 3.99 (1H, sept x d, $J_{\text{HH}} = 6.1\text{Hz}$, $J_{\text{HF}} = 1.3\text{Hz}$, CHN), 5.34 (2H, d, $J = 47.3\text{Hz}$, CH₂F), 7.36 (2H, d, $J = 8.8\text{Hz}$, 2 x CH_{ar}), 7.69 (2H, d, $J = 8.8\text{Hz}$, 2 x CH_{ar}). ¹³C NMR (CDCl₃): $\delta_{E\text{-isomer}}$ 23.7 (2 x CH₃), 52.4 (CHN), 85.7 (d, $J = 175.4\text{Hz}$, CH₂F), 128.4 (2 x CH_{ar}), 129.0 (2 x CH_{ar}), 133.0 (C_{quat}), 135.0 (CCl), 163.0 (d, $J = 18.4\text{Hz}$, C=N). $\delta_{Z\text{-isomer}}$ 24.1 (2 x CH₃), 51.6 (CHN), 75.3 (d, $J = 170.7\text{Hz}$, CH₂F), 128.5 (2 x CH_{ar}), 128.7 (2 x CH_{ar}), 135.9 (C_{quat}), 136.8 (CCl), 159.3 (d, $J = 15.0\text{Hz}$, C=N). ¹⁹F NMR (CDCl₃): $\delta_{E\text{-isomer}}$ -220.5 (t, $J = 47.3\text{Hz}$, CH₂F). $\delta_{Z\text{-isomer}}$ -219.5 (t, $J = 47.3\text{Hz}$, CH₂F). IR (NaCl): ν_{max} 1648, 1634, 1593, 1489, 1093 cm⁻¹. MS m/z (%): 213/15 (M⁺, 6), 198/200 (M⁺-CH₃, 10), 180/82 (30), 138/40(100). Anal. Calcd. for C₁₁H₁₃NCIF: C, 61.83; H, 6.13; N, 6.56. Found: C, 61.67; H, 6.30; N, 6.40.

***N*-(2-Fluoro-1-phenylbutylidene)cyclohexylamine 3f.** Bp 96°C (0.02 mmHg), yield 71%. *E/Z* ratio 100:0. ¹H NMR (CDCl₃): δ 0.91-1.72 (10H, m, 5 x xCH₂), 1.00 (3H, t, $J = 7.6\text{Hz}$, CH₃), 1.74-1.84 (2H, m, CH₂), 3.09 (1H, quint, $J = 7.3\text{Hz}$, NCH), 5.01 (1H, dt, $J_{\text{HF}} = 48.6\text{Hz}$, $J_{\text{HH}} = 6.3\text{Hz}$, CHF), 7.10-7.13 (2H, m, 2 x CH_{ar}), 7.33-7.42 (3H, m, 3 x CH_{ar}). ¹³C NMR (CDCl₃): δ 9.2 (d, $J = 5.3\text{Hz}$, CH₃), 24.4 (2 x CH₂), 25.6 (CH₂), 26.0 (CH₂), 26.3 (CH₂), 33.8 (d, $J = 13.8\text{Hz}$, CH₂CHF), 60.7 (NCH), 97.6 (d, $J = 175.0\text{Hz}$, CHF), 127.2 (2 x CH_{ar}), 128.4 (2 x CH_{ar}), 128.5 (CH_{ar}), 134.8 (C_{quat}), 167.0 (d, $J = 22.3\text{Hz}$, C=N). ¹⁹F NMR (CDCl₃): δ -183.8 (dt, $J = 48.6\text{Hz}$, 21.7Hz, CHF). IR (NaCl): ν_{max} 1647, 1449 cm⁻¹. MS m/z (%): 247 (M⁺, 4), 186 (33), 104 (100). HRMS calcd for C₁₆H₂₃NF (M+H⁺) 247.17308, found 247.17317. Copies of ¹H NMR and ¹³C NMR spectra are included in chapter III of the supporting information file.

***N*-((1*E*)-2-Fluoro-3,4-dihydro-1(2*H*)-naphthalenyldene)isopropylamine 3g.** Bp 94°C (0.3 mmHg), yield 71%. *E/Z* ratio 0:100. ¹H NMR (CDCl₃): δ 1.22 (3H, d, $J = 6.2\text{Hz}$, CH₃), 1.28 (3H, d, $J = 6.2\text{Hz}$, CH₃), 1.87-2.12 (1H, m, CH_aH_bCHF), 2.38-2.51 (1H, m, CH_aH_bCHF), 2.71 (1H, dt, $J = 15.8\text{Hz}$, 4.1Hz, CH_aH_bCH₂CHF), 3.09 (1H, ddd, $J = 15.8\text{Hz}$, 12.1Hz, 4.1Hz, CH_aH_bCH₂CHF), 4.19 (1H, sept x d, $J_{\text{HH}} = 6.2\text{Hz}$, $J_{\text{HF}} = 2.4\text{Hz}$, CHN), 5.69 (1H, dt, $J_{\text{HF}} = 43.8\text{Hz}$, $J_{\text{HH}} = 3.5\text{Hz}$, CHF), 7.14 (1H, d, $J = 7.4\text{Hz}$, CH_{ar}), 7.22-7.33 (2H, m, 2 x CH_{ar}), 8.18 (1H, dd, 7.6Hz, 1.5Hz, CH_{ar}). ¹³C NMR (CDCl₃): δ 23.9 (d, $J = 4.6\text{Hz}$, CH₂), 24.4 (CH₃), 24.5 (CH₃), 29.5 (d, $J = 20.8\text{Hz}$, CH₂), 51.1 (NCH), 81.2 (d, $J = 171.9\text{Hz}$, CHF), 126.6 (CH_{ar}), 126.8 (CH_{ar}), 128.2 (CH_{ar}), 130.0 (CH_{ar}), 133.4 (C_{quat}), 139.4 (C_{quat}), 156.5 (d, $J = 13.8\text{Hz}$, C=N). ¹⁹F NMR (CDCl₃): δ -183.2 (dt, $J = 43.8\text{Hz}$, 7.0Hz, CHF). IR (NaCl): ν_{max} 1629, 1599, 1455 cm⁻¹. MS m/z (%): 205 (M⁺, 18), 190 (77), 117 (100). HRMS calcd for

$C_{13}H_{17}NF$ ($M+H^+$) 206.13395, found 206.13411. Copies of 1H NMR and ^{13}C NMR spectra are included in chapter III of the supporting information file.

4. Synthesis of α -fluoroketones 4

2-Fluoroacetophenone 4a.¹⁴ To a solution of 0.5 g of *N*-(2-fluoro-1-phenylethylidene)isopropylamine **3a** in 20 ml of dichloromethane was added 20 ml of aqueous 2M HCl. The mixture was stirred at room temperature for 2 hours and subsequently, the organic layer was separated. After extraction of the aqueous phase with CH_2Cl_2 (2 x 25 ml), the organic phase was dried ($MgSO_4$), filtered and the solvent evaporated, yielding almost pure 2-fluoroacetophenone **4a**. A further purification was possible via chromatography over a very short plug of silica gel. Flash chromatography (hexane/EtOAc 9:1, R_f = 0.52) yielded 91% of 2-fluoroacetophenone. Spectral data were in accordance with literature data.¹⁴ Copies of 1H NMR and ^{13}C NMR spectra as an indication of purity are included in chapter III of the supporting information file.

2-Fluoropropiophenone 4b.⁸ Flash chromatography (hexane/EtOAc 9:1, R_f = 0.39), yield 85%. Spectral data were in accordance with literature data.⁸ Copies of 1H NMR and ^{13}C NMR spectra as an indication of purity are included in chapter III of the supporting information file.

1-(4-Chlorophenyl)-2-fluoroethanone 4c.¹⁴ Flash chromatography (hexane/EtOAc 9:1, R_f = 0.53), yield 94%. Spectral data were in accordance with literature data.¹⁴ Copies of 1H NMR and ^{13}C NMR spectra as an indication of purity are included in chapter III of the supporting information file.

2-Fluorobutyrophenone 4d.¹⁵ Flash chromatography (hexane/EtOAc 9:1, R_f = 0.40), yield 90%. Spectral data were in accordance with literature data.¹⁵ Copies of 1H NMR and ^{13}C NMR spectra as an indication of purity are included in chapter III of the supporting information file.

2-Fluoro- α -tetralone 4e.⁶ⁱ Flash chromatography (hexane/EtOAc 9:1, R_f = 0.43), yield 86%. Spectral data were in accordance with literature data.⁶ⁱ Copies of 1H NMR and ^{13}C NMR spectra as an indication of purity are included in chapter III of the supporting information file.

5. Synthesis of β -fluorinated amines 5

***N*-(2-Fluoro-1-phenylethyl)-*N*-isopropylamine 5a.** To a solution of 0.5 g (2.79 mmol) of imine **3a** in 25 ml of absolute methanol was added 0.19 g (3.07 mmol, 1.1 equiv.) of

acetic acid and 0.18 g (3.07 mmol, 1.1 equiv.) of sodium cyanoborohydride at 0°C. The mixture was stirred for 15 h while the temperature reached room temperature. After completion of the reaction, the mixture was poured in 50 ml of a saturated aqueous NaHCO₃ solution and was subsequently extracted with diethyl ether (3 x 50 ml). The combined organic layers were washed with brine and dried over MgSO₄. Filtration of the solids and evaporation of the solvent yielded almost pure *N*-(2-fluoro-1-phenylethyl)-*N*-isopropylamine **5a** which could be further purified by flash chromatography (hexane/EtOAc 98:2, R_f = 0.08), yield 72%. ¹H NMR (CDCl₃): δ 1.01 (3H, d, J = 6.3Hz, Me), 1.04 (3H, d, J = 6.3Hz, Me), 2.69 (1H, sept, J = 6.3Hz, NCH), 4.11 (1H, ddd, J_{HF} = 12.3Hz, J_{HH} = 7.8Hz, J_{HH} = 4.2Hz, NCH), 4.35 (1H, ddd, J_{HF} = 47.3Hz, J_{HH} = 8.9Hz, J_{HH} = 7.8Hz, C(H)HF), 4.48 (1H, ddd, J_{HF} = 47.3Hz, J_{HH} = 8.9Hz, J_{HH} = 4.2Hz, C(H)HF), 7.01-7.63 (5H, m, 5 x CH_{ar}). ¹³C NMR (CDCl₃): δ 22.2 (CH₃), 24.3 (CH₃), 45.9 (NCH), 60.2 (d, J = 18.5Hz, NCH), 87.2 (d, J = 175.4Hz, CH₂F), 127.8 (2 x CH_{ar}), 127.9 (CH_{ar}), 128.7 (2 x CH_{ar}), 139.4 (d, J = 6.9Hz, C_{quat}). ¹⁹F NMR (CDCl₃): δ -219.3 (dt, J = 12.3Hz, 47.3Hz, CHF). IR (NaCl): ν_{max} 3331 cm⁻¹. MS (ES+) *m/z* (%): 182 (M+H⁺, 100). HRMS calcd for C₁₁H₁₇NF (M+H⁺) 182.13395, found 182.13422. Copies of ¹H NMR and ¹³C NMR spectra are included in chapter III of the supporting information file.

***N*-(2-Fluoro-1-(4-chlorophenyl)ethyl)-*N*-isopropylamine 5b.** Flash chromatography (hexane/EtOAc/Et₃N 95:4.9:0.1, R_f = 0.16), yield 68%. ¹H NMR (CDCl₃): δ 1.00 (3H, d, J = 6.3Hz, Me), 1.04 (3H, d, J = 6.3Hz, Me), 1.5 (1H, s(b), NH), 2.66 (1H, sept, J = 6.3Hz, NCH), 4.09 (1H, ddd, J_{HF} = 12.6Hz, J_{HH} = 7.7Hz, J_{HH} = 4.3Hz, NCH), 4.32 (1H, ddd, J_{HF} = 46.9Hz, J_{HH} = 8.9Hz, J_{HH} = 7.7Hz, C(H)HF), 4.42 (1H, ddd, J_{HF} = 46.9Hz, J_{HH} = 8.9Hz, J_{HH} = 4.3Hz, C(H)HF), 7.31 (4H, s, 4 x CH_{ar}). ¹³C NMR (CDCl₃): δ 22.1 (CH₃), 24.3 (CH₃), 45.9 (NCH), 59.7 (d, J = 18.5Hz, NCH), 86.9 (d, J = 175.4Hz, CH₂F), 128.8 (2 x CH_{ar}), 129.1 (2 x CH_{ar}), 133.5 (C_{quat}), 138.1 (d, J = 6.9Hz, C_{quat}). ¹⁹F NMR (CDCl₃): δ -219.6 (dt, J = 12.6Hz, 46.9Hz, CHF). IR (NaCl): ν_{max} 3335, 1596, 1490 cm⁻¹. MS (ES+) *m/z* (%): 216/18 (M+H⁺, 100). HRMS calcd for C₁₁H₁₆N³⁵ClF (M+H⁺) 216.09498, found 216.09481.

***N*-Butyl-*N*-(2-fluoro-1-phenylethyl)amine 5c.** Flash chromatography (hexane/EtOAc/Et₃N 95:4.9:0.1, R_f = 0.18), yield 76%. ¹H NMR (CDCl₃): δ 0.87 (3H, t, J = 7.1Hz, Me), 1.26-1.38 (2H, m, CH₂), 1.46 (2H, quint, J = 7.1Hz, CH₂), 1.70 (1H, s(b), NH), 2.49 (2H, t, J = 7.0Hz, NCH₂), 4.01 (1H, ddd, J_{HF} = 13.0Hz, J_{HH} = 8.8Hz, J_{HH} = 4.2Hz, NCH), 4.37 (1H, dt, J_{HF} = 47.4Hz, J_{HH} = 8.8Hz, C(H)HF), 4.45 (1H, ddd, J_{HF} = 47.4Hz, J_{HH} = 8.8Hz, J_{HH} = 4.2Hz, C(H)HF), 7.26-7.36 (5H, m, 5 x CH_{ar}). ¹³C NMR (CDCl₃): δ 14.1 (CH₃), 20.5 (CH₂), 32.4 (CH₂), 47.3 (NCH₂), 63.2 (d, J = 18.5Hz, NCH), 87.2 (d, J = 174.1Hz, CH₂F),

127.8 (2 x CH_{ar}), 128.0 (CH_{ar}), 128.7 (2 x CH_{ar}), 138.9 (d, J = 8.1Hz, C_{quat}). ¹⁹F NMR (CDCl₃): δ -218.4 (dt, J=13.0Hz, 47.4Hz, CHF). IR (NaCl): ν_{max} 3339, 1454 cm⁻¹. MS (ES+): *m/z* (%): 196 (M+H⁺, 100). HRMS calcd for C₁₂H₁₉NF (M+H⁺) 196.14960, found 196.14986.

6. Synthesis of α-fluoroketones 8

2-Fluorocyclohexanone 8a.^{6m} For the synthetic procedure is referred to the synthesis of fluorinated imines **3** and the hydrolysis to fluorinated ketones **4** (see above). Flash chromatography (hexane/EtOAc 9:1, R_f = 0.13), yield 86%. Spectral data were in accordance with literature data.^{6m} Copies of ¹H NMR and ¹³C NMR spectra as an indication of purity are included in chapter III of the supporting information file.

2-Fluoro-5-nonanone 8b.¹⁷ Flash chromatography (hexane/EtOAc 94:6, R_f = 0.52), yield 65%. Spectral data were in accordance with literature data.¹⁷ Copies of ¹H NMR and ¹³C NMR spectra as an indication of purity are included in chapter III of the supporting information file.

1-Fluoro-1-phenylacetone 8c.¹⁶ Flash chromatography (hexane/EtOAc 9:1, R_f = 0.32), yield 61%. Spectral data were in accordance with literature data.¹⁶ Copies of ¹H NMR and ¹³C NMR spectra as an indication of purity are included in chapter III of the supporting information file.

7. Synthesis of α,α-difluoroimines 9

N-(2,2-Difluoro-1-phenylethylidene)isopropylamine 9a. In a flame dried 100 ml flask, a heterogeneous mixture of 5.87 g (18.63 mmol, 3 equiv.) of NFSI (*N*-fluorobenzenesulfonimide) and 1.72 g (12.42 mmol, 2 equiv.) of K₂CO₃ and 8 g of 4Å molecular sieves (dried at 120°C) in 50 ml of acetonitrile and 10 ml of DMF was vigorously stirred under N₂ atmosphere for 15 minutes at room temperature. Subsequently, 1.00 g (6.21 mmol) of freshly distilled *N*-(1-phenylethylidene)isopropylamine **2a** was added dropwise via a syringe. The mixture was stirred for 15 hours. Subsequently, an excess of triethylamine (5 ml) was added at 0°C and stirring was continued for 2 minutes. The reaction mixture was filtered over Celite and the solids were washed 3 times with 20 ml of diethyl ether. Subsequently, the filtrate was poured in 100 ml of aqueous 0.5M NaOH. After separation of the organic layer, the aqueous phase was extracted with 3 x 25 ml of diethyl ether. To remove the DMF, the combined organic phases were washed with a saturated aqueous NaCl-solution (2 x 100 ml) and subsequently dried with MgSO₄/K₂CO₃ (10:1). Filtration of the drying agents and evaporation of the solvents in vacuo yielded difluoroimine **9a** (together with 18%

of 2-fluoroacetophenone). Other imines **9b-9e** were obtained using an analogous procedure and were obtained without significant amounts of the corresponding monofluoroimines or -ketones. Purification of imines **9b-9e** could be established by column chromatography (short column) using dry solvents and silica that was dried at high vacuum (0.01 mmHg) at 100°C for 5h and treated with Et₃N directly after cooling to room temperature. Yield of **9a** 74%. Mp 41°C. ¹H NMR (CDCl₃): δ 1.12 (6H, d, J=6.2Hz, 2xMe), 3.59 (1H, sept, J=6.2Hz, CHMe₂), 6.13 (1H, t, J=55.4Hz, CHF₂), 7.20-7.25 (2H, m, 2xCH_{ar}), 7.41-7.48 (3H, m, 3xCH_{ar}). ¹³C NMR (CDCl₃): δ 2 x 23.2 (2 x CH₃), 52.3 (CH), 115.4 (t, J = 244.0Hz, CHF₂), 127.6 (2 x CH_{ar}), 128.4 (2 x CH_{ar}), 129.2 (CH_{ar}), 131.2 (C_{quat}), 161.1 (t, J = 27.7Hz, C=N). ¹⁹F NMR (CDCl₃): δ -117.8 (2F, d, J=55.4Hz). IR (KBr) ν_{max} 1660 cm⁻¹. MS (ES+) m/z (%): 198 (M+H⁺, 100). Copies of ¹H NMR and ¹³C NMR spectra are included in chapter III of the supporting information file.

N-(2,2-Difluoro-1-phenylpropylidene)isopropylamine 9b. Flash chromatography (hexane/EtOAc/Et₃N 95:4.8:0.2, R_f = 0.6), yield 54%. ¹H NMR (CDCl₃): δ 1.08 (6H, d, J=6.1Hz, CH(Me)₂), 1.86 (3H, t, J=17.9Hz, CH₃), 3.44 (1H, sept, J=6.1Hz, CH(Me)₂), 7.15-7.18 (2H, m, 2xCH_{ar}), 7.39-7.46 (3H, m, 3xCH_{ar}). ¹³C NMR (CDCl₃): δ 21.6 (t, J=26.5Hz, CH₃), 23.4 (CH(Me)₂), 52.6 (CH(Me)₂), 120.9 (t, J=239.4Hz, CF₂), 127.7 (2xCH_{ar}), 128.4 (2xCH_{ar}), 128.9 (CH_{ar}), 133.1 (C_{ar,quat}), 162.5 (t, J=31.2Hz, C=N). ¹⁹F NMR (CDCl₃): δ -92.4 (2F, q, J=17.9Hz). IR (NaCl): ν_{max} 1657 cm⁻¹. MS (ES+): m/z(%): 212 (M+H⁺, 100). Copies of ¹H NMR and ¹³C NMR spectra are included in chapter III of the supporting information file.

N-(2,2-Difluoro-1-phenylbutylidene)isopropylamine 9c. Flash chromatography (hexane/EtOAc/Et₃N 94:4:2, R_f = 0.3), yield 63%. ¹H NMR (CDCl₃): δ 1.07 (3H, t, J=7.5Hz, CH₂Me), 1.09 (6H, d, J=6.1Hz, CH(Me)₂), 2.18 (2H, tq, J_{HF}=16.7Hz, J_{HH}=7.5Hz, CH₂Me), 3.49 (1H, sept, J=6.1Hz, CH(Me)₂), 7.14-7.17 (2H, m, 2xCH_{ar}), 7.38-7.44 (3H, m, 3xCH_{ar}). ¹³C NMR (CDCl₃, int. ref.=77.13ppm): δ 6.5 (t, J=5.2Hz, CH₂Me), 23.5 (CH(Me)₂), 27.8 (t, J=25.4Hz, CH₂CF₂), 52.7 (CH(Me)₂), 121.3 (t, J=242.9Hz, CF₂), 127.6 (2xCH_{ar}), 128.4 (2xCH_{ar}), 128.9 (CH_{ar}), 133.3 (C_{ar,quat}), 162.2 (t, J=30.0Hz, C=N). ¹⁹F NMR (CDCl₃): δ -101.6 (2F, s(br)). IR (NaCl): ν_{max} 1663 cm⁻¹. MS (ES+): m/z(%): 226 (M+H⁺, 100). Copies of ¹H NMR and ¹³C NMR spectra are included in chapter III of the supporting information file.

N-(2,2-Difluoro-1-phenylbutylidene)cyclohexylamine 9d. Flash chromatography (hexane/EtOAc/Et₃N 95:4.8:0.2, R_f = 0.5), yield 63%. ¹H NMR (CDCl₃): δ 0.82-1.29 (4H, m, 4 x CH_{cyclohexyl}), 1.06 (3H, t, J=7.4Hz, CH₃), 1.49-1.59 (4H, m, 4 x CH_{cyclohexyl}), 1.67-1.73 (2H, m, 2 x CH_{cyclohexyl}), 2.17 (2H, tq, J_{HH}=7.4Hz, J_{HF}=16.7Hz, CH₂), 3.12 (1H, quint,

$J=7.1\text{Hz}$, NCH), 7.13-7.18 (2H, m, $2\times\text{CH}_{\text{ar}}$), 7.38-7.43 (3H, m, $3\times\text{CH}_{\text{ar}}$). ^{13}C NMR (CDCl_3 , int. ref.=77.00ppm): δ 6.4 (t, $J=4.6\text{Hz}$, CH_2Me), 24.0 ($2\times\text{CH}_{2,\text{cHex}}$), 25.5 (CH_2, cHex), 27.7 (t, $J=25.4\text{Hz}$, CH_2CF_2), 32.3 (NCH), 121.2 (t, $J=242.3\text{Hz}$, CF_2), 127.5 ($2\times\text{CH}_{\text{ar}}$), 128.3 ($2\times\text{CH}_{\text{ar}}$), 128.7 (CH_{ar}), 133.2 ($\text{C}_{\text{ar,quat}}$), 162.1 (t, $J=30.6\text{Hz}$, $\text{C}=\text{N}$). ^{19}F NMR (CDCl_3): δ -101.3 (2F, s(br)). IR (NaCl): ν_{max} 1660 cm^{-1} . MS (ES⁺): $m/z(\%)$: 266 ($\text{M}+\text{H}^+$, 100). Copies of ^1H NMR and ^{13}C NMR spectra are included in chapter III of the supporting information file.

***N*-(2,2-Difluoro-1-phenylpentylidene)isopropylamine 9e.** Bp 80°C (1 mmHg), yield: 49%. ^1H NMR (CDCl_3): δ 0.98 (3H, t, $J=7.8\text{Hz}$, Me), 1.09 (6H, d, $J=6.3\text{Hz}$, $2\times\text{Me}$), 1.58 (2H, sext, $J=7.8\text{Hz}$, CH_2Me), 2.13 (2H, tt, $J_{\text{HF}}=17.0\text{Hz}$, $J_{\text{HH}}=7.8\text{Hz}$, CH_2CF_2), 3.44 (1H, sept, $J=6.3\text{Hz}$, CHMe_2), 7.12-7.18 (2H, m, $2\times\text{CH}_{\text{ar}}$), 7.38-7.43 (3H, m, $3\times\text{CH}_{\text{ar}}$). ^{13}C NMR (CDCl_3): δ 14.0 (Me), 15.5 (t, $J=4.6\text{Hz}$, CH_2Me), 23.4 ($2\times\text{Me}$), 36.4 (t, $J=24.2\text{Hz}$, CH_2CF_2), 52.6 (CHMe_2), 120.9 (t, $J=242.9\text{Hz}$, CF_2), 127.5 ($2\times\text{CH}_{\text{ar}}$), 128.3 ($2\times\text{CH}_{\text{ar}}$), 128.8 (CH_{ar}), 133.2 ($\text{C}_{\text{ar,quat}}$), 162.3 (t, $J=30.6\text{Hz}$, $\text{C}=\text{N}$). ^{19}F NMR (CDCl_3): δ -99.8 (2F, t, $J=17.0\text{Hz}$). IR (NaCl): ν_{max} 1657 cm^{-1} . MS (ES⁺): $m/z(\%)$: 240 ($\text{M}+\text{H}^+$, 100). Copies of ^1H NMR and ^{13}C NMR spectra are included in chapter III of the supporting information file.

8. Synthesis of β,β -difluoroamines 10

***N*-(2,2-Difluoro-1-phenylethyl)-*N*-isopropylamine 10a.** For the synthetic procedures of the reductions of α,α -difluoroimines **9** towards amines **10** is referred to the synthesis of amines **5** from α -fluoroimines **3**. Amine **10a** was purified by flash chromatography using a short plug of silicagel (hexane/EtOAc 95:5, $R_f = 0.42$), yield 82%. ^1H NMR (CDCl_3): δ 1.03 (3H, d, $J=6.3\text{Hz}$, Me), 1.04 (3H, d, $J=6.3\text{Hz}$, Me), 1.45 (1H, s(br), NH), 2.69 (1H, sept, $J=6.3\text{Hz}$, $\text{CH}(\text{Me})_2$), 4.03 (1H, ddd, $J_{\text{HF}}=13.3\text{Hz}$, $J_{\text{HF}}=10.6\text{Hz}$, $J_{\text{HH}}=4.4\text{Hz}$, CHCHF_2), 5.79 (1H, dt, $J_{\text{HF}}=56.4\text{Hz}$, $J_{\text{HH}}=4.4\text{Hz}$, CHF_2), 7.25-7.40 (5H, m, $5\times\text{CH}_{\text{ar}}$). ^{13}C NMR (CDCl_3 , int. ref.=77.13ppm): δ 21.9 (Me), 24.1 (Me), 45.5 ($\text{CH}(\text{Me})_2$), 62.3 (t, $J=87.0\text{Hz}$, CCF_2), 117.2 (t, $J=245.2\text{Hz}$, CF_2), 128.5 (CH_{ar}), 128.5 ($2\times\text{CH}_{\text{ar}}$), 128.7 ($2\times\text{CH}_{\text{ar}}$), 136.5 ($\text{C}_{\text{ar,quat}}$). ^{19}F NMR (CDCl_3): δ -123.7 (1F, ddd, $J_{\text{FF}}=277.0\text{Hz}$, $J_{\text{HF}}=56.7\text{Hz}$, $J_{\text{HF}}=10.6\text{Hz}$); -125.44 (1F, ddd, $J_{\text{FF}}=277.0\text{Hz}$, $J_{\text{HF}}=56.4\text{Hz}$, $J_{\text{HF}}=13.3\text{Hz}$). IR (NaCl): ν_{max} 3339 cm^{-1} . MS (ES⁺): $m/z(\%)$: 200 ($\text{M}+\text{H}^+$, 100). Copies of ^1H NMR and ^{13}C NMR spectra are included in chapter III of the supporting information file.

***N*-(2,2-Difluoro-1-phenylpropyl)-*N*-isopropylamine 10b.** Purification by acid-base extraction ($\text{Et}_2\text{O}/\text{HCl}/\text{NaOH}$), yield 90%. ^1H NMR (CDCl_3): δ 1.01 (3H, d, $J = 6.3\text{Hz}$, Me), 1.02 (3H, d, $J = 6.3\text{Hz}$, Me), 1.53 (3H, t, $J = 19.0\text{Hz}$, CH_3CF_2), 2.62 (1H, sept, $J = 6.3\text{Hz}$,

NCH), 3.99 (1H, t, $J = 11.6\text{Hz}$, NCH), 7.26-7.38 (5H, m, 5 x CH_{ar}). ^{13}C NMR (CDCl_3 , int. ref.=77.00ppm): δ 20.8 (t, $J = 27.1\text{Hz}$, CH_3), 21.5 (CH_3), 24.1 (CH_3), 45.4 (NCH), 64.9 (t, $J = 25.4\text{Hz}$, CHCF_2), 124.1 (t, $J = 242.9\text{Hz}$, CF_2), 128.0 (CH_{ar}), 128.2 (2 x CH_{ar}), 128.6 (2 x CH_{ar}), 137.6 (C_{quat}). ^{19}F NMR (CDCl_3): δ -98.0 (s(br), CF_2). IR (NaCl): ν_{max} 3337 cm^{-1} . MS (ES+) m/z (%): 214 ($\text{M}+\text{H}^+$, 100). Copies of ^1H NMR and ^{13}C NMR spectra are included in chapter III of the supporting information file.

***N*-(2,2-Difluoro-1-phenylbutyl)-*N*-isopropylamine 10c.** Flash chromatography (hexane/EtOAc/ Et_3N 94:4:2, $R_f = 0.33$), yield 60%. ^1H NMR (CDCl_3): δ 0.96-1.01 (9H, m, 3xMe), 1.52 (1H, s(br), NH), 1.68-2.04 (2H, m, CH_2Me), 2.60 (1H, sept, $J=6.3\text{Hz}$, $\text{CH}(\text{Me})_2$), 3.99 (1H, d, $J_{\text{HF}}=11.5\text{Hz}$, CHCF_2), 7.24-7.41 (5H, m, 5x CH_{ar}). ^{13}C NMR (CDCl_3 , int. ref.=77.13ppm): δ 6.0 (CH_2Me), 21.6 ($\text{CH}(\text{Me})\text{Me}$), 24.3 ($\text{CH}(\text{Me})\text{Me}$), 27.3 (t, $J=25.4\text{Hz}$, CH_2CF_2), 45.4 ($\text{CH}(\text{Me})_2$), 63.8 (t, $J=25.4\text{Hz}$, CHCF_2), 124.9 (t, $J=245.8\text{Hz}$, CF_2), 128.0 (CH_{ar}), 128.4 (2x CH_{ar}), 128.8 (2x CH_{ar}), 137.7 ($\text{C}_{\text{ar,quat}}$). ^{19}F NMR (CDCl_3): δ -109.06 (2F, s(br), CF_2). IR (NaCl): ν_{max} 3340 cm^{-1} . MS (ES+): m/z (%): 228 ($\text{M}+\text{H}^+$, 100). Copies of ^1H NMR and ^{13}C NMR spectra are included in chapter III of the supporting information file.

***N*-Cyclohexyl-*N*-(2,2-difluoro-1-phenylbutyl)amine 10d.** Purification by acid-base extraction ($\text{Et}_2\text{O}/\text{HCl}/\text{NaOH}$), yield 94%. ^1H NMR (CDCl_3): δ 0.98 (3H, t, $J = 7.6\text{Hz}$, CH_2CH_3), 1.01-1.23 (4H, m, 4 x $\text{CH}_{\text{cyclohexyl}}$), 1.43-2.04 (8H, m, 6 x $\text{CH}_{\text{cyclohexyl}}$ and CH_2CF_2), 2.16-2.30 (1H, m, NCH), 4.05 (1H, t, $J = 12.4\text{Hz}$, CHCF_2), 7.28-7.36 (5H, m, 5 x CH_{ar}). ^{13}C NMR (CDCl_3 , int. ref.=77.00ppm): δ 5.8 (t, $J = 5.2\text{Hz}$, CH_3), 24.4 (CH_2), 24.9 (CH_2), 26.0 (CH_2), 27.0 (t, $J=25.4\text{Hz}$, CH_2CF_2), 32.3 (CH_2), 34.6 (CH_2), 53.0 (NCH), 63.0 (t, $J = 24.8\text{Hz}$, CHCF_2), 124.8 (t, $J = 245.8\text{Hz}$, CF_2), 127.8 (CH_{ar}), 128.2 (2 x CH_{ar}), 128.7 (2 x CH_{ar}), 137.8 ($\text{C}_{\text{ar,quat}}$). ^{19}F NMR (CDCl_3): δ -109.10 (2F, s(br)). IR (NaCl): ν_{max} 3338 cm^{-1} . MS (ES+): m/z (%): 268 ($\text{M}+\text{H}^+$, 100). Copies of ^1H NMR and ^{13}C NMR spectra are included in chapter III of the supporting information file.

***N*-(2,2-Difluoro-1-phenylpentyl)-*N*-isopropylamine 10e.** Yield 81%. ^1H NMR (CDCl_3): δ 0.90 (3H, t, $J=7.4\text{Hz}$, Me), 0.99 (3H, d, $J=6.2\text{Hz}$, Me), 1.00 (3H, d, $J=6.2\text{Hz}$, Me), 1.41-1.60 (2H, m, CH_2Me), 1.63-1.97 (2H, m, CH_2CF_2), 2.59 (1H, sept, $J=6.2\text{Hz}$, CHMe_2), 3.97 (1H, t, $J=12.4\text{Hz}$, CH_{ar}), 7.27-7.39 (5H, m, 5x CH_{ar}). ^{13}C NMR (CDCl_3): δ 13.9 (Me), 15.0 (t, $J=4.6\text{Hz}$, CH_2Me), 21.5 (MeCH), 24.2 (MeCH), 35.9 (t, $J=24.2\text{Hz}$, CH_2CF_2), 45.4 (CHMe_2), 64.1 (t, $J=25.4\text{Hz}$, CH_{ar}), 124.6 (t, $J=245.8\text{Hz}$, CF_2), 127.9 (CH_{ar}), 128.3 (2x CH_{ar}), 128.7 (2x CH_{ar}), 137.6 ($\text{C}_{\text{ar,quat}}$). ^{19}F NMR (CDCl_3): δ -106.9 (2F, m). IR (NaCl): ν_{max} 3339 cm^{-1} . MS (ES+): m/z (%): 241 ($\text{M}+\text{H}^+$, 100). Copies of ^1H NMR and ^{13}C NMR spectra are included in chapter III of the supporting information file.

III. Copies of ^1H NMR and ^{13}C NMR spectra of compounds **2e**, **2g**, **2i**, **3a-c**, **3f**,
3g, **4a-e**, **5a-c**, **6b** and **8a-c**

5a