SUPPORTING INFORMATION

A Direct, One-Step Synthesis of Condensed Heterocycles: A Palladium-Catalyzed Coupling Approach

*Farnaz Jafarpour and Mark Lautens**

Davenport Chemical Research Laboratories, Chemistry Department, University of Toronto

Toronto, Ontario, Canada, M5S 3H6

General remarks: The following includes general experimental procedures, specific details for representative reactions, and isolation and spectroscopic information for new compounds. Anhydrous solvents (DMSO, DMF, CH₂Cl₂, CCl₄) were systematically used. DME and MeCN were distilled under nitrogen from CaH₂ and immediately used or stored under 4 Å molecular sieves. Other reagents, metal catalysts (PdCl₂, Pd(OAc)₂) and ligands (PPh₃, TFP) were commercially available and used as received. The cyclization reactions were carried out in an oil bath using Biotage Microwave Vials (2-5 mL). ¹H and ¹³C NMR spectra were recorded at room temperature on a Mercury 300 MHz or a Mercury 400 MHz spectrometers using CDCl₃ as the NMR solvent. ¹H NMR spectra are referenced to tetramethylsilane (0.00 ppm) and ¹³C NMR spectra are referenced from the solvent central peak (77.23 ppm). Chemical shifts are given in ppm. IR spectra were recorded on a Nicolet DX FT IR spectrometer. IR is reported as characteristic bands (cm⁻¹) in their maximal intensity. The letters br mean the signal is broad. High-Resolution Mass Spectra (HRMS) were obtained using a VG 70-250S (double focusing) mass spectrometer at 70 eV unless otherwise noted.

Table of Contents

Synthesis of Iodoarenes	S2
Annulation Reactions	S3
General Procedure for the reaction of bromoenoates and iodoarenes	S3
General Procedure for the reaction of bromoalkyl indoles and iodoarenes	S6
Synthesis of the Key Intermediate	S7
¹H and ¹³C NMR Spectra	S9

Synthesis of Iodoarenes

Iodoarenes **1a-1b**, **4b** and **6b** synthesis were previously reported.¹ Iodoarenes **4a** and **4b** were synthesized as follows:

1-Iodo-3-(3-bromo-propoxy)-benzene **4a**

4a

3-Iodophenol (1.11 g, 5.0 mmol) was dissolved in acetone (4.5 ml). 1,3-dibromopropane (1.00 ml, 10 mmol, 2 eq.) and K₂CO₃ (1.38 g, 10 mmol, 2 eq.) were added and the mixture was stirred at 50°C for 2 days. The mixture was diluted with diethyl ether, washed with water and dried over MgSO₄. Purification by FC (silica, AcOEt /hexane, 10%) afforded the desired product as a colorless liquid (1.2 g, 3.5 mmol, 70%) ; ¹H NMR (300 MHz, CDCl₃) δ 7.26 (m, 2H), 7.00 (t, *J* = 8.0 Hz, 1H), 6.86 (dd, *J* = 8.4, 2.4 Hz, 1H), 4.07 (t, *J* = 6.0 Hz, 2H), 3.58 (t, *J* = 6.3 Hz, 2H), 2.31 (quintet, *J* = 6.3 Hz, 2H) ; ¹³C NMR (75 MHz, CDCl₃) δ 159.4, 131.0, 130.3, 123.9, 114.4, 94.6, 65.7, 32.4, 30.0; IR 2930, 2877, 1583, 1464, 1284, 1240, 1028, 989, 766 cm⁻¹ ; HRMS (EI, *m/z*) calcd. for C₉H₁₀OBrI (M⁺) 339.8960, Found 339.8963.

1-Iodo-3-(4-bromo-butoxy)-benzene **6a**

6a

3-iodophenol (1.11 g, 5.0 mmol) was dissolved in acetone (4.5 ml). 1,4-Dibromobutane (1.32 ml, 10 mmol, 2 eq.) and K₂CO₃ (1.38 g, 10 mmol, 2 eq.) were added and the mixture was stirred at 50°C for 2 days. The mixture was diluted with diethyl ether, washed with water and dried over MgSO₄. Purification by FC (silica, AcOEt /hexane, 10%) afforded the desired product as a colorless liquid (1.3 g, 3.7 mmol, 72%) ; ¹H NMR (300 MHz, CDCl₃) δ 7.24 (m, 2H), 6.971 (t, *J* = 8.0 Hz, 1H), 6.83 (m, 1H), 3.93 (t, *J* = 6.4 Hz, 2H), 3.46 (t, *J* = 6.4 Hz, 2H), 2.03 (m, 2H), 1.91 (m, 2H) ; ¹³C NMR (75 MHz, CDCl₃) δ 159.5, 130.9, 130, 123.7, 114.3, 94.6, 67.1, 33.5, 29.5, 27.9; IR 2939, 2867, 1583, 1465, 1284, 1225, 1038, 989, 765 cm⁻¹ ; HRMS (EI, *m/z*) calcd. for C₁₀H₁₂OBrI (M⁺) 353.9116, Found 353.9112

¹ Pache, S.; Lautens, M. *Org. Lett.* **2003**, 5, 4827-4830.

Synthesis of Bifunctional Acceptors

Bromoenoates **2**,² **8**,³ **17**,⁴ the amide **12**,⁵ and bromoalkyl indoles⁶ synthesis were previously reported.

Annulation Reactions

General Procedure for the reaction of bromoenoates and iodoarenes

A vial equipped with a stir bar was charged with iodoarene (0.20 mmol, 1 eq.), bromoenoate (1 mmol, 5 eq.), norbornene (1 mmol, 5 eq.), Cs₂CO₃ (1.2 mmol, 6 eq.), Pd(OAc)₂ (0.02 mmol, 10 mol%) and PPh₃ (0.04 mmol, 20 mol%) and 2 mL of degassed dry DME was added and the vial was capped. The resulting mixture was heated in an oil bath at 100 °C for 48 h, cooled then filtered through a short plug of silica. Removal of the solvent gave a crude mixture which was purified by flash column chromatography (hexane/CH₂Cl₂ gradient).

Ethyl-2-(1,2,7,8-tetrahydronaphtho[2,1-b]furan-9(6H)-ylidene) acetate **3**

Colourless oil (42 mg, 80%) ; R_f = 0.18 on silica gel (hexane/CH₂Cl₂ 1:1) ; ¹H NMR (300 MHz, CDCl₃) δ 6.94 (d, *J* = 8.1 Hz, 1H), 6.76 (d, *J* = 8.1 Hz, 1H), 6.04 (m, 1H), 4.53 (t, *J* = 8.4, 2H), 4.2 (q, *J* = 7.2 Hz, 2H), 3.38 (t, *J* = 8.4 Hz, 2H), 3.15 (dt, *J* = 6.5, 1.8 Hz, 2H), 2.7 (t, *J* = 6.5 Hz, 2H), 1.81 (quintet, *J* = 6.5 Hz, 2H), 1.31 (t, *J* = 7.2 Hz, 3H) ; ¹³C NMR (100 MHz, CDCl₃) δ 167.3, 159.4, 155.1, 133.3, 132.7, 128.5, 124.4, 116.0, 110.7, 71.4, 60.0, 32.3, 30.1, 28.9, 23.1, 14.6 ; IR 2927, 2846, 1712, 1614, 1462, 1248, 1159 cm⁻¹ ; HRMS (EI, *m/z*) calcd. for C₁₆H₁₈O₃ (M⁺) 258.1256, Found 258.1255.

Ethyl-2-(2,3,8,9-tetrahydro-1H-benzo[f]chromen-10(7H)-ylidene) acetate **5**

² (a) Vedejs, E.; Arnost, M. J.; Hagen, J. P. *J. Org. Chem.* **1979**, *44*, 3230-3238. (b) Amputch, M. A.; Matamoros, R.; Little, R. D. *Tetrahedron* **1994**, *50*, 5591-5614.

³ Somekawa, K.; Okuhira, H.; Sendayama, M.; Suishu, T.; Shimo, T. *J. Org. Chem.* **1992**, *57*, 5708-5712.

⁴ Lautens, M.; Paquin, J.-F.; Piguel S. *J. Org. Chem.* **2002**, *67*, 3972-3974.

⁵ Lautens, M.; Paquin, J.-F.; Piguel S.; Dahlmann, M. *J. Org. Chem.* **2001**, *66*, 8127-8134.

⁶ Bressy, C.; Alberico, D.; Lautens, M. *J. Am. Chem. Soc.* **2005**, *127*, 13148-13149.

Colourless oil (27 mg, 51%) ; R_f = 0.22 on silica gel (hexane/ CH_2Cl_2 1:1) ; ^1H NMR (400 MHz, CDCl_3) δ 6.90 (d, J = 8.2 Hz, 1H), 6.72 (d, J = 8.2 Hz, 1H), 5.92 (m, 1H), 4.19 (m, 4H), 3.11 (dt, J = 6.8, 1.6 Hz, 2H), 2.90 (t, J = 6.0 Hz, 2H), 2.55 (t, J = 6.2, 2H), 1.92 (quintet, J = 5.4 Hz, 2H), 1.75 (quintet, J = 6.4 Hz, 2H), 1.31 (t, J = 7.2 Hz, 3H) ; ^{13}C NMR (100 MHz, CDCl_3) δ 167.0, 154.5, 153.8, 137.0, 133.8, 126.8, 119.4, 118.4, 117.4, 66.3, 60.0, 29.7, 28.4, 25.8, 23.0, 22.1, 14.6 ; IR 2938, 2867, 1710, 1615, 1476, 1155 cm^{-1} ; HRMS (EI, m/z) calcd. for $\text{C}_{17}\text{H}_{20}\text{O}_3$ (M^+) 272.1412, Found 272.1410.

Ethyl-2-(1,2,3,4,9,10-hexahydronaphtho[2,1-b]oxepin-11(8H)-ylidene) acetate **7**

Colourless oil (25 mg, 44%) ; R_f = 0.23 on silica gel (hexane/ CH_2Cl_2 1:1) ; ^1H NMR (400 MHz, CDCl_3) δ 6.89 (s, 2H), 5.76 (t, J = 1.8 Hz, 1H), 4.20 (q, J = 7.2 Hz, 2H), 4.03 (t, J = 5.0 Hz, 2H), 3.14 (dt, J = 7.2, 1.6 Hz, 2H), 2.96 (t, J = 5.4 Hz, 2H), 2.55 (t, J = 6.2, 2H), 1.99 (quintet, J = 5.4 Hz, 2H), 1.77 (quintet, J = 6.4 Hz, 4H), 1.31 (t, J = 7.2 Hz, 3H) ; ^{13}C NMR (100 MHz, CDCl_3) δ 166.9, 160.4, 155.1, 138.1, 137.2, 133.3, 125.9, 121.1, 119.2, 74.1, 60.0, 32.3, 30.1, 29.7, 28.3, 26.2, 21.9, 14.6 ; IR 2924, 2853, 1715, 1614, 1470, 1163 cm^{-1} ; HRMS (EI, m/z) calcd. for $\text{C}_{18}\text{H}_{22}\text{O}_3$ (M^+) 286.1569, Found 286.1570.

Tricyclic compound **9**

Colourless oil (31 mg, 58%) ; R_f = 0.28 on silica gel (hexane/ CH_2Cl_2 1:1) ; ^1H NMR (400 MHz, CDCl_3) δ 6.87 (d, J = 8.0 Hz, 1H), 6.60 (d, J = 8.0 Hz, 1H), 5.78 (s, 1H), 4.53 (t, J = 8.4 Hz, 2H), 4.21 (q, J = 7.2 Hz, 2H), 3.16 (t, J = 8.6 Hz, 2H), 2.91 (br, 2H), 2.64 (t, J = 5.8 Hz, 2H), 1.83 (m, 2H), 1.71 (m, 2H), 1.31 (t, J = 7.2 Hz, 3H) ; ^{13}C NMR (100 MHz, CDCl_3) δ 166.61, 162.20, 158.77, 140.32, 131.77, 128.61, 124.02, 118.21, 108.25, 71.47, 60.06, 34.17, 31.91, 29.89, 29.69, 28.25, 14.51 ; IR 2927, 2854, 1713, 1634, 1455, 1172 cm^{-1} ; HRMS (EI, m/z) calcd. for $\text{C}_{17}\text{H}_{20}\text{O}_3$ (M^+) 272.1412, Found 272.1411.

Ethyl 2-(2,3,7,8,9,10-hexahydrocyclohepta[f]chromen-11(1H)-ylidene)acetate **10**

Colourless oil (25 mg, 45%) ; R_f = 0.23 on silica gel (hexane/ CH_2Cl_2 1:1) ; ^1H NMR (300 MHz, CDCl_3) δ 6.86 (d, J = 8.4 Hz, 1H), 6.63 (d, J = 8.4 Hz, 1H), 5.71 (s, 1H), 4.21 (m, 3H), 4.02 (m, 1H), 3.66 (m, 1H), 2.83 (m, 1H), 2.68 (m, 1H), 2.50 (m, 2H), 1.94 (m, 6H), 1.49 (m, 1H), 1.32 (t, J = 7.0 Hz, 3H) ; ^{13}C NMR (100 MHz, CDCl_3) δ 166.6, 162.3, 153.6, 144.1, 131.3, 127.7, 118.8, 118.0, 115.6, 66.1, 60.1, 34.6, 32.1, 30.3, 28.5, 24.0, 22.8, 14.5 ; IR 2929, 1715, 1636, 1476, 1172 cm^{-1} ; HRMS (EI, m/z) calcd. for $\text{C}_{18}\text{H}_{22}\text{O}_3$ (M^+) 286.1569, Found 286.1573.

Tricyclic compound 11

Colourless oil (21 mg, 35%) ; R_f = 0.24 on silica gel (hexane/ CH_2Cl_2 1:1) ; ^1H NMR (400 MHz, CDCl_3) δ 6.84 (q, J = 8.0 Hz, 2H), 5.63 (s, 1H), 4.20 (m, 3H), 3.84 (m, 1H), 3.66 (m, 1H), 2.79 (m, 2H), 2.69 (m, 1H), 2.58 (m, 1H), 1.94 (m, 6H), 1.73 (m, 1H), 1.54 (m, 2H), 1.32 (t, J = 7.2, 3H) ; ^{13}C NMR (75 MHz, CDCl_3) δ 166.6, 163.2, 159.6, 144.1, 134.8, 132.0, 127.2, 120.0, 118.8, 73.9, 60.1, 34.8, 32.4, 29.7, 29.5, 28.2, 26.3, 14.5 ; IR 2926, 2853, 1714, 1630, 1470, 1172 cm^{-1} ; HRMS (EI, m/z) calcd. for $\text{C}_{19}\text{H}_{24}\text{O}_3$ (M^+) 300.1725, Found 300.1723.

2-(1,2,7,8-Tetrahydronaphtho[2,1-b]furan-9(6H)-ylidene)-1-(pyrrolidin-1-yl)ethanone 13

Colourless oil (28 mg, 50%) ; R_f = 0.57 on silica gel (hexane/ CH_2Cl_2 1:1) ; ^1H NMR (300 MHz, CDCl_3) δ 6.93 (d, J = 8.1 Hz, 1H), 6.73 (d, J = 8.1 Hz, 1H), 6.187 (s, 1H), 4.53 (t, J = 8.4 Hz, 2H), 3.56 (t, J = 6.6 Hz, 2H), 3.47 (t, J = 6.6 Hz, 2H), 3.36 (t, J = 8.4 Hz, 2H), 3.00 (dt, J = 6.6, 1.6 Hz, 2H), 2.74 (t, J = 6.3 Hz, 2H), 1.90 (m, 6H) ; ^{13}C NMR (100 MHz, CDCl_3) δ 166.7, 159.2, 148.1, 133.2, 132.5, 128.7, 123.6, 118.8, 110.0, 71.3, 47.3, 45.7, 32.3, 30.0, 28.7, 26.4, 24.6, 23.5 ; IR 2932, 2871, 1634, 1615, 1463, 1418, 1247 cm^{-1} ; HRMS (EI, m/z) calcd. for $\text{C}_{18}\text{H}_{22}\text{NO}_2$ ($\text{M}^+ + 1$) 284.1645, Found 284.1652.

Ethyl 2-(1,2,6,7-tetrahydrobenzofuro[4,5-d]oxepin-10(9H)-ylidene)acetate **18**

Colourless oil (24 mg, 45%) ; R_f = 0.24 on silica gel (hexane/ CH_2Cl_2 1:1) ; ^1H NMR (400 MHz, CDCl_3) δ 6.92 (d, J = 8.0 Hz, 1H), 6.73 (d, J = 8.0 Hz, 1H), 5.81 (s, 1H), 4.92 (s, 2H), 4.55 (t, J = 8.6 Hz, 2H), 4.20 (q, J = 7.2 Hz, 2H), 3.79 (t, J = 6.4 Hz, 2H), 3.26 (t, J = 8.4 Hz, 2H), 2.82 (t, J = 6.4 Hz, 2H), 1.32 (t, J = 7.2, 3H) ; ^{13}C NMR (75 MHz, CDCl_3) δ 1166.2, 162.8, 160.0, 136.1, 129.0, 127.8, 125.5, 117.3, 109.8, 71.6, 68.8, 68.7, 60.4, 33.1, 30.1, 14.5 ; IR 2955, 1706, 1620, 1465, 1242, 1178 cm^{-1} ; HRMS (EI, m/z) calcd. for $\text{C}_{16}\text{H}_{19}\text{O}_4$ ($\text{M}^+ + 1$) 275.1277, Found 275.1287.

General Procedure for the reaction of bromoalkyl indoles and iodoarenes

A vial equipped with a stir bar was charged with aryl iodide (0.200 mmol, 1.0 equiv), tri-2-furylphosphine (0.044 mmol, 22 mol %), norbornene (0.400 mmol, 2.0 equiv), Cs_2CO_3 (0.400 mmol, 2.0 equiv) and $\text{Pd}(\text{OAc})_2$ (0.020 mmol, 10 mol %). A solution of bromoalkyl indole (0.400 mmol, 2.0 equiv) in CH_3CN (2 mL) was then added and the vial was capped. The resulting mixture was heated in an oil bath at 90 °C for 24 h, cooled then filtered through a short plug of silica. Removal of the solvent gave a crude mixture which was purified by flash column chromatography (hexane/ CH_2Cl_2 gradient).

Annulated Indole **20**

Colourless oil (17 mg, 33%) ; R_f = 0.32 on silica gel (hexane/ CH_2Cl_2 1:1) ; ^1H NMR (400 MHz, CDCl_3) δ 7.65 (d, J = 8.0 Hz, 1H), 7.36 (d, J = 8.4 Hz, 1H), 7.23 (d, J = 8.4 Hz, 1H), 7.11 (t, J = 8.0 Hz, 1H), 7.04 (d, J = 8.0 Hz, 1H), 6.71 (s, 1H), 6.70 (d, J = 8.4 Hz, 1H), 4.70 (t, J = 8.6 Hz, 2H), 4.25 (t, J = 6.4 Hz, 2H), 3.49 (t, J = 8.6 Hz, 2H), 3.15 (t, J = 6.4 Hz, 2H) ; ^{13}C NMR (100 MHz, CDCl_3) δ 159.9, 136.3, 134.6, 128.8, 127.9, 126.1, 124.9, 123.3, 122.3, 121.0, 120.1, 109.2, 108.2, 100.0, 71.5, 40.9, 31.0, 29.2 ; IR 2914, 2846, 1592, 1481, 1460, 1242, 1160 cm^{-1} ; HRMS (EI, m/z) calcd. for $\text{C}_{18}\text{H}_{16}\text{NO}$ ($\text{M}^+ + 1$) 262.1226, Found 272.1237.

Annulated Indole 22

Colourless oil (34 mg, 62%) ; $R_f = 0.36$ on silica gel (hexane/ CH_2Cl_2 1:1) ; ^1H NMR (400 MHz, CDCl_3) δ 7.66 (d, $J = 8.0$ Hz, 1H), 7.37 (d, $J = 8.4$ Hz, 1H), 7.22 (d, $J = 8.0$ Hz, 1H), 7.12 (dt, $J = 7.5, 0.8$ Hz, 1H), 7.04 (d, $J = 8.0$ Hz, 1H), 6.74 (d, $J = 8.0$ Hz, 1H), 6.57 (s, 1H), 4.61 (t, $J = 8.8$ Hz, 2H), 4.10 (t, $J = 6.6$ Hz, 2H), 3.36 (t, $J = 8.6$ Hz, 2H), 2.62 (t, $J = 7.0$ Hz, 2H), 2.23 (quintet, $J = 6.8$ Hz, 2H) ; ^{13}C NMR (100 MHz, CDCl_3) δ 159.4, 138.8, 136.3, 130.6, 130.1, 129.0, 128.0, 126.4, 121.8, 120.9, 119.6, 108.9, 108.8, 101.2, 71.7, 41.0, 31.3, 30.6, 30.5 ; IR 2946, 2866, 1588, 1510, 1454, 1313, 1238, 1003 cm^{-1} ; HRMS (EI, m/z) calcd. for $\text{C}_{19}\text{H}_{17}\text{NO}$ (M^+) 275.1310, Found 275.1304.

Annulated Indole 24

Colourless oil (32 mg, 52%) ; $R_f = 0.45$ on silica gel (hexane/ CH_2Cl_2 1:1) ; ^1H NMR (300 MHz, CDCl_3) δ 7.25 (d, $J = 9.0$ Hz, 1H), 7.12 (d, $J = 2.4$ Hz, 1H), 7.03 (d, $J = 8.1$ Hz, 1H), 6.90 (dd, $J = 9.0, 2.4$ Hz, 1H), 6.73 (d, $J = 7.8$ Hz, 1H), 6.49 (s, 1H), 4.60 (t, $J = 8.7$ Hz, 2H), 4.05 (t, $J = 6.6$ Hz, 2H), 3.87 (s, 3H), 3.35 (t, $J = 8.7$ Hz, 2H), 2.61 (t, $J = 7.2$ Hz, 2H), 2.22 (quintet, $J = 6.7$ Hz, 2H) ; ^{13}C NMR (100 MHz, CDCl_3) δ 159.2, 154.0, 139.0, 131.5, 130.3, 129.9, 128.8, 128.0, 126.1, 112.0, 109.4, 108.5, 102.2, 100.5, 71.5, 56.0, 40.9, 31.2, 30.4, 30.3 ; IR 2939, 1593, 1487, 1455, 1238, 1216 cm^{-1} ; HRMS (EI, m/z) calcd. for $\text{C}_{20}\text{H}_{19}\text{NO}_2$ (M^+) 305.1388, Found 275.1380.

Synthesis of the Key Intermediate

Ethyl 2-(1,2,6,7-tetrahydronaphtho[2,1-b]furan-9-yl)acetate 15

Tetrahydronaphtho[2,1-b]furan **3** (50 mg, 0.2 mmol) was diluted with toluene (2 mL). 4-Toluenesulphonic acid (2 mg, 0.012 mmol, 0.06 eq.) was added and the mixture was stirred

at 110°C for 24 hrs. The mixture was cooled to room temperature, evaporated to dryness, diluted with ethyl acetate and washed successively with 10% hydrochloric acid, 2N aqueous sodium carbonate solution and brine. The organic solution was dried over MgSO₄. Purification by FC (silica, hexane/AcOEt 9:1) afforded the desired product. Colourless liquid (26 mg, 50%) ; R_f = 0.29 on silica gel (hexane/AcOEt 9:1) ; ¹H NMR (300 MHz, CDCl₃) δ 6.91 (d, *J* = 8.1 Hz, 1H), 6.60 (d, *J* = 8.1 Hz, 1H), 6.06 (t, *J* = 4.8 Hz, 1H), 4.50 (t, *J* = 8.7 Hz, 2H), 4.14 (q, *J* = 7.0 Hz, 2H), 3.48 (s, 2H), 3.33 (t, *J* = 8.7 Hz, 2H), 2.66 (t, *J* = 7.6, 2H), 2.21 (m, 2H), 1.22 (t, *J* = 7.0, 3H) ; ¹³C NMR (100 MHz, CDCl₃) δ 172.1, 159.7, 132.1, 131.9, 130.8, 129.7, 127.3, 121.7, 107.7, 70.9, 60.9, 40.8, 31.1, 28.6, 23.8, 14.4 ; IR 2934, 1732, 1580, 1457, 1242 cm⁻¹ ; HRMS (EI, *m/z*) calcd. for C₁₆H₁₈O₃ (M⁺) 258.1256, Found 258.1260.

4a

¹HNMR (300 MHz, CDCl₃)

4a

^{13}C NMR (75 MHz, CDCl_3)

6a

¹HNMR (300 MHz, CDCl₃)

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (100 MHz, CDCl_3)

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (100 MHz, CDCl_3)

^1H NMR (400 MHz, CDCl_3)

¹³C NMR (100 MHz, CDCl₃)

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (100 MHz, CDCl_3)

¹H NMR (300 MHz, CDCl₃)

^{13}C NMR (100 MHz, CDCl_3)

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (100 MHz, CDCl_3)

13

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (100 MHz, CDCl_3)

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (100 MHz, CDCl_3)

¹H NMR (400 MHz, CDCl₃)

^{13}C NMR (75 MHz, CDCl_3)

20

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (100 MHz, CDCl_3)

22

^1H NMR (400 MHz, CDCl_3)

^{13}C NMR (100 MHz, CDCl_3)

^1H NMR (300 MHz, CDCl_3)

^{13}C NMR (100 MHz, CDCl_3)