

## Direct Photocatalyzed Hydrogen Atom Transfer (HAT) for Aliphatic C–H Bonds Elaboration

Luca Capaldo, Davide Ravelli, and Maurizio Fagnoni\*



Cite This: *Chem. Rev.* 2022, 122, 1875–1924




Read Online

ACCESS |

Metrics & More

Article Recommendations

**ABSTRACT:** Direct photocatalyzed hydrogen atom transfer (*d*-HAT) can be considered a method of choice for the elaboration of aliphatic C–H bonds. In this manifold, a photocatalyst (PC<sub>HAT</sub>) exploits the energy of a photon to trigger the homolytic cleavage of such bonds in organic compounds. Selective C–H bond elaboration may be achieved by a judicious choice of the hydrogen abstractor (key parameters are the electronic character and the molecular structure), as well as reaction additives. Different are the classes of PC<sub>S<sub>HAT</sub></sub> available, including aromatic ketones, xanthene dyes (Eosin Y), polyoxometalates, uranyl salts, a metal-oxo porphyrin and a tris(amino)cyclopropenium radical dication. The processes (mainly C–C bond formation) are in most cases carried out under mild conditions with the help of visible light. The aim of this review is to offer a comprehensive survey of the synthetic applications of photocatalyzed *d*-HAT.



### CONTENTS

1. Introduction	1875
2. Formation of C–C Bonds	1881
2.1. Formation of C(sp <sup>3</sup> )–C(sp <sup>3</sup> ) Bonds via Addition onto C=C Bonds	1881
2.1.1. Oxygen- and Sulfur-Containing Compounds as Hydrogen Donors	1881
2.1.2. Nitrogen-Containing Compounds as Hydrogen Donors	1883
2.1.3. Hydrocarbons as Hydrogen Donors	1886
2.2. Formation of C(sp <sup>3</sup> )–C(sp <sup>3</sup> ) Bonds via Addition onto C=X (X = N, O) Bonds	1888
2.3. Formation of C(sp <sup>3</sup> )–C(sp <sup>2</sup> ) Bonds	1889
2.4. Formation of C(sp <sup>3</sup> )–C(sp) Bonds	1892
2.5. Formation of C(sp <sup>2</sup> )–C(sp <sup>3</sup> ) Bonds	1892
2.6. Formation of C(sp <sup>2</sup> )–C(sp <sup>2</sup> ) Bonds	1894
2.7. Formation of C(sp <sup>2</sup> )–C(sp) Bonds	1896
2.8. Functionalization of Carbon Nanostructures	1896
3. Formation of C–Y Bonds (Y ≠ C)	1896
3.1. Formation of C–N Bonds	1897
3.2. Formation of C–O Bonds	1898
3.2.1. Synthesis of Hydroperoxides	1899
3.2.2. Preparation of Alcohol and Ketone Mixtures	1899
3.2.3. Synthesis of Carbonyl Derivatives	1899
3.2.4. Synthesis of Carboxylic Acids and Esters	1901
3.3. Formation of C–S Bonds	1901
3.4. Formation of C–F Bonds	1903
3.4.1. Fluorination of C(sp <sup>3</sup> )–H Bonds	1903

3.4.2. Fluorination of C(sp <sup>2</sup> )–H Bonds	1905
3.5. Formation of C–Cl Bonds	1905
3.6. Formation of C–D Bonds	1906
4. Formation of Other Bonds	1906
4.1. Formation of P–C Bonds	1906
4.2. Formation of Si–X Bonds (X = Cl or C)	1906
4.3. Formation of S–C Bonds	1907
5. Miscellanea	1907
6. Conclusions	1908
Author Information	1909
Corresponding Author	1909
Authors	1909
Notes	1909
Biographies	1909
Acknowledgments	1909
Abbreviations	1909
References	1910
Note Added in Proof	1924
Note Added after ASAP Publication	1924

**Special Issue:** Photochemical Catalytic Processes

**Received:** March 30, 2021

**Published:** August 6, 2021




## 1. INTRODUCTION

The selective manipulation of C–H bonds (especially C(sp<sup>3</sup>)–H bonds) represents a remarkable challenge in synthetic campaigns because organic molecules contain many of these bonds of different nature. The acidity of hydrogens in the proximity of electron-withdrawing groups has been largely exploited for the smooth generation of enolates, versatile nucleophiles to forge C–C bonds.<sup>1,2</sup> Apart from this fundamental reactivity pattern, aliphatic C–H bonds have been referred to as “unfunctional groups”<sup>3</sup> due to their lack of reactivity and the more articulated strategies needed for their functionalization.<sup>4</sup> Accordingly, the vast majority of C–H bond activation strategies in organic synthesis rely on the use of activating or directing groups, either to enable a particular reaction pathway or to improve selectivity as well as efficiency.<sup>5</sup> Even though the use of temporary directing groups, that is functions that are reversibly bound to the substrate to drive selectivity, has been proposed,<sup>6</sup> the direct aliphatic C–H bond elaboration in organic molecules still remains the unfound Holy Grail in chemistry.<sup>3,7–12</sup> Notably, this is an intense area of research, because it is a godsend for late-stage functionalization<sup>13–17</sup> and in function-oriented synthesis<sup>18</sup> thanks to the innate atom-economy related to the direct elaboration of C–H bonds. Moreover, the selective activation of these bonds in structurally complex molecules is of immense value in medicinal chemistry,<sup>14</sup> where small changes in a given structure may have a profound impact on its biological activity and in natural product synthesis.<sup>19,20</sup>

To address this challenging task, different metal-based strategies<sup>4,19,21–24</sup> have been devised: in particular, Fe-,<sup>25,26</sup> Cu-,<sup>27</sup> Mn-,<sup>28–30</sup> Co-,<sup>31,32</sup> Rh-,<sup>33</sup> Ir-,<sup>34,35</sup> Ru-,<sup>36–38</sup> and Pd-based<sup>39–43</sup> catalysts have been successfully tested. Within this frame, one of the most appealing concepts consists in the homolytic cleavage of the C–H bond via a hydrogen atom transfer (HAT) event.<sup>44–46</sup> This consists in the concerted movement of an electron and a proton ( $H^\bullet \equiv H^+ + e^-$ ) from the substrate, *aka* hydrogen donor, to an accepting species (a hydrogen abstractor); all in a single kinetic step (Scheme 1).

**Scheme 1. Homolytic Cleavage of a C–H Bond via a Hydrogen Atom Transfer Step**



The factors that rule this chemistry (and in general the approaches devoted to design a *selective* HAT) may be tentatively classified as depending on the substrate or on the hydrogen abstractor structures as well as “medium dependent”, as summarized in Scheme 2.

One of the main effects belonging to the former class is the bond dissociation energy (BDE) of the C–H bond to be cleaved. In other words, the lower the BDE and the more stable the generated radical, the easier the bond to break (Scheme 3). However, this is just a rule of thumb and applies only under certain conditions. More often, other factors must be carefully evaluated to account for the difference in selectivity observed in the derivatization of a certain substrate (Scheme 2).<sup>47–52</sup>


Taking the case of hydrocarbons as reference, the relative stability of the generated organoradical drives the cleavage of a tertiary C–H bond (e.g., in compound I<sub>a</sub>) over that of a secondary (as in I<sub>b</sub>) or a primary C–H bond (in I<sub>c</sub>). Also, hyperconjugation and conjugation play an important role in steering the selectivity of HAT. For example, hyperconjugative effects may operate in the case of a hydrocarbon containing a cyclopropyl ring (II). In this case, the overlap between the C–C  $\sigma$  bonding orbital of the three membered ring and the neighboring C–H  $\sigma^*$  antibonding orbital directs the C–H activation to the vicinal position (Scheme 2).<sup>56–59</sup> In some instances, the strain release connected with the cleavage of a C–H bond is another driving force for a selective process. Indeed, this factor is rather significant for six-membered carbocyclic structures (see the case of compound III). In substituted cyclohexanes, the hydrogen abstraction from the equatorial C–H bond that is adjacent to a bulkier axial group leads to the release of 1,3-diaxial strain thanks to the planarization of the incipient carbon centered radical in the HAT transition state.<sup>56,60–76</sup>

While depending also on the nature/identity of the hydrogen abstractor, steric hindrance plays an important role as well, as known for the reactivity of *tert*-butylcyclohexane IV.<sup>77</sup> This is another nice example of a “substrate-dependent” effect where the geometry of the molecule forces the activation toward precise reaction sites (e.g., positions 3 and 4) due to simple steric/torsional effects.<sup>75,77</sup>


Turning to substrates containing heteroatoms (e.g., O, N, etc.), the presence of polar functionalities may influence neighboring C–H bonds and their reactivity. Notably, the polarity match (or mismatch) between the character of the C–H bond to be cleaved and the hydrogen abstractor is another factor that hugely affects HAT. For example, an electrophilic species (e.g., an alkoxy radical or a related derivative) strongly prefers to abstract an hydridic (nucleophilic) C–H bond rather than an electron-poor (electrophilic) one of similar strength, a feature that is directly linked to the electronic character of the accessible radical intermediate.<sup>78</sup> This allows the use of solvents having (rather labile) electrophilic C–H bonds (e.g., acetonitrile, acetone, Scheme 3), when an electrophilic hydrogen abstractor operates in the elaboration of nucleophilic C–H bonds.<sup>45,79–82</sup>

The polarity of C–H bonds in the substrate is thus influenced by the presence of electron-withdrawing or electron-donating groups.<sup>56,60–66</sup> This is apparent in the case of cyclopentanone V, where the more labile electrophilic  $\alpha$ -hydrogens are not activated by an electrophilic hydrogen abstractor since a partial positive charge on the incipient C-centered radical makes the corresponding transition state unfavorable. As a result, the  $\beta$ -C–H cleavage occurs instead.<sup>83</sup>

Another common “substrate-depending” effect governing HAT is that exerted by electron-donating functionalities, notably oxygen and nitrogen atoms. The donation of the nonbonding electrons by these atoms activates vicinal C–H bonds through hyperconjugation. Typical is the case of ethers (e.g., VI), acetals, alcohols, amines, and amides, where the heteroatom causes the decrease of the BDE values of the vicinal C–H bond via hyperconjugation and stabilizes the corresponding radical intermediate (see Scheme 2). Furthermore, the presence of heteroatoms may influence the selectivity via stereoelectronic effects,<sup>56,60–66,85</sup> which allows rationalization of the different reactivities of open chain vs cyclic derivatives. By considering the cleavage of the  $\alpha$ -to-N

Scheme 2. Factors Affecting C–H Bond Cleavage<sup>a</sup>

<sup>a</sup>Different factors operating in the selective HAT-based C–H functionalization in organic compounds. In violet the reactive site where the hydrogen is preferentially cleaved during the functionalization (for the explanation of the effects, see the text).

Scheme 3. Bond Dissociation Energies (BDEs) in kcal/mol of Representative Compounds<sup>a</sup>

<sup>a</sup>Bond dissociation energies (BDEs) in kcal/mol of the X–H bonds (in red) in representative compounds. Values taken from ref 53 except where otherwise noted. <sup>b</sup>Value taken from ref 54. <sup>c</sup>Value taken from ref 55.

C–H bond in amines VII<sub>a</sub> and VII<sub>b</sub> as a representative example, the hydrogen abstraction in VII<sub>a</sub> is more effective than in VII<sub>b</sub>. In fact, the process is more efficient when the bond being broken can be eclipsed with the heteroatom lone pair, not a favorable situation in VII<sub>b</sub> due to the rigidity of the molecular scaffold.

Alternatively, “medium-dependent” effects (Scheme 2) can tune the reactivity pattern in chemical transformations occurring via HAT, again altering the reactivity of substrates containing heteroatoms through the activation or deactivation of the α-C–H bonds. Indeed, the solvent itself may function as hydrogen bond donor or acceptor due to its acidity or basicity;<sup>86–90</sup> albeit, the presence of additives with peculiar acid/base properties may have a similar role.<sup>91–107</sup> The activation effect is well illustrated by the selective C–H functionalization occurring in alcohol VIII<sub>a</sub>. Tetrabutylammonium dihydrogen phosphate forms a complex with the substrate via hydrogen bonding with the alcoholic O–H bond, thus increasing the n–σ\* delocalization of the oxygen lone pair and making the α-to-O C–H bond more prone to a HAT process.<sup>91</sup> On the other hand, a deactivation effect can be

induced by an acidic solvent (e.g., a fluorinated alcohol) or by the addition of a Lewis or a Bronsted acid.<sup>56,60–66,108–111</sup> Accordingly, both in the protonated form (IX<sub>a</sub>) and in the complexed form (IX<sub>c</sub>) the α-to-N C–H bond of triethylamine (IX<sub>b</sub>) is less prone to be cleaved due to the reduced availability of the nitrogen lone pair, and in some instances the selectivity is shifted to the β- (or, in general terms, remote) C–H bonds.<sup>112</sup>


Quite recently, supramolecular chemistry has been exploited to induce chemoselectivity in HAT-based processes.<sup>113–115</sup> The ammonium salt X is functionalized by a Mn-oxo complex (Cat1) preferentially at C<sub>6</sub> and C<sub>7</sub>. By employing a more sophisticated catalyst (Cat2) bearing two crown ether moieties able to complex the ammonium salt, it was possible to shift the functionalization toward C<sub>8</sub> and C<sub>9</sub>.

Upon suitable conditions, a moiety embedded in the molecular scaffold can be activated and triggers the hydrogen abstraction at a specific site in an intramolecular fashion thus inducing a remote activation of a C–H bond (*r*-HAT).<sup>41,116–123</sup> Typically, such site-selectivity is granted by the formation of a favorable six-membered cyclic transition

state, which results in the occurrence of a 1,5-HAT step, despite the fact that the 1,*n*-HAT mode ( $n \geq 6$ ) may compete in some cases.<sup>124</sup>

Given the above, a synthetic route based on HAT has to be judiciously planned, starting from the choice of the proper hydrogen abstractor  $X^\bullet$  (Scheme 4).<sup>125</sup> Thermodynamics-wise,

#### Scheme 4. Common Hydrogen Abstractors Used in Synthetic Planning



the newly formed X–H bond has to be stronger than the C–H bond to cleave to provide the driving force for the overall process, despite BDE not being the only parameter to be considered. As shown in Scheme 3, the BDE of  $\alpha$ -to-heteroatom C–H bonds is mostly comprised between 85 and 95 kcal/mol, while primary and secondary C–H bonds in aliphatic hydrocarbons are quite strong (ca. 100 kcal/mol) calling for a highly reactive species ( $X^\bullet$ ) to promote the cleavage event. Different hydrogen abstractors are known to have a radical or a radical-like character, including alkoxy,<sup>123,126–128</sup> aminoxyl,<sup>62,129</sup> amidyl,<sup>130,131</sup> and sulfonamidyl,<sup>132</sup> azidyl,<sup>133</sup> iodanyl,<sup>134</sup> thiyl,<sup>135–139</sup> and even C-centered<sup>117,140–143</sup> radicals or halogen atoms,<sup>144,145</sup> amine radical cations,<sup>91,118,146–148</sup> N-ammonium ylides,<sup>149</sup> dioxiranes,<sup>59,70,150,151</sup> or metal–oxo complexes.<sup>29,152</sup> These hydrogen abstractors may be thermally or photochemically generated.

Recently, photocatalysis has emerged as a powerful synthetic platform in organic chemistry because it allows taming the tremendous amount of energy associated with light to build molecular complexity. It relies on the use of chemical species, namely photocatalysts (PCs), that can convert light into chemical energy for substrate activation.<sup>153–181</sup>

This methodology has been used to trigger HAT and, in particular, all the reports that appeared so far can be classified


in two approaches: indirect hydrogen atom transfer (*i*-HAT) and direct hydrogen atom transfer (*d*-HAT, Scheme 5).

In the former case, the PC ( $PC_{SET}$ ) takes care of absorbing light and, once in the excited state, generates the hydrogen abstractor ( $X^\bullet$ , a radical or radical ion species) via a single-electron transfer (SET) step (Scheme 5, left).<sup>182–185</sup> In the *d*-HAT process, the PC ( $PC_{HAT}$ ) triggers directly the HAT when in the excited state (Scheme 5, right).<sup>182–186</sup> In other words,  $PC_{HAT}^*$  coincides with  $X^\bullet$ .


A common structural motif to the vast majority of  $PC_{SHAT}$  currently known is the presence of an oxo group ( $Z=O$ ), which acquires a peculiar O-centered radical character in the reactive excited state. The structure of the excited  $PC_{HAT}^*$  strictly resembles electrophilic alkoxy radicals (Scheme 5) behaving as excellent hydrogen abstractors to cleave a C–H bond in the chosen substrate. This leads to the formation of the (protonated) reduced form of the PC ( $PC^{\bullet}H$ ). At each catalytic cycle, the spent PC must be recovered back to its original state, so that it can promote over and over again the process, according to the definition of “photocatalyst” offered by the IUPAC: “Catalyst able to produce, upon absorption of light, chemical transformations of the reaction partners. The excited state of the photocatalyst repeatedly interacts with the reaction partners forming reaction intermediates and regenerates itself after each cycle of such interactions”.<sup>187</sup> The actual mechanism of the PC restoration depends on the synthetic application and can involve a back-HAT step or a sequential electron/proton transfer (ET/PT) mechanism toward a chemical species ( $Y$ , Scheme 5) present in the reaction mixture (e.g., a sacrificial hydrogen acceptor) or transiently formed during the process, also dictating the overall redox balance of the synthetic transformation.<sup>188</sup>

Depending on the X-element carrying the oxo moiety,  $PC_{SHAT}$  can be grouped within different families (Figure 1). These comprise the class of carbonyl derivatives ( $Z = C$ ),<sup>189,190</sup> encompassing simple (aromatic) ketones and aldehydes,<sup>191–193</sup>  $\alpha$ -diketones,<sup>194</sup>  $\alpha$ -ketoacids,<sup>195</sup> and (anthra)-quinones,<sup>196–198</sup> as well as the xanthene dye Eosin Y.<sup>199,200</sup> On the other hand, inorganic derivatives including the decatungstate anion  $[W_{10}O_{32}]^{4-}$  ( $Z = W$ )<sup>64,201–206</sup> and the uranyl cation  $[UO_2]^{2+}$  ( $Z = U$ )<sup>207,208</sup> as well as antimony oxo porphyrin complexes ( $Z = Sb$ )<sup>209</sup> have been likewise proposed as  $PC_{SHAT}$ . A notable exception of  $PC_{HAT}$  lacking the oxo group is known, namely the electrogenerated tris(amino)-cyclopropenium (TAC) radical dication.<sup>210</sup>

#### Scheme 5. Photocatalyzed Indirect Hydrogen Atom Transfer (*i*-HAT) vs Direct Hydrogen Atom Transfer (*d*-HAT)







**Figure 1.** Main photocatalysts ( $PC_{S_{HAT}}$ ) used in photocatalyzed HAT: BP, benzophenone; ABP, aminobenzophenone; DMBP, 4,4'-dimethoxybenzophenone; BPSS, disodium benzophenonedisulfonate; DCBP, 4,4'-dichlorobenzophenone; AP, acetophenone; FL, 9-fluorenone; DBS, dibenzosuberone; DTX, 3,6-dimethoxy-9H-thioxanthen-9-one; PQ, 9,10-phenanthroquinone; XA, xanthone; TX, thioxanthone; AQ, anthraquinone; CIAQ, 2-chloroanthraquinone; tBAQ, 2-*tert*-butylanthraquinone; AQ-2-COOH, anthraquinone-2-carboxylic acid; AQ-2-SO<sub>3</sub>Na, anthraquinone-2-sulfonic acid sodium salt; PT, 5,7,12,14-pentacenetetrone; AQ-2,3-diCOOH, anthraquinone-2,3-dicarboxylic acid; PGA, phenylglyoxylic acid; PYD, 1,6-pyrenedione; EY, Eosin Y; DT, decatungstate anion (TBADT, tetrabutylammonium decatungstate; NaDT, sodium decatungstate); Ur, uranyl cation (UrN, uranyl nitrate hexahydrate; UrP, uranyl perchlorate); Sb-Oxo, antimony-oxo tetra-(*p*-methoxyphenyl)-porphyrin.

Apart from structural similarities, the behavior of excited  $PC_{S_{HAT}}$  featuring an oxo moiety shows many analogies. Thus, the capability of carbonyl derivatives, especially (aromatic) ketones, to act as hydrogen abstractor has been known for a long time.<sup>297–302</sup> In particular, the photochemistry of these compounds is dominated by the triplet excited state and this is due to the very efficient intersystem crossing (ISC) from the first-formed singlet state. These triplet states have a lifetime in the microsecond range<sup>303</sup> (Table 1) and, depending on the structure of the carbonyl derivative, may have a  $n\pi^*$  or a  $\pi\pi^*$  character (the former typically accountable for HAT reactivity).<sup>297,304–310</sup> In carbonyl derivatives, the  $PC^{\bullet-H}$  form is a stable, long-living ketyl radical species (Scheme 5),<sup>311</sup> featuring a very weak O–H bond (the calculated BDE of the O–H bond of the ketyl radical derived from acetone is ca. 16 kcal/mol).<sup>312</sup> Accordingly, the restoration of the carbonyl group is the driving force for the recovery of the spent photocatalyst. However, a common drawback when using ketones as  $PC_{S_{HAT}}$  is that ketyl radical may dimerize in solution to form pinacols thus decreasing the efficiency of the reaction.<sup>313</sup> A particular case is that of the excited carbonyl of EY that has some phenoxyl radical character.<sup>221</sup>

Another deeply studied family of  $PC_{S_{HAT}}$  is that of inorganic polyoxometalate (POM) derivatives. The first reports describing a HAT reactivity upon excitation of these metal–oxygen clusters in the presence of organic substrates (mostly alcohols) dates back to the 1980s.<sup>314–318</sup> It was soon realized that tungsten-based POMs, in particular DT, outperform all the other known POMs in terms of HAT reactivity,<sup>201,319</sup> offering a catalytic tool for the elaboration of C–H bonds.<sup>320–331</sup> A common occurrence in photocatalytic systems based on DT is the observation of a typical blue color of the  $PC^{\bullet-H}$  form

( $[W_{10}O_{32}]^{5-}$ , either protonated or not)<sup>332</sup> that accumulates in solution.<sup>333</sup> The studies of DT-based systems by means of time-resolved spectroscopic techniques<sup>334–340</sup> concluded that the state responsible for the HAT reactivity is a relaxed excited state (“wO” having a lifetime of 55 ns in acetonitrile),<sup>334</sup> probably of triplet multiplicity,<sup>341</sup> not directly accessible upon excitation (a so-called dark state).<sup>342,343</sup> Theoretical simulations supported these experimental spectroscopic works.<sup>265,344,345</sup>

Turning to the uranyl cation, despite the very weak absorption in the blue region of the spectrum ( $\epsilon \sim 10 \text{ M}^{-1} \text{ cm}^{-1}$  at  $\lambda = 423 \text{ nm}$ ), visible light irradiation can be adopted to trigger its photochemistry. This transition has been proposed to populate a long-lived ( $\mu\text{s}$  range lifetime) state which contains an extremely reactive oxyl radical, well explaining the HAT reactivity.<sup>208,271,346–350</sup>

A partial oxyl radical character of the triplet excited state has been likewise postulated to be the active species in the hydrogen abstraction operated by Sb-Oxo.<sup>266</sup> In the latter case, Sb<sup>V</sup> (a high-valent oxidation state element of the *p*-block) is used in the dihydroxo form that contains two hydroxyl groups in the axial positions. Upon treatment with a base, one of the two hydroxy groups turns into the desired oxo moiety; the excited state of the so-generated oxo species was exploited for hydrogen abstraction.<sup>266,351</sup> Even in this case the lifetime of the triplet involved is in the microsecond range (Table 1).

It is important to stress, however, that  $PC_{S_{HAT}}$  may be engaged in photocatalytic processes different from HAT (mainly electron transfer, but energy transfer may not be excluded).<sup>186,266,310,352</sup> Accordingly, the real mechanism has to be checked carefully to ascertain if a HAT process is involved rather than an electron transfer followed by proton transfer or

Table 1. Photophysical Properties of Selected PCsHAT

Photocatalyst	$\lambda_{\text{max}}$ (nm)	$\lambda_{\text{use}}$ (nm)	$\Phi_{\text{ISC}}$	$\tau_{\text{T}}$ ( $\mu\text{s}$ )
AP	243, 279, 315 (THF or MeCN) <sup>254</sup>	19 W CFL <sup>225</sup>	$\sim 1$ (computational) <sup>222</sup>	6.6 (H <sub>2</sub> O/MeCN 9:1) <sup>211</sup>
ABP	337 ( <i>i</i> -PrOH), <sup>255</sup> 303 (C <sub>6</sub> H <sub>12</sub> ), <sup>255</sup> 308 (CCl <sub>4</sub> ) <sup>255</sup>	365, UV LED <sup>226</sup>	0.82 (C <sub>6</sub> H <sub>12</sub> ) <sup>212</sup>	3.6 <sup>212</sup>
AQ	325, 370 (DCM) <sup>256</sup>	360, <sup>227</sup> 390, 427 <sup>228</sup>	0.95 (MeCN) <sup>213</sup>	1.62 (MeCN) <sup>213</sup>
BP	347, 415 <sup>257</sup>	350, <sup>229</sup> Hg lamp, <sup>230–233</sup> 350 Rayonet, <sup>234</sup> 366 Rayonet, <sup>235</sup> 365 LED lamp, <sup>236</sup> 18 W CFL, <sup>237</sup> 400 nm LED <sup>238</sup>	1 (MeCN) <sup>214</sup>	19.6 (MeCN) <sup>214</sup>
BPSS	330 <sup>215</sup>	315, 366 <sup>239</sup> sunlight <sup>240</sup>	1 (H <sub>2</sub> O) <sup>215</sup>	352 (H <sub>2</sub> O) <sup>215</sup>
CIAQ	256, 265, 274, 325 (MeOH) <sup>258</sup>	365 LED lamp, <sup>241</sup> Xe lamp <sup>242</sup>		1150 (CF <sub>3</sub> CH <sub>2</sub> OH at 77 K) <sup>216</sup>
DMBP	339 <sup>259</sup>	365 UV fluorescent lamp <sup>243</sup>		3.4–17.6 (CH <sub>2</sub> Cl <sub>2</sub> /CHCl <sub>3</sub> 1:1 at 77 K) <sup>217</sup>
DBS	251, 306, 347 (C <sub>6</sub> H <sub>12</sub> ), <sup>260</sup> 253, 316, 326 (EtOAc), <sup>260</sup> 253, 306, 347 (MeCN), <sup>260</sup> 255, 305, 348 ( <i>i</i> -PrOH), <sup>260</sup> 255, 306, 349 (MeOH) <sup>260</sup>	CFL <sup>244</sup>		980 (C <sub>6</sub> H <sub>12</sub> , Ar), <sup>218</sup> 890 (EtOH, Ar) <sup>218</sup> 740 (MeOH, Ar), <sup>218</sup> 650 (C <sub>6</sub> H <sub>6</sub> , Ar) <sup>218</sup>
PQ	268, 324, 500 (CHCl <sub>3</sub> ), <sup>261</sup> 319, 312, 503 (C <sub>6</sub> H <sub>6</sub> ) <sup>262</sup>	Blue LED strip <sup>245</sup>	1 <sup>223</sup>	2.2 <sup>219</sup>
PGA	350 (dioxane) <sup>263</sup>	household lamps <sup>246</sup>	$\sim 1$ <sup>195</sup>	8 <sup>195</sup>
DTX	354 (MeCN) <sup>220</sup>	365 LED lamp <sup>247,248</sup>	0.93 <sup>219</sup>	862 (MeCN) <sup>220</sup>
EY	541 (DMF) <sup>264</sup>	Blue LED, <sup>249,250</sup> 460 Blue LED, <sup>251</sup> 470 blue LED, <sup>252</sup> 520 Green LED, <sup>251</sup> Xe lamp (400 nm cutoff) <sup>253</sup>	0.32 <sup>224</sup>	$\sim 21$ <sup>221</sup>
FL	382 (C <sub>6</sub> H <sub>12</sub> ), <sup>288</sup> 380 (C <sub>6</sub> H <sub>6</sub> ), <sup>288</sup> 377 (MeCN), <sup>288</sup> 379 (EtOH) <sup>288</sup>	Xe lamp, <sup>272</sup> CFL lamp <sup>273</sup>	0.97 (MeCN) <sup>214</sup>	70 (MeCN) <sup>214</sup>
NaDT/ TBADT	260, 323 (H <sub>2</sub> O), <sup>274</sup> 272, 321 <sup>289</sup>	BLB lamp, <sup>274</sup> 310 Multilamp reactor, <sup>275</sup> 366 LED, <sup>276</sup> Xe lamp, <sup>272</sup> Solar light, <sup>277</sup> 390 LED <sup>278</sup>	0.5 (MeCN) <sup>265</sup>	50 (MeCN) <sup>265</sup>
PT	270, 343 (CHCl <sub>3</sub> ) <sup>290</sup>	365, 425, 279, 390, 427, 455 <sup>282</sup>		
PYD	427 (MeCN/THF 1:1), <sup>291</sup> 266, 277, 398, 424, 448 (ether) <sup>292</sup>	Blue LED <sup>283</sup>		
Sb-Oxo	440 <sup>296</sup>	405, 455 <sup>286</sup>		
TX	380 <sup>220</sup>	405 LED <sup>272</sup>	0.99 (MeCN), <sup>214</sup> 0.76 <sup>268</sup>	8 (MeCN/H <sub>2</sub> O 95:5) <sup>266</sup> 28 (MeCN) <sup>214,267</sup> 760 ns (CD <sub>3</sub> CN), <sup>220</sup> 45 (MeCN) <sup>268</sup>
UrN	414, <sup>293–295</sup>	456, <sup>284</sup> Blue LED <sup>285</sup>	1 <sup>271</sup>	400 (80 K, MeOH) <sup>269</sup>
XA	338 <sup>296</sup>	365 LED, <sup>286</sup> CFL <sup>273,287</sup>	1 (MeCN) <sup>214</sup>	4.8–8.3 (MeCN) <sup>214,270</sup>

even a proton-coupled electron transfer (PCET) mechanism.<sup>163</sup>

In view of the above, the aim of the present review is to offer an overview of the synthetic applications based on photocatalyzed *direct* HAT (*d*-HAT), wherein the excited  $PC_{HAT}$  triggers the HAT step. On the other hand, examples dealing with a photocatalyzed *indirect* HAT (*i*-HAT)<sup>182–185</sup> or *remote* HAT (*r*-HAT)<sup>184</sup> as well as the activation of C–H bonds via a PCET mechanism<sup>353</sup> will be excluded. The threshold that we used throughout the entire work to consider an approach photocatalytic is 20 mol % of catalyst loading. We then considered photocatalytic HAT reactions where the generated radical must be incorporated in the desired compound, so the photogeneration of a thermally active redox agent will not be treated here.<sup>291,354–356</sup> Photoinitiated processes wherein the light-absorbing species undergoes degradation during the process have been likewise excluded.<sup>357–359</sup>

Similarly, the adoption of a *d*-HAT strategy in polymerizations will not be mentioned; however, the reader is invited to refer to seminal works in the field.<sup>360–365</sup> Thus, synthetic applications are preferentially treated here avoiding (when possible) the works simply devoted to mechanistic purposes and where the  $PC_{HAT}$  tested gave a very low yield or a very low reagent consumption.

The following sections have been organized based on the bond being formed during the transformation and found in the final product, while different types of transformations (e.g., dehydrogenation and fragmentation reactions) have been reported in the final part of the review. Thus, under the section “formation of a  $C(sp^3)$ – $C(sp^2)$  bond”, examples wherein a  $C(sp^3)$ -centered radical (formed from the photocatalyzed homolysis of a  $C(sp^3)$ –H bond via a HAT step) will be attached to a  $C(sp^2)$  atom in the final product will be described. Moreover, despite the fact that most of the examples reported in this review deal with the functionalization of C–H bonds, the elaboration of P–H, Si–H, and S–H bonds via a photocatalyzed *d*-HAT step has been mentioned, for the sake of comprehensiveness.

All the schemes have been color-coded so that the bond activated via HAT has been reported in violet, while the bond formed has been highlighted in red.

## 2. FORMATION OF C–C BONDS


### 2.1. Formation of $C(sp^3)$ – $C(sp^3)$ Bonds via Addition onto C=C Bonds

A typical reactivity mode that can be exploited to forge a  $C(sp^3)$ – $C(sp^3)$  bond is the radical addition of nucleophilic radicals onto Michael acceptors. In this scenario, the C-centered radical generated via photocatalyzed HAT is trapped by an electrophilic olefin and the resulting radical adduct is quenched via back hydrogen atom transfer (or sequential electron/proton transfer) from the reduced form of the photocatalyst ( $PC^{\bullet}$ –H; see Scheme 5), thus closing the photocatalytic cycle. This is a very reliable and general protocol as also demonstrated by the vast amount of hydrogen donors that can be profitably employed, with notable examples including alcohols, ethers, dioxolanes, sulfides, amides, nitriles, as well as simple hydrocarbons such as toluenes, allylated derivatives, and even (cyclo)alkanes.

**2.1.1. Oxygen- and Sulfur-Containing Compounds as Hydrogen Donors.** Oxygenated derivatives have been the elective substrates for this reactivity manifold since the earliest

reports on photocatalyzed HAT; this is because of the low BDE of the  $\alpha$ -to-O C–H bond and the relatively stable  $\alpha$ -oxy radical generated. In particular, to the best of our knowledge, alcohols are the first hydrogen donors ever investigated,<sup>366</sup> and the earliest preparative example appeared in 1957 dealing with the photoaddition of isopropanol onto maleic acid **6.1** to give terebic acid **6.2** (Scheme 6).<sup>367</sup> In those days, aromatic ketones

**Scheme 6. Photoaddition of Isopropanol onto Maleic Acid**




(e.g., BP) were the elective class of  $PC_{sHAT}$  to perform the reaction under UV light coming from a Hg lamp.<sup>367</sup> Due to the low cost of isopropanol, this was used as the reaction medium. The mechanism is depicted in Scheme 6 (lower part) and, as mentioned in the introduction to this section, it is quite common in all cases wherein a radical Michael addition takes place. Thus, the excited  $PC_{HAT}$  abstracts a hydrogen atom from the hydrogen donor and the resulting radical adds onto the olefin to give the radical adduct **6.3**. Back hydrogen atom transfer from the reduced form of the PC to **6.3** yielded the hydroxy acid **6.4** (and **6.2** from it by spontaneous lactonization) with the concomitant regeneration of  $PC_{HAT}$ .

The same reaction was later on replicated on a gram scale by using the SOLFIN (SOLar synthesis of FINE chemicals) apparatus as solar light concentrator placed in Almeria (Spain).<sup>240</sup> In this case, BPSS was used in the role of  $PC_{HAT}$ , which was synthesized by sulfonation of parent BP (the sulfonation took place both at the 3- and 4-positions of the aromatic rings). The reason was that the thus-obtained  $PC_{HAT}$  was easily removed at the end of the reaction by extraction with water. Thus, ca. 14 g of terebic acid (**6.2**) was isolated in 75% yield upon 14 h solar light irradiation of an isopropanol/water 1:1 solution of **6.1** in the presence of 10 mol % BPSS.<sup>240</sup> This  $PC_{HAT}$  has been likewise used to trigger the addition of alcohols (isopropanol, ethanol, and methanol) onto  $\alpha,\beta$ -unsaturated aldehydes for the preparation of  $\gamma$ -lactols and  $\gamma$ -lactones upon treatment of the crude lactols with bromine.<sup>239</sup> When maleic or fumaric acids were converted to the corresponding chiral (–)-menthyl diesters, the BP (19 mol %) photocatalyzed addition of isopropanol gave the acyclic diaterebic acid ester (63% yield) with a modest degree (8%) of diastereoselectivity.<sup>368</sup>

Alcohols (in particular, methanol) were used to functionalize carbohydrate enones, such as hex-2-enopyranosid-4-ulose **7.1**, to form branched-chain monosaccharides (Scheme 7).<sup>234,369–371</sup> Irradiation of this  $\alpha$ -enone in MeOH in the


### Scheme 7. Photocatalyzed Functionalization of an $\alpha$ -Enone



presence of BP afforded 1,4-ketoalcohol **7.2** in 66% yield. Interestingly, the incorporation of the alcohol took place from the less-hindered side of the enone in a complete stereo- and regioselective fashion.<sup>370</sup>

Isopropanol was likewise used for the derivatization of 1,3-dioxin-4-ones having a (–)-menthone moiety embedded as chiral auxiliary in the 2-position (**8.1**, Scheme 8). The resulting 1,5-dioxaspiro[5.5]undecane-2-one (**8.2**) was formed, however, in a poor yield (<30% by using 15 mol % BP).<sup>372</sup>


### Scheme 8. Photocatalyzed Synthesis of 1,5-Dioxaspiro[5.5]undecane-2-ones



On the other hand, the adoption of TBADT allowed the activation of isopropanol even by using a low amount of the PC<sub>HAT</sub> (2–4 mol %) in the reaction with acrylonitrile (72% yield).<sup>81,373</sup>

The addition of isopropanol onto a Michael acceptor (e.g., furanone **9.1**, Scheme 9) was likewise carried out under flow

### Scheme 9. Photocatalyzed Functionalization of Furanones under Flow Conditions




conditions<sup>374</sup> by using either an LED-driven microchip reactor,<sup>375</sup> a continuous-flow photoreactor with parallel capillaries,<sup>376</sup> or a multimicrocapillary flow reactor.<sup>243</sup> In all cases, the adduct **9.2** was formed in a less than 10 min irradiation. Of note, DMBP was found to be the best PC<sub>HAT</sub> among the several aromatic ketones tested.<sup>243</sup>

Cyclic alcohols have been rarely used, but the methine hydrogen atom in cyclohexanol (**10.1**, Scheme 10) was selectively abstracted by the excited state of PT and the radical formed was engaged in an allylation reaction to give homoallyl alcohol **10.3**.<sup>290</sup>



The lability of the C–H bonds in position 2- in (2-substituted) 1,3-dioxolanes has been exploited for the

### Scheme 10. Synthesis of Homoallyl Alcohols



generation of dioxolan-2-yl radicals, which moiety was used to formally introduce a masked formyl group. The reaction was initially tested on  $\alpha$ -enones similar to **7.1** by using 1,3-dioxolane as the solvent.<sup>234</sup> Later on, the process was extended to other enones, such as 1-phenyl-2-propen-1-one or chalcone **11.1** (Scheme 11).<sup>227</sup> In that case, AQ was adopted as a

### Scheme 11. Photocatalyzed Incorporation of a Masked Formyl Group



visible-light-absorbing photoorganocatalyst (POC)<sup>154</sup> and, despite the long reaction time needed (ca. 60 h), the final adduct **11.2** was isolated in 85% yield. The same reaction described in Scheme 11 was also performed in a 3D-printed, chemically resistant, nonswelling, and UV–vis transparent postfunctionalized flow reactor by using ABP as an immobilized PC<sub>HAT</sub>; however, an unsatisfactory yield (13%) was reported.<sup>226</sup>


The BP-photocatalyzed addition of 1,3-dioxolane onto 5-alkoxymethyl-2(*SH*)-furanone was used as the key step for the preparation of a bis-tetrahydrofuran ligand for HIV protease inhibitor UIC-94017 (TMC-114).<sup>377</sup> BPSS was likewise used as PC<sub>HAT</sub> to promote the radical addition of 1,3-dioxolane onto  $\alpha,\beta$ -unsaturated aldehydes to give 1,4-monoprotected succinaldehydes upon solar light exposure.<sup>240</sup> The radical 1,3-dioxolanylation of alkenoic acids was also performed by using DTX (10 mol %) as POC.<sup>248</sup> Shifting to a metal-based PC, the adoption of UrN allowed the hydrogen activation in 1,3-dioxolane for the addition onto Michael acceptors.<sup>284</sup>

Along the same line, 2-alkyl-1,3-dioxolanes were exploited as hydrogen donors for the (formal) incorporation of a ketone moiety, but they had to be used as cosolvents. In such a way, 1,4-monoprotected ketoaldehydes were obtained upon radical alkylation of  $\alpha,\beta$ -unsaturated aldehydes.<sup>239</sup>

TBADT was the elective PC<sub>HAT</sub> for the activation of the methylene hydrogens in substituted 1,3-benzodioxoles (e.g., **12.1**) to give the corresponding 2-substituted derivatives by reaction with various Michael acceptors<sup>378</sup> or with styrene<sup>379</sup> (in the latter case in the presence of a disulfide cocatalyst). When the process was carried out on the  $\beta$ -substituted cyclic enone **12.2** (Scheme 12) in the presence of a chiral organocatalyst (i.e., carbazole derivative (*S,S*)-**12.3**), an enantioselective radical conjugate addition took place with formation of **12.4** in 99% yield. Notably, the latter product was formed with e.e. 88% and contains two quaternary carbon




### Scheme 12. Dual-Catalytic Asymmetric Formation of Quaternary Carbons



stereocenters.<sup>380</sup> The reaction is based on an electron relay mechanism. In fact, the carbazole moiety is oxidized by an intramolecular electron transfer with the unstable radical cation formed by radical addition onto the chiral iminium ion intermediate, thus functioning as an electron donor.

Another class of widely used oxygen-based hydrogen donors is that of cyclic ethers, wherein the HAT step occurs at the labile  $\alpha$ -to-O C–H bonds. It is perhaps important to stress here that cyclic ethers cannot be easily activated otherwise; in fact, they are routinely used as inert solvents. Scheme 13

### Scheme 13. Cyclic Ethers as H-Donors in the Functionalization of Butendioate Esters



collects some representative examples concerning the derivatization of butendioate esters **13.1**. TBADT enabled the facile cleavage of the C–H bond adjacent to the oxygen atom both in 1,4-dioxane **13.2**<sup>277</sup> and in oxetane **13.4**.<sup>381</sup> In the former case, sunlight was effectively used to irradiate the solution poured in a glass vessel placed on a window ledge. Despite the long time required (4 days), the reaction did not make use of any external source of artificial energy.<sup>277</sup> The same process was performed upon UV light irradiation under flow conditions in a shorter period.<sup>382</sup>

The generation of radicals from tetrahydrofuran (**14.1**) is useful to compare different PC<sub>S<sub>HAT</sub></sub> in their role and to stress the versatility of the photocatalyzed HAT process (Scheme 14). Thus, **14.1** may be photoactivated by having recourse to several PCs, including aromatic ketones such as TX and FL under visible light LED irradiation,<sup>272</sup> TBADT under solar simulated conditions,<sup>272,383</sup> as well as with Sb-Oxo,<sup>266</sup> UrN,<sup>284</sup> PYD,<sup>283</sup> EY,<sup>221</sup> and CIAQ.<sup>241</sup> In all cases, satisfactory yields of adducts **14.3**, **14.5**, **14.7**, and **14.9** were obtained.

The photocatalyzed addition of THF was also applied to quinones **15.1a–d** (4-benzylidene-2,6-di-*tert*-butylcyclohexa-2,5-dien-1-ones, Scheme 15) under blue LED irradiation by

using UrN (5 mol %) to give 2,6-di-*tert*-butyl-4-[phenyl-(tetrahydrofuran-2-yl)methyl]phenols **15.2a–d**.<sup>384</sup>


In rare instances, the activation of the C–H bond was applied to cyclic carbonates (**16.1a,b**, Scheme 16), where the presence of the carbonyl group did not hamper the C–H cleavage in these substrates. The importance of using TBADT is evident in this case, since the same process promoted by aromatic ketones gave no products **16.3a,b**.<sup>272</sup> Introducing a methyl group in carbonate **16.1b** drove the cleavage to the most labile C–H bond present.

Apart from the case of carbonates, other carbonyl-containing derivatives (ketones, esters, lactones, etc.) can be used as hydrogen donors. However, it is important to remember that in these cases the selectivity of the HAT step is shifted toward remote positions due to the mismatched polarity (Scheme 2). A typical case is the photoactivation of cyclopentanones **17.1a,b** (Scheme 17a). Despite the lability (and the acidity) of the  $\alpha$ -C–H bonds with respect to  $\beta$ -C–H bonds in compound **17.1a**, the former are left untouched under the action of TBADT and a selective  $\beta$ -C–H to C–C bond conversion occurred.<sup>83</sup> A more favorable polar HAT transition state has been invoked in this case to rationalize the observed regioselectivity.<sup>84</sup> The presence of a methyl group in compound **17.1b** made the methine hydrogen sufficiently labile to allow the preparation of compound **17.3b** as the sole product. Notably, the combination of polar and steric effects may direct the selective C–H cleavage in cycloalkanones and lactones. As an example, compound **17.4** underwent a selective  $\beta$ -C–H cleavage since both the hydrogen abstraction from the  $\alpha$ -C–H and the  $\gamma$ -C–H bonds is prevented by polar and steric effects, respectively (Scheme 17b).<sup>385</sup> In particular cases, the regioselective cleavage may be induced even in open chain esters (e.g., **17.7**) exploiting the lability of the methine hydrogen of the isopropyl group and taking advantage of the bulkiness of the *t*Bu group that prevents any other competitive C–H cleavage (Scheme 17c).<sup>385</sup>


Sparse examples have been reported involving the use of sulfides (mainly cyclic derivatives) as hydrogen donors. Thus, tetrahydrothiophene **18.1** (or thioxane) was allylated at the C–H bond adjacent to the S-atom by reaction with allyl sulfone **18.2** (Scheme 18).<sup>290</sup> Despite the easy oxidizability of these sulfides, the adoption of PT avoided any competitive electron transfer reaction. Additionally, both CIAQ<sup>241</sup> and EY<sup>221</sup> were likewise effective PC<sub>S<sub>HAT</sub></sub> to trigger the C–H to C–C bond conversion in tetrahydrothiophene.

**2.1.2. Nitrogen-Containing Compounds as Hydrogen Donors.** A widely used class of nitrogen-containing hydrogen donors is that of amides (often used as the solvent) and carbamates; albeit, often the PC<sub>HAT</sub> has to be used in a (super)stoichiometric amount for their activation.<sup>158</sup> Nevertheless, catalytic amounts of TBADT smoothly promoted the C–H functionalization in amides and carbamates, used only in a 4 equiv excess (Scheme 19). The C–H bonds adjacent to the nitrogen atom in protected pyrrolidine **19.1** were sufficiently labile to be cleaved under photocatalyzed conditions to afford nitrile **19.3** (Scheme 19a), while excess **19.1** could be recovered during the purification.<sup>386</sup> The reaction was found to be effective even under sunlight exposure.<sup>277</sup> A similar C–H activation has been reported by using CIAQ (10 mol %)<sup>241</sup> and PT (5 mol %)<sup>290</sup> as PC<sub>S<sub>HAT</sub></sub>.


As an alternative to typical Michael acceptors, the  $\alpha$ -amidoalkyl radical formed from dimethylformamide **19.4** in the presence of TBADT was trapped by vinyl pyridine **19.5** to

Scheme 14. Different PC<sub>S<sub>HAT</sub></sub> for the Photocatalyzed Cleavage of the C–H Bond in THF

Scheme 15. Photocatalyzed Addition of THF onto Cyclohexa-2,5-dien-1-ones




Scheme 16. TBADT-Mediated Derivatization of Cyclic Carbonates





give adduct **19.6** in a very high yield (82%, Scheme 19b).<sup>387</sup> It is important to note that in the latter case no C–H cleavage of the formyl hydrogen competed (see further section 2.5).

A related C–H functionalization of carbamates to perform a Giese-type alkylation was carried out by combining the action of a POC (BP, 20% mol) with a catalytic amount of Cu(OAc)<sub>2</sub> (2% mol) under UV-A irradiation. In this case, the copper species prevents the otherwise feasible polymerization of

Scheme 17. Selective C–H Cleavage in Ketones and Esters




Scheme 18. Allylation of Tetrahydrothiophene



Michael acceptors, such as unsubstituted acrylates, acrylonitrile, or methyl vinyl ketone.<sup>388</sup>

## Scheme 19. Amides, Carbamates, and Amines as H-Donors




Even EY may be used for the selective, photocatalyzed addition of acetamide onto benzylidenemalononitrile.<sup>221</sup> When using UrN, the C–H cleavage in **19.4** was not selective since competitive hydrogen abstraction from the C(sp<sup>2</sup>)–H bond took place (ca. 1/3 ratio).<sup>284</sup> When *N*-methylacetamide was subjected to a hydrogen abstraction reaction by using DCBP (20 mol %), the resulting  $\alpha$ -amidoalkyl radical was trapped by  $\beta$ -phenyl allyl sulfone to give the corresponding allylated derivative.<sup>389</sup>

The TBADT-photocatalyzed addition of tertiary amides (e.g., **19.7**, Scheme 19c) onto vinyl sulfones under flow conditions was selected for the easy preparation of  $\gamma$ -aminopropylsulfones (**19.9**).<sup>390</sup> The latter conditions allowed scale-up of the process with a substrate concentration up to 0.5 M.<sup>390</sup>

In rare instances, an amine functioned as the hydrogen donor. Indeed, the electron-donor capability of such substrate may engage an electron transfer rather than a hydrogen atom transfer reaction (this is a typical case when using aromatic ketones).<sup>186,352</sup> However, EY was able to functionalize amine **19.10** via enantioselective addition onto  $\alpha,\beta$ -unsaturated *N*-acyl-3,5-dimethylpyrazole **19.11** (Scheme 19d).<sup>391</sup> The asymmetric Giese-type addition of the photogenerated  $\alpha$ -amino radical was promoted by the presence of the chiral rhodium Lewis acid catalyst  $\Delta$ -RhS. As a result, adduct **19.12** was formed in a modest yield but with a high e.e..<sup>391</sup>


Similarly, in one instance a primary amine was derivatized under photocatalyzed conditions (Scheme 20). Thus, the visible light irradiation of a mixture of amine **20.1**, a styrene (**20.2a–d**), and a catalytic amount of EY caused the C–H cleavage of the methine hydrogen in **20.1**, finally affording the

Scheme 20. Photocatalyzed C(sp<sup>3</sup>)–H Alkylation of Amines


hoped-for 2-methyl-4-arylbutan-2-amine derivatives **20.3a–d**.<sup>392</sup> Apart from the mildness of the reaction conditions, this is an important example dealing with the derivatization of vinyl aromatics.<sup>392</sup>

A particular case in the activation of the C–H bond in amines is depicted in Scheme 21 and deals with the introduction of a trifluoromethyl group, which is known to improve the pharmacokinetic properties of drugs.<sup>393</sup> This challenging C(sp<sup>3</sup>)–C(sp<sup>3</sup>) bond formation was made possible by merging NaDT chemistry with copper catalysis and made use of the Togni's reagent **21.2** as the trifluoromethylating agent.<sup>394</sup> The adopted acidic conditions here caused the formation of the ammonium salt of pyrrolidine **21.1**, thus deactivating the C–H bonds adjacent to the nitrogen atom (Scheme 2). Overall, the strategy is based on the addition of the photocatalyzed C-centered radical onto a Cu<sup>II</sup>–CF<sub>3</sub> species. The hoped-for trifluoromethylated product **21.3** (trifluoroacetate salt) was then obtained in 68% isolated yield as a single regioisomer. The same procedure enabled the trifluoromethylation of benzylic C–H bonds and of biologically valuable compounds such as lidocaine, prilocaine,

### Scheme 21. Metallaphotoredox Strategy for the Trifluoromethylation of Amines



Notable examples:



celecoxib, and torsemide. Mechanistic studies are consistent with the involvement of a “Cu- $\text{CF}_3$  complex”.<sup>394</sup>

Other nitrogen-containing hydrogen donors have been reported, such as aliphatic nitriles and alkyipyridines, where the influence of the heteroatom is not so important as in the previous cases; albeit, it still has a role in directing the C–H cleavage event. A representative case is that of adiponitrile **22.1** (Scheme 22a). The electron-withdrawing effect of the cyano group hampers the cleavage of  $\alpha$ -C–H but not of  $\beta$ -C–H bonds. Thus, the TBADT-photocatalyzed reaction between **22.1** and dimethyl maleate **22.2** easily gave tetrafunctionalized adduct **22.3** in a satisfying yield.<sup>395</sup> Related reactions involve the photocatalyzed addition of 4-methylpentanenitrile to phenyl vinyl sulfone (under flow conditions)<sup>382</sup> or to a vinylpyridine.<sup>387</sup>

The same site-selective C–H to C–C bond conversion in nitriles took place when DT was incorporated within the pores of a copper-based metal organic framework (MOF) ( $[\text{Cu}_4(\text{BPY})_6\text{Cl}_2(\text{W}_{10}\text{O}_{32})]\cdot 3\text{H}_2\text{O}$ ; DT-BPY, BPY = 4,4'-bipyridine). This is a rare case where the HAT process is carried out under heterogeneous conditions. The new  $\text{PC}_{\text{HAT}}$  showed high catalytic efficiency, high stability, and good recyclability, allowing use of a lower excess of the aliphatic nitrile substrate (only 5 equiv), thus improving the sustainability of the process (Scheme 22a).<sup>396</sup>


The C–H activation in alkyipyridines is interesting, since the labile benzylic hydrogens are not involved in the process, at variance with the alkylbenzene counterparts (see also section 2.1.3). This is well exemplified by the case of **22.4**, wherein the methine hydrogen was selectively cleaved and the resulting tertiary radical was then trapped by ketone **22.5** to give adduct

**22.6** (Scheme 22b).<sup>397</sup> In the last case, preference of the excited  $\text{PC}_{\text{HAT}}$  to abstract the less acidic (or, in other words, the less electrophilic) hydrogen atom in the investigated alkyipyridine was observed.


**2.1.3. Hydrocarbons as Hydrogen Donors.** In hydrocarbons, it is possible to find quite labile hydrogens that can be easily cleaved under photocatalytic conditions. Hydrocarbons displaying labile benzylic<sup>398</sup> and allylic C–H bonds can be easily cleaved at these sites under photocatalytic conditions. In fact, the BDEs of the most labile C–H bonds in toluene and cyclohexene are 88 and 82 kcal/mol, respectively (see Scheme 3).

The main problem here is the high stability of the radical formed and its reluctance to react with the reaction partner (e.g., a C=C bond) to forge a  $\text{C}(\text{sp}^3)\text{--C}(\text{sp}^3)$  bond. This probably explains why very few processes involving these substrates have been reported. Simple alkylaromatics have been derivatized by using TBADT to perform valuable benzylations. However, only easily reducible olefins, including fumaronitrile, maleic anhydride, and substituted maleic imides, gave good results.<sup>399</sup> The same conjugate radical additions were carried out in a mesoscale flow photoreactor by adopting a water-cooled 500 W medium-pressure Hg-vapor lamp as the light source. The use of this apparatus led to a marked increase of the STY (space time yield) and a reduction of the irradiation time compared with the same processes developed under batch conditions.<sup>382</sup> Other  $\text{PC}_{\text{HAT}}$  were likewise useful for this C–H activation strategy, as collected in Scheme 23.

### Scheme 23. Functionalization of (a) Benzylic and (b,c) Allylic Hydrogens



### Scheme 22. Regioselective Photocatalyzed C–H Cleavage in (a) Aliphatic Nitriles and (b) Alkyipyridines






Thus, the benzylic position in toluene **23.1** was functionalized under EY photocatalysis despite heating at 60 °C being required (Scheme 23a).<sup>221</sup> The activation of allylic hydrogens was also attempted by using DT, albeit not on a preparative scale.<sup>201</sup>

More recently, cyclohexene has been used as the hydrogen donor for the preparation of allylated derivatives **23.5** and **23.7** (Schemes 23b,c). Both EY<sup>221</sup> and CIAQ<sup>241</sup> were employed in the functionalization of very good Michael acceptors **23.2** and **23.6**.

The most challenging reaction for the construction of C(sp<sup>3</sup>)-C(sp<sup>3</sup>) bonds is related to the functionalization of (cyclo)alkanes,<sup>125</sup> due to the high BDE of the C-H bonds involved (ca. 100 kcal/mol, see Scheme 3). Early photocatalytic experiments made use of a high amount of the PC<sub>HAT</sub> to pursue this issue,<sup>158</sup> but the use of DT allowed performing a real photocatalyzed process with only a few mol % loading of the PC<sub>HAT</sub>. Simple symmetric cycloalkanes were the preferred substrates.<sup>81,275,400</sup> As shown in Scheme 24a, cyclohexane **24.1a** easily gave access to the corresponding cycloalkyl radical that was in turn trapped by dinitrile **24.2** to give **24.3** through a C-C bond formation step.<sup>275</sup>

#### Scheme 24. Photocatalyzed C-H Cleavage in (Cyclo)alkanes



Similarly, various 5- to 12-membered cycloalkanes were used to functionalize conjugated enones (**24.6**)<sup>401</sup> even with the help of a chiral spiro phosphoric acid (**24.7**) to promote an asymmetric C-H functionalization (Scheme 24b).<sup>402</sup> A chiral phosphoric acid similar to **24.7** was likewise adopted as a chiral proton-transfer shuttle in the cycloalkane addition onto  $\alpha$ -substituted acrylates (e.g., *N*-acyl dehydroalanine benzyl esters) used as Michael acceptors for the smooth forging of enantioenriched  $\alpha$ -stereogenic esters.<sup>403</sup>


Methylene norbornanone was alkylated in a good yield by using cyclohexane as the hydrogen donor under UrN photocatalysis.<sup>284</sup> Similarly, the activation of nonacidic C-(sp<sup>3</sup>)-H bonds in cyclohexane (or adamantane) was carried out upon UV light irradiation (CIAQ as the PC<sub>HAT</sub>) by using 1,1-bis(phenylsulfonyl)ethylene as the radical trap.<sup>241</sup> The allylation of alkanes has been performed by means of the PT-photocatalyzed addition of cyclohexane, cyclododecane, or adamantane onto 1,2-bis(phenylsulfonyl)-2-propene as the allyl source.<sup>290</sup>

In rare instances, the reaction was applied to substituted cycloalkanes. Thus, the presence of a *t*Bu group in compound **24.1b** exerted a profound effect in steering the hydrogen abstraction process. In fact, the bulkiness of the *t*Bu group completely shielded the hydrogens in positions 1- and 2-, allowing the selective C-H cleavage in positions 3- and 4- (Scheme 24a). The bulkiness of the PC<sub>HAT</sub> and the radical trap **24.2** further helped in reducing the number of possible isomers formed, with only *cis*-3-substituted **24.4** and *trans*-4-substituted **24.5** formed in an overall 70% yield.<sup>77</sup> Interestingly, in the latter case, when BP (1 equiv) was used in place of TBADT, the same product distribution was roughly observed.<sup>77</sup> However, when alkane **24.1b** reacted with acrylonitrile (TBADT as the PC<sub>HAT</sub>), a more complex mixture resulted.<sup>77</sup>

Open-chain alkanes were poorly investigated. A rare case is that reported in Scheme 24c. Despite the fact that compound **24.9** has five different types of hydrogen atoms, only the methine C-H position was effectively cleaved, and the reaction with maleate **24.10** led to diester **24.11** as the sole product.<sup>373</sup> Even in this case, the bulkiness of the *t*Bu group helped in the regioselective cleavage of the C-H bond.

The activation of methane (BDE = 105 kcal/mol) was proved to be feasible by adopting DT photocatalysis. The process required specifically optimized conditions, namely the adoption of flow conditions and application of a high pressure (45 bar), to allow the correct mixing of the reagents (Scheme 25). Unfortunately, the C-H bond was so reluctant to


#### Scheme 25. Photocatalyzed Derivatization of Methane under Flow Conditions




undergo cleavage that acetonitrile competed in the HAT event. Accordingly, during the alkylation of dinitriles **25.1a-c**, a deuterated acetonitrile/water 7:1 mixture was mandatory to obtain a decent yield of methylated derivatives **25.2a-c**.<sup>404</sup> The functionalization of ethane and propane was likewise carried out under milder conditions, with no need of deuterated solvents.<sup>404</sup>

The addition of an alkyl radical onto an electron-poor olefin may ultimately lead to difunctionalization of the double bond thanks to a dual-catalytic approach, as shown in Scheme 26. TBADT was again used as PC<sub>HAT</sub> to generate an alkyl radical from cyclohexane. In this case, however, the adduct radical formed by addition of the cyclohexyl radical onto acrylate ester **26.2** was intercepted by a Ni<sup>0</sup> catalyst to form the alkyl-Ni<sup>I</sup>

## Scheme 26. Photocatalyzed Three-Component Difunctionalization of Alkenes



## Scheme 27. Photocatalyzed Derivatization of Electron-Rich Olefins



intermediate **26.3**. Oxidative addition of selected aryl derivatives **26.1a–d** onto **26.3** led to ester **26.4** in variable amounts depending on the leaving group X on the aromatic ring, with the bromine atom being the best choice. This approach showed a broad substrate scope since it may be applicable to several functionalized tertiary, secondary, and primary alkyl radicals.<sup>405</sup> A related approach was likewise devised by combining a POC (BP) with the same [Ni(dtbbpy)Br<sub>2</sub>] catalyst.<sup>406</sup>

Another recent example where HAT catalysis was merged with metal catalysis involved a Pd-catalyzed allylic alkylation. In this strategy, the alkyl radical was trapped by a Michael acceptor and the resulting carbanion interacted with *in situ* formed  $\pi$ -allylpalladium species that finally released the desired alkylation product.<sup>407</sup> The approach developed was then used for the concise synthesis of ( $\pm$ )-mesembrine.<sup>407</sup>

As a final note to this section, it is worth highlighting that the addition of photogenerated radicals onto olefins different from Michael acceptors (e.g., electron-rich C=C bonds or captodative olefins) intended for the formation of C(sp<sup>3</sup>)–C(sp<sup>3</sup>) bonds has only a few precedents in the literature. These processes, however, took place only in the presence of high PC<sub>HAT</sub> loadings or showed a low conversion of the starting materials.<sup>158,201,326</sup>


A representative example is shown in Scheme 27 where ethylene **27.1** was alkylated under TBADT photocatalysis to give **27.2** as the major product.<sup>326</sup> Such examples have not been presented in detail here due to their limited synthetic significance.

2.2. Formation of C(sp<sup>3</sup>)–C(sp<sup>3</sup>) Bonds via Addition onto C=X (X = N, O) Bonds

The formation of C(sp<sup>3</sup>)–C(sp<sup>3</sup>) bonds can be realized also via the addition of a photogenerated radical onto a C=X (X = N, O) double bond. In order to promote reactivity, the N-atom typically bears an electron-withdrawing S-based substituent,

either S(=O)<sub>2</sub>R or S(=O)R. Thus, N-tosylimines have been reported to act as excellent radical traps in TBADT-triggered alkylations with alkanes, ethers, and DMF. As reported in Scheme 28a, cyclohexane **28.1** (10 equiv) underwent addition

## Scheme 28. Photocatalyzed Addition of Hydrocarbons onto Imine Derivatives




onto the C=N bond of **28.2** to give the hydroalkylated adduct **28.3** in 85% yield in the presence of TBADT (2 mol %) upon irradiation with 400 nm LEDs (16 h). The occurrence of a chain mechanism (at least in part), however, could not be excluded.<sup>408</sup> Similarly, chiral N-sulfinyl imines were smoothly alkylated by adamantane scaffolds in the presence of a catalytic amount of PT (5 mol %) upon irradiation with 390 nm LEDs.

Notably, this strategy allowed the enantioselective synthesis of the saxagliptin core, containing an adamantyl-glycine motif.<sup>280</sup> In another instance, a dual-catalytic system based on **PT** and a chiral Cu-based complex containing a bisoxazoline (**BOX**) ligand allowed the regio- and stereoselective functionalization of benzylic, allylic, and even unactivated hydrocarbons with an imine derivative. As shown in **Scheme 28b**, toluene **28.6** (10 equiv) reacted with **28.7** to give product **28.9** in an excellent yield (93%) and enantioselectivity (e.e. 93%) in the presence of **PT** (2 mol %) and Cu(BF<sub>4</sub>)<sub>2</sub> (10 mol %) and chiral **BOX** ligand **28.8** (11 mol %).<sup>282</sup>

Very recently, the preparation of amines has been realized via a multicomponent carbonyl alkylative amination strategy. The protocol was promoted by **TBADT** (2 mol %) and comprised of *N*-arylamines, aldehydes, and hydrocarbons as starting materials. Slightly different conditions were required depending on the nature of the amine, being either an aniline or a diphenylamine. As shown in **Scheme 29**, the process

**Scheme 29. Multicomponent Synthesis of Secondary Amines**




involved the *in situ* formation of an iminium ion (**29.6<sup>+</sup>**), which acted as the trap of the photogenerated radical. When adopting cyclohexane **29.1**, benzaldehyde **29.2**, and anilines **29.3a-c**, secondary amines **29.4a-c** were obtained in good yields upon irradiation at 390 nm for 24 h, only requiring acetic acid (0.5 equiv) as an additive.<sup>409</sup>

The challenging addition of photogenerated intermediates onto C=O bonds has been realized only in a few instances. One notable example involved a strategy comprised of **TBADT** and a Cr<sup>III</sup> salt, where the role of the latter was to promote the formation of an organochromium compound via interception of the photogenerated radical. Indeed, this approach has been exploited to trigger the alkylation, aminomethylation, and oxymethylation of both aliphatic and aromatic aldehydes. Thus, *N,N*-dimethylacetamide **30.1** reacted with aldehydes **30.2a-c** to give 1,2-aminoalcohol derivatives **30.3a-c** in the presence of **TBADT** (10 mol %) and CrCl<sub>3</sub> (3 equiv) upon irradiation with 390 nm LEDs for 48 h (**Scheme 30**).<sup>410</sup>

### 2.3. Formation of C(sp<sup>3</sup>)-C(sp<sup>2</sup>) Bonds

This section describes the formation of C(sp<sup>3</sup>)-C(sp<sup>2</sup>) bonds between a photocatalytically generated C(sp<sup>3</sup>)-centered radical and suitable reaction partners, which include alkynes via an addition process or vinyl/aryl derivatives via a (formal) substitution or cross-dehydrogenative coupling reaction. Like-

**Scheme 30. Photocatalyzed Functionalization of C=O Bonds**




wise, the addition onto carbon monoxide (CO) or carbon dioxide (CO<sub>2</sub>) will be reported here.

Seminal works in the field focused on the addition of cycloalkyl radicals, obtained from the corresponding hydrocarbons, onto electron-poor alkynes in the presence of aromatic carbonyls. Although these PC<sub>S<sub>HAT</sub></sub> were routinely adopted stoichiometrically, the reaction was demonstrated to work smoothly also in the presence of a catalytic amount of **BP**. Thus, methyl propiolate **31.2** was functionalized by cyclopentane (**31.1**, used as the solvent) to give an *E/Z* mixture of vinylcycloalkanes **31.3** (**Scheme 31**) in a very good yield. **BP** loading could be lowered to 9 mol % without affecting the reaction yield; albeit, a longer irradiation time was required in the latter case.<sup>229</sup> Notably, the employed aromatic ketone could be supported onto a solid material (a polystyrene matrix or silica), rendering the PC<sub>HAT</sub> potentially recyclable. This heterogeneous variant has been shown to work to some extent under natural sunlight irradiation.<sup>411,412</sup> When applied to alcohols as substrates and dimethyl acetylenedicarboxylate as radical trap, this (heterogeneous) methodology opened the way to the generation of  $\alpha$ -hydroxyalkyl radicals and to the preparation of  $\gamma$ -butenolides from them.<sup>413</sup> Very recently, an analogous strategy based on the use of chloroalkynes (and, in selected cases, terminal alkynes) has been reported. **DCBP** (15 mol %) was used as the PC<sub>HAT</sub>, while the substrate scope included alcohols, ethers, amides, and even alkanes. Furthermore, when applied to THF, this process could be performed on the gram scale, without any significant yield decrease. Mechanistic studies revealed that this process occurred with a quantum yield >1, indicating the involvement of a radical chain mechanism.<sup>414</sup> Similarly, the functionalization of chloroalkynes to give functionalized vinyl chlorides has been likewise carried out in the presence of **EY**<sup>221</sup> and **TBADT**,<sup>278</sup> respectively.


Another option to forge a C(sp<sup>3</sup>)-C(sp<sup>2</sup>) bond is to intercept the photogenerated radical with an olefinic reaction partner containing a suitable radicofugal group. Thus, the alkenylation of ethers and amides with a library of vinyl sulfones smoothly occurred in the presence of **DCBP** (20 mol %) upon irradiation with CFL bulbs, wherein the loss of a sulfonyl radical occurred during the process. Thus, 2-pyrrolidone **32.1** (used as the solvent) reacted with sulfones **32.2a-c** to give the expected alkenylated amides **32.3a-c** in good yields and with a marked preference for the formation of the *E*-isomer (**Scheme 32**).<sup>389</sup>

The dehydrogenative coupling between alkanes and aryl alkenes is also possible and has been recently realized thanks to

Scheme 31. Photocatalyzed Addition of Cycloalkanes onto Alkynes




Scheme 32. Photocatalyzed Alkenylation of Amides




a dual-catalytic strategy. This approach relies on the synergistic combination of TBADT photocatalysis with cobaloxime-mediated hydrogen-evolution cross-coupling. The Co-catalyst is responsible for intercepting the radical adduct formed upon addition of the photogenerated radical onto the olefin and then undergoes a photoinduced  $\beta$ -hydride elimination, restoring the original double bond. As depicted in Scheme 33, a series of cycloalkanes (33.1a–d) was alkenylated by styrene 33.2 (10 equiv) in the presence of TBADT (4 mol %), Co(dmgH)(dmgH<sub>2</sub>)Cl<sub>2</sub> (1 mol %; dmgH<sub>2</sub> and dmgH: dimethylglyoxime and its monoanion), and 2,6-lutidine (10 mol %) as the ligand to deliver adducts 33.3a–d in good yields with complete regio- and stereoselectivity. The reaction took place in acetonitrile at 60 °C upon irradiation with a 370 nm LED and could be applied in the late-stage alkenylation of natural products, including steroid derivatives.<sup>415</sup>

Turning to arylation reactions, the preparation of alkylated pyrimidines was realized through the coupling of saturated heterocycles (including oxygen-, nitrogen-, and sulfur-based derivatives) with sulfonylated pyrimidines in the presence of BP (10 mol %). As an example, 5-membered heterocycles 34.1a,b were arylated by 34.2 to give pyrimidine derivatives 34.3a,b in good yields upon irradiation with a medium-pressure Hg lamp via an *ipso*-substitution process (Scheme 34).<sup>416</sup> Similarly, EY (2 mol %) was employed to promote the arylation of THF at the 2-position upon reaction with 2-phenylsulfonylbenzothiazole.<sup>221</sup> In a related instance, the 4-pyridination of cumene at the benzylic position was performed

Scheme 33. Dual-Catalyzed Dehydrogenative (*E*)-Alkenylation of Cycloalkanes

Scheme 34. Photocatalyzed Arylation of Five-Membered Heterocycles





in the presence of a catalytic amount of BP (10 mol %). In the process, the photogenerated radical added onto 4-cyanopyridine, while the desired product was formed upon loss of HCN from the initially formed adduct.<sup>230</sup> In a very recent report, *N*-aminopyridinium salts have been likewise used as radical traps for photogenerated C-centered radicals (AQ as the photocatalyst) and enabled the site-selective C–H pyridylation of unactivated alkanes. Notably, this protocol could be adopted for the late-stage site-selective functionalization of biorelevant compounds.<sup>417</sup>

The merging of HAT photocatalysis with Ni-catalysis opened new avenues on the route toward the arylation of (strong) aliphatic C–H bonds, allowing adoption of aromatic halides (mostly, bromides) as coupling partners. In particular, the Ni-based cocatalyst was responsible for activating the C(sp<sup>2</sup>)–Br bond and intercepting the photogenerated radical. This chemistry was successfully combined with different classes of PC<sub>S<sub>HAT</sub></sub>, including DT<sup>418</sup> and aromatic carbonyls.<sup>419,420</sup> Thus, cyclohexane 35.1 was functionalized by (hetero)aryl bromides 35.2a,b to deliver cross-coupled products 35.3a–b in very good yields upon irradiation with a 390 nm LED (Scheme 35). Of note, this protocol could be applied to the manipulation of natural products (see the case of 35.5).<sup>418</sup> Very recently, a dual-catalytic strategy based on DCBP and a Ni-based complex enabled the construction of C(sp<sup>3</sup>)–C(sp<sup>2</sup>) bonds via the acylation of methylbenzenes with *N*-acylsuccinimides.<sup>421</sup>





### Scheme 35. Arylation of Strong C–H Bonds via a TBADT/Nickel Dual-Catalyzed Strategy



Another opportunity for C(sp<sup>3</sup>)–C(sp<sup>2</sup>) bond construction is represented by photocatalytic cross-dehydrogenative couplings (CDC)<sup>422</sup> between aliphatic H-donors and (hetero)arenes. These processes require the adoption of oxidative conditions to remove the extra electrons, and this has been realized either by having recourse to a chemical oxidant or through electrochemical means. As for the former case, DT photocatalysis has been successfully exploited to trigger the functionalization of alkanes, ethers, and amides with heteroarenes in the presence of a persulfate salt. As an example, this Minisci-type reaction allowed the functionalization of quinaldine (36.2) with DMF (36.1) to give 36.3 as the only product (73% isolated yield) in the presence of TBADT (4 mol %) and K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> (2 equiv.) upon irradiation with simulated solar light (Scheme 36).<sup>423</sup> More recently, a similar strategy has been applied to the preparation of 2-alkylated benzothiazoles under chemical oxidant-free photoelectrochemical conditions.<sup>424</sup>


A different cross-dehydrogenative coupling encompassing the merging between HAT photocatalysis and electrochemistry allowed the regioselective functionalization of ethers with

### Scheme 36. Photocatalyzed Cross-Dehydrogenative Coupling between Amides and Heteroarenes



isoquinolines. This strategy, tagged “electrophotocatalysis”, was based on the use of a trisaminocyclopropenium ion (TAC<sup>+</sup>), which was electrochemically converted to the stable TAC<sup>•2+</sup> species via one-electron oxidation. The latter species then underwent excitation and, once in the excited state, triggered the desired HAT from the chosen ether. Thus, adducts 37.3a–d have been prepared by reaction between THF (37.1) and substituted isoquinolines 37.2a–d in the presence of TAC<sup>+</sup> (perchlorate salt; 1 mol %) upon application of a constant potential ( $E_{\text{cell}} = 1.5 \text{ V}$ ) under irradiation with a CFL (Scheme 37).<sup>210</sup>


### Scheme 37. Electrophotocatalytic Arylation of Ethers Mediated by the Trisaminocyclopropenium (TAC<sup>+</sup>) Ion



Carbon monoxide (CO) is an excellent radical trap and has been frequently exploited to get access to valuable acyl radicals.<sup>425–427</sup> Seminal examples of this chemistry within photocatalytic applications date back to the early 90s, when aromatic carbonyls were adopted to promote the carbonylation of alkanes (mainly, cyclohexane) to afford cyclohexanecarboxaldehyde either under high CO pressure (20–80 atm)<sup>428</sup> or in the presence of metal carbonyl complexes based on Ir, Rh, or Ru.<sup>429</sup> Around the same period, an 8% formation of cyclohexanecarboxaldehyde from cyclohexane and CO (1 atm) in the presence of TBADT was reported.<sup>327</sup> More recently, the functionalization of alkanes with electron-poor olefins under photocatalytic conditions mediated by TBADT was realized in the presence of CO. This allowed the preparation of unsymmetrical ketones in an atom-economical fashion in an overall multicomponent process, where the photogenerated alkyl radical was trapped by CO to form a C(sp<sup>3</sup>)–C(sp<sup>2</sup>) bond and then by the chosen electron-poor olefin. Thus, upon irradiation with a Xe lamp equipped with a Pyrex filter, 5- to 7-membered cycloalkanes 38.1a–c reacted with dibutyl maleate 38.3 under an atmosphere of CO (38.2, 80 atm) in the presence of TBADT (4 mol %) to afford ketones 38.4a–c in good isolated yields (Scheme 38).<sup>430</sup> Later, the same protocol was applied to the regioselective  $\beta$ -acylation of cyclopentanone in the role of H-donor,<sup>83</sup> as well as to the preparation of acyl hydrazides using diisopropyl azodicarboxylate (DIAD) in place of electron-poor olefins.<sup>431</sup>

In one instance, the carboxylation of the allylic position in simple alkenes by CO<sub>2</sub> has been realized in the presence of 3,6-diphenylxanthone and a Cu-based complex. The process has been proposed to occur through a sequence involving two independent steps, where activation of the allylic C–H bond was promoted by the excited carbonyl, while the copper complex operated the desired carboxylation, also restoring the initial ketone. Indeed, it was demonstrated that both the xanthone derivative and the copper complex behaved catalytically in the overall process.<sup>286</sup>


### Scheme 38. Three-Component Photocatalyzed Synthesis of Unsymmetrical Ketones



### 2.4. Formation of C(sp<sup>3</sup>)-C(sp) Bonds

The HAT-photocatalyzed C(sp<sup>3</sup>)-H to C(sp<sup>3</sup>)-C(sp) bond conversion can be related to two different families of processes, namely the introduction of an alkynyl or a cyano group. As for the first instance, the photogenerated radicals have been trapped by suitable alkynylating agents, namely alkynes substituted with a convenient radicofugal group. Indeed, only a handful of examples of this chemistry have been reported, which are based on the use of bromoalkynes,<sup>414</sup> alkynylbenziodoxolones,<sup>238</sup> or alkynylsulfones.<sup>278,389</sup> Either the aromatic ketone DCBP<sup>238,389,414</sup> or DT<sup>278</sup> was used as the PCs<sub>HAT</sub>. As an example, THF (39.1, used as the solvent) reacted with bromoalkynes 39.2a-c in the presence of DCBP (15 mol %) and KOAc (1.5 equiv) to give alkynes 39.3a-c in good isolated yield, independently from the electronic character of the aromatic substituent in 39.2a-c (Scheme 39). In the


### Scheme 39. Photocatalyzed Alkynylation of Ethers by Bromoalkynes



process, the formation of a vinyl bromide intermediate initially takes place, which then undergoes HBr elimination to give the desired alkynylated product aided by the employed base (KOAc).<sup>414</sup>

Turning to cyanation processes, an excellent option to intercept the photogenerated radical is using tosyl cyanide, which allows introduction of the desired cyano group via displacement of the sulfonyl moiety. Thus, BP can be successfully adopted to trigger this transformation in a variety of substrates, including ethers, alkanes, and nitrogen-containing substrates; however, only in the latter case it behaves as a real PC<sub>HAT</sub>. Thus, protected nitrogen-heterocycles 40.1a-c were cyanated by tosyl cyanide 40.2 (2 equiv) to give 40.3a-c in good to excellent yield in the presence of BP (20 mol %) and 2,6-di-*tert*-butylpyridine (4 equiv; functioning as an acid scavenger) upon irradiation with a medium-pressure Hg lamp (Scheme 40). Worthful of notice is the example related to 40.1c, wherein the functionalization of the  $\alpha$ -to-N position occurred

### Scheme 40. BP-Photocatalyzed Cyanation of Nitrogen-Containing Heterocycles




chemoselectively.<sup>231</sup> Similarly, EY (2 mol %) has been adopted in the C-H to C-CN conversion in 1,4-dioxane in the presence of tosyl cyanide.<sup>221</sup> A different strategy is based on the use of a seven-coordinated (chiral) Ur salen complex (2 mol %), which was used for the cyanation of a variety of (substituted) *N,N*-dimethylanilines under oxidative conditions (H<sub>2</sub>O<sub>2</sub>) in the presence of NaCN and AcOH.<sup>432</sup>

### 2.5. Formation of C(sp<sup>2</sup>)-C(sp<sup>3</sup>) Bonds

This section gathers examples enabling the formation of a C(sp<sup>2</sup>)-C(sp<sup>3</sup>) bond via the intermediacy of a photogenerated C(sp<sup>2</sup>)-hybridized radical. Specifically, either aldehydes or formamides (see also section 2.1.2) can be exploited as H-donors in the formation of acyl and carbamoyl radicals, respectively. Seminal works in the area involved the use of BP to trigger the acylation of enones (mainly carbohydrate enones, see also Scheme 7 for the analogous C(sp<sup>3</sup>)-C(sp<sup>3</sup>) bond formation)<sup>234</sup> and  $\alpha,\beta$ -unsaturated esters or acids<sup>235</sup> with aldehydes. An interesting example is reported in Scheme 41,


### Scheme 41. Photocatalyzed Acetylation of Crotonic Acid



showing the hydroacylation of crotonic acid 41.2 with acetaldehyde 41.1 to give 4-oxoalkanoic acid 41.3 (60% yield) in the presence of BP (10 mol %) upon irradiation at 366 nm for 24 h.<sup>235</sup>

More recently, given their excellent reactivity as H-donors, aldehydes have been adopted as substrates in combination with a plethora of PC<sub>S<sub>HAT</sub></sub> to perform the hydroacylation of a huge variety of unsaturated systems. One of the most studied systems involves the functionalization of electron-poor olefins ( $\alpha,\beta$ -unsaturated esters, ketones, and nitriles, as well as vinyl sulfones) triggered by TBADT.<sup>383,433,434</sup> As an example, 2-cyclohexenone 42.1 was smoothly acylated by both hydrocinnamaldehyde 42.2a and *p*-anisaldehyde 42.2b to give interesting 1,4-diketones 42.3a,b in a good yield (Scheme 42). Thus, the optimal TBADT loading was 2 mol %, but contrary to aliphatic aldehydes, a slight excess of 42.1 (1.2 equiv), a longer irradiation time (30 vs 24 h), and an increased light intensity were required in the preparation of 42.3b.<sup>433,434</sup>

### Scheme 42. Cyclohexenone Acylation by Addition of Aliphatic (Upper Part) and Aromatic (Lower Part) Aldehydes




Of note, these reactions have been demonstrated to occur under natural sunlight irradiation by simply exposing the reaction vessel containing the mixture on a window ledge for a few days. In the acylation of dimethyl maleate with heptanal, it was possible to increase the concentration of the starting materials up to 0.5 M, therefore reducing the amount of solvent needed and bringing about important ecological advantages.<sup>277</sup>

The environmental performance of these acylations was further ameliorated making use of continuous flow conditions.<sup>382</sup> The adoption of this operation mode also allowed design of multistep procedures, wherein the photocatalytic C(sp<sup>2</sup>)-C(sp<sup>3</sup>) bond formation was followed by additional thermal steps on the resulting acylated derivatives.<sup>435,436</sup>


An elegant one-pot protocol comprised of two distinct photochemical steps was adopted for the preparation of homoallyl ketones starting from cyclopentanones and electron-poor olefins. The sequence involved an initial Norrish type-I photoinduced fragmentation of the 5-membered ring to give a 4-pentenal derivative followed by the TBADT-photocatalyzed hydroacylation of an electron-poor olefin. Thus, cyclopentanone **43.1** underwent ring opening via photoinduced cleavage of the C<sub>1</sub>-C<sub>2</sub> bond to give aldehyde **43.2**. Next, TBADT (4 mol %) and the chosen electron-poor olefins (e.g., **43.3a,b**) were added to the crude mixture, finally affording the desired adducts (**43.4a,b**) in a good yield upon irradiation for an additional 24 h (Scheme 43).<sup>437</sup>

### Scheme 43. Preparation of Homoallyl Ketones from Cyclopentanones via a Two-Step Photochemical Norrish Type-I Cleavage/Photocatalyzed Hydroacylation Sequence



Apart from TBADT, other PC<sub>S<sub>HAT</sub></sub> have been recently reported to promote the hydroacylation of electron-poor olefins under visible light irradiation, including EY,<sup>221</sup> UrN,<sup>284</sup> and Sb-Oxo.<sup>266</sup> Scheme 44 gathers selected examples describing the hydroacylation of benzylidene malononitrile **44.1**. Thus, EY and UrN allowed the hydroacylation of **44.1** with hexanal **44.2a** and heptanal **44.2b** to give the corresponding adducts **44.3a** and **44.3b** in 84 and 93%

### Scheme 44. Hydroacylation of Benzylidene Malononitrile Triggered by Different PC<sub>S<sub>HAT</sub></sub>



isolated yield, respectively (Schemes 44a,b).<sup>221,284</sup> On the other hand, the preparation of **44.3b** in the presence of Sb-Oxo proceeded with a partial conversion of the starting materials; however, an almost quantitative yield based on remaining starting material (99% brsm) was observed (Scheme 44c).<sup>266</sup>

EY has been further tested for the asymmetric synthesis of 1,4-dicarbonyls. This strategy encompassed a dual-catalytic system, which also involved the use of a chiral rhodium catalyst responsible for coordinating the chosen electron-poor olefin (an unsaturated *N*-acylpyrazole) and driving the radical addition step (for a related example, see Scheme 19d).<sup>391</sup>


Very recently, the formation of a C(sp<sup>2</sup>)-C(sp<sup>3</sup>) bond by the EY-photocatalyzed reaction of aldehydes with *N*-(hetero)-arylsulfonyl propiolamides has been proposed. This transformation led to the preparation of the isothiazolidin-3-one 1,1-dioxide core and proceeded through a cascade involving addition of the photogenerated radical onto the C≡C triple bond of propiolamide, Smiles rearrangement, and 5-*endo*-trig cyclization.<sup>438</sup> As an example, *N*-heterocycles **45.3a-c** have been prepared from aldehydes **45.1a-c** and amide **45.2** upon irradiation with blue light for 48 h in the presence of EY (4 mol %, Scheme 45). The preparation of **45.3c** has been successfully realized on a gram scale in 83% yield, and of note, some of the synthesized compounds may have potential anticancer activity.<sup>438</sup>

Apart from the use of electron-poor olefins, (hetero)-aromatic alkenes can be used as well as radical traps for the photogenerated acyl radicals, with the driving force for the process being the formation of a stabilized benzyl radical. Thus, different combinations of the PC<sub>HAT</sub> and alkenes have been adopted, including vinylpyridines,<sup>387</sup>  $\alpha$ -trifluoromethyl aryl alkenes (in this case, an aliphatic trifluoromethylalkene has been used as well),<sup>439</sup> and aryl alkenes.<sup>379,440</sup> Scheme 46 gathers the case of benzaldehyde **46.1**, that has been adopted for the functionalization of 2-vinylpyridine **46.2a**, CF<sub>3</sub>-substituted alkene **46.2b**, styrene **46.2c** (in all cases TBADT as the PC<sub>HAT</sub>), and *p*-fluorostyrene **46.2d** (EY as the PC<sub>HAT</sub>).


Finally, *N*-tosyl imines (for a related example, see Scheme 28a)<sup>408</sup> and dehydroalanine derivatives<sup>441</sup> have been likewise adopted as acyl radical traps under TBADT-mediated photocatalytic conditions.

Along the same line, a dual-catalytic strategy comprised of TBADT and a Ni-based cocatalyst allowed the asymmetric acyl-carbamoylation of alkenes starting from aldehydes and a

## Scheme 45. Photocatalyzed Preparation of Isothiazolidin-3-one 1,1-Dioxides



## Scheme 46. Photocatalyzed Hydroacylation of Vinyl (Hetero)aromatics




carbamoyl chloride incorporating a C=C double bond. In this case, the Ni-based cocatalyst intercepted the photogenerated acyl radical and triggered the activation of the carbamoyl chloride, supervising the sequence of steps leading to the formation of the final product. Thus, butanal 47.1 reacted with aryl carbamic chlorides 47.2a–c in the presence of TBADT (5 mol %), Ni(OTf)<sub>2</sub> (10 mol %), ligand 47.4 (12 mol %), and K<sub>3</sub>PO<sub>4</sub> (1.1 equiv). The process took place in MeCN upon irradiation with a 390 nm LED for 9 h, delivering oxindoles 47.3a–c in good yields and excellent enantioselectivity (Scheme 47).<sup>442</sup>

A similar reaction system, comprised of TBADT and a Ni-based cocatalyst, enabled the cross-coupling between acyl radicals, photogenerated from aldehydes, and (fluorinated)  $\alpha$ -bromoacetates. As reported in Scheme 48, 1,3-dicarbonyl derivatives 48.3a–c were readily accessed upon reaction between *p*-anisaldehyde 48.1 and esters 48.2a–c.<sup>443</sup>

As mentioned above, apart from aldehydes, formamides can be likewise adopted as H-donors for the generation of carbamoyl radicals. These, in turn, are interesting intermediates for C(sp<sup>2</sup>)–C(sp<sup>3</sup>) bond formation campaigns.<sup>158</sup> Thus, only a handful of PC<sub>S</sub>HAT have been employed to trigger this reactivity, including TBADT<sup>382,383,386,387,390</sup> and UrN.<sup>284</sup> In the latter case, however, this chemistry represented a minor

## Scheme 47. Asymmetric Acyl-Carbamylation of Alkenes



pathway with respect to the preferred  $\alpha$ -to-N C–H cleavage (see also section 2.1.2).<sup>284</sup>


On one hand, TBADT has been shown to cleave chemoselectively the C(=O)–H bond in primary and secondary formamides, while a completely different reactivity has been observed with tertiary formamides, (e.g., DMF, see Scheme 19b).<sup>386</sup> As shown in Scheme 49, methyl crotonate 49.2 has been successfully carbamoylated by formamide 49.1a and *N*-methyl formamide 49.1b in the presence of TBADT (2 mol %) upon irradiation with phosphor-coated lamps centered at 310 nm. Indeed, the former H-donor led to the formation of product 49.3a in a higher yield (76%) than the latter (49.3b, 46% yield). The carbamoylation of electron-poor olefins has been successfully implemented under continuous flow conditions, delivering the desired products in shorter reaction times and increased productivity.<sup>382,390</sup> More recently, the same reactivity has been likewise applied to the carbamoylation of vinylpyridines<sup>387</sup> and styrenes<sup>379</sup> (in the latter case, in the presence of a disulfide cocatalyst). In sharp contrast, UrN enabled the functionalization of the C(=O)–H bond even when using DMF as substrate, being a competitive path to the usual  $\alpha$ -to-N functionalization.<sup>284</sup>

2.6. Formation of C(sp<sup>2</sup>)–C(sp<sup>2</sup>) Bonds


As in the previous section, the formation of C(sp<sup>2</sup>)–C(sp<sup>2</sup>) bonds made use of aldehydes or formamides as radical precursors. The photogenerated radicals, however, are here used in the addition onto C $\equiv$ C triple bonds or in (formal) substitution reactions or cross-dehydrogenative couplings with substituted arenes or alkenes.



## Scheme 48. Synthesis of 1,3-Dicarbonyl Derivatives via a Dual-Catalytic Strategy





## Scheme 49. TBADT-Photocatalyzed Carbamoylation of Electron-Poor Olefins





Early examples described the acylation of quinones in the presence of BP,<sup>232,233</sup> in what has been tagged as “photo-Friedel–Crafts acylation”.<sup>233</sup> Albeit the reaction proceeded to some extent also in the absence of BP, its addition in catalytic quantities allowed an improvement of the product yield. As depicted in Scheme 50, 1,4-naphthoquinone **50.2** reacted with *ortho*-substituted benzaldehydes **50.1a–c** to afford hydroquinones **50.3a–c** in a good yield upon irradiation with a high-pressure mercury lamp for 5 days in the presence of BP (6 mol %).<sup>232</sup>

## Scheme 50. Photocatalyzed Acylation of 1,4-Naphthoquinone



More recently, the construction of the 3-acyl-4-aryl coumarin scaffold has been realized upon photocatalyzed addition of aldehydes (e.g., **51.1a–c**) onto an aromatic ynoate (**51.2**) triggered by an anthraquinone derivative (Scheme 51). Thus,



## Scheme 51. Photocatalyzed Acylation of Ynoates on the Route to Coumarin Scaffolds



the photogenerated acyl radical added regioselectively onto the  $\alpha$ -position of the ynoate to give the vinyl radical **51.4**<sup>•</sup>, which cyclized onto the tethered aromatic ring followed by rearomatization to give coumarins **51.3a–c** in good yields. The reaction was promoted under visible light irradiation by adopting *t*BAQ (10 mol %) and benzoyl peroxide (BPO) as a stoichiometric oxidant. A possible role of the carboxyl radical resulting from BPO decomposition in the desired C–H cleavage has been proposed. Of note, the biological activity of some of the synthesized coumarins has been studied, suggesting their efficacy as potential candidates for new therapeutics.<sup>444</sup>

Very recently, AQ (1 mol %) has been used as the PC<sub>HAT</sub> to promote the C(=O)–H pyridylation of a small library of aldehydes and formamide with *N*-aminopyridinium salts. As an example, Scheme 52 depicts the reaction between aldehydes


## Scheme 52. Photocatalyzed Preparation of Pyridyl Ketones



**52.1a,b** and **52.2**, which took place upon visible light irradiation for 48 h, to deliver acylated pyridines **52.3a,b**. In the process, AQ triggered the formation of acyl radicals **52.4**<sup>•</sup>, which underwent addition onto the pyridinium derivative in a regioselective fashion and ultimately gave the desired products upon deprotonation and release of sulfonamidyl radical **52.5**<sup>•</sup> via N–N bond cleavage. Indeed, the possible involvement of sulfonamidyl radical **52.5**<sup>•</sup> in a chain pathway has been likewise proposed.<sup>417</sup>

Another opportunity for C(sp<sup>2</sup>)–C(sp<sup>2</sup>) bond formation is the cross-coupling between aldehydes and aromatics. Most of the reported routes rely on the adoption of a dual-catalytic strategy comprised of a PC<sub>HAT</sub> (TBADT<sup>228,443</sup> or an aromatic ketone<sup>228</sup>) and a transition-metal cocatalyst (based on Ni<sup>442</sup> or Pd<sup>227</sup>), responsible for intercepting the photogenerated acyl radical and activating the aromatic derivative. Scheme 53a describes the arylation of cyclohexanecarboxaldehyde **53.1** with 4-bromotoluene **53.2**. Interestingly, two different PC<sub>HAT</sub> have been used in the preparation of **53.3**, either TBADT (2 mol %) or AQ (10 mol %), in the presence of the same cocatalyst, viz. Pd(OAc)<sub>2</sub> (5 mol %) complexed with the Xantphos ligand (5 mol %). Notably, while the former system


### Scheme 53. Photocatalyzed Preparation of Aromatic Ketones and Benzamides



required UV light irradiation (390 nm), the latter worked upon visible light exposure (427 nm). Apart from aryl bromides, aryl iodides and triflates have been demonstrated to be competent substrates as well. One example also demonstrated that *N*-methyl formamide **53.4** could be used instead of aldehydes, enabling the carbamoylation of aryl bromide **53.5** to give benzamide **53.6** (Scheme 53b).<sup>228</sup> Very recently, a dual-catalytic approach based on **PQ** and a Pd-based cocatalyst enabled the acylation of arenes via a double C–H activation strategy. The process took place under visible light irradiation in the presence of Ag<sub>2</sub>O as the terminal oxidant.<sup>445</sup> In another instance, the net-oxidative cross-coupling between heptanal and quinaldine in the presence of **TBADT** (4 mol %) and K<sub>2</sub>S<sub>2</sub>O<sub>8</sub> (2 equiv) has been described (see Scheme 36 for a related process).<sup>423</sup> Also in this case, the aldehyde could be substituted with *N*-methyl formamide as the H-donor to unlock quinaldine carbamoylation.<sup>423</sup>

Apart from aromatic derivatives, the formation of C(sp<sup>2</sup>)–C(sp<sup>2</sup>) bonds starting from aldehydes and alkenes has been reported. Only a couple of examples of this chemistry are available in the literature, both making use of dual-catalytic strategies. The first one enabled the cross-dehydrogenative coupling between aldehydes and alkenes, based on the use of **TBADT** and a Co-based cocatalyst. Thus, adopting the same protocol described in Scheme 33, hexanal **54.1** reacted with styrene **54.2** to give the  $\alpha,\beta$ -unsaturated ketone **54.3** in a good yield (Scheme 54). Notably, the process occurred in a

### Scheme 54. Dual-Catalytic Cross-Dehydrogenative Coupling between Aldehydes and Aryl Alkenes




stereoselective fashion, affording exclusively the *E*-isomer.<sup>415</sup>

In one instance, the alkenylation of cyclohexanecarboxaldehyde with  $\beta$ -bromostyrene has been likewise realized through the merging of **TBADT** photocatalysis with Pd-catalysis (see also Scheme 53).<sup>228</sup>

### 2.7. Formation of C(sp<sup>2</sup>)–C(sp) Bonds

Very recently, an example of C(sp<sup>2</sup>)–C(sp) bond formation has been reported, making use of a SOMOphilic alkylation protocol. As shown in Scheme 55, heptanal **55.1** reacted with alkynyl sulfone **55.2** to give ynone **55.3** in a good isolated yield (67%) upon irradiation with a 390 nm LED in the presence of **TBADT** (2 mol %).<sup>278</sup>

### Scheme 55. TBADT-Photocatalyzed Alkylation of Aldehydes



### 2.8. Functionalization of Carbon Nanostructures


Fullerene is one of the most investigated nanomaterials nowadays, and this is mainly due to its versatility in supramolecular chemistry,<sup>446,447</sup> in technological applications,<sup>448,449</sup> as well as in medicinal chemistry and nanotechnology.<sup>450</sup> One of the most convenient ways to edit the fullerene structure, taking advantage of the high number of unsaturations, is by radical addition, wherein fullerene works as a radical sponge.<sup>451</sup> However, the challenging aspect of this chemistry stems from the tendency of C<sub>60</sub> to give a complex mixture of multiadducts. This reactivity could be tamed by having recourse to DT-photocatalyzed HAT;<sup>452–454</sup> albeit, only in one case the PC<sub>HAT</sub> could be used in a 20 mol % loading. Therein, the functionalization of fullerene C<sub>60</sub> with acyl radicals generated from aromatic,  $\alpha,\beta$  unsaturated and aliphatic aldehydes was reported.<sup>455</sup> The mechanism is based on a fast radical–radical anion coupling between the acyl radical and the fullerene radical anion generated concomitantly with the recovery of the PC<sub>HAT</sub>. Intriguingly, it was found that when a pivaloyl or phenylacetyl radical was generated (for both intermediates the decarbonylation rate is >10<sup>5</sup> s<sup>–1</sup>), CO loss occurred already at 10 °C. Notably, the decarbonylation step could be suppressed and the corresponding acylated fullerenes were synthesized with great selectivity by decreasing the temperature down to –40 °C. In fact, when pivalaldehyde (**56.1a**) or phenylacetaldehyde (**56.1b**) was subjected to the optimized reaction conditions at 10 °C, a mixture of **56.3** and **56.4** was obtained in the ratio 1:1.8 or 1:2, respectively. However, by decreasing the temperature, the ratio was completely reversed to 12:1 for **56.1a**, while for **56.1b** the decarbonylation product was not even observed (Scheme 56).

In another instance, the DT-photocatalyzed PEGylation of carbon nanotubes was reported.<sup>456,457</sup> In particular, when a solution of single-walled carbon nanotubes was irradiated in the presence of **TBADT** in a PEG400/MeCN solution, the grafting occurred smoothly in 48 h, albeit with modest efficiency (20 wt %, 1 PEG400 chain every 120 carbon atoms). Interestingly, it was shown that sunlight could be used as an inexpensive light source and afforded the PEGylated carbon nanotubes in a comparable yield.<sup>456,457</sup>

### 3. FORMATION OF C–Y BONDS (Y ≠ C)

Besides the formation of C–C bonds, photocatalyzed HAT has been successfully employed to forge C–Y bonds. Specifically, this section gathers synthetic examples dealing with the formation of C–N, C–O, C–S, C–F, and C–Cl bonds.

### Scheme 56. Functionalization of Fullerene-C60 via Addition of Photogenerated Acyl Radicals




Finally, the adoption of *d*-HAT for the deuteration of organic molecules is presented.

### 3.1. Formation of C–N Bonds

Diisopropyl azodicarboxylate (DIAD) is one of the most widely adopted unsaturated compounds to forge C–N bonds via photocatalyzed HAT. Thus, its remarkable electrophilic character makes DIAD an excellent trap for nucleophilic radicals such as alkyl, acyl,  $\alpha$ -oxyalkyl, and  $\alpha,\alpha$ -dioxyalkyl. Upon addition of these nucleophilic radicals onto the N=N double bond, the corresponding N-centered radical is generated, which is typically entrusted for the recovery of the spent PC<sub>HAT</sub>.

In one instance, DIAD was exploited for the synthesis of hydrazides, as shown in Scheme 57.<sup>431</sup> Therein, TBADT was

### Scheme 57. TBADT-Photocatalyzed Synthesis of Hydrazides




used for the functionalization of (cyclo)alkanes (e.g., cyclohexane 57.1), ethers, and acetals via homolytic C–H bond cleavage. The photogenerated C-centered radicals were trapped by DIAD to deliver the corresponding hydrazides (e.g., 57.3) in good to excellent yields. Interestingly, aldehydes were competent substrates as well, allowing the synthesis of acyl hydrazides through the formation of a C(sp<sup>2</sup>)-N bond.<sup>431</sup>

The synthesis of hydrazides, including adduct 57.3, was also reported under continuous-flow conditions. In such case, the photochemical reactor (*V* = 50 mL) consisted of a water-cooled 500 W medium-pressure Hg-vapor lamp wrapped in FEP (fluorinated ethylene propylene) tubing. When a MeCN solution of THF (58.1), DIAD, and TBADT (0.4 mol %) was flown through the tubing, the labile  $\alpha$ -to-O C(sp<sup>3</sup>)-H bond of 58.1 was cleaved, thus delivering the corresponding  $\alpha$ -oxyalkyl radical and, ultimately, product 58.3 in 73% yield upon

reaction with 58.2 (Scheme 58). Interestingly, this reaction showed an excellent process mass intensity (PMI, i.e. the ratio

### Scheme 58. TBADT-Photocatalyzed Formation of a C–N Bond in Flow



of the total mass of materials to the mass of isolated product) of 9.36 kg kg<sup>-1</sup> and a specific productivity (SP, i.e. mmol of product formed with respect to the energy consumed) of 22 mol W<sup>-1</sup> h<sup>-1</sup>.<sup>382</sup>


A similar approach was reported for the one-pot synthesis of hydroxamic acids from aldehydes.<sup>246</sup> In detail, acyl radicals were generated via HAT by PGA (10 mol %) and trapped by DIAD. The resulting acylhydrazides (e.g., 59.3) were then converted in a one-pot fashion to the corresponding hydroxamic acids via nucleophilic acyl substitution with hydroxylamine. For example, propanal (59.1) was smoothly converted to hydroxamic acid 59.4 in 78% yield over two steps (Scheme 59).

This approach was also extended to the synthesis of amides by employing primary and secondary amines in the place of hydroxylamine.<sup>458</sup> The validity and the robustness of the protocol were demonstrated through the preparation of Moclobemide (~30% yield over two steps), a drug used against depression and social anxiety, starting from 4-chlorobenzaldehyde.<sup>458</sup>


A conceptually different route for the formation of C–N bonds consists in the synthesis of organic azides. This approach is particularly appealing since these compounds are widely used in medicinal chemistry for biorthogonal labeling via the Cu-catalyzed Huisgen reaction. In one instance, photocatalyzed HAT has been used to introduce the N<sub>3</sub> group on leucine in the synthesis of Manzacidin C.<sup>459,460</sup> The first step of the synthetic route consisted in the TBADT-promoted activation of the methine site of the amino acid (60.1); ensuing quenching of the photogenerated radical with sulfonyl azide 60.2 readily afforded the hoped-for azide 60.3 in 49% yield (Scheme 60).

Recently, the azidation of C(sp<sup>3</sup>)-H bonds via HAT has been achieved through an electrophotocatalytic strategy, where DMBP (5.5 mol %) was used as PC<sub>HAT</sub> and a Mn-salt as the cocatalyst.<sup>461</sup> In detail, compound 61.1 was activated via HAT to afford the corresponding (stabilized) benzyl radical. In the meanwhile, the manganese ion (from MnF<sub>2</sub>) was complexed by the added ligand (1,10-phenanthroline) and the azide anion to afford a Mn<sup>III</sup>/L–N<sub>3</sub> complex after anodic oxidation. The interaction between the benzyl radical and the latter species afforded the hoped-for organic azide and the Mn<sup>II</sup> ion, prone to start a new catalytic cycle. It is however important to mention that the reaction proceeded to a certain extent (>40%) even without the photocatalyst, suggesting that a competitive direct anodic oxidation of the N<sub>3</sub><sup>-</sup> anion might be likewise responsible for the observed reactivity. Notably, the reaction could be run on a gram scale and product 61.3 was obtained in 71% yield (1.36 g, Scheme 61).



## Scheme 59. One-Pot Synthesis of Hydroxamic Acids from Aldehydes



## Scheme 60. Azidation of the Methine Site of Leucine




## Scheme 61. Electrophotocatalytic Azidation via a DMBP/Mn Dual-Catalytic System




Very recently, a different concept for the C–H to C–N bond conversion based on radical-polar crossover (RPC) has been disclosed. RPC opens the world of polar chemistry to radicals and enables a whole new range of synthetic possibilities for radical chemistry.<sup>462,463</sup> More specifically, HAT was coupled with a subsequent chemical oxidation of the first-formed C-centered radical to afford a carbocation. The latter intermediate could be conveniently trapped with *N*-heteroaryl-based nucleophiles, thus establishing the targeted C–N bond.<sup>464</sup> Thus, when an acetonitrile solution of tetrahydrofuran (62.1, 6 equiv.), pyrazole (62.2), TBADT (5 mol %), and TBHP (*tert*-butyl hydroperoxide, 3 equiv.) was irradiated with a UV LED (365 nm) for 16 h, product 62.3 was formed in 86% <sup>1</sup>H NMR yield. The reaction was made more efficient by adopting a flow apparatus, which allowed reduction of reaction time to just 1 h giving a similar result (86% <sup>1</sup>H NMR yield, 81% isolated yield), even though 18 equiv. of 62.1

was required (Scheme 62). Notably, the reaction remained efficient when performed on a 10 mmol scale (80% yield) and

Scheme 62. C(sp<sup>3</sup>)–H Heteroarylation via Radical-Polar Crossover

Notable examples:



the synthetic approach could be employed for the late-stage functionalization of biologically active molecules such as acetylated podophyllotoxin (62.4) and stanozolol (62.5).<sup>464</sup>

## 3.2. Formation of C–O Bonds

The formation of C–O bonds is a widely investigated transformation among those that can be carried out via photocatalyzed HAT. In these manifolds, the C-centered radical generated via HAT is typically intercepted by a sacrificial oxidant, such as molecular oxygen (O<sub>2</sub>). The first-formed oxygenated intermediate is a hydroperoxyl radical, which can be isolated as the corresponding hydroperoxide or *in situ* converted to alcohols, ketones, and even carboxylic acids. Besides its operational simplicity, this transformation has remarkable commercial and industrial implications. For



example, when using cyclohexane as the starting material, a mixture of cyclohexanol and cyclohexanone is obtained, the so-called KA-oil (ketone-alcohol mixture).<sup>465</sup> The latter has an immense value in modern industry, since it is used for the synthesis of adipic acid, in turn needed to produce Nylon-6,6.<sup>466</sup>

In view of the above, several research groups have designed photocatalytic systems able to perform this transformation in a sustainable way and carried out meticulous mechanistic studies. Most of them rely on the use of metal-oxo compounds (e.g., the DT or Ur ions) or POCs (e.g., aromatic ketones) under aerobic conditions. Homogeneous conditions are routinely adopted,<sup>202,315,473–477,334,339,467–477</sup> albeit, heterogeneous systems have been reported as well.<sup>289,478,479</sup>

Hereby, the synthetic applications of this chemistry have been reviewed and classified according to the aimed-for product, namely hydroperoxides, alcohols, and ketones or carboxylic acids.

**3.2.1. Synthesis of Hydroperoxides.** A seminal report on the synthesis of hydroperoxides via photocatalyzed HAT was published in the 1980s and focused on the oxidation of isobutane relying on the use of polyoxometalates as PC<sub>S<sub>HAT</sub></sub>.<sup>201,324</sup> In such a case, an acetonitrile solution of the light alkane was irradiated in the presence of TBADT (4 mol %) to afford *tert*-butylhydroperoxide in a quantitative yield (at 55% conversion of isobutane) after 2 h of irradiation with a 1000 W Xe lamp.<sup>324</sup> In another instance, the hydroperoxidation of benzylic C–H bonds by using EY has been reported.<sup>251</sup> The irradiation of an acetonitrile solution of **63.1** (0.2 M) under an O<sub>2</sub>-atmosphere with a blue LED in the presence of the organic dye (2 mol %) delivered compound **63.2** in 80% isolated yield (Scheme 63). Notably, the same reaction was run on a gram scale and 1.18 g of **63.1** was smoothly converted into 0.95 g of **63.2** (68% yield) after 15 h of irradiation.<sup>251</sup>

**3.2.2. Preparation of Alcohol and Ketone Mixtures.** Under certain circumstances, hydroperoxyl radicals undergo spontaneous disproportionation to yield a mixture of the


corresponding alcohol and ketone.<sup>480</sup> This chemistry has been exploited in several instances, especially with cyclohexane as the substrate and TBADT as the PC<sub>HAT</sub>, to obtain a mixture of cyclohexanol and cyclohexanone.<sup>323,481–488</sup>

Thus, when a TBADT solution ( $2 \times 10^{-4}$  M in CH<sub>2</sub>Cl<sub>2</sub>/C<sub>6</sub>H<sub>12</sub>/CH<sub>3</sub>CN 6:3:1) was irradiated at 325 nm with a 250 W Xe lamp (equipped with a grating monochromator), the formation of the KA-oil was observed with a quantum yield of 0.35.<sup>481</sup> It is important to mention that a great influence of the oxygen pressure on the final product distribution was observed. In fact, while the formation of the alcohol was not affected when increasing O<sub>2</sub> pressure from ~0.03 to 1 atm, the percentage of ketone in the mixture increased considerably. Thus, while at 0.03 atm the alcohol/ketone ratio was found to be 7:3, at 1 atm the ratio was fully reversed (ca. 3:7).<sup>481</sup> As stated by the authors, the distribution of the final products depends on a delicate interplay of the involved radical species. In particular, at low oxygen concentration, the reduced photocatalyst interacts with O<sub>2</sub> to afford H<sub>2</sub>O<sub>2</sub> and eventually OH•. Radical recombination of the latter intermediate with the photocatalytically generated cyclohexyl radical affords cyclohexanol. In such a scenario, the formation of the ketone is proposed to be a marginal phenomenon. Conversely, at high pressure of oxygen, different pathways are stimulated, such as radical trapping of cyclohexyl radical by triplet oxygen to get the corresponding hydroperoxyl intermediate. The decomposition of such an intermediate, as also mentioned above, leads to the formation of a mixture of ketone and alcohol through radical chain autoxidation, with a preference for the former.


**3.2.3. Synthesis of Carbonyl Derivatives.** As mentioned in the previous paragraph, when TBADT-photocatalyzed HAT is carried out under an O<sub>2</sub>-rich atmosphere, the selectivity can be diverted toward the formation of the C=O double bond. The direct, 4-electron oxidation of a CH<sub>2</sub> group into ketones has been achieved on a preparative scale by combining the use of TBADT (5 mol %) and O<sub>2</sub> with a flow technology (V = 5 mL; PFA tubing,  $\phi_{in}$  = 750  $\mu$ m).<sup>276</sup> Thus, a MeCN/1 M HCl 2.5:1 solution of cyclohexane (**64.1**, ~0.1 M) and molecular oxygen were pumped through a PFA tubing coiled around a 3D-printed PLA cylinder and the mixture was irradiated with UV LEDs ( $\lambda$  = 365 nm). As a result, a cyclohexanone/cyclohexanol mixture (**64.2/64.2'** 9:1) was obtained in 90% overall yield from cyclohexane, outperforming the results obtained in batch. Interestingly, this methodology could be extended to natural scaffolds (**64.3–64.5**) and, for instance, allowed us to realize the oxidation of artemisinin to 9-artemisitonone **64.3** with great efficiency and selectivity (59%, gram-scale) in a MeCN/DCM mixture (acid was omitted, Scheme 64).

In another instance, the use of NaDT for the remote oxidation of aliphatic amines was reported.<sup>112</sup> Typically, protected amines undergo photocatalyzed HAT only at the  $\alpha$ -to-N position. Building upon previous reports, however, it was envisaged that the protonation of amines could divert the HAT reactivity toward remote C–H bonds<sup>99</sup> (see also the Introduction, Scheme 2). Thus, the use of flow technology enabled the conversion of pyrrolidine **65.1** to (protected) 3-pyrrolidinone **65.3** under mild conditions on a 5 g scale in 46% isolated yield in the presence of 1.5 equiv of sulfuric acid and molecular oxygen as the oxidant (Scheme 65). The initially formed ketoamine **65.2** was not stable enough for purification, so isolation was performed after derivatization, for example


**Scheme 63. Photocatalyzed Benzylic Hydroperoxidation under Visible Light Irradiation**




### Scheme 64. Molecular Oxygen as the Oxidant in the Selective CH<sub>2</sub> to C=O Conversion



Notable examples:



### Scheme 65. Gram-Scale Remote Oxidation of Pyrrolidine



through Boc-protection to give **65.3**. The reaction could also be conducted in batch by adopting hydrogen peroxide as the oxidant.<sup>112</sup>


The direct oxidation of labile benzylic C–H bonds for the synthesis of carbonyl compounds was reported to take place under visible light. For example, UrN was used for the benzylic oxidation of indane and isochroman, respectively, to indanone and isochromanone in high yields (up to 87 and 80%) upon irradiation at 420 nm.<sup>489</sup> Similarly, UrP was employed for the oxidation of toluene to benzaldehyde, with minor amounts of benzyl alcohol detected.<sup>467</sup>

Ketones can be smoothly prepared upon photooxidation of the corresponding alcohols via photocatalyzed HAT. Accordingly, TBADT was used to convert benzyl alcohols to the corresponding ketones in high yields on a 0.1 mmol scale.<sup>490,491</sup> Molecular oxygen was used as the oxidant and was bubbled with a constant flow (20 mL min<sup>-1</sup>) in the solution. Thus, when an acetonitrile solution of 1-(4-methylphenyl)ethanol in the presence of 1 mol % of the PC<sub>HAT</sub> was irradiated for 90 min, 82% conversion to the corresponding acetophenone was observed (>99% yield). Notably, when the very same PC<sub>HAT</sub> was immobilized on TiO<sub>2</sub>, a competition between hydrogen atom transfer and single-electron transfer was observed and the conversion rate was halved (35% in 90 min).<sup>490</sup> It is important to mention that, albeit homogeneous TBADT showed higher reaction rates, the heterogenized photocatalyst showed better chemoselectivity toward the  $\alpha$ -to-O C–H bond when two distinguishable benzylic sites were present in the substrate, as demonstrated for 1-(4-ethylphenyl)ethanol.<sup>490</sup>

In another instance, TBADT immobilized on silica (10% w/w) promoted the oxygen-assisted chemoselective photooxidation of 1-pentanol or 3-pentanol to pentanal or 3-pentanone, respectively.<sup>492</sup> When the alcohol (5 mM) was irradiated under O<sub>2</sub> for 1 h at  $\lambda > 290$  nm in the presence of the heterogeneous PC<sub>HAT</sub> (8 g L<sup>-1</sup>), 90% mass balance was observed. Of note, no overoxidation products were detected in this case, contrary to what was observed under the same conditions when using homogeneous TBADT. The performances of homogeneous and heterogeneous TBADT were then directly compared, and the role of the matrix (silica) was evoked to explain the differences in reactivity. Indeed, the oxidation of 1-pentanol occurred 4-times faster when the PC<sub>HAT</sub> was supported on SiO<sub>2</sub> due to the positive effect of adsorption phenomena. Later on, this heterogeneous PC<sub>HAT</sub> was also used for the oxidation of diols with similar results in terms of chemoselectivity.<sup>493</sup> Similarly, silica-heterogenized Ur has been used for the oxidation of alcohols.<sup>479</sup>

Intriguingly, a conceptually similar approach was reported for the oxidation of benzyl alcohols (and of aliphatic alcohols to some extent) by NaDT on a preparative scale.<sup>494</sup> In detail, zirconia was used as a solid support for this PC<sub>HAT</sub> (~19% w/w) and the transformation occurred by irradiating a 0.2 M solution of the starting alcohol with a 400 W high-pressure Hg lamp using a cutoff filter ( $\lambda > 320$  nm) for less than 2 h in most cases, delivering the expected ketone in >90% yield. However, when benzyl alcohols bearing strongly electron-withdrawing substituents were used, longer irradiation times were required (up to 4 h). As an example, the oxidation of 1-phenylethanol (**66.1**) occurred smoothly on a 2 mmol scale to give the corresponding acetophenone (**66.2**) in 94% yield after only 1 h of irradiation (Scheme 66). Notably, the use of the

### Scheme 66. Heterogenized NaDT-Photocatalyzed Oxidation of Benzylic Alcohols




corresponding homogeneous PC<sub>HAT</sub> led to decreased yields (68%). The photocatalyzed oxidation of **66.1** could be conducted on a 100 mmol scale in 3.25 h to deliver the expected ketone in 92% isolated yield. When the same methodology was applied to nonbranched benzylic alcohols, aldehydes were formed. Thus, benzyl alcohol was smoothly converted to benzaldehyde in 88% yield after 2 h of irradiation.

The photooxidation of benzyl alcohols to aldehydes under aerobic conditions has also been reported using water as a benign solvent.<sup>495</sup> In particular, a tandem transformation was reported where this first step was triggered by AQN-2-SO<sub>3</sub>Na (10 mol %), followed by an asymmetric aldol reaction with ketones in the presence of a chiral pyrrolidine-based organocatalyst. Prolonged irradiation (48 h) and substoichiometric acetic acid (50 mol %) were needed to obtain full conversion of the starting materials. For example, (4-bromophenyl)methanol (**67.1**) was a competent substrate in the presence of cyclohexanone (**67.2**, 5 equiv), delivering product **67.3** in 71% yield with exquisite diastereo- and enantioselectivity. Benzyl alcohols bearing strong electron-donating substituents failed to give the expected products due

to the diminished electrophilicity of the intermediate aldehyde (Scheme 67).

### Scheme 67. One-Pot Asymmetric Aldol Reaction Starting from Benzyl Alcohols under Aerobic Conditions




The same PC<sub>HAT</sub> was later used to promote the oxidation of both hydrocarbons and alcohols to the corresponding carbonyl compounds. For example, it was found that, upon irradiation with a 200 W white LED lamp, the water-soluble AQN-2-SO<sub>3</sub>Na (3 mol %) could be used to promote efficiently the oxidation of toluene to benzaldehyde (GC yield: 81%).<sup>496</sup> It was observed that the transformation proceeded through the rate-determining formation of benzyl alcohol.<sup>496</sup> In another instance, the use of EY as POC extended this transformation to primary alcohols, besides easily oxidizable benzyl alcohols, for the synthesis of quinazolinones in a one-pot fashion.<sup>253</sup> As an example, when ethanol (68.1) was used as the substrate in DMSO in the presence of EY (1 mol %) and the mixture was irradiated for 72 h with a 50 W Xe lamp (cutoff > 400 nm) under O<sub>2</sub> atmosphere, acetaldehyde was formed *in situ* and conveniently trapped by *o*-aminobenzamide 68.2 to give the expected product 68.3 in 66% yield (Scheme 68). Remarkably, this procedure was adopted for the synthesis of 2-alkylquinazolinones, including the sedative-hypnotic drugs methaqualone 68.4 and mecloqualone 68.5.<sup>253</sup>

#### 3.2.4. Synthesis of Carboxylic Acids and Esters.


Photocatalyzed HAT was likewise exploited for the oxidation of methyl aromatics to the corresponding benzoic acids.<sup>497</sup> As an example, when 4-*tert*-butyltoluene 69.1 was reacted under aerobic conditions in the presence of CIAQ, a mixture of the corresponding benzoic and perbenzoic acids was obtained. It was found that the use of either a base or an acid could steer the selectivity toward the acid. Accordingly, when a 0.3 M methyl acetate solution of 69.1 was irradiated in the presence of CIAQ (8 mol %), the corresponding benzoic acid 69.2 was isolated in 97% and 99% yields when using K<sub>2</sub>CO<sub>3</sub> or TFA as an additive, respectively (Scheme 69). A stepwise mechanism was unveiled, where the alkyl aromatic is first converted to the aldehyde via a hydroperoxide intermediate, which is in turn hydrated and then transformed into the acid by a similar photooxidative process.<sup>497</sup>

Later on, this strategy was used for the direct aerobic photooxidative synthesis of aromatic methyl esters starting from methyl aromatics.<sup>242</sup> In detail, when methanol was adopted as the reaction solvent, the transient aldehyde was converted to the corresponding hemiacetal and, eventually, to the methyl

### Scheme 68. EY-Photocatalyzed Synthesis of Alkyl Quinazolinones via Oxidation of Primary Alcohols



### Scheme 69. Anthraquinone-Photocatalyzed Benzylic Oxidation of Methylarenes



ester. For example, 69.1 was converted to 69.3 in 89% isolated yield by irradiating a solution containing AQ-2,3-diCOOH (10 mol %) for 24 h in methanol (Scheme 69). Unfortunately, this strategy could not be extended to the synthesis of other alkyl carboxylates, since other alcohols, such as ethanol or propanol, were found to undergo uncontrolled oxidation when used as solvents.


Intriguingly, when 69.1 was reacted in the presence of AQ-2-COOH (5 mol %) in pure aerated ethyl acetate without any additive, the perbenzoic acid 69.4 was formed predominantly (88% overall <sup>1</sup>H NMR yield, perbenzoic:benzoic acid ratio 2.8:1). This product could be in turn used in a one-pot fashion to epoxidize 5-decene to give 69.5 in 86% isolated yield when MeCN was used as cosolvent (Scheme 69).<sup>498</sup>

#### 3.3. Formation of C–S Bonds

Photocatalyzed HAT was also employed to forge C–S bonds via addition of the photogenerated C-centered radicals onto a weak S–S single bond (see also section 4.3). In one instance, PQ was chosen as visible light-triggered PC<sub>HAT</sub> (5 mol %) to promote the metal- and oxidant-free thioesterification of

aldehydes.<sup>245</sup> Thus, it was found that thiosulfonate **70.2** used in slight excess (1.2 equiv) was a suitable trap for the acyl radical generated via HAT from **70.1** (0.2 M) because of the high polarization of the S–SO<sub>2</sub> bond (Scheme 70).

### Scheme 70. Photocatalyzed Thioesterification of Aldehydes




Consequently, the generated sulfonyl radical accounted for the closure of the photocatalytic cycle, yielding product **70.3** in 95% yield after 14 h of irradiation under blue light. The presence of a base (Na<sub>2</sub>CO<sub>3</sub>, 0.5 equiv) was found crucial to neutralize the sulfinic acid developed during the reaction. Notably, this protocol could be extended to several structurally diverse aldehydes and thiosulfonate *S*-esters and could be adopted for the functionalization of complex biologically active molecules, such as probenecid and ursodeoxycholic aldehyde derivatives to give **70.4** and **70.5**, respectively.<sup>245</sup>


Very recently, aldehydes functioned as substrates for a difluoromethylthiolation strategy triggered by photocatalyzed HAT (Scheme 71).<sup>499</sup> Both aliphatic and aromatic aldehydes were activated by the excited state of TBADT (4 mol %) to form the corresponding acyl radicals that were in turn trapped by compound **71.2**, used in excess. Indeed, the expected difluoromethylthiolated product could be isolated in good yields on a 0.2 mmol scale. As an example, 2-naphthaldehyde **71.1** was converted to thioester **71.3** in 86% yield. As reported in Scheme 71, this methodology was exploited to functionalize biologically relevant molecules, such as *L*-menthol (product **71.4**) and dehydrocholic acid derivatives (product **71.5**).

A conceptually similar procedure was reported for the trifluoromethylthiolation of aromatic aldehydes.<sup>500</sup> For example, TBADT (4 mol %) was used to cleave the formyl C–H bond in *p*-anisaldehyde (**72.1**) and the acyl radical was readily trapped by *N*-(trifluoromethylthio)phthalimide (**72.2**) to afford the expected product (**72.3**, Scheme 72). Notably, the protocol proved robust and enabled this transformation on a gram-scale (5 mmol), as well as the functionalization of


### Scheme 71. Difluoromethylthiolation of Aldehydes




Notable examples:



### Scheme 72. Trifluoromethylthiolation of Aldehydes



Notable examples:




fenbufen and ibuprofen to deliver products **72.4** and **72.5**, respectively. It is perhaps important to mention that the authors found some limitations with substrates containing electron-withdrawing groups or substituents in the *ortho* position. They proposed the steric bulkiness of TBADT as the main reason for this diminished reactivity.<sup>500</sup>

Recently, a completely different strategy for the formation of C–S bonds has been disclosed. In this work, the authors made use of photocatalyzed HAT to generate C-centered radicals that were readily trapped by a SO<sub>2</sub>-surrogate (1,4-diazabicyclo[2.2.2]octane bis(sulfur dioxide), DABSO) to afford sulfonyl radicals. The latter intermediates were then exploited for a Ni-mediated enantioselective radical addition onto electrophilic olefins (Scheme 73).<sup>501</sup> As an example, when adamantane (**73.1**) was irradiated in the presence of PT (5 mol %) under blue light (455 nm), the tertiary radical was generated. This intermediate was readily trapped by DABSO (**73.3**, 1.5 equiv) to forge the aimed-for C–S bond and afford a stable sulfonyl radical (e.g., **73.9**). In the meanwhile, an asymmetric Ni complex was formed upon coordination of Ni<sup>II</sup> by chiral ligand **73.4**; this Ni complex was responsible for imparting asymmetry on the following step via coordination of **73.2**. Finally, the sulfonyl radical added onto **73.2** to eventually afford the expected product **73.5** (77% yield, e.e. 90%, r.r. >



### Scheme 73. Three-Component Asymmetric Sulfonation via HAT




50:1).<sup>501</sup> This protocol was likewise amenable to activated benzylic C–H bonds and to functionalize celestolide (to give 73.6) and a synthetic intermediate of differin (73.7).

### 3.4. Formation of C–F Bonds

**3.4.1. Fluorination of C(sp<sup>3</sup>)–H Bonds.** The introduction of a fluorine atom onto unactivated sites of organic molecules, such as strong aliphatic C(sp<sup>3</sup>)–H bonds, is a hot topic in medicinal and pharmaceutical sciences. In fact, fluorination makes a molecule more lipophilic, which results in ameliorated biological absorption and distribution. This transformation is of great importance in materials science too, where the incorporation of the halogen endows materials with a remarkable hydrophobicity.<sup>502</sup> Photocatalyzed HAT has been widely used to promote this highly desirable transformation.<sup>503</sup> This chemistry mainly relies on fluorinating agents containing a weak N–F bond (BDE = 61–63 kcal mol<sup>−1</sup>), such as *N*-fluorobenzenesulfonimide and Selectfluor.<sup>504</sup>


In one of the earliest examples of this chemistry, the use of aromatic ketones for the mono- and *gem*-difluorination of benzylic C(sp<sup>3</sup>)–H bonds in the presence of Selectfluor was reported.<sup>273</sup> In particular, when an acetonitrile solution of 1-(*tert*-butyl)-4-methylbenzene (74.1, 0.08 M) was irradiated with a CFL lamp in the presence of FL (5 mol %) and 2 equiv of Selectfluor (74.2a), the monofluorination at the benzylic site occurred in 82% yield to give compound 74.3 in 6 h (Scheme 74). However, when the reaction was performed in the presence of XA (5 mol %) and Selectfluor II (74.2b, 3 equiv) as PC<sub>HAT</sub> and fluorinating agent, respectively, the *gem*-difluorinated product 74.4 was isolated in 72% yield after 16 h. From the mechanistic standpoint, the obtained organoradical was fluorinated by reaction with 74.2 to deliver the (di)fluorinated product and a *N*-centered radical cation, which is entrusted for the closure of the photocatalytic cycle.

### Scheme 74. Diarylketone-Photocatalyzed Selective Benzylic Mono- and Difluorinations



More recently, the above-mentioned procedure for the monofluorination of alkylaromatics was used to develop a one-pot, transition metal-free cross-dehydrogenative arylation of unactivated benzylic C–H bonds (Scheme 75).<sup>505</sup> In

### Scheme 75. One-Pot Arylation of Unactivated Benzylic C–H Bonds




particular, FL was used to trigger the halogenation step, while the so-formed benzyl fluorides (75.2) were used as electrophilic partners in a nucleophilic substitution with electron-rich (hetero)arenes (e.g. 75.3 to give 75.4).<sup>505</sup>


In a related example, the same monofluorination reaction was performed under flow conditions by circulating an acetonitrile solution of 76.1 through a transparent FEP tubing coiled around a glass cylinder (V = 28 mL, flow rate = 1 mL·min<sup>−1</sup>) containing a 105 W CFL.<sup>287</sup> Under these conditions, it was found that XA (5 mol %) could perform the monofunctionalization of the benzylic site to get 76.3 in 89% yield with a residence time of 28 min (temperature of the system = 40 °C). This reaction manifold was adapted also to the fluorination of drugs, such as celestolide and ester-protected ibuprofen (Scheme 76). As for celestolide, milder conditions were required (flow rate = 3 mL min<sup>−1</sup>, t<sub>r</sub> = 9 min, temperature: 25 °C) to get the corresponding fluorinated compound in 88% yield and the reaction could be conveniently scaled up with comparable efficiency (87% yield). Contrarily, in the case of ibuprofen, “harsher” conditions were needed (flow rate = 1 mL min<sup>−1</sup>, t<sub>r</sub> = 28 min, temperature: 60 °C) to get the expected product in 80% yield, with the secondary benzylic C–H position being almost exclusively functionalized (>90% selectivity).

In another case, the benzylic fluorination of alkyl aromatics was achieved in the presence of TBADT and NFSI (77.2) as the fluorinating agent.<sup>506</sup> Thus, ibuprofen methyl ester 77.1


## Scheme 76. Photocatalyzed Benzylic Fluorination under Flow Conditions



Notable examples:



## Scheme 77. TBADT-Photocatalyzed Benzylic Fluorination in Flow




was smoothly functionalized at the less-hindered benzylic position in 75% yield. It is important to stress that, when a flow apparatus was adopted ( $V = 1.4 \text{ mL}$ , FEP tubing wrapped around a blacklight blue lamp;  $\lambda = 365 \text{ nm}$ ), the reaction was significantly sped up, from 24 h required in batch to only 5 h (70% yield, Scheme 77). Hydrolysis of the fluorinated ester gave fluoroibuprofen, whose metabolic stability in human and rat microsomes was evaluated to be somewhat higher compared to the parent drug.

Similarly, **DBS** was used to promote the selective fluorination of phenylalanine-like residues in amino acids and short chain peptides at the benzylic site. In this case, Selectfluor was employed as the fluorinating agent while a 14 W CFL was used as the light source.<sup>244</sup>



Photocatalyzed HAT can be conveniently used also for the fluorination of strong aliphatic C–H bonds. In one case, the fluorination of cycloalkanes took place making use of aromatic ketones such as **AP**, which was effective in the C–H to C–F bond conversion upon irradiation with a commercially available CFL (Scheme 78). For example, the fluorination of cyclohexane, cycloheptane, and cyclooctane **78.1a–c** occurred in 90%, 85%, and 82% <sup>19</sup>F NMR yield, respectively.<sup>225</sup>

Another convenient PC<sub>HAT</sub> that can be used for the fluorination of strong, unactivated C–H bonds is **TBADT** (Scheme 79).<sup>507</sup> In particular, this polyoxometalate has been exploited for the fluorination of a wide array of compounds, including biologically valuable molecules such as (protected) amino acids and sclareolide (79.1). Thus, when an acetonitrile solution of **79.1** was irradiated for 17 h in the presence of NFSI (79.2, 1.2 equiv) and **TBADT** (2 mol %) at 365 nm (15

## Scheme 78. Photocatalyzed Preparation of Fluorocycloalkanes



## Scheme 79. Photocatalytic Fluorination of Unactivated C–H Bonds




W black light bulbs), a mixture of the 8- and 9-fluorinated adducts was obtained (68% overall yield, ratio **79.3/79.4** 5.8:1). Interestingly, the selectivity enabled by this system is analogous to that previously reported adopting manganese porphyrins in the presence of fluoride anion.<sup>508</sup>

In this context, a similar strategy was adopted for the functionalization of the methine site in leucine. Thus, **NaDT** was found useful for the selective fluorination of  $\gamma$ -leucine methyl ester (**80.1**), a key intermediate on the route to odanacatib, a potent inhibitor of cathepsin K. Intriguingly, this transformation was carried out under flow conditions and could be easily scaled up, to get ~45 g of product (90% yield, Scheme 80).<sup>509</sup>

The same procedure was used to achieve the site-selective <sup>18</sup>F-fluorination of unprotected amino acids<sup>274</sup> and peptides,<sup>510</sup> which has an intrinsic value for positron emission tomography (PET) imaging. Later on, the same fluorination concept was adopted for the labeling of ZJ-43 analogues, potent binders for PSMA (prostate specific membrane antigen), that are overexpressed in the case of prostate cancer. Parent ZJ-43 could not be efficiently labeled (<12%) with <sup>18</sup>F; however, the addition of pendant ammonium groups was found to accelerate the functionalization (>46% yield).<sup>511</sup>


Quite recently, the **UrN** photocatalyst has been used for the fluorination of cycloalkanes under visible light irradiation. Thus, a deuterated acetonitrile solution of cyclooctane (**81.1**) was irradiated in the presence of **UrN** (1 mol %) and 1.5 equiv of NFSI (**81.2**) for 16 h under inert atmosphere to afford

## Scheme 80. TBADT-Photocatalyzed Fluorination of Leucine in Flow




fluorocyclooctane (**81.3**) in 95% <sup>1</sup>H NMR yield (Scheme 81).<sup>285</sup>

## Scheme 81. Visible Light Fluorination of Cyclooctane



**3.4.2. Fluorination of C(sp<sup>2</sup>)-H Bonds.** The fluorination of aldehydes via the activation of the formyl C(sp<sup>2</sup>)-H bond has been sparsely reported.<sup>512</sup> Here, the capability of DT to generate acyl radicals from aldehydes (**82.1**) was exploited. These intermediates were in turn trapped by NFSI to get the corresponding acyl fluorides (**82.3**, Scheme 82). Since acyl

## Scheme 82. Synthesis of Acyl Fluorides from Aldehydes



fluorides were found to be unstable in several cases, they were treated *in situ* with benzylamine to get the corresponding amides (**82.4**). For example, when an acetonitrile solution of benzaldehyde **82.1** (0.3 M) was irradiated at 365 nm in the presence of NaDT (2.5 mol %) and NFSI (**82.2**, 1.2 equiv), the corresponding acyl fluoride **82.3** was formed in 79% <sup>19</sup>F NMR yield. After irradiation, the addition of 2 equiv of benzylamine allowed isolation of benzamide **82.4** in 79% yield. This reactivity was extended to several aliphatic aldehydes as well. Notably, **82.4** could be obtained in 53% isolated yield starting from benzyl alcohol in the presence of a higher amount of NFSI (2.5 equiv) in a one-pot fashion.<sup>512</sup>


## 3.5. Formation of C-Cl Bonds

Similarly to what was shown in the previous section for C-F bond formation, C-Cl bonds can also be formed via photocatalyzed HAT. In these cases, electrophilic reactants

containing a weak N-Cl bond are exploited, with one typical example being *N*-chlorosuccinimide (NCS).

In one instance, the halogenation of α-to-boron C(sp<sup>3</sup>)-H bonds in benzyl *N*-methyliminodiacetyl (MIDA) boronates was reported.<sup>513</sup> Thus, the authors started off by studying the bromination of said bonds and found an effective photochemical way to promote reactivity. In particular, the direct irradiation of a solution containing the boronate ester and *N*-bromosuccinimide (NBS) under inert atmosphere with a 13 W CFL led to the expected products. However, this protocol failed when replacing NBS with NCS to perform chlorination of C-H bonds; intriguingly, the authors found that the addition of 5 mol % of FL restored reactivity (Scheme 83).


## Scheme 83. Chlorination of Benzyl MIDA Boronates



Upon mechanistic investigation, they found that FL acted as a photocatalyst for HAT. Interestingly, by replacing NCS with Selectfluor, the authors were able to achieve the fluorination of α-to-boron C(sp<sup>3</sup>)-H bonds.<sup>513</sup>

In another instance, a similar chlorine source was used for the chlorination of (+)-sclareolide (**84.1**) under flow conditions (Scheme 84).<sup>514</sup> Interestingly, the authors designed

## Scheme 84. Chlorination of (+)-Sclareolide via a Schlenk-in-Flow Approach



a “Schlenk-in-flow” technique for the safe handling of oxygen- and moisture-sensitive reagents, where argon is used instead of solvent to drive reagents through the tubing. Doing so, distribution phenomena are suppressed, which allowed saving both reagents and solvent. The photocatalyzed chlorination of **84.1** was one of their benchmark reactions: when an acetonitrile solution of **84.1** was pumped through the

photoreactor ( $V_R = 10$  mL,  $t_R = 100$  min) in the presence of **TBADT** (10 mol %) and **84.2** (1.5 equiv) under UV light (365 nm), 38% of the chlorinated product **84.3** was obtained on a gram scale.


### 3.6. Formation of C–D Bonds

Deuterium labeling has a range of applications, including the investigation of reaction mechanisms, drugs metabolism and distribution, as well as in spectroscopy. Accordingly, it is no surprise that photocatalyzed HAT was investigated as a direct and unique platform to access the formation of the C–D (D: deuterium) bond by the so-called hydrogen isotope exchange (HIE).<sup>515–518</sup>

In this frame, two similar approaches have been published very recently. In both cases, the photogenerated radicals (**TBADT** as the  $PC_{HAT}$ ) were quenched by an aromatic thiol (triisopropylbenzenethiol, TIPSH) present in catalytic amounts and involved in an acid/base equilibrium with  $D_2O$ . While in one instance the deuteration of the formyl C–H bond in both aromatic and aliphatic aldehydes was achieved,<sup>519</sup> in the other one the methodology was extended to the deuteration of benzylic sites and tertiary  $C(sp^3)$ –H bonds, besides aldehydes.<sup>520</sup>

In the former case, 4-bromobenzaldehyde **85.1** was deuterated in 95% yield (D incorporation: 96%) by means of the **TBADT**/TIPSH system (used in 4 and 40 mol %, respectively) in DCM/ $D_2O$  1:1 upon irradiation for 24 h (Scheme 85a).<sup>519</sup> In the latter case, the deuteration could be

**Scheme 85. Photocatalyzed Hydrogen Isotope Exchange (HIE)**



extended to benzylic sites and tertiary  $C(sp^3)$ –H bonds simply by adding a phase transfer catalyst (TBAB, tetrabutylammonium bromide). For example, **85.2** and **85.3** were functionalized in 92% (D incorporation: 91%, 24 h) and 99% (D incorporation: 81%, 48 h) yield, respectively (Scheme 85b).<sup>520</sup>


## 4. FORMATION OF OTHER BONDS

Photocatalyzed HAT has been likewise employed for the generation of heteroradicals, including P-, Si-, and S-centered radicals.

### 4.1. Formation of P–C Bonds

As for P–C bond formation, the enantioselective conversion of hydrophosphine oxide **86.1** to the phosphine oxide **86.3** via **EY**-photocatalyzed HAT has been reported (Scheme 86).<sup>391</sup> In particular, when a solution of amide **86.2** (1 M) in MTBE/

**Scheme 86. Enantioselective Formation of a P–C Bond**



$H_2O \sim 3:1$  was irradiated with a blue LED in the presence of **EY** (1 mol %) and **86.1** as the H-donor, **86.3** was formed in 71% yield (e.e. 68%). As mentioned before (see Scheme 19d), the enantioselectivity was entrusted to a chiral Rh complex (4 mol %). Unfortunately, the reaction required a long time to go to completion (3 days).

Very recently, **EY** was used to promote the formation of a P–C bond starting from hydrophosphine oxide as H-donor and *N*-(hetero)arylsulfonyl propiolamides. As already mentioned above (see Scheme 45 in section 2.5), this transformation proceeded through a cascade involving 1,4-addition of the P-centered radical onto the  $C\equiv C$  triple bond of propiolamide, Smiles rearrangement, and 5-*endo*-trig cyclization.<sup>438</sup> In a very recent report, *N*-aminopyridinium salts have been used as radical traps for photogenerated P-centered radicals (**AQ** as the photocatalyst), resulting in the pyridylation of a handful of hydrophosphine oxides.<sup>417</sup>


### 4.2. Formation of Si–X Bonds (X = Cl or C)

Photocatalyzed HAT has been used also for the formation of Si–X bonds (X = Cl, C). This strategy requires that a Si–H bond is photocatalytically cleaved, which is easily predicted by the enhanced electropositivity of silicon compared to carbon. This feature contributes to make the H atom to be abstracted more hydridic.<sup>249</sup> The forging of the Si–Cl bond gives access to chlorosilanes, currently used for the protection of alcohols and acids. The reaction occurred under visible light irradiation (**EY** as the  $PC_{HAT}$ ) and relied upon DCM in the double role of chlorinating agent and solvent. The range of applicability of this transformation is remarkably broad, and almost all chlorosilanes were isolated in quantitative yield. Notably, flow chemistry allowed successfully performing the process up to the gram scale. For example, when hydrosilane **87.1** (0.2 M) was irradiated with **EY** (1 mol %) in DCM with an 18 W blue LED strip, the corresponding chlorinated product **87.2** was obtained quantitatively after 4 h. By means of the flow technique (HPFA tubing,  $V = 7$  mL, flow rate  $\sim 0.1$  mL  $\cdot$  min<sup>−1</sup>), the same yield was achieved with a residence time of 1.1 h and 1.18 g of **87.2** was produced after 4 h (Scheme 87).<sup>249</sup>


As for the formation of the Si–C bond, a handful of reports describe the use of **TBADT** for the activation of the Si–H bond (Scheme 88).<sup>278,379,387,521</sup> Thus, when an acetonitrile solution of methyl-diphenylsilane **88.1** was irradiated with UV light (310 nm,  $10 \times 15$  W phosphor-coated lamps) for 24 h in the presence of **TBADT** (2 mol %), the expected Si-centered radical was generated via HAT.<sup>521</sup> This intermediate was readily trapped by dimethyl maleate (**88.2**) to afford the corresponding silylated compound in excellent isolated yield (90%), even though it should be noted that 28% yield was observed in the absence of the  $PC_{HAT}$ . The use of transient absorption spectroscopy and steady-state EPR experiments served to prove the nature of the activation step. Interestingly, when silanes containing more labile Si–H bonds (e.g., tris(trimethylsilyl)silane) were used, the reaction proceeded



Scheme 87. Photocatalyzed Synthesis of Chlorosilanes



Scheme 88. TBADT-Photocatalyzed Hydrosilylation of Electron-Poor Olefins




even in the absence of the photocatalyst due to radical chain processes. On the other hand, when trialkylsilanes were used, the competitive activation of the  $\alpha$ -to-Si C–H and Si–H bonds was observed, thus generating a mixture of C- and Si-functionalized products.<sup>521</sup>

#### 4.3. Formation of S–C Bonds

The formation of S–C bonds can be achieved through the addition of S-centered radicals generated via HAT onto electron-rich olefins such as styrenes. In one case, a thiol–ene process has been reported by using BP (10 mol %) as the PC<sub>HAT</sub>, which allowed formation of sulfide **89.3** in 87% yield after 12 h of irradiation of a mixture of thiophenol (**89.1**) and styrene (**89.2**) with a 18 W CFL (Scheme 89).<sup>237</sup>

Quite recently, an asymmetric manifold for the formation of the S–C bond has been proposed.<sup>391</sup> Therein, a sulfinic acid was activated via HAT by EY to get the corresponding S-centered radical. The latter intermediate was trapped by  $\alpha,\beta$ -unsaturated amides through a radical addition manifold to forge the desired S–C bond. Notably, a Ru-based chiral Lewis

Scheme 89. BP-Photocatalyzed S–C Bond Formation



acid was used to coordinate the amide with a dual objective: favoring the radical addition step and introducing asymmetry.<sup>391</sup>

Finally, given the importance of the trifluoromethylthio (–SCF<sub>3</sub>) moiety in medicinal chemistry, a methodology based on HAT has been recently proposed for the trifluoromethylation of  $\beta$ -ketodithioesters under visible light irradiation.<sup>252</sup> Thus, when methyl 3-oxo-3-(3,4,5-trimethoxyphenyl)propanedithioate was irradiated with blue light in the presence of EY (2 mol %) and sodium triflate, the corresponding trifluoromethylthiolated product was obtained as a single diastereomer (*Z*) in 89% yield after 15 h. It is perhaps worth mentioning that the reaction worked to some extent (37% yield) also in the absence of the PC<sub>HAT</sub>.


## 5. MISCELLANEA

This section gathers examples that do not fit into any of the categories above. In particular, net-oxidative processes based on a HAT step such as dehydrogenation and dehydroformylation reactions, as well as reactions based on a C–C oxidative cleavage, are described.

Seminal examples of such processes have been reported in the mid 1980s/early 1990s and were based on the use of polyoxometalates as PC<sub>S<sub>HAT</sub></sub>, with particular focus on the DT anion; albeit, the reactions were usually stopped at low conversion of the starting mixtures.<sup>201</sup> Thus, the dehydrogenation of a library of organic substrates (mostly alkanes and alcohols) based on photocatalyzed HAT has been reported.<sup>322,325,330,522,523</sup> Indeed, specific conditions have been found to deliver the expected products with a high quantum yield (approaching unity),<sup>523</sup> as well as to favor the formation of nonthermodynamic alkenes for alkanes.<sup>330</sup> Furthermore, it was demonstrated that photocatalyzed HAT can also be exploited to trigger the epimerization of C–H bonds (e.g., in decalone derivatives).<sup>325,522</sup>

More recently, a dual-catalytic system consisting of TBADT (0.4 mol %) and cobaloxime pyridine chloride ([Co(dmgH)<sub>2</sub>(pyr)Cl], COPC; 0.4 mol %) was used for the preparative dehydrogenation of alkanes.<sup>524</sup> This methodology was baptized “cooperative HAT”, since it involved the combination of photocatalyzed HAT and thermal HAT.<sup>525</sup> In detail, the PC<sub>HAT</sub> was responsible for the activation of a strong C(sp<sup>3</sup>)–H bond, for example in cyclooctane **90.1**, in what has been called a “hard” HAT step (Scheme 90a). Subsequently, the Co-based complex exerted an “easy” HAT

Scheme 90. TBADT/Cobaloxime Dual-Catalytic Approaches for the (a) Dehydrogenation of Alkanes and (b) Dehydroformylation of Aldehydes



from the  $\alpha$ -position with respect to the radical site. The two catalytic cycles met when the reduced DT and the spent cocatalyst interacted to release a molecule of hydrogen gas. Ultimately, cyclooctene (**90.2**) was obtained in 19% NMR yield upon irradiation for 48 h. The very same procedure was also successfully applied to secondary alcohols, delivering the corresponding ketones in good yields.<sup>524</sup>

Later on, a dehydroformylation procedure via HAT has been developed.<sup>526</sup> In particular, the same dual-catalytic system has been proposed to realize the decomposition of an aldehyde to an alkene, hydrogen gas, and carbon monoxide. Thus, aldehydes containing a nonenolizable quaternary  $\alpha$ -carbon, such as 2,2-dimethyl-3-phenylpropanal **90.3**, were tested at first (Scheme 90b). TBADT (4 mol %) generated an acyl radical under UV light irradiation, that spontaneously underwent decarbonylation to give a tertiary radical. The latter intermediate was in turn intercepted by cobaloxime (COPC, 4 mol %), that triggered a further dehydrogenation step, with a preference toward the less substituted olefin (terminal olefin **90.4**). Aldehydes with an enolizable carbon  $\alpha$  to the carbonyl group, however, were not competent substrates in this reaction.<sup>526</sup>

Another photocatalytic approach for the deconstructive cleavage of organic compounds via HAT dealt with the aerobic photooxidative cleavage of 1,3-diketones to give carboxylic acids.<sup>527</sup> In such a case, CIAQ (10 mol %) was used to trigger a H-abstraction in benzoylacetones, such as 1-phenyl-1,3-butanedione **91.1** (Scheme 91a). Thus, when an acetone


mechanism underlying this transformation is quite complicated and proceeds through the formation of transient hydroperoxides and  $\alpha$ -hydroxycarbonyl and  $\alpha$ -dicarbonyl compounds.<sup>527</sup> In a related instance, similar conditions were applied for the aerobic oxidative cleavage of cyclic acetals (Scheme 91b).<sup>528</sup> As an example, irradiation of 1,3-dioxolane **91.3** in ethyl acetate solution in the presence of AQ-2-COOH (10 mol %) followed by quenching with sodium thiosulfate allowed synthesizing the corresponding ester **91.4** in good yield (69%).

## 6. CONCLUSIONS

It is apparent from the examples described in this review that the photocatalyzed hydrogen atom transfer approach has had and will have in future years a primary role in synthetic planning, in both academic and industrial settings. The versatility of the method is witnessed by the different types of bonds that can be forged, even though most of them are C–C bonds (see Table 2). In the majority of the examples reviewed, the reactions are carried out under mild conditions with no need of heating or aggressive conditions. In many instances, in the reactions developed all of the atoms of the reagents are incorporated in the desired products (100% atom economy).

Table 2 gives an idea of the importance of the different classes of PC<sub>S<sub>HAT</sub></sub> used in connection with the bond they allow to build. Decatungstate salts (either NaDT or TBADT) are by far the most employed PC<sub>S<sub>HAT</sub></sub>, and they are involved in all of the processes described. The capability of DT-based PC<sub>S<sub>HAT</sub></sub> to cleave even very strong bonds, such as the C–H bond in methane (BDE = 105 kcal/mol), showcases their impressive potentialities. Moreover, dual-catalytic approaches with Ni, Pd, Cu, and Co have extended even further the boundaries of the chemical space that can be explored with them. This colorless anion is robust, cheap, and easy to prepare; however, its chemistry is mainly restricted by the need for highly energetic UV light (<400 nm). Accordingly, in recent years a lot of work has been devoted to the discovery of visible light absorbing PC<sub>S<sub>HAT</sub></sub>. Albeit colored polyoxometalates that can compete with DT are so far lacking, other classes of colored PC<sub>S<sub>HAT</sub></sub> have been tested, including oxo-porphyrins (Sb-Oxo), uranyl salts (Ur), or MOFs. Photoorganocatalysts appear as promising derivatives worthy to be further investigated. As for aromatic ketones, early examples dealt only with the use of colorless BP, but nowadays slightly colored diaryl ketones

### Scheme 91. Aerobic Photooxidative Cleavage of (a) 1,3-Diketones and (b) 1,3-Dioxolanes



solution (0.06 M) was irradiated with four 22 W CFLs in the presence of oxygen and a base, the corresponding benzoic acid **91.2** was obtained in 72% yield after 20 h. The proposed

Table 2. Use of the Diverse Classes of PC<sub>S<sub>HAT</sub></sub> in the Forging of New Bonds

Bond formed/PC <sub>S<sub>HAT</sub></sub>	Aromatic ketones	Anthraquinones	EY	DT	Ur	Sb-Oxo	TAC <sup>*2+</sup>
C(sp <sup>3</sup> )–C(sp <sup>3</sup> )	✓	✓	✓	✓	✓	✓	
C(sp <sup>3</sup> )–C(sp <sup>2</sup> )	✓		✓	✓			✓
C(sp <sup>3</sup> )–C(sp)	✓		✓	✓	✓		
C(sp <sup>2</sup> )–C(sp <sup>3</sup> )	✓		✓	✓	✓	✓	
C(sp <sup>2</sup> )–C(sp <sup>2</sup> )		✓		✓			
C(sp <sup>2</sup> )–C(sp)				✓			
C–N	✓			✓			
C–O	✓	✓	✓	✓	✓		
C–S	✓		✓	✓			
C–Cl	✓			✓			
C–F	✓			✓		✓	
C–D				✓			
Other bonds P–C, Si–Cl, Si–C, S–C	✓		✓	✓			

(e.g., FL, DTX, PT) or  $\alpha$ -diketones (PQ) are preferred. Anthraquinone derivatives could have a more important role in the future; albeit, they have been used only in selected cases so far. EY is close to the ideal candidate as PC<sub>HAT</sub> due to its wide availability and the performance showed, but its competitive role as PC<sub>SET</sub> or singlet oxygen sensitizer must be carefully considered to avoid undesired side reactions. The use of alternative hydrogen abstractors, such as the excited state of the electrogenerated tris(amino)cyclopropenium radical dication (TAC<sup>•2+</sup>), is still in its infancy as it is the photoelectrochemistry approach.<sup>529</sup>

As apparent from Table 2, the forging of C(sp<sup>3</sup>)-C(sp<sup>3</sup>) bonds is widely investigated and a lot of conditions have been tested, making use of almost all of the classes of PC<sub>S<sub>HAT</sub></sub> available in conjunction with other catalysts, in turn resulting in interesting dual-catalytic applications. The Giese radical addition is the archetypal reactivity mode in this group.

However, in C-C bond formation limited efforts have been devoted to the formation of C(sp<sup>2</sup>)-C(sp<sup>2</sup>) and even less to C(sp<sup>2</sup>)-C(sp) bonds. The same holds for the construction of C-N and C-D bonds.

One of the main drawbacks of HAT reactions is that the homolytic cleavage of C-H bonds involves “nucleophilic” rather than “electrophilic” hydrogens, and it is quite rare to find a process that points to the generation (and ensuing reactions) of an electrophilic radical.<sup>180</sup> However, quite recently, a new methodology based on radical polar crossover was exploited to reverse the polarity of said nucleophilic radical by generating a carbocation upon chemical oxidation.<sup>464</sup> This concept finally made photocatalyzed HAT suitable to explore new chemical spaces, namely the world of polar chemistry. The possibility to choose the nucleophile to trap the carbocation sets up a fertile ground to develop new synthetic methodologies based on HAT.

The final comment is on selectivity in the hydrogen abstraction. As mentioned in the Introduction, there are several factors that may affect and direct the cleavage of a C-H bond out of all those present in the hydrogen donor. Dated examples typically dealt with hydrogen donors where all C-H bonds (e.g., in cyclohexane) were equivalent or the hydrogen abstraction was merely governed by the BDE (e.g., in tetrahydrofuran). Conversely, in recent times, very selective HAT reactions have been reported especially in the late-stage functionalization of complex molecules. This demonstrates that mastering the knowledge of all the factors explained in the Introduction allows chemists to develop surprisingly selective processes. Accordingly, further efforts toward a better understanding of these factors is needed to increase the potentialities of HAT reactions in the future. Finally, efforts on the development of hydrogen abstractors having peculiar shapes or a marked bulkiness as well as on the study of the cooperative effects in tuning the selectivity of the reaction are then crucial in this respect.

## AUTHOR INFORMATION

### Corresponding Author

**Maurizio Fagnoni** – PhotoGreen Lab, Department of Chemistry, University of Pavia, 27100 Pavia, Italy; [orcid.org/0000-0003-0247-7585](https://orcid.org/0000-0003-0247-7585); Email: [fagnoni@unipv.it](mailto:fagnoni@unipv.it)

## Authors

**Luca Capaldo** – Flow Chemistry Group, Van't Hoff Institute for Molecular Sciences (HIMS), University of Amsterdam, 1098 XH Amsterdam, The Netherlands; [orcid.org/0000-0001-7114-267X](https://orcid.org/0000-0001-7114-267X)

**Davide Ravelli** – PhotoGreen Lab, Department of Chemistry, University of Pavia, 27100 Pavia, Italy; [orcid.org/0000-0003-2201-4828](https://orcid.org/0000-0003-2201-4828)

Complete contact information is available at: <https://pubs.acs.org/10.1021/acs.chemrev.1c00263>

## Notes

The authors declare no competing financial interest.

## Biographies

Luca Capaldo was born in Milan, Italy, in 1991. After an Erasmus Traineeship in the De Cola group at the University of Strasbourg and a visiting period as a Ph.D. student in the Yoon Group at the University of Wisconsin—Madison, he received his doctorate in Chemical and Pharmaceutical Sciences at the University of Pavia in 2019 (advisor: Prof. M. Fagnoni). After a 2-year postdoctoral fellowship in Pavia (PI: Prof. D. Ravelli), Luca joined the Flow Chemistry Group at the University of Amsterdam as a Marie Curie Fellow (advisor: Prof. T. Noël) to develop novel synthetic methodologies based on photocatalyzed HAT in flow.

Davide Ravelli is currently Assistant Professor at the Department of Chemistry of the University of Pavia, where he also obtained his Ph.D. in Chemistry in 2012 (Prof. A. Albini as the supervisor). His main research interests focus on the generation of radical intermediates through photocatalyzed hydrogen atom transfer and their application in sustainable organic synthesis. In 2017, he received the Ciamician medal and the Vincenzo Caglioti international award.

Maurizio Fagnoni is currently a Full Professor in Organic Chemistry at the PhotoGreen Lab (Department of Chemistry, University of Pavia, Italy). His academic and professional background is in organic photochemistry, and his activity has always been focused on the exploration of the use of photochemistry and photocatalysis for organic synthesis. The photochemical generation of intermediates, for example, radicals and cations and radical ions by photochemical means, is the main topic of his research. Particular attention has been given to the significance of such mild synthetic procedures in the frame of the increasing interest for sustainable/green chemistry. He was the recipient in 2019 of the “Elsevier Lectureship Award” from the Japanese Photochemical Association. He was recently coeditor of the book *Photoorganocatalysis in Organic Synthesis* (World Scientific, 2019).

## ACKNOWLEDGMENTS

This review is in honor of Prof. Ilhyong Ryu on the occasion of his 70th birthday. L.C. acknowledges European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 101023615 (project name: HAT-TRICK).

## ABBREVIATIONS

ABP	aminobenzophenone
AP	acetophenone
AQ	9,10-anthraquinone
AQ-2-COOH	anthraquinone-2-carboxylic acid
AQ-2,3-diCOOH	anthraquinone-2,3-dicarboxylic acid
AQ-2-SO <sub>3</sub> Na	anthraquinone-2-sulfonic acid sodium salt



tBAQ	2- <i>tert</i> -butylanthraquinone
BLB	Black light bulb
Boc	<i>tert</i> -butoxycarbonyl
BOX	bisoxazoline
BP	benzophenone
BPO	benzoyl peroxide
BPSS	disodium benzophenondisulfonate
Brsm	based on remaining starting material
CFL	compact fluorescent lamp
CIAQ	2-chloroantraquinone
COPC	cobaloxime pyridine chloride
DBS	dibenzosuberone
DCBP	4,4'-dichlorobenzophenone
DCM	dichloromethane
DDQ	2,3-dichloro-5,6-dicyano-1,4-benzoquinone
DIAD	diisopropyl azodicarboxylate
DIPEA	<i>N,N</i> -diisopropylethylamine
DMBP	4,4'-dimethoxybenzophenone
DT	decatungstate anion
DT-BPY	[Cu <sub>4</sub> (BPY) <sub>6</sub> Cl <sub>2</sub> (W <sub>10</sub> O <sub>32</sub> )]·3H <sub>2</sub> O
DTX	3,6-dimethoxy-9 <i>H</i> -thioxanthen-9-one
EY	Eosin Y
FL	9-fluorenone
FPT	freeze pump–thaw
HAT	hydrogen atom transfer
h.p.	high pressure
l.p.	low pressure
MFC	mass flow controller
m.p.	medium pressure
MTBE	<i>tert</i> -butyl methyl ether
NaDT	sodium decatungstate
PC	photocatalyst
P.E.	petroleum ether
POC	photoorganocatalyst
PGA	phenylglyoxylic acid
PSMA	Prostate specific membrane antigen
PQ	9,10-phenanthrenequinone
PT	5,7,12,14-pentacenetetrone
PTSA	<i>p</i> -toluenesulfonic acid.
PYD	1,6-pyrenedione
Sb-Oxo	antimony-oxo tetra-( <i>p</i> -methoxyphenyl)-porphyrin
SOLFIN	solar synthesis of fine chemicals
TBAB	tetrabutyl ammonium bromide
TBADT	tetrabutylammonium decatungstate
TBDPS	<i>tert</i> -butyldiphenylsilyl
TEA	triethylamine
TFA	trifluoroacetic acid
TIPSH	triisopropylbenzenethiol
Ts	tosyl
TX	thioxanthone
Ur	uranyl cation
UrN	uranyl nitrate hexahydrate
UrP	uranyl perchlorate
XA	xanthone

## REFERENCES

- Braun, M. *Modern Enolate Chemistry*; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, Germany, 2015.
- Stolz, D.; Kazmaier, U. Metal Enolates as Synthons in Organic Chemistry. In *PATAI'S Chemistry of Functional Groups*; John Wiley & Sons, Ltd: Chichester, UK, 2010.
- Goldman, A. S.; Goldberg, K. I. Organometallic C–H Bond Activation: An Introduction. In *Activation and Functionalization of C–H Bonds*; ACS Symposium Series 2004; Vol. 885, pp 1–43.
- Hartwig, J. F.; Larsen, M. A. Undirected, Homogeneous C–H Bond Functionalization: Challenges and Opportunities. *ACS Cent. Sci.* **2016**, *2*, 281–292.
- Sambigioglio, C.; Schönbauer, D.; Blicke, R.; Dao-Huy, T.; Pototschnig, G.; Schaaf, P.; Wiesinger, T.; Zia, M. F.; Wencel-Delord, J.; Besset, T.; Maes, B. U. W.; Schnürch, M. A Comprehensive Overview of Directing Groups Applied in Metal-Catalysed C–H Functionalisation Chemistry. *Chem. Soc. Rev.* **2018**, *47*, 6603–6743.
- Rasheed, O. K.; Sun, B. Advances in Development of C–H Activation/Functionalization Using a Catalytic Directing Group. *ChemistrySelect* **2018**, *3*, 5689–5708.
- Crabtree, R. H. Introduction to Selective Functionalization of C–H Bonds. *Chem. Rev.* **2010**, *110*, 575–575.
- Davies, H. M. L.; Du Bois, J.; Yu, J.-Q. C–H Functionalization in Organic Synthesis. *Chem. Soc. Rev.* **2011**, *40*, 1855–1856.
- Doyle, M. P.; Goldberg, K. I. C–H Functionalization (special issue). *Acc. Chem. Res.* **2012**, *45*, 777–777.
- Rouquet, G.; Chatani, N. Catalytic Functionalization of C(sp<sup>2</sup>)–H and C(sp<sup>3</sup>)–H Bonds by Using Bidentate Directing Groups. *Angew. Chem., Int. Ed.* **2013**, *52*, 11726–11743.
- Hartwig, J. F. Evolution of C–H Bond Functionalization from Methane to Methodology. *J. Am. Chem. Soc.* **2016**, *138*, 2–24.
- Kärkäs, M. D. Electrochemical Strategies for C–H Functionalization and C–N Bond Formation. *Chem. Soc. Rev.* **2018**, *47*, 5786–5865.
- Wencel-Delord, J.; Glorius, F. C–H Bond Activation Enables the Rapid Construction and Late-Stage Diversification of Functional Molecules. *Nat. Chem.* **2013**, *5*, 369–375.
- Cernak, T.; Dykstra, K. D.; Tyagarajan, S.; Vachal, P.; Krska, S. W. The Medicinal Chemist's Toolbox for Late Stage Functionalization of Drug-like Molecules. *Chem. Soc. Rev.* **2016**, *45*, 546–576.
- Moir, M.; Danon, J. J.; Reekie, T. A.; Kassiou, M. An Overview of Late-Stage Functionalization in Today's Drug Discovery. *Expert Opin. Drug Discovery* **2019**, *14*, 1137–1149.
- Cannalire, R.; Pelliccia, S.; Sancineto, L.; Novellino, E.; Tron, G. C.; Giustiniano, M. Visible Light Photocatalysis in the Late-Stage Functionalization of Pharmaceutically Relevant Compounds. *Chem. Soc. Rev.* **2021**, *50*, 766–897.
- Dénès, F. Intermolecular Radical C–H Bond Activation: A Powerful Tool for Late Stage Functionalization. *Chimia* **2020**, *74*, 23–32.
- Wender, P. A.; Verma, V. A.; Paxton, T. J.; Pillow, T. H. Function-Oriented Synthesis, Step Economy, and Drug Design. *Acc. Chem. Res.* **2008**, *41*, 40–49.
- McMurray, L.; O'Hara, F.; Gaunt, M. J. Recent Developments in Natural Product Synthesis Using Metal-Catalysed C–H Bond Functionalisation. *Chem. Soc. Rev.* **2011**, *40*, 1885.
- Qiu, Y.; Gao, S. Trends in Applying C–H Oxidation to the Total Synthesis of Natural Products. *Nat. Prod. Rep.* **2016**, *33*, 562–581.
- Crabtree, R. H. Alkane C–H Activation and Functionalization with Homogeneous Transition Metal Catalysts: A Century of Progress - a New Millennium in Prospect. *J. Chem. Soc. Dalton Trans.* **2001**, 2437–2450.
- C–H Activation and Functionalization of Alkanes and Arenes. In *Organometallic Chemistry and Catalysis*; Springer Berlin Heidelberg: Berlin, Heidelberg, 2007; pp 409–429.
- Caplin, M. J.; Foley, D. J. Emergent Synthetic Methods for the Modular Advancement of sp<sup>3</sup>-Rich Fragments. *Chem. Sci.* **2021**, *12*, 4646–4660.
- Reed-Berendt, B. G.; Latham, D. E.; Dambatta, M. B.; Morrill, L. C. Borrowing Hydrogen for Organic Synthesis. *ACS Cent. Sci.* **2021**, *7*, 570–585.
- Chen, M. S.; White, M. C. A Predictably Selective Aliphatic C–H Oxidation Reaction for Complex Molecule Synthesis. *Science* **2007**, *318*, 783–787.



- (26) Cowley, R. E.; Eckert, N. A.; Vaddadi, S.; Figg, T. M.; Cundari, T. R.; Holland, P. L. Selectivity and Mechanism of Hydrogen Atom Transfer by an Isolable Imidoiron(III) Complex. *J. Am. Chem. Soc.* **2011**, *133*, 9796–9811.
- (27) Yang, Y.; Gao, W.; Wang, Y.; Wang, X.; Cao, F.; Shi, T.; Wang, Z. Recent Advances in Copper Promoted Inert C(sp<sup>3</sup>)-H Functionalization. *ACS Catal.* **2021**, *11*, 967–984.
- (28) Shen, D.; Miao, C.; Wang, S.; Xia, C.; Sun, W. Efficient Benzylic and Aliphatic C–H Oxidation with Selectivity for Methylene Sites Catalyzed by a Bioinspired Manganese Complex. *Org. Lett.* **2014**, *16*, 1108–1111.
- (29) Liu, W.; Groves, J. T. Manganese Catalyzed C–H Halogenation. *Acc. Chem. Res.* **2015**, *48*, 1727–1735.
- (30) Li, G.; Dilger, A. K.; Cheng, P. T.; Ewing, W. R.; Groves, J. T. Selective C–H Halogenation with a Highly Fluorinated Manganese Porphyrin. *Angew. Chem., Int. Ed.* **2018**, *57*, 1251–1255.
- (31) Baccalini, A.; Vergura, S.; Dolui, P.; Zanoni, G.; Maiti, D. Recent Advances in Cobalt-Catalyzed C–H Functionalizations. *Org. Biomol. Chem.* **2019**, *17*, 10119–10141.
- (32) Liu, Y.; You, T.; Wang, H.-X.; Tang, Z.; Zhou, C.-Y.; Che, C.-M. Iron- and Cobalt-Catalyzed C(sp<sup>3</sup>)-H Bond Functionalization Reactions and Their Application in Organic Synthesis. *Chem. Soc. Rev.* **2020**, *49*, 5310–5358.
- (33) Rej, S.; Chatani, N. Rhodium-Catalyzed C(sp<sup>2</sup>)- or C(sp<sup>3</sup>)-H Bond Functionalization Assisted by Removable Directing Groups. *Angew. Chem., Int. Ed.* **2019**, *58*, 8304–8329.
- (34) Choi, J.; Goldman, A. S. Ir-Catalyzed Functionalization of C–H Bonds; Andersson, P. G., Ed.; *Topics in Organometallic Chemistry*; Springer Berlin Heidelberg: Berlin, Heidelberg, 2011; Vol. 34, pp 139–167.
- (35) Li, X.; Ouyang, W.; Nie, J.; Ji, S.; Chen, Q.; Huo, Y. Recent Development on Cp\*Ir(III)-Catalyzed C–H Bond Functionalization. *ChemCatChem* **2020**, *12*, 2358–2384.
- (36) Thirunavukkarasu, V. S.; Kozhushkov, S. I.; Ackermann, L. C–H Nitrogenation and Oxygenation by Ruthenium Catalysis. *Chem. Commun.* **2014**, *50*, 29–39.
- (37) Dana, S.; Yadav, M. R.; Sahoo, A. K. Ruthenium-Catalyzed C–N and C–O Bond-Forming Processes from C–H Bond Functionalization; *Topics in Organometallic Chemistry*; Springer, Cham, 2015; Vol. 55, pp 189–215.
- (38) Bryant, J. R.; Mayer, J. M. Oxidation of C–H Bonds by [(Bpy)<sub>2</sub>(Py)Ru<sup>IV</sup>O]<sup>2+</sup> Occurs by Hydrogen Atom Abstraction. *J. Am. Chem. Soc.* **2003**, *125*, 10351–10361.
- (39) Lyons, T. W.; Sanford, M. S. Palladium-Catalyzed Ligand-Directed C–H Functionalization Reactions. *Chem. Rev.* **2010**, *110*, 1147–1169.
- (40) Neufeldt, S. R.; Sanford, M. S. Controlling Site Selectivity in Palladium-Catalyzed C–H Bond Functionalization. *Acc. Chem. Res.* **2012**, *45*, 936–946.
- (41) He, G.; Wang, B.; Nack, W. A.; Chen, G. Syntheses and Transformations of  $\alpha$ -Amino Acids via Palladium-Catalyzed Auxiliary-Directed sp<sup>3</sup> C–H Functionalization. *Acc. Chem. Res.* **2016**, *49*, 635–645.
- (42) He, C.; Whitehurst, W. G.; Gaunt, M. J. Palladium-Catalyzed C(sp<sup>3</sup>)-H Bond Functionalization of Aliphatic Amines. *Chem.* **2019**, *5*, 1031–1058.
- (43) Das, J.; Guin, S.; Maiti, D. Diverse Strategies for Transition Metal Catalyzed Distal C(sp<sup>3</sup>)-H Functionalizations. *Chem. Sci.* **2020**, *11*, 10887–10909.
- (44) *Hydrogen-Transfer Reactions*; Hynes, J. T., Klinman, J. P., Limbach, H., Schowen, R. L., Eds.; Wiley-VCH: Weinheim, 2007.
- (45) Lai, W.; Li, C.; Chen, H.; Shaik, S. Hydrogen-Abstraction Reactivity Patterns from A to Y: The Valence Bond Way. *Angew. Chem., Int. Ed.* **2012**, *51*, 5556–5578.
- (46) Mayer, J. M. Understanding Hydrogen Atom Transfer: From Bond Strengths to Marcus Theory. *Acc. Chem. Res.* **2011**, *44*, 36–46.
- (47) Tedder, J. M. The Importance of Polarity, Bond Strength and Steric Effects in Determining the Site of Attack and the Rate of Free Radical Substitution in Aliphatic Compounds. *Tetrahedron* **1982**, *38*, 313–329.
- (48) Newhouse, T.; Baran, P. S. If C–H Bonds Could Talk: Selective C–H Bond Oxidation. *Angew. Chem., Int. Ed.* **2011**, *50*, 3362–3374.
- (49) Salamone, M.; Bietti, M. Tuning Reactivity and Selectivity in Hydrogen Atom Transfer from Aliphatic C–H Bonds to Alkoxy Radicals: Role of Structural and Medium Effects. *Acc. Chem. Res.* **2015**, *48*, 2895–2903.
- (50) Costas, M.; Bietti, M. Uncovering the Complexity of the Simplest Atom Transfer Reaction. *Acc. Chem. Res.* **2018**, *51*, 2601–2602.
- (51) Bietti, M. Activation and Deactivation Strategies Promoted by Medium Effects for Selective Aliphatic C–H Bond Functionalization. *Angew. Chem., Int. Ed.* **2018**, *57*, 16618–16637.
- (52) Milan, M.; Salamone, M.; Costas, M.; Bietti, M. The Quest for Selectivity in Hydrogen Atom Transfer Based Aliphatic C–H Bond Oxygenation. *Acc. Chem. Res.* **2018**, *51*, 1984–1995.
- (53) Luo, Y.-R. *Comprehensive Handbook of Chemical Bond Energies*; CRC Press: Boca Raton, 2007.
- (54) Ghosh, M.; Singh, K. K.; Panda, C.; Weitz, A.; Hendrich, M. P.; Collins, T. J.; Dhar, B. B.; Sen Gupta, S. Formation of a Room Temperature Stable Fe<sup>V</sup>(O) Complex: Reactivity Toward Unactivated C–H Bonds. *J. Am. Chem. Soc.* **2014**, *136*, 9524–9527.
- (55) Kariofillis, S. K.; Shields, B. J.; Tekle-Smith, M. A.; Zacuto, M. J.; Doyle, A. G. Nickel/Photoredox-Catalyzed Methylation of (Hetero)Aryl Chlorides Using Trimethyl Orthoformate as a Methyl Radical Source. *J. Am. Chem. Soc.* **2020**, *142*, 7683–7689.
- (56) Chen, M. S.; White, M. C. Combined Effects on Selectivity in Fe-Catalyzed Methylene Oxidation. *Science* **2010**, *327*, 566–571.
- (57) Proksch, E.; de Meijere, A. Oxidation of Cyclopropyl Hydrocarbons with Ozone. *Angew. Chem., Int. Ed. Engl.* **1976**, *15*, 761–762.
- (58) Volker Dehmlow, E.; Heiligenstädt, N. Dimethyldioxirane Oxidations of Some Cyclopropanes. *Tetrahedron Lett.* **1996**, *37*, 5363–5364.
- (59) D'Accolti, L.; Dinoi, A.; Fusco, C.; Russo, A.; Curci, R. Oxyfunctionalization of Non-Natural Targets by Dioxiranes. 5. Selective Oxidation of Hydrocarbons Bearing Cyclopropyl Moieties 1. *J. Org. Chem.* **2003**, *68*, 7806–7810.
- (60) Roberts, B. P. Polarity-Reversal Catalysis of Hydrogen-Atom Abstraction Reactions: Concepts and Applications in Organic Chemistry. *Chem. Soc. Rev.* **1999**, *28*, 25–35.
- (61) Salamone, M.; Basili, F.; Bietti, M. Reactivity and Selectivity Patterns in Hydrogen Atom Transfer from Amino Acid C–H Bonds to the Cumyloxy Radical: Polar Effects as a Rationale for the Preferential Reaction at Proline Residues. *J. Org. Chem.* **2015**, *80*, 3643–3650.
- (62) DiLabio, G. A.; Franchi, P.; Lanzalunga, O.; Lapi, A.; Lucarini, F.; Lucarini, M.; Mazzonna, M.; Prasad, V. K.; Ticconi, B. Hydrogen Atom Transfer (HAT) Processes Promoted by the Quinolinimide-N-Oxyl Radical. A Kinetic and Theoretical Study. *J. Org. Chem.* **2017**, *82*, 6133–6141.
- (63) Salamone, M.; Ortega, V. B.; Martin, T.; Bietti, M. Hydrogen Atom Transfer from Alkanols and Alkanediols to the Cumyloxy Radical: Kinetic Evaluation of the Contribution of  $\alpha$ -C–H Activation and  $\beta$ -C–H Deactivation. *J. Org. Chem.* **2018**, *83*, 5539–5545.
- (64) Ravelli, D.; Fagnoni, M.; Fukuyama, T.; Nishikawa, T.; Ryu, I. Site-Selective C–H Functionalization by Decatungstate Anion Photocatalysis: Synergistic Control by Polar and Steric Effects Expands the Reaction Scope. *ACS Catal.* **2018**, *8*, 701–713.
- (65) Bietti, M.; Cucinotta, E.; DiLabio, G. A.; Lanzalunga, O.; Lapi, A.; Mazzonna, M.; Romero-Montalvo, E.; Salamone, M. Evaluation of Polar Effects in Hydrogen Atom Transfer Reactions from Activated Phenols. *J. Org. Chem.* **2019**, *84*, 1778–1786.
- (66) Ticconi, B.; Mazzonna, M.; Lanzalunga, O.; Lapi, A. Oxidation of  $\alpha$ -Amino Acids Promoted by the Phthalimide N-Oxyl Radical: A Kinetic and Product Study. *Tetrahedron* **2019**, *75*, 3579–3585.

- (67) Chen, K.; Baran, P. S. Total Synthesis of Eudesmane Terpenes by Site-Selective C–H Oxidations. *Nature* **2009**, *459*, 824–828.
- (68) Gómez, L.; García-Bosch, I.; Company, A.; Benet-Buchholz, J.; Polo, A.; Sala, X.; Ribas, X.; Costas, M. Stereospecific C–H Oxidation with H<sub>2</sub>O<sub>2</sub> Catalyzed by a Chemically Robust Site-Isolated Iron Catalyst. *Angew. Chem., Int. Ed.* **2009**, *48*, 5720–5723.
- (69) Prat, I.; Gómez, L.; Canta, M.; Ribas, X.; Costas, M. An Iron Catalyst for Oxidation of Alkyl C–H Bonds Showing Enhanced Selectivity for Methylenic Sites. *Chem. - Eur. J.* **2013**, *19*, 1908–1913.
- (70) Zou, L.; Paton, R. S.; Eschenmoser, A.; Newhouse, T. R.; Baran, P. S.; Houk, K. N. Enhanced Reactivity in Dioxirane C–H Oxidations via Strain Release: A Computational and Experimental Study. *J. Org. Chem.* **2013**, *78*, 4037–4048.
- (71) Moteki, S. A.; Usui, A.; Zhang, T.; Solorio Alvarado, C. R.; Maruoka, K. Site-Selective Oxidation of Unactivated C(sp<sup>3</sup>)–H Bonds with Hypervalent Iodine(III) Reagents. *Angew. Chem., Int. Ed.* **2013**, *52*, 8657–8660.
- (72) Schmidt, V. A.; Quinn, R. K.; Brusoe, A. T.; Alexanian, E. J. Site-Selective Aliphatic C–H Bromination Using N -Bromoamides and Visible Light. *J. Am. Chem. Soc.* **2014**, *136*, 14389–14392.
- (73) Salamone, M.; Ortega, V. B.; Bietti, M. Enhanced Reactivity in Hydrogen Atom Transfer from Tertiary Sites of Cyclohexanes and Decalins via Strain Release: Equatorial C–H Activation vs Axial C–H Deactivation. *J. Org. Chem.* **2015**, *80*, 4710–4715.
- (74) Milan, M.; Bietti, M.; Costas, M. Highly Enantioselective Oxidation of Nonactivated Aliphatic C–H Bonds with Hydrogen Peroxide Catalyzed by Manganese Complexes. *ACS Cent. Sci.* **2017**, *3*, 196–204.
- (75) Salamone, M.; Martin, T.; Milan, M.; Costas, M.; Bietti, M. Electronic and Torsional Effects on Hydrogen Atom Transfer from Aliphatic C–H Bonds: A Kinetic Evaluation via Reaction with the Cumyloxy Radical. *J. Org. Chem.* **2017**, *82*, 13542–13549.
- (76) Chen, K.; Eschenmoser, A.; Baran, P. S. Strain Release in C–H Bond Activation? *Angew. Chem., Int. Ed.* **2009**, *48*, 9705–9708.
- (77) Dondi, D.; Ravelli, D.; Fagnoni, M.; Mella, M.; Molinari, A.; Maldotti, A.; Albini, A. Regio- and Stereoselectivity in the Decatungstate Photocatalyzed Alkylation of Alkenes by Alkylcyclohexanes. *Chem. - Eur. J.* **2009**, *15*, 7949–7957.
- (78) De Vleeschouwer, F.; Van Speybroeck, V.; Waroquier, M.; Geerlings, P.; De Proft, F. Electrophilicity and Nucleophilicity Index for Radicals. *Org. Lett.* **2007**, *9*, 2721–2724.
- (79) Li, C.; Danovich, D.; Shaik, S. Blended Hydrogen Atom Abstraction and Proton-Coupled Electron Transfer Mechanisms of Closed-Shell Molecules. *Chem. Sci.* **2012**, *3*, 1903–1918.
- (80) Tanielian, C.; Cougnon, F.; Seghrouchni, R. Acetone, a Substrate and a New Solvent in Decatungstate Photocatalysis. *J. Mol. Catal. A: Chem.* **2007**, *262*, 164–169.
- (81) Dondi, D.; Fagnoni, M.; Albini, A. Tetrabutylammonium Decatungstate-Photosensitized Alkylation of Electrophilic Alkenes: Convenient Functionalization of Aliphatic C–H Bonds. *Chem. - Eur. J.* **2006**, *12*, 4153–4163.
- (82) Naguib, Y. M. A.; Steel, C.; Cohen, S. G.; Young, M. A. Photoreduction of Benzophenone by Acetonitrile: Correlation of Rates of Hydrogen Abstraction from RH with the Ionization Potentials of the Radicals R•. *J. Phys. Chem.* **1987**, *91*, 3033–3036.
- (83) Okada, M.; Fukuyama, T.; Yamada, K.; Ryu, I.; Ravelli, D.; Fagnoni, M. Sunlight Photocatalyzed Regioselective  $\beta$ -Alkylation and Acylation of Cyclopentanones. *Chem. Sci.* **2014**, *5*, 2893–2898.
- (84) Ueda, M.; Kitano, A.; Matsubara, H. A computational study of site-selective hydrogen abstraction by sulfate radical anion. *Org. Biomol. Chem.* **2021**, *19*, 4775–4782.
- (85) Milan, M.; Carboni, G.; Salamone, M.; Costas, M.; Bietti, M. Tuning Selectivity in Aliphatic C–H Bond Oxidation of N -Alkylamides and Phthalimides Catalyzed by Manganese Complexes. *ACS Catal.* **2017**, *7*, 5903–5911.
- (86) Tanko, J. M.; Suleman, N. K. Solvent Effects in the Reactions of Neutral Free Radicals. In *Energetics of Organic Free Radicals*; Springer Netherlands: Dordrecht, 1996; pp 224–293.
- (87) Bietti, M.; Salamone, M. Kinetic Solvent Effects on Hydrogen Abstraction Reactions from Carbon by the Cumyloxy Radical. The Role of Hydrogen Bonding. *Org. Lett.* **2010**, *12*, 3654–3657.
- (88) Bietti, M.; Martella, R.; Salamone, M. Understanding Kinetic Solvent Effects on Hydrogen Abstraction Reactions from Carbon by the Cumyloxy Radical. *Org. Lett.* **2011**, *13*, 6110–6113.
- (89) Salamone, M.; Giammarioli, I.; Bietti, M. Kinetic Solvent Effects on Hydrogen Abstraction Reactions from Carbon by the Cumyloxy Radical. The Importance of Solvent Hydrogen-Bond Interactions with the Substrate and the Abstracting Radical. *J. Org. Chem.* **2011**, *76*, 4645–4651.
- (90) Salamone, M.; Mangiacapra, L.; Bietti, M. Kinetic Solvent Effects on the Reactions of the Cumyloxy Radical with Tertiary Amides. Control over the Hydrogen Atom Transfer Reactivity and Selectivity through Solvent Polarity and Hydrogen Bonding. *J. Org. Chem.* **2015**, *80*, 1149–1154.
- (91) Jeffrey, J. L.; Terrett, J. A.; MacMillan, D. W. C. O-H Hydrogen Bonding Promotes H-Atom Transfer from C–H Bonds for C-Alkylation of Alcohols. *Science* **2015**, *349*, 1532–1536.
- (92) Dewanji, A.; Mück-Lichtenfeld, C.; Studer, A. Radical Hydrodeiodination of Aryl, Alkenyl, Alkynyl, and Alkyl Iodides with an Alcoholate as Organic Chain Reductant through Electron Catalysis. *Angew. Chem., Int. Ed.* **2016**, *55*, 6749–6752.
- (93) Salamone, M.; Giammarioli, I.; Bietti, M. Tuning Hydrogen Atom Abstraction from the Aliphatic C–H Bonds of Basic Substrates by Protonation. Control over Selectivity by C–H Deactivation. *Chem. Sci.* **2013**, *4*, 3255–3262.
- (94) Milan, M.; Salamone, M.; Bietti, M. Hydrogen Atom Transfer from 1,*n*-Alkanediamines to the Cumyloxy Radical. Modulating C–H Deactivation Through Acid–Base Interactions and Solvent Effects. *J. Org. Chem.* **2014**, *79*, 5710–5716.
- (95) Howell, J. M.; Feng, K.; Clark, J. R.; Trzepakowski, L. J.; White, M. C. Remote Oxidation of Aliphatic C–H Bonds in Nitrogen-Containing Molecules. *J. Am. Chem. Soc.* **2015**, *137*, 14590–14593.
- (96) Lee, M.; Sanford, M. S. Platinum-Catalyzed, Terminal-Selective C(sp<sup>3</sup>)–H Oxidation of Aliphatic Amines. *J. Am. Chem. Soc.* **2015**, *137*, 12796–12799.
- (97) Bietti, M.; Forcina, V.; Lanzalunga, O.; Lapi, A.; Martin, T.; Mazzonna, M.; Salamone, M. Kinetic Study of the Reaction of the Phthalimide- N -Oxyl Radical with Amides: Structural and Medium Effects on the Hydrogen Atom Transfer Reactivity and Selectivity. *J. Org. Chem.* **2016**, *81*, 11924–11931.
- (98) Mbofana, C. T.; Chong, E.; Lawniczak, J.; Sanford, M. S. Iron-Catalyzed Oxyfunctionalization of Aliphatic Amines at Remote Benzylic C–H Sites. *Org. Lett.* **2016**, *18*, 4258–4261.
- (99) Salamone, M.; Carboni, G.; Bietti, M. Fine Control over Site and Substrate Selectivity in Hydrogen Atom Transfer-Based Functionalization of Aliphatic C–H Bonds. *J. Org. Chem.* **2016**, *81*, 9269–9278.
- (100) Lee, M.; Sanford, M. S. Remote C(sp<sup>3</sup>)–H Oxygenation of Protonated Aliphatic Amines with Potassium Persulfate. *Org. Lett.* **2017**, *19*, 572–575.
- (101) Mack, J. B. C.; Gipson, J. D.; Du Bois, J.; Sigman, M. S. Ruthenium-Catalyzed C–H Hydroxylation in Aqueous Acid Enables Selective Functionalization of Amine Derivatives. *J. Am. Chem. Soc.* **2017**, *139*, 9503–9506.
- (102) Dantignana, V.; Milan, M.; Cussó, O.; Company, A.; Bietti, M.; Costas, M. Chemoselective Aliphatic C–H Bond Oxidation Enabled by Polarity Reversal. *ACS Cent. Sci.* **2017**, *3*, 1350–1358.
- (103) Gaster, E.; Kozuch, S.; Pappo, D. Selective Aerobic Oxidation of Methylarenes to Benzaldehydes Catalyzed by N -Hydroxyphthalimide and Cobalt(II) Acetate in Hexafluoropropan-2-ol. *Angew. Chem., Int. Ed.* **2017**, *56*, S912–S915.
- (104) Bietti, M.; Lanzalunga, O.; Lapi, A.; Martin, T.; Mazzonna, M.; Polin, M.; Salamone, M. Aerobic Oxidation of 4-Alkyl-N,N-Dimethylbenzylamines Catalyzed by N-Hydroxyphthalimide: Protonation-Driven Control over Regioselectivity. *J. Org. Chem.* **2017**, *82*, 5761–5768.

- (105) Pipitone, L. M.; Carboni, G.; Sorrentino, D.; Galeotti, M.; Salamone, M.; Bietti, M. Enhancing Reactivity and Site-Selectivity in Hydrogen Atom Transfer from Amino Acid C–H Bonds via Deprotonation. *Org. Lett.* **2018**, *20*, 808–811.
- (106) Cianfanelli, M.; Olivo, G.; Milan, M.; Klein Gebbink, R. J. M.; Ribas, X.; Bietti, M.; Costas, M. Enantioselective C–H Lactonization of Unactivated Methylenes Directed by Carboxylic Acids. *J. Am. Chem. Soc.* **2020**, *142*, 1584–1593.
- (107) Borrell, M.; Gil-Caballero, S.; Bietti, M.; Costas, M. Site-Selective and Product Chemoselective Aliphatic C–H Bond Hydroxylation of Polyhydroxylated Substrates. *ACS Catal.* **2020**, *10*, 4702–4709.
- (108) Salamone, M.; Mangiacapra, L.; DiLabio, G. A.; Bietti, M. Effect of Metal Ions on the Reactions of the Cumyloxyl Radical with Hydrogen Atom Donors. Fine Control on Hydrogen Abstraction Reactivity Determined by Lewis Acid–Base Interactions. *J. Am. Chem. Soc.* **2013**, *135*, 415–423.
- (109) Salamone, M.; Carboni, G.; Mangiacapra, L.; Bietti, M. Binding to Redox-Inactive Alkali and Alkaline Earth Metal Ions Strongly Deactivates the C–H Bonds of Tertiary Amides toward Hydrogen Atom Transfer to Reactive Oxygen Centered Radicals. *J. Org. Chem.* **2015**, *80*, 9214–9223.
- (110) Salamone, M.; Mangiacapra, L.; Carboni, G.; Bietti, M. Hydrogen Atom Transfer from Tertiary Alkanamides to the Cumyloxyl Radical. The Role of Substrate Structure on Alkali and Alkaline Earth Metal Ion Induced C–H Bond Deactivation. *Tetrahedron* **2016**, *72*, 7757–7763.
- (111) Martin, T.; Salamone, M.; Bietti, M. Hydrogen Atom Transfer from 1,2- and 1,3-Diols to the Cumyloxyl Radical. The Role of Structural Effects on Metal-Ion Induced C–H Bond Deactivation. *Chem. Commun.* **2019**, *55*, 5227–5230.
- (112) Schultz, D. M.; Lévesque, F.; DiRocco, D. A.; Reibarkh, M.; Ji, Y.; Joyce, L. A.; Dropinski, J. F.; Sheng, H.; Sherry, B. D.; Davies, I. W. Oxyfunctionalization of the Remote C–H Bonds of Aliphatic Amines by Decatungstate Photocatalysis. *Angew. Chem., Int. Ed.* **2017**, *56*, 15274–15278.
- (113) Olivo, G.; Farinelli, G.; Barbieri, A.; Lanzalunga, O.; Di Stefano, S.; Costas, M. Supramolecular Recognition Allows Remote, Site-Selective C–H Oxidation of Methylenic Sites in Linear Amines. *Angew. Chem., Int. Ed.* **2017**, *56*, 16347–16351.
- (114) Olivo, G.; Capocasa, G.; Lanzalunga, O.; Di Stefano, S.; Costas, M. Enzyme-like Substrate-Selectivity in C–H Oxidation Enabled by Recognition. *Chem. Commun.* **2019**, *55*, 917–920.
- (115) Olivo, G.; Capocasa, G.; Ticconi, B.; Lanzalunga, O.; Di Stefano, S.; Costas, M. Predictable Selectivity in Remote C–H Oxidation of Steroids: Analysis of Substrate Binding Mode. *Angew. Chem., Int. Ed.* **2020**, *59*, 12703–12708.
- (116) Stateman, L.; Nakafuku, K.; Nagib, D. Remote C–H Functionalization via Selective Hydrogen Atom Transfer. *Synthesis* **2018**, *50*, 1569–1586.
- (117) Sarkar, S.; Cheung, K. P. S.; Gevorgyan, V. C–H Functionalization Reactions Enabled by Hydrogen Atom Transfer to Carbon-Centered Radicals. *Chem. Sci.* **2020**, *11*, 12974–12993.
- (118) Kumar, G.; Pradhan, S.; Chatterjee, I. N-Centered Radical Directed Remote C–H Bond Functionalization via Hydrogen Atom Transfer. *Chem. - Asian J.* **2020**, *15*, 651–672.
- (119) Costas, M. Remote Oxidation of Aliphatic C–H Bonds with Biologically Inspired Catalysts. In *Remote C–H Bond Functionalizations*; Maiti, D., Guin, S., Eds.; Wiley-VCH: Weinheim, 2021; pp 383–421.
- (120) Sambiagio, C.; Maes, B. U. W. Non-Directed Functionalization of Distal C(sp<sup>3</sup>)-H Bonds. In *Remote C–H Bond Functionalizations*; Maiti, D., Guin, S., Eds.; Wiley-VCH: Weinheim, 2021; pp 343–382.
- (121) Li, W.; Zhu, C. Radically Initiated Distal C(sp<sup>3</sup>)-H Functionalization. In *Remote C–H Bond Functionalizations*; Maiti, D., Guin, S., Eds.; Wiley-VCH: Weinheim, 2021; pp 315–341.
- (122) Li, Y.; Zhang, Q.; Shi, B. Directing Group Assisted Distal C(sp<sup>3</sup>)-H Functionalization of Aliphatic Substrates. In *Remote C–H Bond Functionalizations*; Maiti, D., Guin, S., Eds.; Wiley-VCH: Weinheim, 2021; pp 279–314.
- (123) Čeković, Z. Reactions of  $\delta$ -Carbon Radicals Generated by 1,5-Hydrogen Transfer to Alkoxy Radicals. *Tetrahedron* **2003**, *59*, 8073–8090.
- (124) Nechab, M.; Mondal, S.; Bertrand, M. P. 1,n -Hydrogen-Atom Transfer (HAT) Reactions in Which n  $\neq$  5: An Updated Inventory. *Chem. - Eur. J.* **2014**, *20*, 16034–16059.
- (125) Fokin, A. A.; Schreiner, P. R. Selective Alkane Transformations via Radicals and Radical Cations: Insights into the Activation Step from Experiment and Theory. *Chem. Rev.* **2002**, *102*, 1551–1594.
- (126) Tsui, E.; Wang, H.; Knowles, R. R. Catalytic Generation of Alkoxy Radicals from Unfunctionalized Alcohols. *Chem. Sci.* **2020**, *11*, 11124–11141.
- (127) Wu, X.; Zhu, C. Recent Advances in Alkoxy Radical-Promoted C–C and C–H Bond Functionalization Starting from Free Alcohols. *Chem. Commun.* **2019**, *55*, 9747–9756.
- (128) Guo, J.-J.; Hu, A.; Zuo, Z. Photocatalytic Alkoxy Radical-Mediated Transformations. *Tetrahedron Lett.* **2018**, *59*, 2103–2111.
- (129) Melone, L.; Punta, C. N-Hydroxyphthalimide (NHPI)-Organocatalyzed Aerobic Oxidations: Advantages, Limits, and Industrial Perspectives. In *Liquid Phase Aerobic Oxidation Catalysis: Industrial Applications and Academic Perspectives*; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, Germany, 2016; pp 253–265.
- (130) Quinn, R. K.; Konst, Z. A.; Michalak, S. E.; Schmidt, Y.; Szklarski, A. R.; Flores, A. R.; Nam, S.; Horne, D. A.; Vanderwal, C. D.; Alexanian, E. J. Site-Selective Aliphatic C–H Chlorination Using N-Chloroamides Enables a Synthesis of Chlorolissoclimide. *J. Am. Chem. Soc.* **2016**, *138*, 696–702.
- (131) Czaplowski, W. L.; Na, C. G.; Alexanian, E. J. C–H Xanthylation: A Synthetic Platform for Alkane Functionalization. *J. Am. Chem. Soc.* **2016**, *138*, 13854–13857.
- (132) Tanaka, H.; Sakai, K.; Kawamura, A.; Oisaki, K.; Kanai, M. Sulfonamides as New Hydrogen Atom Transfer (HAT) Catalysts for Photoredox Allylic and Benzylic C–H Arylations. *Chem. Commun.* **2018**, *54*, 3215–3218.
- (133) Ryder, A. S. H.; Cunningham, W. B.; Ballantyne, G.; Mules, T.; Kinsella, A. G.; Turner-Dore, J.; Alder, C. M.; Edwards, L. J.; McKay, B. S. J.; Grayson, M. N.; Cresswell, A. J. Photocatalytic  $\alpha$ -Tertiary Amine Synthesis via C–H Alkylation of Unmasked Primary Amines. *Angew. Chem., Int. Ed.* **2020**, *59*, 14986–14991.
- (134) Lu, M.; Zhang, T.; Tan, D.; Chen, C.; Zhang, Y.; Huang, M.; Cai, S. Visible-Light-Promoted Oxidative Alkylarylation of N-Aryl/Benzoyl Acrylamides Through Direct C–H Bond Functionalization. *Adv. Synth. Catal.* **2019**, *361*, 4237–4242.
- (135) Li, W.; Duan, Y.; Zhang, M.; Cheng, J.; Zhu, C. A Photoredox Catalyzed Radical–Radical Coupling Reaction: Facile Access to Multi-Substituted Nitrogen Heterocycles. *Chem. Commun.* **2016**, *52*, 7596–7599.
- (136) Jin, J.; MacMillan, D. W. C. Alcohols as Alkylating Agents in Heteroarene C–H Functionalization. *Nature* **2015**, *525*, 87–90.
- (137) Hager, D.; MacMillan, D. W. C. Activation of C–H Bonds via the Merger of Photoredox and Organocatalysis: A Coupling of Benzylic Ethers with Schiff Bases. *J. Am. Chem. Soc.* **2014**, *136*, 16986–16989.
- (138) Cuthbertson, J. D.; MacMillan, D. W. C. The Direct Arylation of Allylic sp<sup>3</sup> C–H Bonds via Organic and Photoredox Catalysis. *Nature* **2015**, *519*, 74–77.
- (139) Qvortrup, K.; Rankic, D. A.; MacMillan, D. W. C. A General Strategy for Organocatalytic Activation of C–H Bonds via Photoredox Catalysis: Direct Arylation of Benzylic Ethers. *J. Am. Chem. Soc.* **2014**, *136*, 626–629.
- (140) Lu, P.; Hou, T.; Gu, X.; Li, P. Visible-Light-Promoted Conversion of Alkyl Benzyl Ether to Alkyl Ester or Alcohol via O- $\alpha$ -sp<sup>3</sup> C–H Cleavage. *Org. Lett.* **2015**, *17*, 1954–1957.
- (141) Hou, T.; Lu, P.; Li, P. Visible-Light-Mediated Benzylic sp<sup>3</sup> C–H Bond Functionalization to C–Br or C–N Bond. *Tetrahedron Lett.* **2016**, *57*, 2273–2276.



- (142) Fadeyi, O. O.; Mousseau, J. J.; Feng, Y.; Allais, C.; Nuhant, P.; Chen, M. Z.; Pierce, B.; Robinson, R. Visible-Light-Driven Photocatalytic Initiation of Radical Thiol–Ene Reactions Using Bismuth Oxide. *Org. Lett.* **2015**, *17*, 5756–5759.
- (143) Keylor, M. H.; Park, J. E.; Wallentin, C.-J.; Stephenson, C. R. J. Photocatalytic Initiation of Thiol–Ene Reactions: Synthesis of Thiomorpholin-3-Ones. *Tetrahedron* **2014**, *70*, 4264–4269.
- (144) Kariofillis, S. K.; Doyle, A. G. Synthetic and Mechanistic Implications of Chlorine Photoelimination in Nickel/Photoredox C(sp<sup>3</sup>)–H Cross-Coupling. *Acc. Chem. Res.* **2021**, *54*, 988–1000.
- (145) Xu, P.; Chen, P.; Xu, H. Scalable Photoelectrochemical Dehydrogenative Cross-Coupling of Heteroarenes with Aliphatic C–H Bonds. *Angew. Chem., Int. Ed.* **2020**, *59*, 14275–14280.
- (146) McMillan, A. J.; Sieńkowska, M.; Di Lorenzo, P.; Gransbury, G. K.; Chilton, N. F.; Salamone, M.; Ruffoni, A.; Bietti, M.; Leonori, D. Practical and Selective sp<sup>3</sup> C–H Bond Chlorination via Aminium Radicals. *Angew. Chem., Int. Ed.* **2021**, *60*, 7132–7139.
- (147) Shaw, M. H.; Shurtleff, V. W.; Terrett, J. A.; Cuthbertson, J. D.; MacMillan, D. W. C. Native Functionality in Triple Catalytic Cross-Coupling: sp<sup>3</sup> C–H Bonds as Latent Nucleophiles. *Science* **2016**, *352*, 1304–1308.
- (148) Xiao, W.; Wang, X.; Liu, R.; Wu, J. Quinuclidine and Its Derivatives as Hydrogen-Atom-Transfer Catalysts in Photoinduced Reactions. *Chin. Chem. Lett.* **2021**, *32*, 1847–1856.
- (149) Saito, M.; Kawamata, Y.; Meanwell, M.; Navratil, R.; Chiodi, D.; Carlson, E.; Hu, P.; Chen, L.; Udyavara, S.; Kingston, C.; Tanwar, M.; Tyagi, S.; McKillican, B. P.; Gichinga, M. G.; Schmidt, M. A.; Eastgate, M. D.; Lambert, M.; He, C.; Tang, T.; Malapit, C. A.; Sigman, M. S.; Minter, S. D.; Neurock, M.; Baran, P. S. N-Ammonium Ylide Mediators for Electrochemical C–H Oxidation. *J. Am. Chem. Soc.* **2021**, *143*, 7859–7867.
- (150) González-Núñez, M. E.; Castellano, G.; Andreu, C.; Royo, J.; Báguena, M.; Mello, R.; Asensio, G. Influence of Remote Substituents on the Equatorial/Axial Selectivity in the Monooxygenation of Methylene C–H Bonds of Substituted Cyclohexanes. *J. Am. Chem. Soc.* **2001**, *123*, 7487–7491.
- (151) Mello, R.; Fiorentino, M.; Fusco, C.; Curci, R. Oxidations by Methyl(Trifluoromethyl)Dioxirane. 2. Oxyfunctionalization of Saturated Hydrocarbons. *J. Am. Chem. Soc.* **1989**, *111*, 6749–6757.
- (152) Nam, W. High-Valent Iron(IV)–Oxo Complexes of Heme and Non-Heme Ligands in Oxygenation Reactions. *Acc. Chem. Res.* **2007**, *40*, 522–531.
- (153) Stephenson, C.; Yoon, T.; MacMillan, D. W. C. *Visible Light Photocatalysis in Organic Chemistry*; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, Germany, 2018.
- (154) Fagnoni, M.; Protti, S.; Ravelli, D. *Photoorganocatalysis in Organic Synthesis*; Catalytic Science Series; World Scientific (Europe), 2019; Vol. 18.
- (155) Albini, A.; Fagnoni, M. *Photochemically-Generated Intermediates in Synthesis*; John Wiley & Sons, Inc.: Hoboken, NJ, 2013.
- (156) *Chemical Photocatalysis*; König, B., Ed.; De Gruyter: Berlin, Boston, 2013.
- (157) Palmisano, G.; Augugliaro, V.; Pagliaro, M.; Palmisano, L. Photocatalysis: A Promising Route for 21st Century Organic Chemistry. *Chem. Commun.* **2007**, 3425–3437.
- (158) Fagnoni, M.; Dondi, D.; Ravelli, D.; Albini, A. Photocatalysis for the Formation of the C–C Bond. *Chem. Rev.* **2007**, *107*, 2725–2756.
- (159) Ravelli, D.; Dondi, D.; Fagnoni, M.; Albini, A. Photocatalysis. A Multi-Faceted Concept for Green Chemistry. *Chem. Soc. Rev.* **2009**, *38*, 1999–2011.
- (160) Ravelli, D.; Fagnoni, M.; Albini, A. Photoorganocatalysis. What For? *Chem. Soc. Rev.* **2013**, *42*, 97–113.
- (161) Staveness, D.; Bosque, I.; Stephenson, C. R. J. Free Radical Chemistry Enabled by Visible Light-Induced Electron Transfer. *Acc. Chem. Res.* **2016**, *49*, 2295–2306.
- (162) Douglas, J. J.; Sevrin, M. J.; Stephenson, C. R. J. Visible Light Photocatalysis: Applications and New Disconnections in the Synthesis of Pharmaceutical Agents. *Org. Process Res. Dev.* **2016**, *20*, 1134–1147.
- (163) Gentry, E. C.; Knowles, R. R. Synthetic Applications of Proton-Coupled Electron Transfer. *Acc. Chem. Res.* **2016**, *49*, 1546–1556.
- (164) Reiser, O. Shining Light on Copper: Unique Opportunities for Visible-Light-Catalyzed Atom Transfer Radical Addition Reactions and Related Processes. *Acc. Chem. Res.* **2016**, *49*, 1990–1996.
- (165) Goddard, J.-P.; Ollivier, C.; Fensterbank, L. Photoredox Catalysis for the Generation of Carbon Centered Radicals. *Acc. Chem. Res.* **2016**, *49*, 1924–1936.
- (166) Skubi, K. L.; Blum, T. R.; Yoon, T. P. Dual Catalysis Strategies in Photochemical Synthesis. *Chem. Rev.* **2016**, *116*, 10035–10074.
- (167) Levin, M. D.; Kim, S.; Toste, F. D. Photoredox Catalysis Unlocks Single-Electron Elementary Steps in Transition Metal Catalyzed Cross-Coupling. *ACS Cent. Sci.* **2016**, *2*, 293–301.
- (168) Xie, J.; Jin, H.; Hashmi, A. S. K. The Recent Achievements of Redox-Neutral Radical C–C Cross-Coupling Enabled by Visible-Light. *Chem. Soc. Rev.* **2017**, *46*, 5193–5203.
- (169) Lee, K. N.; Ngai, M.-Y. Recent Developments in Transition-Metal Photoredox-Catalysed Reactions of Carbonyl Derivatives. *Chem. Commun.* **2017**, 53, 13093–13112.
- (170) Matsui, J. K.; Lang, S. B.; Heitz, D. R.; Molander, G. A. Photoredox-Mediated Routes to Radicals: The Value of Catalytic Radical Generation in Synthetic Methods Development. *ACS Catal.* **2017**, *7*, 2563–2575.
- (171) Twilton, J.; Le, C.; Zhang, P.; Shaw, M. H.; Evans, R. W.; MacMillan, D. W. C. The Merger of Transition Metal and Photocatalysis. *Nat. Rev. Chem.* **2017**, *1*, 0052.
- (172) Zhang, L.; Meggers, E. Steering Asymmetric Lewis Acid Catalysis Exclusively with Octahedral Metal-Centered Chirality. *Acc. Chem. Res.* **2017**, *50*, 320–330.
- (173) Savateev, A.; Antonietti, M. Heterogeneous Organocatalysis for Photoredox Chemistry. *ACS Catal.* **2018**, *8*, 9790–9808.
- (174) Revathi, L.; Ravindar, L.; Fang, W.-Y.; Rakesh, K. P.; Qin, H.-L. Visible Light-Induced C–H Bond Functionalization: A Critical Review. *Adv. Synth. Catal.* **2018**, *360*, 4652–4698.
- (175) Silvi, M.; Melchiorre, P. Enhancing the Potential of Enantioselective Organocatalysis with Light. *Nature* **2018**, *554*, 41–49.
- (176) Sambiagio, C.; Noël, T. Flow Photochemistry: Shine Some Light on Those Tubes! *Trends Chem.* **2020**, *2*, 92–106.
- (177) Guillemard, L.; Wencel-Delord, J. When Metal-Catalyzed C–H Functionalization Meets Visible-Light Photocatalysis. *Beilstein J. Org. Chem.* **2020**, *16*, 1754–1804.
- (178) Crespi, S.; Fagnoni, M. Generation of Alkyl Radicals: From the Tyranny of Tin to the Photon Democracy. *Chem. Rev.* **2020**, *120*, 9790–9833.
- (179) Prentice, C.; Morrisson, J.; Smith, A. D.; Zysman-Colman, E. Recent Developments in Enantioselective Photocatalysis. *Beilstein J. Org. Chem.* **2020**, *16*, 2363–2441.
- (180) Lenardon, G. V. A.; Nicchio, L.; Fagnoni, M. Photogenerated Electrophilic Radicals for the Umpolung of Enolate Chemistry. *J. Photochem. Photobiol., C* **2021**, *46*, 100387.
- (181) Sideri, I. K.; Voutyritsa, E.; Kokotos, C. G. Photoorganocatalysis, Small Organic Molecules and Light in the Service of Organic Synthesis: The Awakening of a Sleeping Giant. *Org. Biomol. Chem.* **2018**, *16*, 4596–4614.
- (182) Protti, S.; Fagnoni, M.; Ravelli, D. Photocatalytic C–H Activation by Hydrogen-Atom Transfer in Synthesis. *ChemCatChem* **2015**, *7*, 1516–1523.
- (183) Capaldo, L.; Ravelli, D. Hydrogen Atom Transfer (HAT): A Versatile Strategy for Substrate Activation in Photocatalyzed Organic Synthesis. *Eur. J. Org. Chem.* **2017**, *2017*, 2056–2071.
- (184) Capaldo, L.; Quadri, L. L.; Ravelli, D. Photocatalytic Hydrogen Atom Transfer: The Philosopher’s Stone for Late-Stage Functionalization? *Green Chem.* **2020**, *22*, 3376–3396.



- (185) Cao, H.; Tang, X.; Tang, H.; Yuan, Y.; Wu, J. Photoinduced Intermolecular Hydrogen Atom Transfer Reactions in Organic Synthesis. *Chem. Catal* **2021**, DOI: 10.1016/j.checcat.2021.04.008.
- (186) Hoffmann, N. Electron and Hydrogen Transfer in Organic Photochemical Reactions. *J. Phys. Org. Chem.* **2015**, *28*, 121–136.
- (187) *IUPAC Compendium of Chemical Terminology*; Nič, M., Jiráč, J., Kořata, B., Jenkins, A., McNaught, A., Eds.; IUPAC: Research Triangle Park, NC, 2009.
- (188) Capaldo, L.; Ravelli, D. The Dark Side of Photocatalysis: One Thousand Ways to Close the Cycle. *Eur. J. Org. Chem.* **2020**, *2020*, 2783–2806.
- (189) Dantas, J. A.; Correia, J. T. M.; Paixão, M. W.; Corrêa, A. G. Photochemistry of Carbonyl Compounds: Application in Metal-Free Reactions. *ChemPhotoChem.* **2019**, *3*, 506–520.
- (190) Zhu, D.-L.; Young, D. J.; Li, H.-X. Carbonyl-Photoredox/Metal Dual Catalysis: Applications in Organic Synthesis. *Synthesis* **2020**, *52*, 3493–3510.
- (191) Perez-Prieto, J.; Galian, R.; Miranda, M. Diaryl Ketones as Photoactivators. *Mini-Rev. Org. Chem.* **2006**, *3*, 117–135.
- (192) Kamijo, S. Ketones and Aldehydes. In *Photoorganocatalysis in Organic Synthesis*; Fagnoni, M., Protti, S., Ravelli, D., Eds.; World Scientific (Europe), 2019; pp 1–37.
- (193) Kamijo, S. C–H Activation via Radical Processes Using Photo-Excited Ketones. *Topics in Organometallic Chemistry*; Springer, Cham, 2018; Vol. 54, pp 71–92.
- (194) Das, A.; Thomas, K. R. J. Facile Thiol–Ene Click Protocol Using Benzil as Sensitizer and White LED as Light Source. *Eur. J. Org. Chem.* **2020**, *2020*, 7214
- (195) Kuhn, H. J.; Goerner, H. Triplet State and Photodecarboxylation of Phenylglyoxylic Acid in the Presence of Water. *J. Phys. Chem.* **1988**, *92*, 6208–6219.
- (196) Itoh, A. Quinones. In *Photoorganocatalysis in Organic Synthesis*; Fagnoni, M., Protti, S., Ravelli, D., Eds.; World Scientific (Europe), 2019; pp 39–70.
- (197) Cervantes-González, J.; Vosburg, D. A.; Mora-Rodriguez, S. E.; Vázquez, M. A.; Zepeda, L. G.; Villegas Gómez, C.; Lagunas-Rivera, S. Anthraquinones: Versatile Organic Photocatalysts. *Chem-CatChem* **2020**, *12*, 3811–3827.
- (198) de Lucas, N. C.; Elias, M. M.; Firme, C. L.; Corrêa, R. J.; Garden, S. J.; Netto-Ferreira, J. C.; Nicodem, D. E. A Combined Laser Flash Photolysis, Density Functional Theory and Atoms in Molecules Study of the Photochemical Hydrogen Abstraction by Pyrene-4,5-Dione. *J. Photochem. Photobiol., A* **2009**, *201*, 1–7.
- (199) Xiao, W.-J.; Hu, X.-Q.; Chen, J.-R. Oxygen Heterocycles: Eosin Derivatives. In *Photoorganocatalysis in Organic Synthesis*; Fagnoni, M., Protti, S., Ravelli, D., Eds.; World Scientific (Europe), 2019; pp 243–286.
- (200) Yan, D.-M.; Chen, J.-R.; Xiao, W.-J. New Roles for Photoexcited Eosin Y in Photochemical Reactions. *Angew. Chem., Int. Ed.* **2019**, *58*, 378–380.
- (201) Hill, C. L. Introduction of Functionality into Unactivated Carbon-Hydrogen Bonds. Catalytic Generation and Nonconventional Utilization of Organic Radicals. *Synlett* **1995**, *1995*, 127–132.
- (202) Tzirakis, M. D.; Lykakis, I. N.; Orfanopoulos, M. Decatungstate as an Efficient Photocatalyst in Organic Chemistry. *Chem. Soc. Rev.* **2009**, *38*, 2609.
- (203) N. Lykakis, I.; Evgenidou, E.; Orfanopoulos, M. Photocatalysis and Polyoxo-Anion Decatungstate in Organic Chemistry: A Manifold Concept for Green Chemistry. *Curr. Org. Chem.* **2012**, *16*, 2400–2414.
- (204) Ravelli, D.; Protti, S.; Fagnoni, M. Decatungstate Anion for Photocatalyzed “Window Ledge” Reactions. *Acc. Chem. Res.* **2016**, *49*, 2232–2242.
- (205) Lazzaroni, S.; Ravelli, D.; Protti, S.; Fagnoni, M.; Albini, A. Photochemical Synthesis: Using Light to Build C–C Bonds under Mild Conditions. *C. R. Chim.* **2017**, *20*, 261–271.
- (206) Yuan, X.; Yang, G.; Yu, B. Photoinduced Decatungstate-Catalyzed C–H Functionalization. *Youji Huaxue* **2020**, *40*, 3620–3632.
- (207) Cowie, B. E.; Purkis, J. M.; Austin, J.; Love, J. B.; Arnold, P. L. Thermal and Photochemical Reduction and Functionalization Chemistry of the Uranyl Dication, [U<sup>VI</sup>O<sub>2</sub>]<sup>2+</sup>. *Chem. Rev.* **2019**, *119*, 10595–10637.
- (208) Li, Y.; Su, J.; Mitchell, E.; Zhang, G.; Li, J. Photocatalysis with Visible-Light-Active Uranyl Complexes. *Sci. China: Chem.* **2013**, *56*, 1671–1681.
- (209) Shiragami, T.; Matsumoto, J.; Inoue, H.; Yasuda, M. Antimony Porphyrin Complexes as Visible-Light Driven Photocatalyst. *J. Photochem. Photobiol., C* **2005**, *6*, 227–248.
- (210) Huang, H.; Strater, Z. M.; Lambert, T. H. Electrophotocatalytic C–H Functionalization of Ethers with High Regioselectivity. *J. Am. Chem. Soc.* **2020**, *142*, 1698–1703.
- (211) Adam, W.; Arnold, M. A.; Nau, W. M.; Pischel, U.; Saha-Möller, C. R. A Comparative Photomechanistic Study (Spin Trapping, EPR Spectroscopy, Transient Kinetics, Photoproducts) of Nucleoside Oxidation (DG and 8-OxodG) by Triplet-Excited Acetophenones and by the Radicals Generated from  $\alpha$ -Oxy-Substituted Derivatives through No. *J. Am. Chem. Soc.* **2002**, *124*, 3893–3904.
- (212) Ghoneim, N.; Monbelli, A.; Pilloud, D.; Suppan, P. Photochemical Reactivity of Para-Aminobenzophenone in Polar and Non-Polar Solvents. *J. Photochem. Photobiol., A* **1996**, *94*, 145–148.
- (213) Serpa, C.; Arnaut, L. G. Does Molecular Size Matter in Photoinduced Electron Transfer Reactions? *J. Phys. Chem. A* **2000**, *104*, 11075–11086.
- (214) Timpe, H.-J.; Kronfeld, K.-P. Light-Induced Polymer and Polymerization Reactions XXXIII: Direct Photoinitiation of Methyl Methacrylate Polymerization by Excited States of Ketones. *J. Photochem. Photobiol., A* **1989**, *46*, 253–267.
- (215) Favaro, G.; Romani, A. Acid–Base Properties of Disodium 3,3′-Disulfonatobenzophenone (DSB) in the Ground and Excited States. *J. Chem. Soc., Faraday Trans.* **1993**, *89*, 699–702.
- (216) Hamanoue, K.; Nakayama, T.; Kajiwar, Y.; Yamaguchi, T.; Teranishi, H. The Lowest Triplet States of Anthraquinone and Chloroanthraquinones: The 1-chloro, 2-chloro, 1,5-dichloro, and 1,8-dichloro Compounds. *J. Chem. Phys.* **1987**, *86*, 6654–6659.
- (217) Bergamini, G.; Ceroni, P.; Maestri, M.; Balzani, V.; Lee, S.-K.; Vögtle, F. Forward (Singlet–Singlet) and Backward (Triplet–Triplet) Energy Transfer in a Dendrimer with Peripheral Naphthalene Units and a Benzophenone Core. *Photochem. Photobiol. Sci.* **2004**, *3*, 898–905.
- (218) Kavlakoglu, E.; Yaman, A.; Bayrakçeken, F. Photophysical Properties of SH-Dibenzo(a,d)cyclohepten-5-One in Different Solutions. *Spectrochim. Acta, Part A* **2000**, *56*, 1117–1121.
- (219) Fukuzumi, S.; Itoh, S.; Komori, T.; Suenobu, T.; Ishida, A.; Fujitsuka, M.; Ito, O. Photochemical Reactions of Coenzyme PQQ (Pyrroloquinolinequinone) and Analogues with Benzyl Alcohol Derivatives via Photoinduced Electron Transfer. *J. Am. Chem. Soc.* **2000**, *122*, 8435–8443.
- (220) Elliott, L. D.; Kayal, S.; George, M. W.; Booker-Milburn, K. Rational Design of Triplet Sensitizers for the Transfer of Excited State Photochemistry from UV to Visible. *J. Am. Chem. Soc.* **2020**, *142*, 14947–14956.
- (221) Fan, X.; Rong, J.; Wu, H.; Zhou, Q.; Deng, H.; Tan, J. D.; Xue, C.; Wu, L.; Tao, H.; Wu, J. Eosin Y as a Direct Hydrogen-Atom Transfer Photocatalyst for the Functionalization of C–H Bonds. *Angew. Chem., Int. Ed.* **2018**, *57*, 8514–8518.
- (222) Huix-Rotlant, M.; Burghardt, I.; Ferré, N. Population of Triplet States in Acetophenone: A Quantum Dynamics Perspective. *C. R. Chim.* **2016**, *19*, 50–56.
- (223) Bohning, J. J.; Weiss, K. A Kinetic Study of the Photochemical Reaction of Phenanthrenequinone with Olefins. *J. Am. Chem. Soc.* **1966**, *88*, 2893–2898.
- (224) Shen, T.; Zhao, Z.-G.; Yu, Q.; Xu, H.-J. Photosensitized Reduction of Benzil by Heteroatom-Containing Anthracene Dyes. *J. Photochem. Photobiol., A* **1989**, *47*, 203–212.

- (225) Xia, J.-B.; Zhu, C.; Chen, C. Visible Light-Promoted Metal-Free  $\text{sp}^3\text{-C-H}$  Fluorination. *Chem. Commun.* **2014**, *50*, 11701–11704.
- (226) Hansen, A.; Renner, M.; Griesbeck, A. G.; Büsgen, T. From 3D to 4D Printing: A Reactor for Photochemical Experiments Using Hybrid Polyurethane Acrylates for Vat-Based Polymerization and Surface Functionalization. *Chem. Commun.* **2020**, *56*, 15161–15164.
- (227) Manfrotto, C.; Mella, M.; Freccero, M.; Fagnoni, M.; Albin, A. Photochemical Synthesis of 4-Oxobutanol Acetals and of 2-Hydroxycyclobutanone Ketals. *J. Org. Chem.* **1999**, *64*, 5024–5028.
- (228) Wang, L.; Wang, T.; Cheng, G.-J.; Li, X.; Wei, J.-J.; Guo, B.; Zheng, C.; Chen, G.; Ran, C.; Zheng, C. Direct C–H Arylation of Aldehydes by Merging Photocatalyzed Hydrogen Atom Transfer with Palladium Catalysis. *ACS Catal.* **2020**, *10*, 7543–7551.
- (229) Doohan, R. A.; Hannan, J. J.; Geraghty, N. W. A. The Photomediated Reaction of Alkynes with Cycloalkanes. *Org. Biomol. Chem.* **2006**, *4*, 942–952.
- (230) Hoshikawa, T.; Inoue, M. Photoinduced Direct 4-Pyridination of C( $\text{sp}^3$ )–H Bonds. *Chem. Sci.* **2013**, *4*, 3118–3123.
- (231) Hoshikawa, T.; Yoshioka, S.; Kamijo, S.; Inoue, M. Photoinduced Direct Cyanation of C( $\text{sp}^3$ )–H Bonds. *Synthesis* **2013**, *45*, 874–887.
- (232) Kraus, G. A.; Liu, P. Benzophenone-Mediated Conjugate Additions of Aromatic Aldehydes to Quinones. *Tetrahedron Lett.* **1994**, *35*, 7723–7726.
- (233) Oelgemöller, M.; Schiel, C.; Fröhlich, R.; Mattay, J. The “Photo-Friedel–Crafts Acylation” of 1,4-Naphthoquinones. *Eur. J. Org. Chem.* **2002**, *2002*, 2465–2474.
- (234) Fraser-Reid, B.; Anderson, R. C.; Hicks, D. R.; Walker, D. L. Synthetic Applications of the Photochemically Induced Addition of Oxycarbonyl Species to  $\alpha$ -Enones. Part II. The Addition of Ketals, Aldehydes, and Polyfunctional Species. *Can. J. Chem.* **1977**, *55*, 3986–3995.
- (235) Cerfontain, H.; Van Noort, P. C. M. Facile Photochemical Synthesis of Some 4-Oxoalkanoic Acids and Esters. *Synthesis* **1980**, *1980*, 490–492.
- (236) Masuda, Y.; Tsuda, H.; Murakami, M.  $\text{C}_1$  Oxidation/ $\text{C}_2$  Reduction Isomerization of Unprotected Aldoses Induced by Light/Ketone. *Angew. Chem., Int. Ed.* **2020**, *59*, 2755–2759.
- (237) Singh, M.; Yadav, A. K.; Yadav, L. D. S.; Singh, R. K. P. Visible Light Photocatalysis with Benzophenone for Radical Thiol-Ene Reactions. *Tetrahedron Lett.* **2017**, *58*, 2206–2208.
- (238) Matsumoto, K.; Nakajima, M.; Nemoto, T. Visible Light-Induced Direct  $\text{S}_0 \rightarrow \text{T}_n$  Transition of Benzophenone Promotes C( $\text{sp}^3$ )–H Alkynylation of Ethers and Amides. *J. Org. Chem.* **2020**, *85*, 11802–11811.
- (239) Dondi, D.; Caprioli, I.; Fagnoni, M.; Mella, M.; Albin, A. A Convenient Route to 1,4-Monoprotected Dialdehydes, 1,4-Ketoaldehydes,  $\gamma$ -Lactols and  $\gamma$ -Lactones through Radical Alkylation of  $\alpha,\beta$ -Unsaturated Aldehydes in Organic and Organic-Aqueous Media. *Tetrahedron* **2003**, *59*, 947–957.
- (240) Dondi, D.; Protti, S.; Albin, A.; Carpio, S. M.; Fagnoni, M. Synthesis of  $\gamma$ -Lactols,  $\gamma$ -Lactones and 1,4-Monoprotected Succinaldehydes under Moderately Concentrated Sunlight. *Green Chem.* **2009**, *11*, 1653–1659.
- (241) Kamijo, S.; Takao, G.; Kamijo, K.; Tsuno, T.; Ishiguro, K.; Murafuji, T. Alkylation of Nonacidic C( $\text{sp}^3$ )–H Bonds by Photoinduced Catalytic Michael-Type Radical Addition. *Org. Lett.* **2016**, *18*, 4912–4915.
- (242) Tada, N.; Ikebata, Y.; Nobuta, T.; Hirashima, S.; Miura, T.; Itoh, A. Direct Aerobic Photo-Oxidative Syntheses of Aromatic Methyl Esters from Methyl Aromatics Using Anthraquinone-2,3-Dicarboxylic Acid as Organophotocatalyst. *Photochem. Photobiol. Sci.* **2012**, *11*, 616–619.
- (243) Yavorsky, A.; Shvydkiv, O.; Hoffmann, N.; Nolan, K.; Oelgemöller, M. Parallel Microflow Photochemistry: Process Optimization, Scale-up, and Library Synthesis. *Org. Lett.* **2012**, *14*, 4342–4345.
- (244) Bume, D. D.; Pitts, C. R.; Jokhai, R. T.; Lectka, T. Direct, Visible Light-Sensitized Benzylic C–H Fluorination of Peptides Using Dibenzosuberone: Selectivity for Phenylalanine-like Residues. *Tetrahedron* **2016**, *72*, 6031–6036.
- (245) Zhang, Y.; Ji, P.; Hu, W.; Wei, Y.; Huang, H.; Wang, W. Organocatalytic Transformation of Aldehydes to Thioesters with Visible Light. *Chem. - Eur. J.* **2019**, *25*, 8225–8228.
- (246) Papadopoulos, G. N.; Kokotos, C. G. Photoorganocatalytic One-Pot Synthesis of Hydroxamic Acids from Aldehydes. *Chem. - Eur. J.* **2016**, *22*, 6964–6967.
- (247) Tripathi, C. B.; Ohtani, T.; Corbett, M. T.; Ooi, T. Photoredox Ketone Catalysis for the Direct C–H Imidation and Acyloxylation of Arenes. *Chem. Sci.* **2017**, *8*, 5622–5627.
- (248) Zhu, K.; Ohtani, T.; Tripathi, C. B.; Uraguchi, D.; Ooi, T. Formal Hydroformylation of  $\alpha,\beta$ -Unsaturated Carboxylic Acids under Photoexcited Ketone Catalysis. *Chem. Lett.* **2019**, *48*, 715–717.
- (249) Fan, X.; Xiao, P.; Jiao, Z.; Yang, T.; Dai, X.; Xu, W.; Tan, J. Da; Cui, G.; Su, H.; Fang, W.; Wu, J. Neutral-Eosin-Y-Photocatalyzed Silane Chlorination Using Dichloromethane. *Angew. Chem., Int. Ed.* **2019**, *58*, 12580–12584.
- (250) Wang, H.; Li, W.; Zeng, K.; Wu, Y.; Zhang, Y.; Xu, T.; Chen, Y. Photocatalysis Enables Visible-Light Uncaging of Bioactive Molecules in Live Cells. *Angew. Chem., Int. Ed.* **2019**, *58*, 561–565.
- (251) Inoa, J.; Patel, M.; Dominici, G.; Eldabagh, R.; Patel, A.; Lee, J.; Xing, Y. Benzylic Hydroperoxidation via Visible-Light-Induced C( $\text{sp}^3$ )–H Activation. *J. Org. Chem.* **2020**, *85*, 6181–6187.
- (252) Soni, S.; Pali, P.; Ansari, M. A.; Singh, M. S. Visible-Light Photocatalysis of Eosin Y: HAT and Complementing MS-CPET Strategy to Trifluoromethylation of  $\beta$ -Ketodithioesters with Langlois’ Reagent. *J. Org. Chem.* **2020**, *85*, 10098–10109.
- (253) Xia, Q.; Shi, Z.; Yuan, J.; Bian, Q.; Xu, Y.; Liu, B.; Huang, Y.; Yang, X.; Xu, H. Visible-Light-Enabled Selective Oxidation of Primary Alcohols through Hydrogen-Atom Transfer and Its Application in the Synthesis of Quinazolinones. *Asian J. Org. Chem.* **2019**, *8*, 1933–1941.
- (254) König, B.; Ramm, S.; Bubenitschek, P.; Jones, P. G.; Hopf, H.; Knieriem, B.; Meijere, A. De. [2.2](4,7)Isobenzofuranophanes - Synthesis, Characterisation, and Reactivity. *Chem. Ber.* **1994**, *127*, 2263–2266.
- (255) Zhang, X.; Cunningham, M. M.; Walker, R. A. Solvent Polarity at Polar Solid Surfaces: The Role of Solvent Structure. *J. Phys. Chem. B* **2003**, *107*, 3183–3195.
- (256) Ohta, A.; Hattori, K.; Kusumoto, Y.; Kawase, T.; Kobayashi, T.; Naito, H.; Kitamura, C. Effects of Alkoxy Substitution on the Optical Properties of 9,10-Anthraquinone and Anthracene: 2,3,6,7-Tetrapropoxy-Substituted vs. 2,6-Dipropoxy-Substituted Derivatives. *Chem. Lett.* **2012**, *41*, 674–676.
- (257) Chen, S.-C.; Fang, T.-S. Observation of a Novel Emission from an Exciplex of Triplet 4-Phenylbenzophenone with Triethylamine. *Chem. Phys. Lett.* **2007**, *450*, 65–70.
- (258) Peters, R. H.; Sumner, H. H. 430. Spectra of Anthraquinone Derivatives. *J. Chem. Soc.* **1953**, 2101–2110.
- (259) Kuzmanich, G.; Simoncelli, S.; Gard, M. N.; Spänig, F.; Henderson, B. L.; Guldi, D. M.; Garcia-Garibay, M. A. Excited State Kinetics in Crystalline Solids: Self-Quenching in Nanocrystals of 4,4'-Disubstituted Benzophenone Triplets Occurs by a Reductive Quenching Mechanism. *J. Am. Chem. Soc.* **2011**, *133*, 17296–17306.
- (260) Rajendiran, N.; Sankaranarayanan, R. K.; Venkatesh, G. Excimer Emission in Inclusion Complexes of Dibenzofuran and 5-Dibenzosuberone with  $\alpha$ - and  $\beta$ -Cyclodextrins. *Bull. Chem. Soc. Jpn.* **2014**, *87*, 797–808.
- (261) Gautrot, J. E.; Hodge, P.; Helliwell, M.; Raftery, J.; Cupertino, D. Synthesis of Electron-Accepting Polymers Containing Phenanthra-9,10-Quinone Units. *J. Mater. Chem.* **2009**, *19*, 4148–4156.
- (262) Togashi, D. M.; Nicodem, D. E. Photophysical Studies of 9,10-Phenanthrenequinones. *Spectrochim. Acta, Part A* **2004**, *60*, 3205–3212.
- (263) Fischer, G.; Oehme, G.; Schellenberger, A. Zur Theorie Der  $\alpha$ -Ketosäuren. *Tetrahedron* **1971**, *27*, 5683–5696.



- (264) Pirenne, V.; Kurtay, G.; Voci, S.; Bouffier, L.; Sojic, N.; Robert, F.; Bassani, D. M.; Landais, Y. Eosin-Mediated Alkylsulfonyl Cyanation of Olefins. *Org. Lett.* **2018**, *20*, 4521–4525.
- (265) Waele, V. De; Poizat, O.; Fagnoni, M.; Bagno, A.; Ravelli, D. Unraveling the Key Features of the Reactive State of Decatungstate Anion in Hydrogen Atom Transfer (HAT) Photocatalysis. *ACS Catal.* **2016**, *6*, 7174–7182.
- (266) Capaldo, L.; Ertl, M.; Fagnoni, M.; Knör, G.; Ravelli, D. Antimony–Oxo Porphyrins as Photocatalysts for Redox-Neutral C–H to C–C Bond Conversion. *ACS Catal.* **2020**, *10*, 9057–9064.
- (267) Ferreira, G. C.; Schmitt, C. C.; Neumann, M. G. Dependence of the Thioxanthone Triplet-Triplet Absorption Spectrum with Solvent Polarity and Aromatic Ring Substitution. *J. Braz. Chem. Soc.* **2006**, *17*, 905–909.
- (268) Iyer, A.; Clay, A.; Jockusch, S.; Sivaguru, J. Evaluating Brominated Thioxanthenes as Organo-Photocatalysts. *J. Phys. Org. Chem.* **2017**, *30*, No. e3738.
- (269) Pant, D. D.; Tripathi, H. B. Spectra, Lifetimes and Energy Transfer for Uranyl Species in Solution. *J. Lumin.* **1974**, *8*, 492–501.
- (270) Scaiano, J. C. Solvent Effects in the Photochemistry of Xanthone. *J. Am. Chem. Soc.* **1980**, *102*, 7747–7753.
- (271) Ghosh, R.; Mondal, J. A.; Ghosh, H. N.; Palit, D. K. Ultrafast Dynamics of the Excited States of the Uranyl Ion in Solutions. *J. Phys. Chem. A* **2010**, *114*, 5263–5270.
- (272) Raviola, C.; Ravelli, D. Efficiency and Selectivity Aspects in the C–H Functionalization of Aliphatic Oxygen Heterocycles by Photocatalytic Hydrogen Atom Transfer. *Synlett* **2019**, *30*, 803–808.
- (273) Xia, J.-B.; Zhu, C.; Chen, C. Visible Light-Promoted Metal-Free C–H Activation: Diarylketone-Catalyzed Selective Benzylic Mono- and Difluorination. *J. Am. Chem. Soc.* **2013**, *135*, 17494–17500.
- (274) Nodwell, M. B.; Yang, H.; Čolović, M.; Yuan, Z.; Merckens, H.; Martin, R. E.; Bénard, F.; Schaffer, P.; Britton, R. 18 F-Fluorination of Unactivated C–H Bonds in Branched Aliphatic Amino Acids: Direct Synthesis of Oncological Positron Emission Tomography Imaging Agents. *J. Am. Chem. Soc.* **2017**, *139*, 3595–3598.
- (275) Dondi, D.; Fagnoni, M.; Molinari, A.; Maldotti, A.; Albini, A. Polyoxotungstate Photoinduced Alkylation of Electrophilic Alkenes by Cycloalkanes. *Chem. - Eur. J.* **2004**, *10*, 142–148.
- (276) Laudadio, G.; Govaerts, S.; Wang, Y.; Ravelli, D.; Koolman, H. F.; Fagnoni, M.; Djuric, S. W.; Noël, T. Selective C(sp<sup>3</sup>)–H Aerobic Oxidation Enabled by Decatungstate Photocatalysis in Flow. *Angew. Chem., Int. Ed.* **2018**, *57*, 4078–4082.
- (277) Protti, S.; Ravelli, D.; Fagnoni, M.; Albini, A. Solar Light-Driven Photocatalyzed Alkylations. Chemistry on the Window Ledge. *Chem. Commun.* **2009**, *47*, 7351–7353.
- (278) Capaldo, L.; Ravelli, D. Decatungstate as Direct Hydrogen Atom Transfer Photocatalyst for SOMophilic Alkynylation. *Org. Lett.* **2021**, *23*, 2243–2247.
- (279) Baxter, I.; Cameron, D. W.; Titman, R. B. Formation of Some Linear Polycyclic Diquinones via Novel Dimerization. *J. Chem. Soc. D* **1969**, 563–564.
- (280) Weigel, W. K.; Dang, H. T.; Yang, H.-B.; Martin, D. B. C. Synthesis of Amino-Diamondoid Pharmacophores via Photocatalytic C–H Aminoalkylation. *Chem. Commun.* **2020**, *56*, 9699–9702.
- (281) Yang, H.-B.; Feceu, A.; Martin, D. B. C. Catalyst-Controlled C–H Functionalization of Adamantanes Using Selective H-Atom Transfer. *ACS Catal.* **2019**, *9*, 5708–5715.
- (282) Li, Y.; Lei, M.; Gong, L. Photocatalytic Regio- and Stereoselective C(sp<sup>3</sup>)–H Functionalization of Benzylic and Allylic Hydrocarbons as Well as Unactivated Alkanes. *Nat. Catal.* **2019**, *2*, 1016–1026.
- (283) Zhang, Y.; Yang, X.; Wu, J.; Huang, D. Aerobic C–H Functionalization Using Pyrenedione as the Photocatalyst. *Synthesis* **2020**, *52*, 2512–2520.
- (284) Capaldo, L.; Merli, D.; Fagnoni, M.; Ravelli, D. Visible Light Uranyl Photocatalysis: Direct C–H to C–C Bond Conversion. *ACS Catal.* **2019**, *9*, 3054–3058.
- (285) West, J. G.; Bedell, T. A.; Sorensen, E. J. The Uranyl Cation as a Visible-Light Photocatalyst for C(sp<sup>3</sup>)–H Fluorination. *Angew. Chem., Int. Ed.* **2016**, *55*, 8923–8927.
- (286) Ishida, N.; Masuda, Y.; Uemoto, S.; Murakami, M. A Light/Ketone/Copper System for Carboxylation of Allylic C–H Bonds of Alkenes with CO<sub>2</sub>. *Chem. - Eur. J.* **2016**, *22*, 6524–6527.
- (287) Cantillo, D.; de Frutos, O.; Rincón, J. A.; Mateos, C.; Kappe, C. O. A Continuous-Flow Protocol for Light-Induced Benzylic Fluorinations. *J. Org. Chem.* **2014**, *79*, 8486–8490.
- (288) Yatsushashi, T.; Nakajima, Y.; Shimada, T.; Inoue, H. Photophysical Properties of Intramolecular Charge-Transfer Excited Singlet State of Aminofluorenone Derivatives. *J. Phys. Chem. A* **1998**, *102*, 3018–3024.
- (289) Tzirakis, M.; Lykakis, I.; Panagiotou, G.; Bourikas, K.; Lycourghiotis, A.; Kordulis, C.; Orfanopoulos, M. Decatungstate Catalyst Supported on Silica and  $\gamma$ -Alumina: Efficient Photocatalytic Oxidation of Benzyl Alcohols. *J. Catal.* **2007**, *252*, 178–189.
- (290) Kamijo, S.; Kamijo, K.; Maruoka, K.; Murafuji, T. Aryl Ketone Catalyzed Radical Allylation of C(sp<sup>3</sup>)–H Bonds under Photoirradiation. *Org. Lett.* **2016**, *18*, 6516–6519.
- (291) Zhang, Y.; Yang, X.; Tang, H.; Liang, D.; Wu, J.; Huang, D. Pyrenediones as Versatile Photocatalysts for Oxygenation Reactions with in Situ Generation of Hydrogen Peroxide under Visible Light. *Green Chem.* **2020**, *22*, 22–27.
- (292) Moriconi, E. J.; Rakoczy, B.; O'Connor, W. F. Oxidation-Reduction Potentials and Absorption Spectra of Polycyclic Aromatic Quinones 1. *J. Org. Chem.* **1962**, *27*, 2772–2776.
- (293) Chaumont, A.; Klimchuk, O.; Gaillard, C.; Billard, I.; Ouadi, A.; Hennig, C.; Wipff, G. Perrhenate Complexation by Uranyl in Traditional Solvents and in Ionic Liquids: A Joint Molecular Dynamics/Spectroscopic Study. *J. Phys. Chem. B* **2012**, *116*, 3205–3219.
- (294) Meinrath, G.; Kato, Y.; Kimura, T.; Yoshida, Z. Stokes Relationship in Absorption and Fluorescence Spectra of U(VI) Species. *Radiochim. Acta* **1998**, *82*, 115–120.
- (295) Karbowiak, M.; Fourest, B.; Hubert, S.; Moulin, C. Complexation of U(VI) with Iodate Ions: Determination of Stability Constants by Using Spectroscopic Methods and Capillary Zone Electrophoresis. *Radiochim. Acta* **2003**, *91*, 505–512.
- (296) Williams, J. D.; Nakano, M.; Gérardy, R.; Rincón, J. A.; de Frutos, O.; Mateos, C.; Monbaliu, J.-C. M.; Kappe, C. O. Finding the Perfect Match: A Combined Computational and Experimental Study toward Efficient and Scalable Photosensitized [2 + 2] Cycloadditions in Flow. *Org. Process Res. Dev.* **2019**, *23*, 78–87.
- (297) Wagner, P. J.; Hammond, G. S. Properties and Reactions of Organic Molecules in Their Triplet States. In *Adv. Photochem.*; 1968; pp 21–156.
- (298) Formosinho, S. J. Photochemical Hydrogen Abstractions as Radiationless Transitions. Part 2. - Thioketones, Quinones, Aza-Aromatics, Olefins and Azobenzenes. *J. Chem. Soc., Faraday Trans. 2* **1976**, *72*, 1332–1339.
- (299) Formosinho, S. J. Photochemical Hydrogen Abstractions as Radiationless Transitions. Part 1. - Ketones, Aldehydes and Acids. *J. Chem. Soc., Faraday Trans. 2* **1976**, *72*, 1313–1331.
- (300) Anpo, M.; Kubokawa, Y. Reactivity of Excited Triplet Alkyl Ketones in Solution. I. Quenching and Hydrogen Abstraction of Triplet Acetone. *Bull. Chem. Soc. Jpn.* **1977**, *50*, 1913–1916.
- (301) Walling, C.; Gibian, M. J. Hydrogen Abstraction Reactions by the Triplet States of Ketones 1. *J. Am. Chem. Soc.* **1965**, *87*, 3361–3364.
- (302) Wagner, P.; Park, B.-S. *Organic Photochemistry*; Padwa, A., Ed.; CRC Press: Routledge, 2017.
- (303) Wu, Y.; Kim, D.; Teets, T. S. Photophysical Properties and Redox Potentials of Photosensitizers for Organic Photoredox Transformations. *Synlett* **2021**, DOI: 10.1055/a-1390-9065.
- (304) Wagner, P. J.; Kemppainen, A. E.; Schott, H. N. Photo-reactivity of p-Methoxyphenyl Ketones. Evidence for Hydrogen Abstraction from Equilibrium Concentrations of Upper n, $\pi^*$  Triplets. *J. Am. Chem. Soc.* **1970**, *92*, 5280–5281.

- (305) Wagner, P. J.; Truman, R. J.; Scaiano, J. C. Substituent Effects on Hydrogen Abstraction by Phenyl Ketone Triplets. *J. Am. Chem. Soc.* **1985**, *107*, 7093–7097.
- (306) Wagner, P. J.; Truman, R. J.; Puchalski, A. E.; Wake, R. Extent of Charge Transfer in the Photoreduction of Phenyl Ketones by Alkylbenzenes. *J. Am. Chem. Soc.* **1986**, *108*, 7727–7738.
- (307) Wagner, P. J.; Thomas, M. J.; Puchalski, A. E. Photoreactivity of  $\alpha$ -Fluorinated Phenyl Alkyl Ketones. *J. Am. Chem. Soc.* **1986**, *108*, 7739–7744.
- (308) Ortica, F.; Romani, A.; Favaro, G. Light-Induced Hydrogen Abstraction from Isobutanol by Thienyl Phenyl, Dithienyl, and Thienyl Pyridyl Ketones. *J. Phys. Chem. A* **1999**, *103*, 1335–1341.
- (309) Chandra, A. K. The Perturbational Treatment of the Process of Hydrogen Abstraction by Ketones. *J. Photochem.* **1979**, *11*, 347–360.
- (310) Fréneau, M.; Lefebvre, C.; Gómez Fernández, M. A.; Richard, C.; Hoffmann, N. Photochemical Reactivity of Phenyl (Methyl-Tetrazolyl) Ketone - Hydrogen Atom Transfer vs. Electron Transfer. *New J. Chem.* **2019**, *43*, 17151–17158.
- (311) Péter, Á.; Agasti, S.; Knowles, O.; Pye, E.; Procter, D. J. Recent Advances in the Chemistry of Ketyl Radicals. *Chem. Soc. Rev.* **2021**, *50*, 5349–5365.
- (312) Yayla, H.; Knowles, R. Proton-Coupled Electron Transfer in Organic Synthesis: Novel Homolytic Bond Activations and Catalytic Asymmetric Reactions with Free Radicals. *Synlett* **2014**, *25*, 2819–2826.
- (313) Cardarelli, A. M.; Fagnoni, M.; Mella, M.; Albini, A. Hydrocarbon Activation. Synthesis of  $\beta$ -Cycloalkyl (Di)Nitriles through Photosensitized Conjugate Radical Addition. *J. Org. Chem.* **2001**, *66*, 7320–7327.
- (314) Papaconstantinou, E.; Dimotikali, D.; Politou, A. Photochemistry of Heteropoly Electrolytes. The 18-Molybdodiphosphate. *Inorg. Chim. Acta* **1980**, *43*, 155–158.
- (315) Papaconstantinou, E. Photocatalytic Oxidation of Organic Compounds Using Heteropoly Electrolytes of Molybdenum and Tungsten. *J. Chem. Soc., Chem. Commun.* **1982**, 12–13.
- (316) Darwent, J. R. Photocatalytic Hydrogen Evolution from Alcohols Using Dodecawolfram-silicic Acid Colloidal Platinum. *J. Chem. Soc., Chem. Commun.* **1982**, 798–799.
- (317) Dimotikali, D.; Papaconstantinou, E. Photochemistry of Heteropoly Electrolytes: The 1:12 Tungstates. *Inorg. Chim. Acta* **1984**, *87*, 177–180.
- (318) Ward, M. D.; Brazdil, J. F.; Grasselli, R. K. Photocatalytic Alcohol Dehydrogenation Using Ammonium Heptamolybdate. *J. Phys. Chem.* **1984**, *88*, 4210–4213.
- (319) Hill, C. L.; Kozik, M.; Winkler, J.; Hou, Y.; Prosser-McCartha, C. M. Polyoxometalates in Catalytic Photochemical Hydrocarbon Functionalization and Photomicroolithography. In *Photosensitive Metal—Organic Systems*; Kutal, C., Serpone, N., Eds.; American Chemical Society, 1993; pp 243–259.
- (320) Yamase, T.; Takabayashi, N.; Kaji, M. Solution Photochemistry of Tetrakis(Tetrabutylammonium) Decatungstate(VI) and Catalytic Hydrogen Evolution from Alcohols. *J. Chem. Soc., Dalton Trans.* **1984**, *5*, 793–799.
- (321) Yamase, T.; Usami, T. Photocatalytic Dimerization of Olefins by Decatungstate(VI),  $[W_{10}O_{32}]^{4-}$ , in Acetonitrile and Magnetic Resonance Studies of Photoreduced Species. *J. Chem. Soc., Dalton Trans.* **1988**, 183–190.
- (322) Hill, C. L.; Bouchard, D. A. Catalytic Photochemical Dehydrogenation of Organic Substrates by Polyoxometalates. *J. Am. Chem. Soc.* **1985**, *107*, 5148–5157.
- (323) Hill, C. L.; Bouchard, D. A.; Kadkhodayan, M.; Williamson, M. M.; Schmidt, J. A.; Hilsinki, E. F. Catalytic Photochemical Oxidation of Organic Substrates by Polyoxometalates. Picosecond Spectroscopy, Photochemistry, and Structural Properties of Charge-Transfer Complexes between Heteropolytungstic Acids and Dipolar Organic Compounds. *J. Am. Chem. Soc.* **1988**, *110*, 5471–5479.
- (324) Chambers, R. C.; Hill, C. L. Excited States of Polyoxometalates as Oxidatively Resistant Initiators of Hydrocarbon Autoxidation. Selective Production of Hydroperoxides. *Inorg. Chem.* **1989**, *28*, 2509–2511.
- (325) Combs-Walker, L. A.; Hill, C. L. Use of Excited-State and Ground-State Redox Properties of Polyoxometalates for Selective Transformation of Unactivated Carbon-Hydrogen Centers Remote from the Functional Group in Ketones. *J. Am. Chem. Soc.* **1992**, *114*, 938–946.
- (326) Jaynes, B. S.; Hill, C. L. Selective Ethylation and Vinylation of Alkanes via Polyoxotungstate Photocatalyzed Radical Addition Reactions. *J. Am. Chem. Soc.* **1993**, *115*, 12212–12213.
- (327) Jaynes, B. S.; Hill, C. L. Radical Carbonylation of Alkanes via Polyoxotungstate Photocatalysis. *J. Am. Chem. Soc.* **1995**, *117*, 4704–4705.
- (328) Zheng, Z.; Hill, C. L. Alkanes to Nitriles and  $\alpha$ -Iminoesters. Polyoxotungstate Photocatalytic Radical Chain Initiation. *Chem. Commun.* **1998**, 2467–2468.
- (329) Prosser-McCartha, C. M.; Hill, C. L. Direct Selective Acylation of an Unactivated Carbon-Hydrogen Bond in a Caged Hydrocarbon. Approach to Systems for Carbon-Hydrogen Bond Functionalization That Proceed Catalytically and Selectively at High Substrate Conversion. *J. Am. Chem. Soc.* **1990**, *112*, 3671–3673.
- (330) Renneke, R. F.; Pasquali, M.; Hill, C. L. Polyoxometalate Systems for the Catalytic Selective Production of Nonthermodynamic Alkenes from Alkanes. Nature of Excited-State Deactivation Processes and Control of Subsequent Thermal Processes in Polyoxometalate Photoredox Chemistry. *J. Am. Chem. Soc.* **1990**, *112*, 6585–6594.
- (331) Chambers, R. C.; Hill, C. L. Redox Catalysis Involving Substrate Photooxidation with Catalyst Regeneration by Substrate Reduction. Simultaneous Oxidative C–H Bond Cleavage and Reductive C–S Bond Cleavage in Thioethers Catalyzed by Polyoxotungstate,  $W_{10}O_{32}^{4-}$ . *J. Am. Chem. Soc.* **1990**, *112*, 8427–8433.
- (332) Chemseddine, A.; Sanchez, C.; Livage, J.; Launay, J. P.; Fournier, M. Electrochemical and Photochemical Reduction of Decatungstate: A Reinvestigation. *Inorg. Chem.* **1984**, *23*, 2609–2613.
- (333) Kothe, T.; Martschke, R.; Fischer, H. Photoreactions of the Decatungstate Anion  $W_{10}O_{32}^{4-}$  with Organic Substrates in Solution Studied by EPR and Kinetic Absorption Spectroscopy: An Example for the Persistent Radical Effect. *J. Chem. Soc., Perkin Trans. 2* **1998**, 503–508.
- (334) Tanielian, C. Decatungstate Photocatalysis. *Coord. Chem. Rev.* **1998**, *178–180*, 1165–1181.
- (335) Ermolenko, L. P.; Delaire, J. A.; Giannotti, C. Laser Flash Photolysis Study of the Mechanism of Photooxidation of Alkanes Catalysed by Decatungstate Anion. *J. Chem. Soc., Perkin Trans. 2* **1997**, 25–30.
- (336) Tanielian, C.; Duffy, K.; Jones, A. Kinetic and Mechanistic Aspects of Photocatalysis by Polyoxotungstates: A Laser Flash Photolysis, Pulse Radiolysis, and Continuous Photolysis Study. *J. Phys. Chem. B* **1997**, *101*, 4276–4282.
- (337) Duncan, D. C.; Fox, M. A. Early Events in Decatungstate Photocatalyzed Oxidations: A Nanosecond Laser Transient Absorbance Reinvestigation. *J. Phys. Chem. A* **1998**, *102*, 4559–4567.
- (338) Tanielian, C.; Schweitzer, C.; Seghrouchni, R.; Esch, M.; Mechin, R. Polyoxometalate Sensitization in Mechanistic Studies of Photochemical Reactions: The Decatungstate Anion as a Reference Sensitizer for Photoinduced Free Radical Oxygenations of Organic Compounds. *Photochem. Photobiol. Sci.* **2003**, *2*, 297–305.
- (339) Tanielian, C.; Lykakis, I. N.; Seghrouchni, R.; Cougnon, F.; Orfanopoulos, M. Mechanism of Decatungstate Photocatalyzed Oxygenation of Aromatic Alcohols Part I. Continuous Photolysis and Laser Flash Photolysis Studies. *J. Mol. Catal. A: Chem.* **2007**, *262*, 170–175.
- (340) Texier, I.; Delaire, J. A.; Giannotti, C. Reactivity of the Charge Transfer Excited State of Sodium Decatungstate at the Nanosecond Time Scale. *Phys. Chem. Chem. Phys.* **2000**, *2*, 1205–1212.
- (341) Yamase, T.; Ohtaka, K. Photochemistry of Polyoxovanadates. Part 1. Formation of the Anion-Encapsulated Polyoxovanadate  $[V_{15}O_{36}(CO_3)]^{7-}$  and Electron-Spin Polarization of  $\alpha$ -Hydroxyalkyl



- Radicals in the Presence of Alcohols. *J. Chem. Soc., Dalton Trans.* **1994**, 53, 2599–2608.
- (342) Duncan, D. C.; Netz, T. L.; Hill, C. L. Early-Time Dynamics and Reactivity of Polyoxometalate Excited States. Identification of a Short-Lived LMCT Excited State and a Reactive Long-Lived Charge-Transfer Intermediate Following Picosecond Flash Excitation of  $[W_{10}O_{32}]^{4-}$  in Acetonitrile. *Inorg. Chem.* **1995**, 34, 4640–4646.
- (343) Texier, I.; Delouis, J. F.; Delaire, J. A.; Giannotti, C.; Plaza, P.; Martin, M. M. Dynamics of the First Excited State of the Decatungstate Anion Studied by Subpicosecond Laser Spectroscopy. *Chem. Phys. Lett.* **1999**, 311, 139–145.
- (344) Jen, S. F.; Anderson, A. B.; Hill, C. L. Alkane Reactions with Photoactivated Decatungstate in Neutral and Acid Solution: Molecular Orbital Theory. *J. Phys. Chem.* **1992**, 96, 5658–5662.
- (345) Ravelli, D.; Dondi, D.; Fagnoni, M.; Albin, A.; Bagno, A. Electronic and EPR Spectra of the Species Involved in  $[W_{10}O_{32}]^{4-}$  Photocatalysis. A Relativistic DFT Investigation. *Phys. Chem. Chem. Phys.* **2013**, 15, 2890–2896.
- (346) McGlynn, S. P.; Smith, J. K. The Electronic Structure, Spectra, and Magnetic Properties of Actinyl Ions. *J. Mol. Spectrosc.* **1961**, 6, 164–187.
- (347) Rabinowitch, E.; Belford, R. L. *Spectroscopy and Photochemistry of Uranyl Compounds*; Pergamon, The MacMillan Company: New York, 1964.
- (348) Burrows, H. D.; Kemp, T. J. The Photochemistry of the Uranyl Ion. *Chem. Soc. Rev.* **1974**, 3, 139–165.
- (349) Hill, R. J.; Kemp, T. J.; Allen, D. M.; Cox, A. Absorption Spectrum, Lifetime and Photoreactivity towards Alcohols of the Excited State of the Aqueous Uranyl Ion ( $UO_2^{+2}$ ). *J. Chem. Soc., Faraday Trans. 1* **1974**, 70, 847–857.
- (350) Jørgensen, C. K.; Reisfeld, R. Uranyl Photophysics. In *Topics in Inorganic and Physical Chemistry. Structure and Bonding*; 1982; Vol. 50, pp 121–171.
- (351) Knör, G. Bionic Catalyst Design: A Photochemical Approach to Artificial Enzyme Function. *ChemBioChem* **2001**, 2, 593–596.
- (352) Hoffmann, N. Photochemical Electron and Hydrogen Transfer in Organic Synthesis: The Control of Selectivity. *Synthesis* **2016**, 48, 1782–1802.
- (353) Morton, C. M.; Zhu, Q.; Ripberger, H.; Troian-Gautier, L.; Toa, Z. S. D.; Knowles, R. R.; Alexanian, E. J. C–H Alkylation via Multisite-Proton-Coupled Electron Transfer of an Aliphatic C–H Bond. *J. Am. Chem. Soc.* **2019**, 141, 13253–13260.
- (354) Cui, L.; Tada, N.; Okubo, H.; Miura, T.; Itoh, A. Efficient Synthesis of Gem-Dihydroperoxides with Molecular Oxygen and Anthraquinone under Visible Light Irradiation with Fluorescent Lamp. *Green Chem.* **2011**, 13, 2347–2350.
- (355) Tada, N.; Cui, L.; Okubo, H.; Miura, T.; Itoh, A. An Efficient Synthesis of Gem-Dihydroperoxides with Molecular Oxygen and Anthracene under Light Irradiation. *Adv. Synth. Catal.* **2010**, 352, 2383–2386.
- (356) Cui, L.; Furuhashi, S.; Tachikawa, Y.; Tada, N.; Miura, T.; Itoh, A. Efficient Generation of Hydrogen Peroxide by Aerobic Photooxidation of 2-Propanol Using Anthraquinone-2-Carboxylic Acid and One-Pot Epoxidation of  $\alpha,\beta$ -Unsaturated Ketones. *Tetrahedron Lett.* **2013**, 54, 162–165.
- (357) Voutyritsa, E.; Garreau, M.; Kokotou, M. G.; Triandafillidi, I.; Waser, J.; Kokotos, C. G. Photochemical Functionalization of Heterocycles with EBX Reagents: C–H Alkynylation versus Deconstructive Ring Cleavage. *Chem. - Eur. J.* **2020**, 26, 14453–14460.
- (358) Papadopoulos, G. N.; Kokotou, M. G.; Spiliopoulou, N.; Nikitas, N. F.; Voutyritsa, E.; Tzaras, D. I.; Kaplaneris, N.; Kokotos, C. G. Phenylglyoxylic Acid: An Efficient Initiator for the Photochemical Hydrogen Atom Transfer C–H Functionalization of Heterocycles. *ChemSusChem* **2020**, 13, 5934–5944.
- (359) Voutyritsa, E.; Kokotos, C. G. Green Metal-Free Photochemical Hydroacylation of Unactivated Olefins. *Angew. Chem., Int. Ed.* **2020**, 59, 1735–1741.
- (360) Yagci, Y.; Jockusch, S.; Turro, N. J. Photoinitiated Polymerization: Advances, Challenges, and Opportunities. *Macromolecules* **2010**, 43, 6245–6260.
- (361) Christmann, J.; Allonas, X.; Ley, C.; Croutxé-Barghorn, C. The Role of Ketyl Radicals in Free Radical Photopolymerization: New Experimental and Theoretical Insights. *Polym. Chem.* **2019**, 10, 1099–1109.
- (362) Sanai, Y.; Kagami, S.; Kubota, K. Cross-Linking Photopolymerization of Monoacrylate Initiated by Benzophenone. *J. Polym. Sci., Part A: Polym. Chem.* **2018**, 56, 1545–1553.
- (363) Tretinnikov, O. N.; Pilipenko, V. V.; Prikhodchenko, L. K. Benzophenone-Initiated Grafting Photopolymerization of Acrylic Acid on the Surface of Polyethylene from the Monomer Aqueous Solution without Its Deaeration. *Polym. Sci., Ser. B* **2012**, 54, 427–433.
- (364) Stache, E. E.; Kottisch, V.; Fors, B. P. Photocontrolled Radical Polymerization from Hydridic C–H Bonds. *J. Am. Chem. Soc.* **2020**, 142, 4581–4585.
- (365) Lalevé, J.; Blanchard, N.; Tehfe, M.-A.; Fouassier, J. P. Decatungstate ( $W_{10}O_{32}^{4-}$ )/Silane: A New and Promising Radical Source Under Soft Light Irradiation. *Macromol. Rapid Commun.* **2011**, 32, 838–843.
- (366) Bolland, J. L.; Cooper, H. R. The Photo-Sensitized Oxidation of Ethanol. *Proc. R. Soc. London. Ser. A. Math. Phys. Sci.* **1954**, 225, 405–426.
- (367) Schenck, G. O.; Koltzenburg, G.; Grossmann, H. Durch Benzophenon Photosensibilisierte Synthese Der Terebinsäure. *Angew. Chem.* **1957**, 69, 177–178.
- (368) Vaßen, R.; Runsink, J.; Scharf, H.-D. Chirale Induktion Bei Photochemischen Reaktionen, VI. Asymmetrische Steuerung Der Addition von Semipinacol-Radikalen an Chirale Malein- Und Fumarsäureester. *Chem. Ber.* **1986**, 119, 3492–3497.
- (369) Fraser-Reid, B.; Holder, N. L.; Yunker, M. B. Ground and Excited State 1,4-Addition Reactions of Some Carbohydrate Enones. *J. Chem. Soc., Chem. Commun.* **1972**, 1286–1287.
- (370) Fraser-Reid, B.; Holder, N. L.; Hicks, D. R.; Walker, D. L. Synthetic Applications of the Photochemically Induced Addition of Oxycarbonyl Species to  $\alpha$ -Enones. Part I. The Addition of Simple Alcohols. *Can. J. Chem.* **1977**, 55, 3978–3985.
- (371) Benko, Z.; Fraser-Reid, B.; Mariano, P. S.; Beckwith, A. L. J. Conjugate Addition of Methanol to  $\alpha$ -Enones: Photochemistry and Stereochemical Details. *J. Org. Chem.* **1988**, 53, 2066–2072.
- (372) Graalfs, H.; Fröhlich, R.; Wolff, C.; Mattay, J. Diastereoselective Addition of Radicals to Chiral 1,3-Dioxin-4-Ones. *Eur. J. Org. Chem.* **1999**, 1999, 1057–1073.
- (373) Fukuyama, T.; Yamada, K.; Nishikawa, T.; Ravelli, D.; Fagnoni, M.; Ryu, I. Site-Selectivity in TBADT-Photocatalyzed C(sp<sup>3</sup>)-H Functionalization of Saturated Alcohols and Alkanes. *Chem. Lett.* **2018**, 47, 207–209.
- (374) Oelgemöller, M.; Hoffmann, N. Photochemically Induced Radical Reactions with Furanones. *Pure Appl. Chem.* **2015**, 87, 569–582.
- (375) Shvydkiv, O.; Yavorsky, A.; Nolan, K.; Youssef, A.; Rigué, E.; Hoffmann, N.; Oelgemöller, M. Photosensitized Addition of Isopropanol to Furanones in a 365 Nm UV-LED Microchip. *Photochem. Photobiol. Sci.* **2010**, 9, 1601–1603.
- (376) Yavorsky, A.; Shvydkiv, O.; Nolan, K.; Hoffmann, N.; Oelgemöller, M. Photosensitized Addition of Isopropanol to Furanones in a Continuous-Flow Dual Capillary Microreactor. *Tetrahedron Lett.* **2011**, 52, 278–280.
- (377) Ghosh, A. K.; Leshchenko, S.; Noetzel, M. Stereoselective Photochemical 1,3-Dioxolane Addition to 5- Alkoxyethyl-2(5 H)-Furanone: Synthesis of Bis-Tetrahydrofuran Ligand for HIV Protease Inhibitor UIC-94017 (TMC-114). *J. Org. Chem.* **2004**, 69, 7822–7829.
- (378) Ravelli, D.; Albin, A.; Fagnoni, M. Smooth Photocatalytic Preparation of 2-Substituted 1,3-Benzodioxoles. *Chem. - Eur. J.* **2011**, 17, 572–579.

- (379) Prieto, A.; Taillefer, M. Visible-Light Decatungstate/Disulfide Dual Catalysis for the Hydro-Functionalization of Styrenes. *Org. Lett.* **2021**, *23*, 1484–1488.
- (380) Murphy, J. J.; Bastida, D.; Paria, S.; Fagnoni, M.; Melchiorre, P. Asymmetric Catalytic Formation of Quaternary Carbons by Iminium Ion Trapping of Radicals. *Nature* **2016**, *532*, 218–222.
- (381) Ravelli, D.; Zoccolillo, M.; Mella, M.; Fagnoni, M. Photocatalytic Synthesis of Oxetane Derivatives by Selective C–H Activation. *Adv. Synth. Catal.* **2014**, *356*, 2781–2786.
- (382) Bonassi, F.; Ravelli, D.; Protti, S.; Fagnoni, M. Decatungstate Photocatalyzed Acylations and Alkylations in Flow via Hydrogen Atom Transfer. *Adv. Synth. Catal.* **2015**, *357*, 3687–3695.
- (383) Ravelli, D.; Montanaro, S.; Zema, M.; Fagnoni, M.; Albini, A. A Tin-Free, Radical Photocatalyzed Addition to Vinyl Sulfones. *Adv. Synth. Catal.* **2011**, *353*, 3295–3300.
- (384) Yu, J.; Zhao, C.; Zhou, R.; Gao, W.; Wang, S.; Liu, K.; Chen, S.; Hu, K.; Mei, L.; Yuan, L.; Chai, Z.; Hu, H.; Shi, W. Visible-Light-Enabled C–H Functionalization by a Direct Hydrogen Atom Transfer Uranyl Photocatalyst. *Chem. - Eur. J.* **2020**, *26*, 16521–16529.
- (385) Yamada, K.; Fukuyama, T.; Fujii, S.; Ravelli, D.; Fagnoni, M.; Ryu, I. Cooperative Polar/Steric Strategy in Achieving Site-Selective Photocatalyzed C(sp<sup>3</sup>)–H Functionalization. *Chem. - Eur. J.* **2017**, *23*, 8615–8618.
- (386) Angioni, S.; Ravelli, D.; Emma, D.; Dondi, D.; Fagnoni, M.; Albini, A. Tetrabutylammonium Decatungstate (Chemo)Selective Photocatalyzed, Radical C–H Functionalization in Amides. *Adv. Synth. Catal.* **2008**, *350*, 2209–2214.
- (387) Capaldo, L.; Fagnoni, M.; Ravelli, D. Vinylpyridines as Building Blocks for the Photocatalyzed Synthesis of Alkylpyridines. *Chem. - Eur. J.* **2017**, *23*, 6527–6530.
- (388) Abadie, B.; Jardel, D.; Pozzi, G.; Toullec, P.; Vincent, J. Dual Benzophenone/Copper-Photocatalyzed Giese-Type Alkylation of C(sp<sup>3</sup>)–H Bonds. *Chem. - Eur. J.* **2019**, *25*, 16120–16127.
- (389) Paul, S.; Guin, J. Radical C(sp<sup>3</sup>)–H Alkenylation, Alkynylation and Allylation of Ethers and Amides Enabled by Photocatalysis. *Green Chem.* **2017**, *19*, 2530–2534.
- (390) Bonciolini, S.; Di Filippo, M.; Baumann, M. A Scalable Continuous Photochemical Process for the Generation of Amino-propylsulfones. *Org. Biomol. Chem.* **2020**, *18*, 9428–9432.
- (391) Kuang, Y.; Wang, K.; Shi, X.; Huang, X.; Meggers, E.; Wu, J. Asymmetric Synthesis of 1,4-Dicarbonyl Compounds from Aldehydes by Hydrogen Atom Transfer Photocatalysis and Chiral Lewis Acid Catalysis. *Angew. Chem., Int. Ed.* **2019**, *58*, 16859–16863.
- (392) Srivastava, V.; Singh, P. K.; Singh, P. P. Photocatalysed Eosin Y Mediated C(sp<sup>3</sup>)–H Alkylation of Amine Substrates via Direct HAT. *Tetrahedron Lett.* **2019**, *60*, 1333–1336.
- (393) Meanwell, N. A. Fluorine and Fluorinated Motifs in the Design and Application of Bioisosteres for Drug Design. *J. Med. Chem.* **2018**, *61*, 5822–5880.
- (394) Sarver, P. J.; Bacauanu, V.; Schultz, D. M.; DiRocco, D. A.; Lam, Y.; Sherer, E. C.; MacMillan, D. W. C. The Merger of Decatungstate and Copper Catalysis to Enable Aliphatic C(sp<sup>3</sup>)–H Trifluoromethylation. *Nat. Chem.* **2020**, *12*, 459–467.
- (395) Yamada, K.; Okada, M.; Fukuyama, T.; Ravelli, D.; Fagnoni, M.; Ryu, I. Photocatalyzed Site-Selective C–H to C–C Conversion of Aliphatic Nitriles. *Org. Lett.* **2015**, *17*, 1292–1295.
- (396) Shi, D.; He, C.; Sun, W.; Ming, Z.; Meng, C.; Duan, C. A Photosensitizing Decatungstate-Based MOF as Heterogeneous Photocatalyst for the Selective C–H Alkylation of Aliphatic Nitriles. *Chem. Commun.* **2016**, *52*, 4714–4717.
- (397) Fukuyama, T.; Nishikawa, T.; Yamada, K.; Ravelli, D.; Fagnoni, M.; Ryu, I. Photocatalyzed Site-Selective C(sp<sup>3</sup>)–H Functionalization of Alkylpyridines at Non-Benzyl Positions. *Org. Lett.* **2017**, *19*, 6436–6439.
- (398) Oliva, M.; Coppola, G. A.; Van der Eycken, E. V.; Sharma, U. K. Photochemical and Electrochemical Strategies towards Benzylic C–H Functionalization: A Recent Update. *Adv. Synth. Catal.* **2021**, *363*, 1810–1834.
- (399) Qrareya, H.; Ravelli, D.; Fagnoni, M.; Albini, A. Decatungstate Photocatalyzed Benzoylation of Alkenes with Alkylaromatics. *Adv. Synth. Catal.* **2013**, *355*, 2891–2899.
- (400) Wen, Z.; Maheshwari, A.; Sambiagio, C.; Deng, Y.; Laudadio, G.; Van Aken, K.; Sun, Y.; Gemoets, H. P. L.; Noël, T. Optimization of a Decatungstate-Catalyzed C(sp<sup>3</sup>)–H Alkylation Using a Continuous Oscillatory Millistructured Photoreactor. *Org. Process Res. Dev.* **2020**, *24*, 2356–2361.
- (401) Dondi, D.; Cardarelli, A. M.; Fagnoni, M.; Albini, A. Photomediated Synthesis of  $\beta$ -Alkylketones from Cycloalkanes. *Tetrahedron* **2006**, *62*, 5527–5535.
- (402) Dai, Z.-Y.; Nong, Z.-S.; Wang, P.-S. Light-Mediated Asymmetric Aliphatic C–H Alkylation with Hydrogen Atom Transfer Catalyst and Chiral Phosphoric Acid. *ACS Catal.* **2020**, *10*, 4786–4790.
- (403) Dai, Z.-Y.; Nong, Z.-S.; Song, S.; Wang, P.-S. Asymmetric Photocatalytic C(sp<sup>3</sup>)–H Bond Addition to  $\alpha$ -Substituted Acrylates. *Org. Lett.* **2021**, *23*, 3157–3161.
- (404) Laudadio, G.; Deng, Y.; van der Wal, K.; Ravelli, D.; Nunõ, M.; Fagnoni, M.; Guthrie, D.; Sun, Y.; Noël, T. C(sp<sup>3</sup>)–H Functionalizations of Light Hydrocarbons Using Decatungstate Photocatalysis in Flow. *Science* **2020**, *369*, 92–96.
- (405) Xu, S.; Chen, H.; Zhou, Z.; Kong, W. Three-Component Alkene Difunctionalization by Direct and Selective Activation of Aliphatic C–H Bonds. *Angew. Chem., Int. Ed.* **2021**, *60*, 7405–7411.
- (406) Campbell, M. W.; Yuan, M.; Polites, V. C.; Gutierrez, O.; Molander, G. A. Photochemical C–H Activation Enables Nickel-Catalyzed Olefin Dicarbofunctionalization. *J. Am. Chem. Soc.* **2021**, *143*, 3901–3910.
- (407) Shen, Y.; Dai, Z.-Y.; Zhang, C.; Wang, P.-S. Palladium-Catalyzed Allylic Alkylation via Photocatalytic Nucleophile Generation. *ACS Catal.* **2021**, *11*, 6757–6762.
- (408) Supranovich, V. I.; Levin, V. V.; Dilman, A. D. Radical Addition to N-Tosylimines via C–H Activation Induced by Decatungstate Photocatalyst. *Org. Lett.* **2019**, *21*, 4271–4274.
- (409) Qi, X.-K.; Guo, L.; Yao, L.-J.; Gao, H.; Yang, C.; Xia, W. Multicomponent Synthesis of  $\alpha$ -Branched Tertiary and Secondary Amines by Photocatalytic Hydrogen Atom Transfer Strategy. *Org. Lett.* **2021**, *23*, 4473–4477.
- (410) Yahata, K.; Sakurai, S.; Hori, S.; Yoshioka, S.; Kaneko, Y.; Hasegawa, K.; Akai, S. Coupling Reaction between Aldehydes and Non-Activated Hydrocarbons via the Reductive Radical-Polar Cross-over Pathway. *Org. Lett.* **2020**, *22*, 1199–1203.
- (411) Doohan, R. A.; Geraghty, N. W. A. A Comparative Analysis of the Functionalisation of Unactivated Cycloalkanes Using Alkynes and Either Sunlight or a Photochemical Reactor. *Green Chem.* **2005**, *7*, 91–96.
- (412) Geraghty, N. W.; Hannan, J. J. Functionalisation of Cycloalkanes: The Photomediated Reaction of Cycloalkanes with Alkynes. *Tetrahedron Lett.* **2001**, *42*, 3211–3213.
- (413) Geraghty, N. W. A.; Herson, E. M. The Reaction of Photochemically Generated  $\alpha$ -Hydroxyalkyl Radicals with Alkynes: A Synthetic Route to  $\gamma$ -Butenolides. *Tetrahedron Lett.* **2009**, *50*, 570–573.
- (414) Adak, T.; Hoffmann, M.; Witzel, S.; Rudolph, M.; Drew, A.; Hashmi, A. S. K. Visible Light-Enabled sp<sup>3</sup>-C–H Functionalization with Chloro- and Bromoalkynes: Chemoselective Route to Vinylchlorides or Alkynes. *Chem. - Eur. J.* **2020**, *26*, 15573–15580.
- (415) Cao, H.; Kuang, Y.; Shi, X.; Wong, K. L.; Tan, B. B.; Kwan, J. M. C.; Liu, X.; Wu, J. Photoinduced Site-Selective Alkenylation of Alkenes and Aldehydes with Aryl Alkenes. *Nat. Commun.* **2020**, *11*, 1956.
- (416) Kamijo, S.; Kamijo, K.; Murafuji, T. Synthesis of Alkylated Pyrimidines via Photoinduced Coupling Using Benzophenone as a Mediator. *J. Org. Chem.* **2017**, *82*, 2664–2671.
- (417) Lee, W.; Jung, S.; Kim, M.; Hong, S. Site-Selective Direct C–H Pyridylation of Unactivated Alkanes by Triplet Excited Anthraquinone. *J. Am. Chem. Soc.* **2021**, *143*, 3003–3012.

- (418) Perry, I. B.; Brewer, T. F.; Sarver, P. J.; Schultz, D. M.; DiRocco, D. A.; MacMillan, D. W. C. Direct Arylation of Strong Aliphatic C–H Bonds. *Nature* **2018**, *560*, 70–75.
- (419) Shen, Y.; Gu, Y.; Martin, R.  $\text{sp}^3$  C–H Arylation and Alkylation Enabled by the Synergy of Triplet Excited Ketones and Nickel Catalysts. *J. Am. Chem. Soc.* **2018**, *140*, 12200–12209.
- (420) Dewanji, A.; Krach, P. E.; Rueping, M. The Dual Role of Benzophenone in Visible-Light/Nickel Photoredox-Catalyzed C–H Arylations: Hydrogen-Atom Transfer and Energy Transfer. *Angew. Chem., Int. Ed.* **2019**, *58*, 3566–3570.
- (421) Ren, C.; Wang, T.; Zhang, Y.; Peng, D.; Liu, X.; Wu, Q.; Liu, X.; Luo, S. Photoinduced Activation of Unactivated C( $\text{sp}^3$ )-H Bonds and Acylation Reactions. *ChemistrySelect* **2021**, *6*, 2523–2528.
- (422) Li, C.-J. Cross-Dehydrogenative Coupling (CDC): Exploring C–C Bond Formations beyond Functional Group Transformations. *Acc. Chem. Res.* **2009**, *42*, 335–344.
- (423) Quattrini, M. C.; Fujii, S.; Yamada, K.; Fukuyama, T.; Ravelli, D.; Fagnoni, M.; Ryu, I. Versatile Cross-Dehydrogenative Coupling of Heteroaromatics and Hydrogen Donors via Decatungstate Photocatalysis. *Chem. Commun.* **2017**, *53*, 2335–2338.
- (424) Capaldo, L.; Quadri, L. L.; Merli, D.; Ravelli, D. Photoelectrochemical Cross-Dehydrogenative Coupling of Benzothiazoles with Strong Aliphatic C–H Bonds. *Chem. Commun.* **2021**, *57*, 4424–4427.
- (425) Chatgililoglu, C.; Crich, D.; Komatsu, M.; Ryu, I. Chemistry of Acyl Radicals. *Chem. Rev.* **1999**, *99*, 1991–2070.
- (426) Sumino, S.; Fusano, A.; Fukuyama, T.; Ryu, I. Carbonylation Reactions of Alkyl Iodides through the Interplay of Carbon Radicals and Pd Catalysts. *Acc. Chem. Res.* **2014**, *47*, 1563–1574.
- (427) Singh, J.; Sharma, S.; Sharma, A. Photocatalytic Carbonylation Strategies: A Recent Trend in Organic Synthesis. *J. Org. Chem.* **2021**, *86*, 24–48.
- (428) Boese, W. T.; Goldman, A. S. Alkane Carbonylation Photocatalyzed by Aromatic Ketones under High CO Pressure. *Tetrahedron Lett.* **1992**, *33*, 2119–2122.
- (429) Boese, W. T.; Goldman, A. S. Photochemical Cyclohexane Carbonylation Cocatalyzed by  $\text{d}^8$  Transition Metal Carbonyls and Aromatic Ketones and Aldehydes. *J. Am. Chem. Soc.* **1992**, *114*, 350–351.
- (430) Ryu, I.; Tani, A.; Fukuyama, T.; Ravelli, D.; Fagnoni, M.; Albini, A. Atom-Economical Synthesis of Unsymmetrical Ketones through Photocatalyzed C–H Activation of Alkanes and Coupling with CO and Electrophilic Alkenes. *Angew. Chem., Int. Ed.* **2011**, *50*, 1869–1872.
- (431) Ryu, I.; Tani, A.; Fukuyama, T.; Ravelli, D.; Montanaro, S.; Fagnoni, M. Efficient C–H/C–N and C–H/C–CO–N Conversion via Decatungstate-Photoinduced Alkylation of Diisopropyl Azodicarboxylate. *Org. Lett.* **2013**, *15*, 2554–2557.
- (432) Azam, M.; Al-Resayes, S. I.; Trzesowska-Kruszynska, A.; Kruszynski, R.; Kumar, P.; Jain, S. L. Seven-Coordinated Chiral Uranyl(VI) Salen Complex as Effective Catalyst for C–H Bond Activation of Dialkylanilines under Visible Light. *Polyhedron* **2017**, *124*, 177–183.
- (433) Esposti, S.; Dondi, D.; Fagnoni, M.; Albini, A. Acylation of Electrophilic Olefins through Decatungstate-Photocatalyzed Activation of Aldehydes. *Angew. Chem., Int. Ed.* **2007**, *46*, 2531–2534.
- (434) Ravelli, D.; Zema, M.; Mella, M.; Fagnoni, M.; Albini, A. Benzoyl Radicals from (Hetero)Aromatic Aldehydes. Decatungstate Photocatalyzed Synthesis of Substituted Aromatic Ketones. *Org. Biomol. Chem.* **2010**, *8*, 4158–4164.
- (435) Fagnoni, M.; Bonassi, F.; Palmieri, A.; Protti, S.; Ravelli, D.; Ballini, R. Flow Synthesis of Substituted  $\gamma$ -Lactones by Consecutive Photocatalytic/Reductive Reactions. *Adv. Synth. Catal.* **2014**, *356*, 753–758.
- (436) Garbarino, S.; Protti, S.; Gabrielli, S.; Fagnoni, M.; Palmieri, A.; Ravelli, D. Multi-Step Continuous Flow Synthesis of  $\beta/\gamma$ -Substituted Ketones. *ChemPhotoChem* **2018**, *2*, 847–850.
- (437) Okada, M.; Yamada, K.; Fukuyama, T.; Ravelli, D.; Fagnoni, M.; Ryu, I. Photocatalytic One-Pot Synthesis of Homoallyl Ketones via a Norrish Type I Reaction of Cyclopentanones. *J. Org. Chem.* **2015**, *80*, 9365–9369.
- (438) Yan, J.; Cheo, H. W.; Teo, W. K.; Shi, X.; Wu, H.; Idres, S. B.; Deng, L.-W.; Wu, J. A Radical Smiles Rearrangement Promoted by Neutral Eosin Y as a Direct Hydrogen Atom Transfer Photocatalyst. *J. Am. Chem. Soc.* **2020**, *142*, 11357–11362.
- (439) Fan, P.; Zhang, C.; Lan, Y.; Lin, Z.; Zhang, L.; Wang, C. Photocatalytic Hydroacylation of Trifluoromethyl Alkenes. *Chem. Commun.* **2019**, *55*, 12691–12694.
- (440) Liu, H.; Xue, F.; Wang, M.; Tang, X.; Wu, J. Neutral-Eosin Y-Catalyzed Regioselective Hydroacylation of Aryl Alkenes under Visible-Light Irradiation. *Synlett* **2021**, *32*, 406–410.
- (441) Wang, X.; Chen, Y.; Song, H.; Liu, Y.; Wang, Q. Synthesis of Unnatural  $\alpha$ -Amino Acids via Photoinduced Decatungstate-Catalyzed Giese Reactions of Aldehydes. *Org. Lett.* **2021**, *23*, 2199–2204.
- (442) Fan, P.; Lan, Y.; Zhang, C.; Wang, C. Nickel/Photocatalyzed Asymmetric Acyl-Carbamoylation of Alkenes. *J. Am. Chem. Soc.* **2020**, *142*, 2180–2186.
- (443) Fan, P.; Zhang, C.; Zhang, L.; Wang, C. Acylation of Aryl Halides and  $\alpha$ -Bromo Acetates with Aldehydes Enabled by Nickel/TBADT Cocatalysis. *Org. Lett.* **2020**, *22*, 3875–3878.
- (444) Kawaai, K.; Yamaguchi, T.; Yamaguchi, E.; Endo, S.; Tada, N.; Ikari, A.; Itoh, A. Photoinduced Generation of Acyl Radicals from Simple Aldehydes, Access to 3-Acyl-4-Arylcoumarin Derivatives, and Evaluation of Their Antiandrogenic Activities. *J. Org. Chem.* **2018**, *83*, 1988–1996.
- (445) Wang, H.; Li, T.; Hu, D.; Tong, X.; Zheng, L.; Xia, C. Acylation of Arenes with Aldehydes through Dual C–H Activations by Merging Photocatalysis and Palladium Catalysis. *Org. Lett.* **2021**, *23*, 3772–3776.
- (446) Mateo-Alonso, A.; Guldi, D. M.; Paolucci, F.; Prato, M. Fullerenes: Multitask Components in Molecular Machinery. *Angew. Chem., Int. Ed.* **2007**, *46*, 8120–8126.
- (447) Vougioukalakis, G. C.; Roubelakis, M. M.; Orfanopoulos, M. Open-Cage Fullerenes: Towards the Construction of Nanosized Molecular Containers. *Chem. Soc. Rev.* **2010**, *39*, 817–844.
- (448) Thompson, B. C.; Fréchet, J. M. J. Polymer–Fullerene Composite Solar Cells. *Angew. Chem., Int. Ed.* **2008**, *47*, 58–77.
- (449) Guldi, D. M.; Illescas, B. M.; Atienza, C. M.; Wielopolski, M.; Martín, N. Fullerene for Organic Electronics. *Chem. Soc. Rev.* **2009**, *38*, 1587–1597.
- (450) Nakamura, E.; Isobe, H. Functionalized Fullerenes in Water. The First 10 Years of Their Chemistry, Biology, and Nanoscience. *Acc. Chem. Res.* **2003**, *36*, 807–815.
- (451) Krusic, P. J.; Wasserman, E.; Keizer, P. N.; Morton, J. R.; Preston, K. F. Radical Reactions of  $\text{C}_{60}$ . *Science* **1991**, *254*, 1183–1185.
- (452) Tzirakis, M. D.; Orfanopoulos, M. Decatungstate-Mediated Radical Reactions of  $\text{C}_{60}$  with Substituted Toluenes and Anisoles: A New Photochemical Functionalization Strategy for Fullerenes. *Org. Lett.* **2008**, *10*, 873–876.
- (453) Tzirakis, M. D.; Orfanopoulos, M. Photochemical Addition of Ethers to  $\text{C}_{60}$ : Synthesis of the Simplest [60]Fullerene/Crown Ether Conjugates. *Angew. Chem., Int. Ed.* **2010**, *49*, 5891–5893.
- (454) Tzirakis, M. D.; Alberti, M. N.; Orfanopoulos, M. Hydroxyalkylation of [60]Fullerene: Free Radical Addition of Alcohols to  $\text{C}_{60}$ . *Chem. Commun.* **2009**, *46*, 8228–8230.
- (455) Tzirakis, M. D.; Orfanopoulos, M. Acyl Radical Reactions in Fullerene Chemistry: Direct Acylation of [60]Fullerene through an Efficient Decatungstate-Photomediated Approach. *J. Am. Chem. Soc.* **2009**, *131*, 4063–4069.
- (456) Ravelli, D.; Montanaro, S.; Tomasi, C.; Galinetto, P.; Quartarone, E.; Merli, D.; Mustarelli, P.; Fagnoni, M. One-Step Decatungstate-Photomediated PEGylation of Single-Walled Carbon Nanotubes. *ChemPlusChem* **2012**, *77*, 210–216.
- (457) Ravelli, D.; Merli, D.; Quartarone, E.; Profumo, A.; Mustarelli, P.; Fagnoni, M. PEGylated Carbon Nanotubes: Preparation, Properties and Applications. *RSC Adv.* **2013**, *3*, 13569–13582.



- (458) Papadopoulos, G. N.; Kokotos, C. G. One-Pot Amide Bond Formation from Aldehydes and Amines via a Photoorganocatalytic Activation of Aldehydes. *J. Org. Chem.* **2016**, *81*, 7023–7028.
- (459) Zwick, C. R.; Renata, H. Remote C–H Hydroxylation by an  $\alpha$ -Ketoglutarate-Dependent Dioxygenase Enables Efficient Chemoenzymatic Synthesis of Manzacidin C and Proline Analogs. *J. Am. Chem. Soc.* **2018**, *140*, 1165–1169.
- (460) Zwick, C. R.; Renata, H. Evolution of Biocatalytic and Chemocatalytic C–H Functionalization Strategy in the Synthesis of Manzacidin C. *J. Org. Chem.* **2018**, *83*, 7407–7415.
- (461) Niu, L.; Jiang, C.; Liang, Y.; Liu, D.; Bu, F.; Shi, R.; Chen, H.; Chowdhury, A. D.; Lei, A. Manganese-Catalyzed Oxidative Azidation of C(sp<sup>3</sup>)–H Bonds under Electrophotocatalytic Conditions. *J. Am. Chem. Soc.* **2020**, *142*, 17693–17702.
- (462) Pitzer, L.; Schwarz, J. L.; Glorius, F. Reductive Radical-Polar Crossover: Traditional Electrophiles in Modern Radical Reactions. *Chem. Sci.* **2019**, *10*, 8285–8291.
- (463) Wiles, R. J.; Molander, G. A. Photoredox-Mediated Net-Neutral Radical/Polar Crossover Reactions. *Isr. J. Chem.* **2020**, *60*, 281–293.
- (464) Wan, T.; Capaldo, L.; Laudadio, G.; Nyuchev, A.; Rincon, J.; Garcia-Losada, P.; Mateos Gutierrez, C.; Frederick, M. O.; Nuno, M.; Noël, T. Decatungstate-mediated C(sp<sup>3</sup>)–H Heteroarylation via Radical-Polar Crossover in Batch and Flow. *Angew. Chem., Int. Ed.* **2021**, DOI: 10.1002/anie.202104682.
- (465) Chen, C.; Wang, Q.; Xiong, Z. Method for preparing KA oil through cyclohexane oxidation. CN105315128A, 2016-02-10.
- (466) Alini, S.; Babini, P. The Industrial Oxidation of KA Oil to Adipic Acid. In *Handbook of Advanced Methods and Processes in Oxidation Catalysis*; Duprez, D., Cavani, F., Eds.; Imperial College Press, 2014; pp 320–333.
- (467) Mao, Y.; Bakac, A. Photocatalytic Oxidation of Toluene to Benzaldehyde by Molecular Oxygen. *J. Phys. Chem.* **1996**, *100*, 4219–4223.
- (468) Ohkubo, K.; Hirose, K.; Fukuzumi, S. Photooxygenation of Alkanes by Dioxygen with p-Benzoquinone Derivatives with High Quantum Yields. *Photochem. Photobiol. Sci.* **2016**, *15*, 731–734.
- (469) Wells, C. F. Hydrogen Transfer to Quinones. Part 1. - Kinetics of the Deactivation of the Photo-Excited Quinone. *Trans. Faraday Soc.* **1961**, *57*, 1703–1718.
- (470) Wells, C. F. Hydrogen Transfer to Quinones. Part 2. - Reactivities of Alcohols, Ethers and Ketones. *Trans. Faraday Soc.* **1961**, *57*, 1719–1731.
- (471) Clark, K. P.; Stonehill, H. I. Photochemistry and Radiation Chemistry of Anthraquinone-2-Sodium-Sulphonate in Aqueous Solution. Part 1. - Photochemical Kinetics in Aerobic Solution. *J. Chem. Soc., Faraday Trans. 1* **1972**, *68*, 577–590.
- (472) Clark, K. P.; Stonehill, H. I. Photochemistry and Radiation Chemistry of 9,10-Anthraquinone-2-Sodium Sulphonate in Aqueous Solution. Part 2. - Photochemical Products. *J. Chem. Soc., Faraday Trans. 1* **1972**, *68*, 1676–1686.
- (473) Lykakis, I. N.; Tanielian, C.; Seghrouchni, R.; Orfanopoulos, M. Mechanism of Decatungstate Photocatalyzed Oxygenation of Aromatic Alcohols Part II. Kinetic Isotope Effects Studies. *J. Mol. Catal. A: Chem.* **2007**, *262*, 176–184.
- (474) Lykakis, I. N.; Orfanopoulos, M. Deuterium Kinetic Isotope Effects in Homogeneous Decatungstate Catalyzed Photooxygenation of 1,1-Diphenylethane and 9-Methyl-9H-Fluorene: Evidence for a Hydrogen Abstraction Mechanism. *Tetrahedron Lett.* **2005**, *46*, 7835–7839.
- (475) Lykakis, I. N.; Vougioukalakis, G. C.; Orfanopoulos, M. Homogeneous Decatungstate-Catalyzed Photooxygenation of Tetra-substituted Alkenes: A Deuterium Kinetic Isotope Effect Study. *J. Org. Chem.* **2006**, *71*, 8740–8747.
- (476) Lykakis, I.; Orfanopoulos, M. Decatungstate-Catalyzed Photooxygenation of S-2-Phenylbutane and Cumene via a Free Carbon-Radical Intermediate. *Curr. Org. Chem.* **2009**, *13*, 1737–1745.
- (477) Nomiya, K.; Sugie, Y.; Miyazaki, T.; Miwa, M. Catalysis by Heteropolyacids - IX. Photocatalytic Oxidation of Isopropyl Alcohol to Acetone under Oxygen Using Tetrabutylammonium Decatungstate. *Polyhedron* **1986**, *5*, 1267–1271.
- (478) Carraro, M.; Gardan, M.; Scorrano, G.; Drioli, E.; Fontananova, E.; Bonchio, M. Solvent-Free, Heterogeneous Photooxygenation of Hydrocarbons by Hyflon? Membranes Embedding a Fluorous-Tagged Decatungstate. *Chem. Commun.* **2006**, 4533–4535.
- (479) Nieweg, J. A.; Lemma, K.; Trewyn, B. G.; Lin, V. S.-Y.; Bakac, A. Mesoporous Silica-Supported Uranyl: Synthesis and Photo-reactivity. *Inorg. Chem.* **2005**, *44*, 5641–5648.
- (480) Hermans, I.; Nguyen, T. L.; Jacobs, P. A.; Peeters, J. Autoxidation of Cyclohexane: Conventional Views Challenged by Theory and Experiment. *ChemPhysChem* **2005**, *6*, 637–645.
- (481) Maldotti, A.; Amadelli, R.; Carassiti, V.; Molinari, A. Catalytic Oxygenation of Cyclohexane by Photoexcited (nBu<sub>4</sub>N)<sub>4</sub>W<sub>10</sub>O<sub>32</sub>: The Role of Radicals. *Inorg. Chim. Acta* **1997**, *256*, 309–312.
- (482) Maldotti, A.; Molinari, A.; Bergamini, P.; Amadelli, R.; Battioni, P.; Mansuy, D. Photocatalytic Oxidation of Cyclohexane by (nBu<sub>4</sub>N)<sub>4</sub>W<sub>10</sub>O<sub>32</sub>Fe(III)Porphyrins Integrated Systems. *J. Mol. Catal. A: Chem.* **1996**, *113*, 147–157.
- (483) Wu, W.; Fu, Z.; Tang, S.; Zou, S.; Wen, X.; Meng, Y.; Sun, S.; Deng, J.; Liu, Y.; Yin, D. (nBu<sub>4</sub>N)<sub>4</sub>W<sub>10</sub>O<sub>32</sub>-Catalyzed Selective Oxygenation of Cyclohexane by Molecular Oxygen under Visible Light Irradiation. *Appl. Catal., B* **2015**, *164*, 113–119.
- (484) Giannotti, C.; Richter, C. Photocatalyzed Oxidation of Cyclohexane by W<sub>10</sub>O<sub>32</sub><sup>4-</sup> Irradiation with Natural Sunlight. *Int. J. Photoenergy* **1999**, *1*, 69–73.
- (485) Fornal, E.; Giannotti, C. Photocatalyzed Oxidation of Cyclohexane with Heterogenized Decatungstate. *J. Photochem. Photobiol., A* **2007**, *188*, 279–286.
- (486) Molinari, A.; Amadelli, R.; Andreotti, L.; Maldotti, A. Heterogeneous Photocatalysis for Synthetic Purposes: Oxygenation of Cyclohexane with H<sub>3</sub>PW<sub>12</sub>O<sub>40</sub> and (nBu<sub>4</sub>N)<sub>4</sub>W<sub>10</sub>O<sub>32</sub> Supported on Silica. *J. Chem. Soc., Dalton Trans.* **1999**, 1203–1204.
- (487) Maldotti, A.; Molinari, A.; Varani, G.; Lenarda, M.; Storaro, L.; Bigi, F.; Maggi, R.; Mazzacani, A.; Sartori, G. Immobilization of (n-Bu<sub>4</sub>N)<sub>4</sub>W<sub>10</sub>O<sub>32</sub> on Mesoporous MCM-41 and Amorphous Silicas for Photocatalytic Oxidation of Cycloalkanes with Molecular Oxygen. *J. Catal.* **2002**, *209*, 210–216.
- (488) Molinari, A.; Amadelli, R.; Mazzacani, A.; Sartori, G.; Maldotti, A. Tetralkylammonium and Sodium Decatungstate Heterogenized on Silica: Effects of the Nature of Cations on the Photocatalytic Oxidation of Organic Substrates. *Langmuir* **2002**, *18*, 5400–5405.
- (489) Arnold, P. L.; Purkis, J. M.; Rutkauskaitė, R.; Kovacs, D.; Love, J. B.; Austin, J. Controlled Photocatalytic Hydrocarbon Oxidation by Uranyl Complexes. *ChemCatChem* **2019**, *11*, 3786–3790.
- (490) Symeonidis, T. S.; Tamiolakis, I.; Armatas, G. S.; Lykakis, I. N. Green Photocatalytic Organic Transformations by Polyoxometalates vs. Mesoporous TiO<sub>2</sub> Nanoparticles: Selective Aerobic Oxidation of Alcohols. *Photochem. Photobiol. Sci.* **2015**, *14*, 563–568.
- (491) Lykakis, I. N.; Tanielian, C.; Orfanopoulos, M. Decatungstate Photocatalyzed Oxidation of Aryl Alkanols. Electron Transfer or Hydrogen Abstraction Mechanism? *Org. Lett.* **2003**, *5*, 2875–2878.
- (492) Molinari, A.; Bratovcic, A.; Magnacca, G.; Maldotti, A. Matrix Effects on the Photocatalytic Oxidation of Alcohols by [nBu<sub>4</sub>N]<sub>4</sub>W<sub>10</sub>O<sub>32</sub> Incorporated into Sol–Gel Silica. *Dalt. Trans.* **2010**, *39*, 7826–7833.
- (493) Maldotti, A.; Molinari, A.; Bigi, F. Selective Photooxidation of Diols with Silica Bound W<sub>10</sub>O<sub>32</sub><sup>4-</sup>. *J. Catal.* **2008**, *253*, 312–317.
- (494) Farhadi, S.; Momeni, Z. Zirconia-Supported Sodium Decatungstate (Na<sub>4</sub>W<sub>10</sub>O<sub>32</sub>/ZrO<sub>2</sub>): An Efficient, Green and Recyclable Photocatalyst for Selective Oxidation of Activated Alcohols to Carbonyl Compounds with O<sub>2</sub>. *J. Mol. Catal. A: Chem.* **2007**, *277*, 47–52.
- (495) Fujiya, A.; Nobuta, T.; Yamaguchi, E.; Tada, N.; Miura, T.; Itoh, A. Aerobic Photooxidative Direct Asymmetric Aldol Reactions of Benzyl Alcohols Using Water as the Solvent. *RSC Adv.* **2015**, *5*, 39539–39543.



- (496) Zhang, W.; Gacs, J.; Arends, I. W. C. E.; Hollmann, F. Selective Photooxidation Reactions Using Water-Soluble Anthraquinone Photocatalysts. *ChemCatChem* **2017**, *9*, 3821–3826.
- (497) Tada, N.; Hattori, K.; Nobuta, T.; Miura, T.; Itoh, A. Facile Aerobic Photooxidation of Methyl Group in the Aromatic Nucleus in the Presence of an Organocatalyst under VIS Irradiation. *Green Chem.* **2011**, *13*, 1669–1671.
- (498) Taguchi, M.; Nagasawa, Y.; Yamaguchi, E.; Tada, N.; Miura, T.; Itoh, A. One-Pot Epoxidation of Alkenes Using Aerobic Photoperoxidation of Toluenes. *Tetrahedron Lett.* **2016**, *57*, 230–232.
- (499) Dong, J.; Yue, F.; Wang, X.; Song, H.; Liu, Y.; Wang, Q. Light-Mediated Difluoromethylthiolation of Aldehydes with a Hydrogen Atom Transfer Photocatalyst. *Org. Lett.* **2020**, *22*, 8272–8277.
- (500) Wang, X.; Dong, J.; Liu, Y.; Song, H.; Wang, Q. Decatungstate as a Direct Hydrogen Atom Transfer Photocatalyst for Synthesis of Trifluoromethylthioesters from Aldehydes. *Chin. Chem. Lett.* **2021**, DOI: 10.1016/j.ccl.2021.03.070.
- (501) Cao, S.; Hong, W.; Ye, Z.; Gong, L. Photocatalytic Three-Component Asymmetric Sulfonylation via Direct C(sp<sup>3</sup>)-H Functionalization. *Nat. Commun.* **2021**, *12*, 2377.
- (502) Yerien, D. E.; Bonesi, S.; Postigo, A. Fluorination Methods in Drug Discovery. *Org. Biomol. Chem.* **2016**, *14*, 8398–8427.
- (503) Yakubov, S.; Barham, J. P. Photosensitized Direct C–H Fluorination and Trifluoromethylation in Organic Synthesis. *Beilstein J. Org. Chem.* **2020**, *16*, 2151–2192.
- (504) Rueda-Becerril, M.; Chatalova Sazepin, C.; Leung, J. C. T.; Okbinoglu, T.; Kennepohl, P.; Paquin, J.-F.; Sammis, G. M. Fluorine Transfer to Alkyl Radicals. *J. Am. Chem. Soc.* **2012**, *134*, 4026–4029.
- (505) Spencer, A. R. A.; Grainger, R.; Panigrahi, A.; Lepper, T. J.; Bentkowska, K.; Larrosa, I. Transition Metal-Free Cross-Dehydrogenative Arylation of Unactivated Benzylic C–H Bonds. *Chem. Commun.* **2020**, *56*, 14479–14482.
- (506) Nodwell, M. B.; Bagai, A.; Halperin, S. D.; Martin, R. E.; Knust, H.; Britton, R. Direct Photocatalytic Fluorination of Benzylic C–H Bonds with N-Fluorobenzenesulfonimide. *Chem. Commun.* **2015**, *51*, 11783–11786.
- (507) Halperin, S. D.; Fan, H.; Chang, S.; Martin, R. E.; Britton, R. A Convenient Photocatalytic Fluorination of Unactivated C–H Bonds. *Angew. Chem., Int. Ed.* **2014**, *53*, 4690–4693.
- (508) Liu, W.; Huang, X.; Cheng, M.-J.; Nielsen, R. J.; Goddard, W. A.; Groves, J. T. Oxidative Aliphatic C–H Fluorination with Fluoride Ion Catalyzed by a Manganese Porphyrin. *Science* **2012**, *337*, 1322–1325.
- (509) Halperin, S. D.; Kwon, D.; Holmes, M.; Regalado, E. L.; Campeau, L.-C.; DiRocco, D. A.; Britton, R. Development of a Direct Photocatalytic C–H Fluorination for the Preparative Synthesis of Odanacatib. *Org. Lett.* **2015**, *17*, 5200–5203.
- (510) Yuan, Z.; Nodwell, M. B.; Yang, H.; Malik, N.; Merckens, H.; Bénard, F.; Martin, R. E.; Schaffer, P.; Britton, R. Site-Selective, Late-Stage C–H <sup>18</sup>F-Fluorination on Unprotected Peptides for Positron Emission Tomography Imaging. *Angew. Chem., Int. Ed.* **2018**, *57*, 12733–12736.
- (511) Yuan, Z.; Yang, H.; Malik, N.; Čolović, M.; Weber, D. S.; Wilson, D.; Bénard, F.; Martin, R. E.; Warren, J. J.; Schaffer, P.; Britton, R. Electrostatic Effects Accelerate Decatungstate-Catalyzed C–H Fluorination Using [<sup>18</sup>F]- and [<sup>19</sup>F]NFSI in Small Molecules and Peptide Mimics. *ACS Catal.* **2019**, *9*, 8276–8284.
- (512) Meanwell, M.; Lehmann, J.; Eichenberger, M.; Martin, R. E.; Britton, R. Synthesis of Acyl Fluorides via Photocatalytic Fluorination of Aldehydic C–H Bonds. *Chem. Commun.* **2018**, *54*, 9985–9988.
- (513) Yang, L.; Tan, D.; Fan, W.; Liu, X.; Wu, J.; Huang, Z.; Li, Q.; Wang, H. Photochemical Radical C–H Halogenation of Benzyl N-Methyliminodiacetyl (MIDA) Boronates: Synthesis of A-Functionalized Alkyl Boronates. *Angew. Chem., Int. Ed.* **2021**, *60*, 3454–3458.
- (514) Kleoff, M.; Schwan, J.; Christmann, M.; Heretsch, P. A Modular, Argon-Driven Flow Platform for Natural Product Synthesis and Late-Stage Transformations. *Org. Lett.* **2021**, *23*, 2370–2374.
- (515) Zhang, Y.; Ji, P.; Dong, Y.; Wei, Y.; Wang, W. Deuteration of Formyl Groups via a Catalytic Radical H/D Exchange Approach. *ACS Catal.* **2020**, *10*, 2226–2230.
- (516) Loh, Y. Y.; Nagao, K.; Hoover, A. J.; Hesk, D.; Rivera, N. R.; Colletti, S. L.; Davies, I. W.; MacMillan, D. W. C. Photoredox-Catalyzed Deuteration and Tritiation of Pharmaceutical Compounds. *Science* **2017**, *358*, 1182–1187.
- (517) Zhou, R.; Li, J.; Cheo, H. W.; Chua, R.; Zhan, G.; Hou, Z.; Wu, J. Visible-Light-Mediated Deuteration of Silanes with Deuterium Oxide. *Chem. Sci.* **2019**, *10*, 7340–7344.
- (518) Zhou, R.; Ma, L.; Yang, X.; Cao, J. Recent Advances in Visible-Light Photocatalytic Deuteration Reactions. *Org. Chem. Front.* **2021**, *8*, 426–444.
- (519) Dong, J.; Wang, X.; Wang, Z.; Song, H.; Liu, Y.; Wang, Q. Formyl-Selective Deuteration of Aldehydes with D<sub>2</sub>O via Synergistic Organic and Photoredox Catalysis. *Chem. Sci.* **2020**, *11*, 1026–1031.
- (520) Kuang, Y.; Cao, H.; Tang, H.; Chew, J.; Chen, W.; Shi, X.; Wu, J. Visible Light Driven Deuteration of Formyl C–H and Hydridic C(sp<sup>3</sup>)-H Bonds in Feedstock Chemicals and Pharmaceutical Molecules. *Chem. Sci.* **2020**, *11*, 8912–8918.
- (521) Qrareya, H.; Dondi, D.; Ravelli, D.; Fagnoni, M. Decatungstate-Photocatalyzed Si–H/C–H Activation in Silyl Hydrides: Hydrosilylation of Electron-Poor Alkenes. *ChemCatChem* **2015**, *7*, 3350–3357.
- (522) Hill, C. L.; Renneke, R. F.; Combs, L. Anaerobic Functionalization of Remote Unactivated Carbon-Hydrogen Bond by Polyoxometalates. *Tetrahedron* **1988**, *44*, 7499–7507.
- (523) Renneke, R. F.; Hill, C. L. Selective Photochemical Dehydrogenation of Saturated Hydrocarbons with Quantum Yields Approaching Unity. *Angew. Chem., Int. Ed. Engl.* **1988**, *27*, 1526–1527.
- (524) West, J. G.; Huang, D.; Sorensen, E. J. Acceptorless Dehydrogenation of Small Molecules through Cooperative Base Metal Catalysis. *Nat. Commun.* **2015**, *6*, 10093.
- (525) Kattamuri, P. V.; West, J. G. Cooperative Hydrogen Atom Transfer: From Theory to Applications. *Synlett* **2021**, *32*, 1179.
- (526) Abrams, D. J.; West, J. G.; Sorensen, E. J. Toward a Mild Dehydroformylation Using Base-Metal Catalysis. *Chem. Sci.* **2017**, *8*, 1954–1959.
- (527) Tachikawa, Y.; Cui, L.; Matsusaki, Y.; Tada, N.; Miura, T.; Itoh, A. Aerobic Photooxidative Cleavage of 1,3-Diketones to Carboxylic Acids Using 2-Chloroanthraquinone. *Tetrahedron Lett.* **2013**, *54*, 6218–6221.
- (528) Yamaguchi, T.; Kudo, Y.; Hirashima, S.; Yamaguchi, E.; Tada, N.; Miura, T.; Itoh, A. Facile and Efficient Synthesis of Hydroxyalkyl Esters from Cyclic Acetals through Aerobic Photo-Oxidation Using Anthraquinone-2-Carboxylic Acid. *Tetrahedron Lett.* **2015**, *56*, 1973–1975.
- (529) Hardwick, T.; Ahmed, N. C–H Functionalization via Electrophotocatalysis and Photoelectrochemistry: Complementary Synthetic Approach. *ACS Sustainable Chem. Eng.* **2021**, *9*, 4324–4340.
- (530) Kim, K.; Lee, S.; Hong, S. H. Direct C(sp<sup>3</sup>)-H Cyanation Enabled by a Highly Active Decatungstate Photocatalyst. *Org. Lett.* **2021**, *23*, 5501–5505.
- (531) Parodi, A.; Jorea, A.; Fagnoni, M.; Ravelli, D.; Samorì, C.; Torri, C.; Galletti, P. Bio-based crotonic acid from polyhydroxybutyrate: synthesis and photocatalyzed hydroacylation. *Green Chem.* **2021**, *23*, 3420–3427.
- (532) Murugesan, V.; Ganguly, A.; Karthika, A.; Rasappan, R. C–H Alkylation of Aldehydes by Merging TBADT Hydrogen Atom Transfer with Nickel Catalysis. *Org. Lett.* **2021**, *23*, 5389–5393.
- (533) Malliaros, N. G.; Kellner, I. D.; Drewello, T.; Orfanopoulos, M. Decatungstate-Photocatalyzed Addition of Lactones to [60]-Fullerene. *J. Org. Chem.* **2021**, *86*, 9876–9882.
- (534) Sarver, P. J.; Bissonnette, N. B.; MacMillan, D. W. C. Decatungstate-Catalyzed C(sp<sup>3</sup>)-H Sulfonylation: Rapid Access to Diverse Organosulfur Functionality. *J. Am. Chem. Soc.* **2021**, *143*, 9737–9743.

(535) Schirmer, T. E.; Rolka, A. B.; Karl, T. A.; Holzhausen, F.; König, B. Photocatalytic C–H Trifluoromethylthiolation by the Decatungstate Anion. *Org. Lett.* **2021**, *23*, 5729–5733.

#### NOTE ADDED IN PROOF

The photocatalyzed cyanation of alkanes has been reported (compare section 2.4)<sup>530</sup> The photocatalyzed acylation of crotonic acid<sup>531</sup> and the alkylation of aldehydes with pyridinium salts<sup>532</sup> to give ketones have been published (compare section 2.5). The decatungstate-photocatalyzed functionalization of [60]fullerene with lactones was devised (compare section 2.8).<sup>533</sup> The photocatalyzed C(sp<sup>3</sup>)–H sulfonylation<sup>534</sup> or C(sp<sup>3</sup>)–H/C(sp<sup>2</sup>)–H trifluoromethylthiolation<sup>535</sup> for the forging of C–S bonds have been obtained (compare section 3.3).

#### NOTE ADDED AFTER ASAP PUBLICATION

This paper was originally published ASAP on August 6, 2021, with Table 1 absent from the PDF version. The revised version was posted August 16, 2021.