SUPPORTING INFORMATION

Direct preparation of nucleoside vinyl disulfides from 2-(trimethylsilyl)ethyl sulfides, an access to vinylthiols.

Béatrice Gerland,[†] Jérôme Désiré,[†] Michel Lepoivre,[‡] Jean-Luc Décout^{†*}

[†]Département de Pharmacochimie Moléculaire UMR 5063 CNRS/Université Joseph Fourier-Grenoble I, ICMG FR CNRS 2607, BP 53, 38041 Grenoble Cedex 9, France and [‡]Institut de Biochimie Biophysique Moléculaire et Cellulaire, UMR 8619 CNRS, Université de Paris Sud, 91405 Orsay, France.

SYNTHESIS

2'-Deoxy-2'-(2-(trimethylsilyl)ethyl)thiothymidine **2**

To a suspension of 2,2'-anhydrothymidine (5.0 g, 22 mmol) and anhydrous potassium carbonate (11 g, 79 mmol) in dry DMF (110 mL) was added 2-(trimethylsilyl)ethanethiol (3.5 g, 26 mmol). The resulting solution was stirred under argon at 120°C for 3 h, filtrated, evaporated, and the residue obtained was purified by chromatography on silica gel in dichloromethane-methanol (95:5) to give compound **2** as a white solid (6.9 g, 19 mmol, 88%); mp 71-73°C; ¹H NMR (400 MHz, CDCl₃) δ 8.73 (1 H, s, NH); 7.25 (1 H, d, J = 8.0 Hz, 6-H), 5.46 (1 H, d, J = 9.2 Hz, 1'-H), 4.36 (1 H, m, 3'-H), 4.24 (1 H, m, 4'-H), 3.99 (2 H, m, 2'-H + 5'-H), 3.81 (1 H, m, 5'-H), 2.60 (2 H, m, S-CH₂), 1.96 (3 H, s, 5-CH₃), 0.85 (2 H, m, CH₂-Si), -0.02 (9 H, s, Si(CH₃)₃); ¹³C NMR (100 MHz, CDCl₃) δ 163.4 (C2), 150.5 (C4), 138.9 (C6), 111.4 (C5), 93.3 (C1'), 86.6 (C4'), 71.5 (C3'), 63.1 (5'-CH₂), 52.8 (C2'), 28.4 (S-CH₂), 18.1 (CH₂-Si), 12.4 (CH₃), -1.8 (Si(CH₃)₃); LRMS (DCI, NH₃-isobutane) m/z 375 [M+H]⁺, 392 [M+H+NH₃]⁺; elemental analysis calculated for C₁₅H₂₆N₂O₄SSi, 0.75 H₂O: C 47.53, H 7.05, N 7.39, S 8.46, found C 47.23, H 7.03, N 7.29, S 8.04.

2',3'-Didehydro-2',3'-dideoxy-2'-(2-(trimethylsilyl)ethyl)thiouridine **3**

To a solution of 2'-deoxy-2'-(2-(trimethylsilyl)ethyl)thiouridine **1** (5.00 g, 13.8 mmol) in anhydrous pyridine (50 mL), at 0°C under argon, was added 4,4'-dimethoxytrityl chloride (5.20 g, 15.3 mmol) and the resulting solution was stirred 24 h at RT, then evaporated. The obtained residue was

dissolved in dichloromethane (100 mL), washed with water (100 mL), and the organic layer was dried over MgSO_4 . After concentration the residue was chromatographed on silica gel in a mixture dichloromethane-methanol (98:2) with 2% de triethylamine to give the tritylated compound as a white solid (7.23 g, 10.9 mmol, 91%).

To this compound (7.10 g, 10.7 mmol) in anhydrous pyridine (50 mL) was added mesyl chloride (1 mL; 12.9 mmol). The solution was stirred at 0°C for 1 h then at RT for 15 h. Water (5 mL) was added then the solution was evaporated. The residue was dissolved in dichloromethane (100 mL), washed with water (50 mL), dried over MgSO_4 and evaporated to dryness.

The mesylated compound obtained (8.36 g, 11.3 mmol) in dichloromethane (20 mL) was treated with dichloroacetic acid (80 mL, 2% in dichloromethane) for 4 h. The solution was neutralized with aqueous sodium hydrogenocarbonate (5%; 80 mL), then the aqueous layer was extracted twice with dichloromethane (100 mL). The combined extracts were dried over MgSO_4 and evaporated. The residue was chromatographed on silica gel in dichloromethane-methanol (98:2) to give the deprotected and mesylated product (3.72 g, 7.24 mmol, 71% for 3 steps).

To this compound (3.72 g, 7.24 mmol) in acetonitrile (50 mL) was added potassium carbonate (6 g, 43 mmol), and then the mixture for stirred at 90°C for 3 days, evaporated and chromatographed on silica gel in dichloromethane-methanol (98:2) to yield **3** as a white solid (2.55 g, 7.44 mmol, 85%, or 60% for 4 steps); mp: 72-74°C; ^1H NMR (400 MHz, CDCl_3) δ 9.30 (1 H, s, NH), 7.61 (1 H, d, J = 8 Hz, 6-H), 6.94 (1 H, m, 1'-H), 5.81 (1 H, m, 3'-H), 5.70 (1 H, d, J = 8 Hz, 5-H), 4.99 (1 H, m, 4'-H), 3.94-3.74 (2 H, m, 5'-H $\times$ 2), 2.85 (2 H, m, S-CH $_2$), 0.94 (2 H, m, CH $_2$ -Si), 0.09 (9 H, s, Si(CH $_3$) $_3$); ^{13}C NMR (100 MHz, CDCl_3) δ 163.7 (C2), 150.8 (C4), 141.3 (C6), 134.9 (C2'), 123.6 (C3'), 102.5 (C5), 90.2 (C1'), 87.5 (C4'), 63.3 (5'-CH $_2$), 28.4 (S-CH $_2$), 16.2 (CH $_2$ -Si), -1.9 (Si(CH $_3$) $_3$); LRMS (DCI, NH_3 -isobutane) m/z 343 $[\text{M}+\text{H}]^+$; elemental analysis calculated for $\text{C}_{14}\text{H}_{22}\text{N}_2\text{O}_4\text{SSi}$, 0.5 H_2O : C 47.84, H 6.60, N 7.97, S 9.12; found: C 47.92, H 6.76, N 7.96, S 9.86.

2',3'-Didehydro-2',3'-dideoxy-2'-(2-(trimethylsilyl)ethyl)thiothymidine **4**

To a solution of 2'-deoxy-2'-(2-(trimethylsilyl)ethyl)thiothymidine **2** (0.75 g, 2.00 mmol) in anhydrous pyridine (5 mL), at 0°C under argon, was added 4,4'-dimethoxytrityl chloride (1.9 g, 2.4 mmol) and the resulting solution was stirred 24 h at RT, then evaporated. The obtained residue was dissolved in dichloromethane (20 mL), washed with water (20 mL), and the organic layer was dried over MgSO_4 . After evaporation, the residue was chromatographed on silica gel in a mixture dichloromethane-methanol (98:2) with 2% de triethylamine to give the tritylated compound as a white solid (1.20 g, 1.35 mmol, 90%).

To this compound (1.20 g, 1.35 mmol) in anhydrous pyridine (10 mL) was added mesyl chloride (125 μL , 1.62 mmol). The solution was stirred at 0°C for 1 h then at RT for 15 h. Water (1 mL) was added then the solution was evaporated. The residue was dissolved in dichloromethane (20 mL), washed with water (20 mL), dried over MgSO_4 and evaporated to dryness.

The mesylated compound in dichloromethane (10 mL) was treated with dichloroacetic acid (30 mL, 2% in dichloromethane) for 4 h. The solution was neutralized with aqueous sodium hydrogenocarbonate (5%, 20 mL), then the aqueous layer was extracted twice with dichloromethane (100 mL). The combined extracts were dried over MgSO_4 and evaporated. The residue was chromatographed on silica gel in dichloromethane-methanol (98:2) to give the deprotected and mesylated product (0.63 g, 0.66 mmol).

To this compound in acetonitrile (15 mL) was added potassium carbonate (0.53 g, 3.85 mmol), and then the mixture for stirred at 90°C for 3 days, evaporated and chromatographed on silica gel in dichloromethane-methanol (98:2) to yield **4** as a white solid (453 mg, 1.25 mmol, 84%; 64% for 4 steps); mp: 59-61°C; ¹H NMR (400 MHz, CDCl₃) δ 8.73 (1 H, s, NH), 7.38 (1 H, s, 6-H), 6.94 (1 H, m, 1'-H), 5.81 (1 H, m, 3'-H), 4.98 (1 H, m, 4'-H), 3.97-3.74 (2 H, m, 5'-H), 2.85 (2 H, m, S-CH₂), 1.87 (3 H, s, 5-CH₃), 0.94 (2 H, m, CH₂-Si), 0.07 (9 H, s, Si(CH₃)₃); ¹³C NMR (100 MHz, CDCl₃) δ 164.0 (C2), 150.6 (C4), 136.9 (C6), 135.2 (C2'), 123.5 (C3'), 110.9 (C5), 90.3 (C1'), 87.4 (C4'), 63.5 (5'-CH₂), 28.5 (S-CH₂), 16.3 (CH₂-Si), 12.4 (CH₃), -1.8 (Si(CH₃)₃); LRMS (DCI, NH₃-isobutane): m/z 357 [M+H]⁺; elemental analysis calculated for C₁₅H₂₄N₂O₄SSi: C 50.54, H 6.79, N 7.86, S 8.99; found: C 50.58, H 6.62, N 7.48, S 8.87.

General procedures for preparing the unsaturated mixed disulfides **5-10**

To a stirred solution of **3** or **4** in the appropriated anhydrous solvent (see table below, dichloromethane DCM, or dichloroethane DCE) was added the sulfenyl chloride (2- or 4-nitrobenzenesulfenyl, trichloromethanesulfenyl chlorides, (2NBSCl, 4NBSCl, TCMSCl) 3 eq.). The solution was stirred at RT or reflux under argon, then evaporated, and the obtained residue was purified by chromatography on silica gel in dichloromethane-methanol (98:2 to 90:10) to give the expected disulfide **5-10**.

Compound	R'	Solvent	t (h)	T	Yield (%)
5	4NB	DCM	4	RT	66
6	2NB	DCE	48	rfx	71
7	TCM	DCM	48	rfx	78
8	4NB	DCM	4	RT	71
9	2NB	DCE	48	rfx	76
10	TCM	DCM	48	rfx	72

2',3'-Didehydro-2',3'-dideoxy-2'-(4-nitrophenyldithio)uridine **5**

5 was obtained as a yellow solid (152 mg, 0.38 mmol, 66%) from **3** (200 mg, 0.58 mmol) in dichloromethane (6 mL) with 4NBSCl (333 mg, 1.75 mmol); mp: 105-109°C; ^1H NMR (400 MHz, CDCl_3) δ 8.12 (1 H, s, NH), 8.18 (2 H, m, Ar), 7.68 (3 H, m, 6-H, Ar), 7.01 (1 H, m, 1'-H), 6.48 (1 H, ls, 3'-H), 5.70 (1 H, d, J = 8.4 Hz, 5-H), 4.99 (1 H, ls, 4'-H), 3.92 (1 H, m, 5'-H), 3.80 (1 H, m, 5'-H); ^{13}C NMR (100 MHz, CDCl_3) δ 162.9 (C2), 150.3 (C4), 146.8 (C-NO₂), 143.9 (C-SS), 140.8 (C6), 135.1 (C2'), 131.8 (C3'), 126.6 (2 C, Ar), 126.3 (2 C, Ar), 102.8 (C5), 89.7 (C1'), 87.0 (C4'), 63.0 (5'-CH₂); LRMS (DCI, NH_3 -isobutane): m/z 396 $[\text{M}+\text{H}]^+$, 413 $[\text{M}+\text{NH}_3]^+$; HRMS (electrospray) calculated for $\text{C}_{15}\text{H}_{13}\text{N}_3\text{O}_6\text{NaS}_2$: $[\text{M}+\text{Na}]^+$ 418.0143; found: 418.0140; elemental analysis calculated for $\text{C}_{15}\text{H}_{13}\text{N}_3\text{O}_6\text{S}_2$, 0.5 MeOH: C 45.26, H 3.67, N 10.21; found C 45.27, H 3.84, N 10.15.

2',3'-Didehydro-2',3'-dideoxy-2'-(2-nitrophenyldithio)uridine **6**

6 was obtained as a yellow solid (82 mg, 0.21 mmol, 71%) from **3** (100 mg, 0.29 mmol) in dichloroethane (3 mL) with 2NBSCl (166 mg, 0.88 mmol); mp: 71-73°C; ^1H NMR (400 MHz, CDCl_3) δ 8.31 (2 H, m, NH + Ar), 8.02 (1 H, m, Ar), 7.68 (2 H, m, Ar, J = 8.0 Hz, 6-H), 7.43 (1 H, m, Ar), 6.98 (1 H, m, 1'-H), 6.40 (1 H, m, 3'-H), 5.66 (1 H, d, J = 8.4 Hz, 5-H), 4.97 (1 H, m, 4'-H), 3.94-3.77 (2 H, m, 5'-H $\times$ 2); ^{13}C NMR (100 MHz, CDCl_3) δ 162.6 (C2), 150.1 (C4), 145.4 (C-NO₂), 140.7 (C6), 135.1 (C-SS), 134.3 (1 C, Ar), 133.1 (C3'), 132.4 (C2'), 127.0, 126.9, 126.2 (3 C, Ar), 102.8 (C5), 89.9 (C1'), 87.0 (C4'), 63.1 (5'-CH₂); LRMS (DCI, NH_3 -isobutane): m/z 396 $[\text{M}+\text{H}]^+$, 413 $[\text{M}+\text{NH}_3]^+$; HRMS (electrospray) calculated for $\text{C}_{15}\text{H}_{13}\text{N}_3\text{O}_6\text{NaS}_2$: $[\text{M}+\text{Na}]^+$ 418.0143; found: 418.0132.

2',3'-Didehydro-2',3'-dideoxy-2'-(trichloromethyldithio)uridine **7**

7 was obtained as a white solid (45 mg, 0.11 mmol, 78%) from **3** (50 mg, 0.15 mmol) in dichloromethane (3 mL) with TCMSCl (48 μL , 0.44 mmol); mp: 149°C (decomposition); ^1H NMR (400 MHz, CD_3OD) δ 9.23 (1 H, s, NH), 7.75 (1 H, d, J = 8 Hz, 6-H), 7.15 (1 H, m, 1'-H), 6.68 (1 H, m, 3'-H), 5.74 (1 H, d, J = 8 Hz, 5-H), 5.04 (1 H, m, 4'-H), 3.97 (1 H, dd, J = 2.4 Hz, J = 12.8 Hz, 5'-H), 3.84 (1 H, dd, J = 2.4 Hz, J = 12.4 Hz, 5'-H); ^{13}C NMR (100 MHz, CDCl_3) δ 164.6 (C2), 150.6 (C4), 141.2 (C6), 136.0 (C3'), 131.3 (C2'), 102.9 (C5), 99.9 (CCl_3), 90.0 (C1'), 87.4 (C4'), 62.9 (5'-CH₂); LRMS (DCI, NH_3 -isobutane): m/z 393 $[\text{M}+\text{H}]^+$; HRMS (electrospray) calculated for $\text{C}_{10}\text{H}_9\text{N}_2\text{O}_4^{35}\text{Cl}_3\text{NaS}_2$: $[\text{M}+\text{Na}]^+$ 412.8967; found: 412.8961.

2',3'-Didehydro-2',3'-dideoxy-2'-(4-nitrophenyldithio)thymidine **8**

8 was obtained as a yellow solid (32 mg, 0.08 mmol, 71%) from **4** (40 mg, 0.11 mmol) in dichloromethane (3 mL) with 4NBSCl (80 mg, 0.42 mmol); mp: 61-63°C; ^1H NMR (400 MHz, CDCl_3) δ 8.18 (2 H, m, Ar), 7.60 (2 H, m, Ar), 7.35 (1H, s, 6-H), 6.91 (1 H, ls, 1'-H), 6.39 (1 H, ls, 3'-H), 4.92 (1 H, ls, 4'-H), 3.84 (1 H, m, 5'-H), 3.73 (1 H, m, 5'-H), 1.83 (3 H, s, 5-CH₃); ^{13}C NMR (100 MHz, CDCl_3) δ 162.5 (C2), 150.7 (C4), 146.7 (C-NO₂), 144.2 (C-SS), 135.1 (C6), 133.1 (C2'), 132.4 (C3'), 126.7 (2 C, Ar), 124.2 (2 C, Ar), 102.5 (C5), 91.2 (C1'), 86.9 (C4'), 63.2 (5'-CH₂), 13.1 (5-CH₃); LRMS (DCI, NH_3 -isobutane): m/z 432 $[\text{M}+\text{Na}]^+$; HRMS (electrospray) calculated for $\text{C}_{16}\text{H}_{15}\text{N}_3\text{O}_6\text{NaS}_2$: $[\text{M}+\text{Na}]^+$ 432.0300; found: 432.0295; calculated for $\text{C}_{16}\text{H}_{14}\text{N}_3\text{O}_6\text{Na}_2\text{S}_2$: $[\text{M}-\text{H}+2\text{Na}]^+$ 454.0119; found: 454.0120.

2',3'-Didehydro-2',3'-dideoxy-2'-(2-nitrophenyldithio)thymidine **9**

9 was obtained as a yellow solid (42 mg, 0.10 mmol, 76%) from **4** (50 mg, 0.14 mmol) in dichloroethane (3 mL) with 2NBSCl (80 mg, 0.42 mmol); mp: 72-74°C; ^1H NMR (400 MHz, CDCl_3) δ 8.62 (1 H, s, NH), 8.30 (1 H, m, Ar), 8.01 (1 H, m, Ar), 7.70 (1 H, m, Ar), 7.44 (1 H, m, Ar), 7.28 (1 H, ls, 6-H), 7.43 (1 H, m, Ar), 6.81 (1 H, m, 1'-H), 6.30 (1 H, m, 3'-H), 4.90 (1 H, m, 4'-H), 3.81 (1H, dd, $J = 2.8$ Hz; $J = 12.4$ Hz; 5'-H), 3.69 (1 H, m, dd, $J = 3.2$ Hz; $J = 12.4$ Hz; 5'-H), 1.86 (3H, s, 5-CH₃); ^{13}C NMR (100 MHz, CDCl_3) δ 162.3 (C2), 150.7 (C4), 145.2 (C-NO₂), 135.3 (C-SS), 135.1 (C6), 134.4 (1 C, Ar), 132.7 (C3'), 131.9 (C2'), 127.1, 126.9, 126.1 (3 C, Ar), 110.4 (C5), 89.9 (C1'), 86.9 (C4'), 63.3 (5'-CH₂), 13.0 (5-CH₃); LRMS (DCI, NH_3 -isobutane): m/z 410 $[\text{M}+\text{H}]^+$; HRMS (electrospray) calculated for $\text{C}_{16}\text{H}_{15}\text{N}_3\text{O}_6\text{NaS}_2$: $[\text{M}+\text{Na}]^+$ 432.0300; found: 432.0295; calculated for $\text{C}_{16}\text{H}_{15}\text{N}_3\text{O}_6\text{KS}_2$: $[\text{M}+\text{K}]^+$ 448.0039; found: 448.0059.

2',3'-Didehydro-2',3'-dideoxy-2'-(trichloromethyldithio)thymidine **10**

10 was obtained as a white solid (41 mg, 0.10 mmol, 72%) from **4** (50 mg, 0.14 mmol) in dichloromethane (3 mL) with TCMSCl (45 μ L, 0.42 mmol); mp: 149°C (decomposition); ^1H NMR (400 MHz, CDCl_3) δ 8.57 (1 H, s, NH), 7.43 (1 H, ls, 6-H), 7.05 (1 H, m, 1'-H), 6.62 (1 H, m, 3'-H), 5.01 (1 H, m, 4'-H), 3.92 (1 H, dd, $J = 2.4$ Hz, $J = 12.4$ Hz, 5'-H), 3.82 (1 H, dd, $J = 2.8$ Hz, $J = 12.4$ Hz, 5'-H), 1.94 (3 H, s, 5- CH_3); ^{13}C NMR (100 MHz, CDCl_3) δ 162.7 (C2), 150.8 (C4), 135.2 (C3'), 135.1 (C6), 131.8 (C2'), 110.6 (C5), 100.1 (CCl_3), 91.4 (C1'), 86.8 (C4'), 63.3 (5'- CH_2), 13.1 (5- CH_3); LRMS (DCI, NH_3 -isobutane): m/z 407 $[\text{M}+\text{H}]^+$; HRMS (electrospray) calculated for $\text{C}_{11}\text{H}_{11}\text{N}_2\text{O}_4^{35}\text{Cl}_3\text{NaS}_2$: $[\text{M}+\text{Na}]^+$: 426.9123; found: 426.9117.

General procedures for preparing the mixed disulfides **11-13** and **17-19**

To a stirred solution of **1** or **14** in freshly distilled methanol was added the sulfonyl chloride (4NBSCl, or 2NBSCl, or TCMSCl, 3 equiv.). The solution was stirred at RT under argon for 4 h, then evaporated, and the obtained residue was purified by chromatography on silica gel in dichloromethane-methanol (98:2 to 90:10) to give the expected disulfide **11-13** or **17-19**.

Compound	R	Yield (%)
11	4NB	92
12	2NB	96
13	TCM	84
17	4NB	69
18	2NB	42
19	TCM	86

2'-Deoxy-2'-(4-nitrophenyldithio)uridine **11**

11 was obtained as a yellow solid (421 mg, 1.02 mmol, 92%) from **1** (400 mg, 1.11 mmol) in methanol (8 mL) with 4NBSCl (632 mg, 3.33 mmol); mp: 192-194°C; ^1H NMR (400 MHz, CD_3OD) δ 8.18 (2 H, d, Ar), 7.78 (1 H, d, $J = 8.2$ Hz, 6-H), 7.66 (2 H, d, Ar), 6.35 (1 H, d, $J = 9.2$

Hz, 1'-H), 5.56 (1 H, d, $J = 8.4$ Hz, 5-H), 4.48 (1 H, m, 3'-H), 4.02 (1 H, m, 4'-H), 3.79 (1 H, m, 2'-H), 3.70 (2 H, m, 5'-H $\times 2$); ^{13}C NMR (100 MHz, CD_3OD) δ 163.9 (C2), 150.8 (C4), 146.5 (C-NO $_2$), 145.7 (C-SS), 140.6 (C6), 125.9 (2 C, Ar), 123.7 (2 C, Ar), 101.9 (C5), 87.9 (C1'), 86.8 (C4'), 72.9 (C3'), 61.5 (C2'), 60.1 (5'-CH $_2$); LRMS (DCI, NH_3 -isobutane): m/z 414 $[\text{M}+\text{H}]^+$, 431 $[\text{M}+\text{H}+\text{NH}_3]^+$; HRMS (electrospray) calculated for $\text{C}_{15}\text{H}_{15}\text{N}_3\text{O}_7\text{NaS}_2$: $[\text{M}+\text{Na}]^+$ 436.0249; found: 436.0247; elemental analysis calculated for $\text{C}_{15}\text{H}_{15}\text{N}_3\text{O}_7\text{S}_2$: C 43.58, H 3.66, N 10.16, S 15.51; found: 43.84, H 3.84, N 9.79, S 15.48.

2'-Deoxy-2'-(2-nitrophenyldithio)uridine **12**

12 was obtained as a yellow solid (220 mg, 0.53 mmol, 96%) from **1** (200 mg, 0.55 mmol) in methanol (4 mL) with 2NBSCl (316 mg, 1.66 mmol); mp: 175-178°C; ^1H NMR (400 MHz, DMSO) δ 11.14 (1 H, s, NH), 8.21 (1 H, d, Ar), 8.13 (1 H, d, Ar), 7.79 (1 H, m, Ar), 7.62 (1 H, d, $J = 8.1$ Hz, 6-H), 7.49 (1 H, m, Ar), 6.23 (1 H, d, $J = 9.2$ Hz, 1'-H), 6.17 (1 H, m, 3'-OH), 5.42 (1 H, d, $J = 8.0$ Hz, 5-H), 5.03 (1 H, m, 3'-H), 4.35 (1 H, m, 5'-OH), 3.91 (1 H, m, 4'-H), 3.69 (1 H, dd, $J = 5.2$ Hz, $J = 9.1$ Hz, 2'-H), 3.51 (2H, m, 5'-H $\times 2$); ^{13}C NMR (100 MHz, DMSO) δ 162.8 (C2), 151.0 (C4), 145.3 (C-NO $_2$), 140.3 (C6), 136.0 (C-SS), 135.3, 127.7, 127.5, 126.4 (4 C, Ar), 102.7 (C5), 87.3 (C1'), 86.9 (C4'), 72.7 (C3'), 61.6 (5'-CH $_2$), 58.3 (C2'); LRMS (DCI, NH_3 -isobutane): m/z 414 $[\text{M}+\text{H}]^+$, 431 $[\text{M}+\text{H}+\text{NH}_3]^+$; HRMS (electrospray) calculated for $\text{C}_{15}\text{H}_{15}\text{N}_3\text{O}_7\text{KS}_2$: $[\text{M}+\text{K}]^+$ 451.9989; found: 452.0008.

2'-Deoxy-2'-(trichloromethyldithio)uridine **13**

13 was obtained as a white solid (48 mg, 0.12 mmol, 84%) from **1** (50 mg, 0.14 mmol) in methanol (1.5 mL) with TCMSCl (44 μL , 0.42 mmol); mp: 109-111°C; ^1H NMR (400 MHz, CD_3OD) δ 8.01 (1 H, d, $J = 8$ Hz, 6-H), 6.38 (1 H, d, $J = 8.8$ Hz, 1'-H), 5.78 (1 H, d, $J = 8$ Hz, 5-H), 4.54 (1 H, m, 3'-H), 4.25 (1 H, dd, $J = 5.6$ Hz, $J = 8.8$ Hz, 2'-H), 4.09 (1 H, m, 4'-H), 3.78 (2 H, m, 5'-H $\times 2$); ^{13}C NMR (100 MHz, CD_3OD) δ 164.4 (C2), 151.1 (C4), 141.0 (C6), 102.2 (C5), 100.2 (CCl $_3$), 88.6 (C1'), 87.3 (C4'), 72.8 (C3'), 61.6 (5'-CH $_2$), 59.0 (C2'); LRMS (DCI, NH_3 -isobutane): m/z 411 $[\text{M}+\text{H}]^+$; HRMS (electrospray) calculated for $\text{C}_{10}\text{H}_{11}\text{N}_2\text{O}_5^{35}\text{Cl}_3\text{NaS}_2$: $[\text{M}+\text{Na}]^+$ 430.9072; found: 430.9081; elemental analysis calculated for $\text{C}_{10}\text{H}_{11}\text{N}_2\text{Cl}_3\text{O}_5\text{S}_2$, 0.75 MeOH: C 29.77, H 3.25, N 6.46, S 14.79; found: C 30.15, H 3.05, N 6.61, S 14.65.

N-4-Benzoyl-3'-(*O*-mesyl)-5'-(*O*-*tert*-butyldiphenylsilyl)-2'-(2-(trimethylsilyl)ethyl)thiocytidine **15**

To a solution of **14** (1.00 g, 2.78 mmol) in dry pyridine (15 mL) was added at 0°C under argon *tert*butyldiphenylsilyl chloride (870 μ l, 3.34 mmol). The solution was stirred at RT for 48 h, evaporated, and the residue was dissolved in dichloromethane (100 mL) and washed with water (50 mL). The organic extract was dried over MgSO₄ and evaporated to be purified by chromatography on silica gel in dichloromethane-methanol (95:5) to give the 5' protected compound (1.37 g, 2.29 mmol, 83%) as a white powder.

To a solution of this compound (1.30 g, 2.20 mmol) in dry DMF (15 mL) was added benzoic anhydride (497 mg, 2.2 mmol). The mixture was refluxed for 24 h, evaporated, dissolved in dichloromethane (80 mL) and washed twice with water (50 mL). The combined organic extracts were dried over MgSO₄ and evaporated.

Mesyl chloride (194 μ L, 2.5 mmol) was added at 0°C under argon to the crude product obtained dissolved in pyridine (15 mL). The mixture was stirred at RT for 15 h, then water (15 mL) was added and after 30 min the solvents were evaporated. The residue was dissolved in dichloromethane (60 mL), washed with water (50 mL), and the organic extract dried over MgSO₄ to be purified by chromatography on silica gel in dichloromethane-methanol (98:2) to give **15** as a white foam (1.57 g, 2.01 mmol, 91% for 2 steps): ¹H NMR (400 MHz, CDCl₃) δ 8.71 (1 H, ls, NH), 8.41 (1 H, d, J = 7.4 Hz, 6-H), 7.68 (5 H, m, Ar), 7.56-7.43 (10 H, m, Ar), 7.27 (1 H, d, J = 7.6 Hz, 5-H), 6.43 (1 H, d, J = 7.6 Hz, 1'-H), 5.34 (1 H, m, 3'-H), 4.43 (1 H, m, 4'-H), 4.09-3.95 (2 H, dd, J = 1.9 Hz, J = 12.0 Hz, 5'-Hx2), 3.64 (1 H, m, 2'-H), 3.15 (3 H, s, CH₃, Ms), 2.65 (2 H, m, S-CH₂), 1.16 (9 H, s, *t*Bu), 0.80 (2 H, m, Si-CH₂), 0.04 (9 H, s, Si(CH₃)₃); ¹³C NMR (100 MHz, CDCl₃) δ 169.5 (CO), 162.3 (C4), 150.1 (C2), 144.1 (C6), [135.6, 135.3, 133.2, 132.2, 131.6, 130.4, 130.3, 129.0, 128.3, 128.2, 127.6] (Ar), 97.9 (C5), 89.2 (C1'), 84.2 (C4'), 79.3 (C3'), 62.9 (5'-CH₂), 52.0 (C2'), 38.7 (CH₃ Ms), 27.8 (S-CH₂), 27.1 ((CH₃)₃), 17.5 (C(CH₃)₃), 17.5 (CH₂-Si), -1.9 (Si(CH₃)₃); LRMS (FAB⁺, NAB+NaCl): *m/z* 780 [M+H]⁺, 802 [M+Na]⁺; elemental analysis calculated for C₃₈H₄₉N₃O₇S₂Si₂: C 58.51, H 6.33, N 5.39, S 8.22; found: C 58.26, H 6.45, N 5.32, S 8.45.

2',3'-Didehydro-2',3'-dideoxy-2'-(2-(trimethylsilyl)ethyl)thiocytidine **16**

To a solution of **15** (1.40 g, 1.79 mmol) in ethanol (5 mL) was added concentrated aqueous ammonia (30%, 10 mL), then the solution was stirred at RT for 24 h. After evaporation the residue was chromatographed on silica gel in dichloromethane-methanol (95:5) to give a white solid (1.15 g, 1.71 mmol, 95%).

To this compound (1.00 g, 1.47 mmol) in acetonitrile (10 mL) was added potassium carbonate (1.24 g, 8.82 mmol) and the resulting mixture was stirred at RT under argon for 48 h. After filtration and evaporation, the residue was chromatographed on silica gel in dichloromethane-methanol (95:5) to give the unsaturated compound (0.51 g, 1.06 mmol, 72%) as a white foam.

To this compound (150 mg, 0.31 mmol) was added a methanolic solution of ammonium fluoride (800 μ L, 0.5 M, 0.41 mmol). The resulting solution was refluxed for 6 h, then allowed to cool at RT overnight before concentration. The product was then chromatographed on silica gel in dichloromethane-methanol (90:10) to give **16** (81 mg, 0.24 mmol, 76%) as a white solid (29% from **14** for 6 steps); mp: 93°C (decomposition); ^1H NMR (400 MHz, CD_3OD) δ 7.86 (1 H, d, J = 7.6 Hz, 6-H), 6.99 (1 H, m, 1'-H), 5.94 (1 H, ls, 3'-H), 5.89 (1 H, d, J = 7.6 Hz, 5-H), 4.90 (1 H, m, 4'-H), 3.73 (2 H, m, 5'-H $\times$ 2), 2.89 (2 H, m, S-CH $_2$), 0.95 (2 H, m, CH $_2$ -Si), 0.07 (9 H, s, Si(CH $_3$) $_3$); ^{13}C NMR (100 MHz, CD_3OD) δ 166.2 (C4), 157.3 (C2), 142.1 (C6), 135.4 (C2'), 123.9 (C3'), 95.1 (C5), 90.6 (C1'), 87.4 (C4'), 62.8 (5'-CH $_2$), 27.6 (S-CH $_2$), 15.9 (CH $_2$ -Si), -3.3 (Si(CH $_3$) $_3$); LRMS (DCI, NH_3 -isobutane): m/z 342 $[\text{M}+\text{H}]^+$; elemental analysis calculated for $\text{C}_{30}\text{H}_{43}\text{N}_3\text{O}_4\text{SSi}$, 0.5 H_2O : C 47.97, H 6.90, N 11.99, S 9.15; found: 47.77, H 6.90, N 11.71, S 9.68.

2'-Deoxy-2'-(4-nitrophenyldithio)cytidine **17**

17 was obtained after crystallization in methanol as a yellow solid (158 mg, 0.38 mmol, 69%) from **14** (200 mg, 0.56 mmol) in methanol (6 mL) with 4NBSCl (316 mg, 1.66 mmol); mp: 167°C (decomposition); ^1H NMR (400 MHz, DMSO) δ 8.13 (2 H, m, Ar), 7.65 (2 H, m, Ar), 7.60 (1 H, d, J = 7.6 Hz, 6-H), 7.16 (2 H, ls, 2 NH), 6.27 (1 H, d, J = 9.2 Hz, 1'-H), 6.13 (1 H, d, J = 5.2 Hz, 3'-OH), 5.57 (1 H, d, J = 7.6 Hz, 5-H), 5.05 (1 H, t, J = 5.2 Hz, 5'-OH), 4.33 (1 H, m, 3'-H), 3.89 (1 H, m, 4'-H), 3.70 (1 H, m, 2'-H), 3.51 (2 H, ls, 5'-H $\times$ 2); ^{13}C NMR (100 MHz, DMSO) δ 165.7 (C4), 155.6 (C2), 146.3 (C-NO $_2$), 146.1 (C-SS), 141.6 (C6), 126.2 (2 C Ar), 124.4 (2 C Ar), 95.1 (C5), 88.6 (C1'), 86.7 (C4'), 72.7 (C3'), 61.8 (5'-CH $_2$), 59.7 (C2'); LRMS (FAB $^+$, NAB): m/z 413 $[\text{M}+\text{H}]^+$; HRMS (electrospray) calculated for $\text{C}_{15}\text{H}_{17}\text{N}_4\text{O}_6\text{S}_2$: $[\text{M}+\text{H}]^+$ 413.0589; found: 413.0592.

2'-Deoxy-2'-(2-nitrophenyldithio)cytidine **18**

18 was obtained as a yellow solid (43 mg, 0.10 mmol, 42%) from **14** (100 mg, 0.28 mmol) in methanol (4 mL) with 2NBSCl (158 mg, 0.83 mmol); mp: 172°C (decomposition); ^1H NMR (400 MHz, CD_3OD) δ 8.21 (2 H, m, Ar), 7.73 (2 H, m, Ar + 6-H, J = 7.5 Hz), 7.44 (1 H, m, Ar), 6.42 (1 H, d, J = 9.3 Hz, 1'-H), 5.70 (1 H, d, J = 7.4 Hz, 5-H), 4.47 (1 H, m, 3'-H), 4.03 (1 H, ls, 4'-H),

3.70-3.62 (3 H, m, 5'-H $\times$ 2 + 2'-H); ^{13}C NMR (100 MHz, CD_3OD) δ 165.1 (C4), 157.4 (C2), 145.3 (C-NO $_2$), 141.3 (C6), 135.9 (C-SS), 134.4 (1 C Ar), 127.0, 126.6, 125.5 (3 C Ar), 96.1 (C5), 88.8 (C1'), 86.9 (C4'), 72.7 (C3'), 61.6 (5'-CH $_2$), 59.2 (C2'); LRMS (FAB $^+$, NAB): m/z 413 [M+H] $^+$; HRMS (electrospray) calculated for $\text{C}_{15}\text{H}_{16}\text{N}_4\text{O}_6\text{NaS}_2$: [M+Na] $^+$ 435.0409; found: 435.0404; elemental analysis calculated for $\text{C}_{15}\text{H}_{16}\text{N}_4\text{O}_6\text{S}_2$, 0.75 MeOH: C 42.30, H 4.14, N 13.15; found: C 42.55, H 4.11, N 12.69.

2'-Deoxy-2'-(trichloromethyldithio)cytidine **19**

19 was obtained as a white solid (53 mg, 0.13 mmol, 86%) from **14** (40 mg, 0.11 mmol) in methanol (1 mL) with TCMSCl (38 μL , 0.33 mmol); mp: 151°C (decomposition); ^1H NMR (400 MHz, CD_3OD) δ 7.99 (1 H, d, J = 7.6 Hz, 6-H), 6.43 (1 H, d, J = 8.8 Hz, 1'-H), 5.98 (1 H, d, J = 7.6 Hz, 5-H), 4.55 (1 H, m, 3'-H), 4.25 (1 H, dd, J = 5.2 Hz, J = 8.8 Hz, 2'-H), 4.09 (1 H, m, 4'-H), 3.78 (2 H, m, 5'-H $\times$ 2); ^{13}C NMR (100 MHz, CD_3OD) δ 165.8 (C4), 156.8 (C2), 141.9 (C6), 100.3 (CCl_3), 95.6 (C5), 89.7 (C1'), 87.1 (C4'), 72.8 (C3'), 61.7 (5'-CH $_2$), 59.5 (C2'); LRMS (DCI, NH_3 -isobutane): m/z 410 [M+H] $^+$; HRMS (electrospray) calculated for $\text{C}_{10}\text{H}_{12}\text{N}_3\text{O}_4\text{Na}^{35}\text{Cl}_3\text{S}_2$: [M+Na] $^+$ 429.9232; found: 429.9237.

Enethiol **20** and bis(2',3'-didehydro-2',3'-dideoxyuridin-2'-yl) disulfide

A solution of **5** (1 mg, 0.03 mmol) in MeOD (0.6 mL) in an NMR tube was degazed with argon and then DTT (6.3 mg, 0.09 mmol) was added, and the solution degazed again with argon. After 30 min at RT a ^1H NMR spectrum was recorded showing the presence of starting material and a new product for which 3'-H was found at 6.55 ppm (3'-H of intermediate **20**). After few hours, only one product was observed by ^1H NMR for which 3'-H was found at 6.22 ppm (3'-H of symmetrical disulfide).

Enethiol **20**

^1H NMR (400 MHz, CD_3OD) δ 7.90 (1 H, d, J = 8.0 Hz, 6-H), 7.06 (1 H, m, 1'-H), 6.55 (1 H, m, 3'-H), 5.72 (1 H, d, J = 8.4 Hz, 5-H), 4.93 (1 H, s, 4'-H), 3.73 (2 H, m, 5'-H $\times$ 2).

Symmetrical disulfide

^1H NMR (400 MHz, CD_3OD) δ 7.88 (1H, d, J = 8.0 Hz, 6-H), 6.83 (1 H, m, 1'-H), 6.22 (1 H, m, 3'-H), 5.75 (1 H, d, J = 8.0 Hz, 5-H), 4.91 (1H, s, 4'-H), 3.71 (2 H, m, 5'-H $\times$ 2); LRMS (FAB $^+$, NBA): m/z = 242 [M/2+H] $^+$.

2',3'-Didehydro-2',3'-dideoxy-2'-methylthiouridine **21**

Argon was bubbled into a stirred solution of **5** (30 mg, 0.08 mmol) in dry methanol. After 5 min DTT (59 mg, 0.38 mmol) was added and the solution was stirred for 30 min at RT under argon. NH₄HCO₃ (30 mg, 0.38 mmol) and methyl iodide (190 μ L, 3.0 mmol) were successively added. The mixture was stirred at RT for 1 h, evaporated, and chromatographed on silica gel in dichloromethane-methanol (95:5) to give **21** as a white foam (16 mg, 0.06 mmol, 82%); ¹H NMR (400 MHz, CDCl₃) δ 7.60 (1 H, d, J = 8.0 Hz, 6-H), 6.93 (1 H, m, 1'-H), 5.76 (1 H, s, 3'-H), 5.73 (1 H, d, J = 7.6 Hz, 5-H), 4.99 (1 H, m, 4'-H), 3.94-3.76 (2 H, m, J = 3.2 Hz, J = 11.6 Hz, 5'-Hx 2), 2.41 (Me); ¹³C NMR (100 MHz, CDCl₃) δ 162.8 (C2), 150.5 (C4), 140.7 (C6), 136.4 (C3'), 122.3 (C2'), 102.8 (C5), 90.2 (C1'), 87.2 (C4'), 63.6 (5'-CH₂), 15.1 (Me). LRMS (DCI, NH₃-isobutane): m/z 257 [M+H]⁺. HRMS calculated for C₁₀H₁₂N₂O₄NaS: [M+Na]⁺ 279.0416; found: 279.0416.

¹H and ¹³C NMR SPECTRA