Direct Synthesis of Sulfonamides and Activated Sulfonate Esters from Sulfonic Acids

Stephen Caddick, Jonathan D. Wilden and Duncan B. Judd*

Supporting Information

N-Allyl-4-methyl-benzenesulfonamide (10a)

To a solution of triphenylphosphine oxide (1.11 g, 4.0 mmol) in degassed DCM (20 mL) was added trifluoromethanesulfonic anhydride (0.35 mL, 0.56 g, 1.8 mmol) and the solution allowed to stir for 15 minutes. The pyridine salt of *p*-toluenesulfonic acid (508 mg, 1.8 mmol) was added and the solution allowed to stir for 30 minutes. A mixture of triethylamine (0.5 mL, 3.6 mmol) and allylamine (0.5 mL) in DCM (10 mL) was added dropwise at 0 °C over a period of 5 minutes. The resulting pale yellow solution was allowed to warm to RT and diluted with DCM (20 mL). The reaction mixture was washed with 2M HCl (3 x 30 mL), saturated sodium bicarbonate solution (3 x 30 mL) and water (30 mL). The organic fraction was dried (MgSO₄), filtered, the filtrate concentrated *in vacuo* and the crude residue purified by flash column chromatography (20-40% ether / petroleum ether 40-60 °C) to yield the product as a pale yellow solid (586 mg, 1.73 mmol, 96%). Recrystallisation from DCM / hexane gave the product as white crystals. Data in agreement with literature values.¹

MP	63-66 °C (DCM / hexane) Lit. 63-65 (aq. MeOH).
¹H NMR	δ _H (300 MHz, CDCl ₃) 7.69 (2H, d, <i>J</i> = 8.3 Hz, 2 x ArH), 7.23 (2H, d, <i>J</i> = 8.2 Hz, 2 x ArH), 5.63 (1H, ddt, <i>J</i> = 17.1, 10.2, 5.8 Hz, CH ₂ CH=CH ₂), 5.08 (1H, app. dq, <i>J</i> = 17.1, 1.4 Hz, =CHH), 4.99 (1H, app. dq, <i>J</i> = 10.2, 1.3 Hz, =CHH), 4.96 (1H, obs. NH), 3.48 (2H, app. tt, <i>J</i> = 6.0, 1.4 Hz, CH ₂ CH=CH ₂), 2.34 (3H, s, ArCH ₃).
¹³C NMR	δ _C (75 MHz, CDCl ₃) 143.9 (s, C (Ar)), 137.3 (s, C (Ar)), 133.4 (d, CH=CH ₂), 130.1 (d, 2 x CH (Ar)), 127.5 (d, 2 x CH (Ar)), 118.0 (t, CH=CH ₂), 46.1 (t, CH ₂ NH), 21.9 (q, ArCH ₃).
FTIR	(CH ₂ Cl ₂ Solution, cm ⁻¹) 3287 br. s, 3056 s, 2987 m, 1647 w, 1598 m, 1495 w, 1495 m, 1422 s, 1328 s, 1265 s, 1161 s, 1094 s.

N-Allyl-ethanesulfonamide (10b)

Prepared as for **10a** to give the product as a colourless oil (242 mg, 1.62 mmol, 90%). Data consistent with literature values.²

¹H NMR	δ _H (300 MHz, CDCl ₃) 5.79 (1H, ddt, <i>J</i> = 17.1, 10.2, 5.8 Hz, CH ₂ CH=CH ₂), 5.22 (1H, app. dq, <i>J</i> = 17.1, 1.5 Hz, =CHH), 5.11 (1H, app. dq, <i>J</i> = 10.3, 1.3 Hz, =CHH), 5.10 (1H, obs. NH), 3.65 (2H, app. tt, <i>J</i> = 6.0, 1.5 Hz, CH ₂ CH=CH ₂), 2.97 (2H, q, <i>J</i> = 7.4 Hz, CH ₃ CH ₂ SO ₂), 1.28 (3H, t, <i>J</i> = 7.4 Hz, CH ₃ CH ₂ SO ₂).
¹³C NMR	δ _C (75 MHz, CDCl ₃) 134.2 (d, CH=CH ₂), 117.7 (t, CH=CH ₂), 45.9 (t, CH ₂ NH), 8.5 (q, CH ₃).
FTIR	(CH ₂ Cl ₂ Solution, cm ⁻¹) 3571 br. m, 3291 br. m, 1642 w, 1430 m, 1317 s, 1282 m, 1142 s, 1069 m.

Pyridine-3-sulfonic acid allylamide

Prepared as for **10a** to give the product as an off-white solid (289 mg, 1.46 mmol, 81%).

MP	102-104 °C (DCM / hexane)
¹H NMR	δ _H (300 MHz, CDCl ₃) 9.07 (1H, d, <i>J</i> = 2.1 Hz, ArH), 8.80 (1H, dd, <i>J</i> = 4.8, 1.4 Hz, ArH), 8.17 (1H, app. dt, <i>J</i> = 8.1, 1.9 Hz, ArH), 7.48 (1H, dd, <i>J</i> = 8.1, 4.9 Hz, ArH), 5.72 (1H, ddt, <i>J</i> = 17.1, 10.2, 5.8 Hz, CH ₂ CH=CH ₂), 5.39 (1H, br. t, <i>J</i> = 6.0 Hz, NH), 5.18 (1H, app. dq, <i>J</i> = 17.1, 1.4 Hz, CH=CHH), 5.11 (1H, app. dq, <i>J</i> = 10.2, 1.2 Hz, CH=CHH), 3.67 (2H, app tt, <i>J</i> = 6.0, 1.5 Hz, CH ₂ CH=CH ₂).
¹³C NMR	δ _C (75 MHz, CDCl ₃) 153.5 (d, CH (Ar)), 148.3 (d, CH (Ar)), 137.4 (s, C (Ar)), 135.3 (d, CH=), 133.0 (d, CH (Ar)), 124.2 (d, CH (Ar)), 118.6 (t, =CH ₂), 46.1 (t, CH ₂ NH).
FTIR	(CH ₂ Cl ₂ Solution, cm ⁻¹) 3056 m, 2855 m, 2715 m, 1645 w, 1583 m, 1495 w, 1417 m, 1324 s, 1265 s, 1165 s, 1108 m.
HRMS	(ES +ve) [MH] ⁺ , C ₈ H ₁₀ N ₂ O ₂ S Requires 199.0536. Found 199.0521.

N-Allyl methanesulfonamide

Prepared as for **10a** to give the product as a colourless oil (207 mg, 1.53 mmol, 85%). Data consistent with literature values.³

¹H NMR	δ _H (300 MHz, CDCl ₃) 5.86 (1H, ddt, <i>J</i> = 17.0, 10.2, 5.7 Hz, CH ₂ CH=CH ₂), 5.26 (1H, app. dq, <i>J</i> = 17.0, 1.5 Hz, CH=CHH), 5.19 (1H, obs., NH), 5.15 (1H, app. dq, <i>J</i> = 10.2, 1.3 Hz, CH=CHH), 3.70 (2H, app. tt, <i>J</i> = 5.9, 1.5 Hz, CH ₂ NH), 2.92 (3H, s, CH ₃ SO ₂).
¹³C NMR	δ _C (75 MHz, CDCl ₃) 134.0 (d, CH=CH ₂), 118.0 (t, CH=CH ₂), 45.9 (t, CH ₂ NH), 41.1 (q, CH ₃ SO ₂).
FTIR	(CH ₂ Cl ₂ Solution, cm ⁻¹) 3584 br. m, 3292 br. s, 3019 w, 2934 w, 1646 w, 1432 m, 1413 m, 1314 vs, 1150 vs, 1066 m.

Thiophene-2-sulfonic acid allylamide

Prepared as for **10a** to give the product as an off-white solid (300 mg, 1.48 mmol, 82%). Data consistent with literature values.⁴

MP	53-55 °C (DCM / hexane) Lit. 54-56 °C (H ₂ O).
¹H NMR	δ _H (300 MHz, CDCl ₃) 7.65-7.58 (2H, m, 2 x ArH), 7.10 (1H, dd, <i>J</i> = 5.0, 3.8 Hz, ArH), 5.75 (1H, ddt, <i>J</i> = 17.1, 10.3, 5.7 Hz, CH ₂ CH=CH ₂), 5.20 (1H, app. dq, <i>J</i> = 17.1, 1.4 Hz, CH=CHH), 5.12 (1H, app. dq, <i>J</i> = 10.3, 1.4 Hz, CH=CHH), 4.91 (1H, br. t, <i>J</i> = 5.8 Hz, NH), 3.68 (2H, app. tt, <i>J</i> = 5.9, 1.4 Hz, CH ₂ NH).
¹³C NMR	δ _C (75 MHz, CDCl ₃) 141.2 (s, C (Ar)), 133.0 (d, CH (Ar)), 132.7 (d, CH=CH ₂), 132.4 (d, CH (Ar)), 127.9 (d, CH (Ar)), 118.3 (t, CH=CH ₂), 46.4 (t, CH ₂ NH).
FTIR	(CH ₂ Cl ₂ Solution, cm ⁻¹) 3290 br. s, 3094 m, 3058 m, 2987 w, 1646 w, 1508 w, 1406 s, 1336 br. s, 1266 s, 1227 s, 1159 s, 1067 s, 1019 s.

5-Dimethylamino-naphthalene-1-sulfonic acid allylamide

Prepared as for **10a** to give the product as a green solid (250 mg, 0.86 mmol, 48%).

MP	84-86 °C (DCM / hexane)
¹H NMR	δ _H (300 MHz, CDCl ₃) 8.55 (1H, d, <i>J</i> = 8.5 Hz, ArH), 8.31 (1H, d, <i>J</i> = 8.7 Hz, ArH), 8.26 (1H, dd, <i>J</i> = 7.3, 1.2 Hz, ArH), 7.61-7.48 (2H, m, 2 x ArH), 7.19 (1H, d, <i>J</i> = 7.7 Hz, ArH), 5.63 (1H, ddt, <i>J</i> = 17.1, 10.2, 5.8 Hz, CH ₂ CH=CH ₂), 5.09 (1H, app. dq, <i>J</i> = 17.1, 1.5 Hz, CH=CHH), 5.00 (1H, app. dq, <i>J</i> = 10.2, 1.4 Hz, CH=CHH), 4.87 (1H, br. t, <i>J</i> = 6.0 Hz, NH), 3.54 (2H, app. tt, <i>J</i> = 6.0, 1.4 Hz, CH ₂ NH), 2.89 (6H, s, N(CH ₃) ₂).
¹³C NMR	δ _C (75 MHz, CDCl ₃) 152.4 (s, C(Ar)), 135.0 (s, C(Ar)), 133.4 (d, CH=CH ₂), 131.0 (d, CH (Ar)), 130.3 (s, C(Ar)), 130.1 (d, CH (Ar)), 130.0 (s, C(Ar)), 128.8 (d, CH (Ar)), 123.6 (d, CH (Ar)), 119.1 (d, CH (Ar)), 118.0 (t, CH=CH ₂), 115.6 (d, CH (Ar)), 46.2 (t, CH ₂ NH), 45.8 (q, 2 x CH ₃).
FTIR	(CH ₂ Cl ₂ Solution, cm ⁻¹) 3584 br. w, 3299 s, 2943 m, 2833 m, 2789 m, 1646 w, 1612 w, 1588 m, 1574 m, 1504 w, 1455 m, 1409 m, 1318 s, 1231 w, 1161 s, 1144 s.
HRMS	(ES +ve) [M] ⁺ C ₁₅ H ₁₉ N ₂ O ₂ S Requires 291.1162. Found 291.1170.

3-Nitrobenzenesulfonic acid allylamide

Prepared as for **10a** to give the product as a white solid (401 mg, 1.66 mmol, 92%). Data consistent with literature values.⁵

MP	71-74 °C (DCM / hexane) Lit.
¹H NMR	δ _H (300 MHz, CDCl ₃) 8.71 (1H, t, <i>J</i> = 2.0 Hz, ArH), 8.44 (1H, ddd, <i>J</i> = 8.2, 2.2, 1.0 Hz, ArH), 8.22 (1H, br. dt, <i>J</i> = 7.9, 1.4 Hz, ArH), 7.76 (1H, t, <i>J</i> = 8.0 Hz, ArH), 5.71 (1H, ddt, <i>J</i> = 17.0, 10.2, 5.8 Hz, CH ₂ CH=CH ₂), 5.18 (1H, app. dq, <i>J</i> = 17.1, 1.4 Hz, CH=CHH), 5.12 (1H, app. dq, <i>J</i> = 10.2, 1.3 Hz, CH=CHH), 5.11 (1H, obs, NH), 3.69 (2H, app. tt, <i>J</i> = 6.0, 1.5 Hz, CH ₂ NH).
¹³C NMR	δ _C (75 MHz, CDCl ₃) 148.7 (s, C(Ar)), 142.8 (s, C(Ar)), 133.1 (d, CH=CH ₂), 132.8 (d, CH (Ar)), 131.0 (d, CH (Ar)), 127.6 (d, CH (Ar)), 122.7 (d, CH (Ar)), 118.7 (t, CH=CH ₂), 46.2 (t, CH ₂ NH).
FTIR	(CH ₂ Cl ₂ Solution, cm ⁻¹) 3379 br. m, 3055 m, 2987 m, 1646 w, 1607 w, 1536 s, 1422 m, 1353 s, 1265 s, 1171 m, 1127 w, 1071 w.

N-Allyl-4-methoxy-benzenesulfonamide

Prepared as for **10a** to give the product as a white solid (360 mg, 1.58 mmol, 88%). Data consistent with literature values.⁶

MP	46-49 °C (Ether / hexane) Lit. 45-47 (MeOH).
¹H NMR	δ_{H} (300 MHz, CDCl ₃) 7.81 (2H, d, J = 8.9 Hz, 2 x ArH), 6.98 (2H, d, J = 8.9 Hz, 2 x ArH), 5.71 (1H, ddt, J = 17.0, 10.2, 5.8 Hz, CH ₂ CH=CH ₂), 5.16 (1H, app. dq, J = 17.1, 1.4 Hz, CH ₂ CH=CHH), 5.08 (1H, app. dq, J = 10.2, 1.3 Hz, CH ₂ CH=CHH), 4.75 (1H, br. t, J = 6.0 Hz, NH), 3.87 (3H, s, OCH ₃), 3.57 (2H, br. t, J = 6.0 Hz, CH ₂ NH).
¹³C NMR	δ_{C} (75 MHz, CDCl ₃) 163.3 (s, C(Ar)), 133.4 (d, CH=CH ₂), 131.8 (s, C(Ar)), 129.7 (d, 2 x CH (Ar)), 118.1 (t, CH=CH ₂), 114.7 (d, CH=CH ₂), 56.0 (q, OCH ₃), 46.1 (t, CH ₂ NH).
FTIR	(CH ₂ Cl ₂ Solution, cm ⁻¹) 3284 br. m, 3055 m, 2987 w, 2842 w, 2305 w, 1646 w, 1598 s, 1579 s, 1498 s, 1421 s, 1329 s, 1264 vs, 1157 vs, 1096 m.

Pentafluorophenyl Tosylate (11a)

To a solution of triphenylphosphine (1.1 g, 4.0 mmol, 2.1 eq.) in freshly distilled DCM (20 mL) at 0 °C under nitrogen was added trifluoromethanesulfonic anhydride (508 mg, 0.30 mL, 1.8 mmol, 1.0 eq.) and the resulting clear solution was allowed to stir for 20 minutes. Pyridinium *p*-toluenesulfonate was then added in a single portion and the mixture allowed to stir for 20 minutes or until all of the sulfonic acid salt had dissolved. A pre-mixed solution of pentafluorophenol (368 mg, 2.0 mmol, 1.1 eq) and triethylamine (0.47 mL, 2.0 mmol) in DCM (5 mL) was then added *via* syringe over a period of 10 minutes. The reaction was allowed to warm to RT then diluted with DCM (*ca.* 30 mL) and washed with 2M sodium carbonate solution (3 x 30 mL), 2M HCl (3 x 30 mL) water (2 x 30 mL), separated, dried (MgSO₄) and filtered. The filtrate was concentrated *in vacuo* and the crude product was purified by flash column chromatography (20% ether / petroleum ether 40-60 °C) to yield the product as a white solid (584 mg, 1.73 mmol, 96%). Recrystallisation from DCM-Hexane gave white needles. Data consistent with literature values.⁷

MP	66-69 °C (DCM / hexane). Lit. 64-65 °C (EtOH).
¹H NMR	δ_{H} (300 MHz, CDCl ₃) 7.86 (2H, d, J = 8.3 Hz, 2 x ArH), 7.41 (2H, d, J = 8.3 Hz, 2 x ArH), 2.51 (3H, s, ArCH ₃).
¹³C NMR	δ_{C} (75 MHz, CDCl ₃) 147.2 (s, C (Ar)), 132.1 (s, C (Ar)), 130.6 (d, 2 x CH (Ar)), 129.0 (d, 2 x CH (Ar)), 22.2 (q, CH ₃).
FTIR	(Thin film, cm ⁻¹) 3055 w, 2927 m, 1653 w, 1578 w, 1520 vs, 1457 m, 1410 m, 1265 s, 1199 m, 1057 s.

Pentafluorophenyl ethanesulfonate (11b)

Prepared as for **11a** to give the product as a colourless oil (428 mg, 1.55 mmol, 86%).

¹H NMR	δ_{H} (300 MHz, CDCl ₃) 3.50 (2H, q, J = 7.4 Hz, CH ₂ SO ₃), 1.63 (3H, t, J = 7.4 Hz, CH ₃ CH ₂)
¹³C NMR	δ_{C} (75 MHz, CDCl ₃) 47.9 (t, CH ₃ CH ₂ SO ₃), 8.6 (q, CH ₃ CH ₂ SO ₃).
FT-IR	(Thin film, cm ⁻¹) 2981 m, 2872 m, 1651 w, 1520 vs, 1472 m, 1460 m, 1318 w, 1293 w, 1239 w, 1182 s, 1141 s, 1116 m, 1022 s, 997 vs.

Pyridine-3-sulfonic acid pentafluorophenyl ester (11c)

Prepared as for **11a** to give the product as an off white solid (474 mg, 1.46 mmol, 81%).

MP	54-56 °C (DCM / hexane).
¹H NMR	δ _H (300 MHz, CDCl ₃) 9.17 (1H, br. d, <i>J</i> = 1.9 Hz, ArH), 8.99 (1H, dd, <i>J</i> = 4.8, 1.5 Hz, ArH), 8.29 (1H, ddd, <i>J</i> = 8.1, 2.3, 1.7 Hz, ArH), 7.60 (1H, ddd, <i>J</i> = 8.2, 4.8, 0.9 Hz, ArH).
¹³C NMR	δ _C (75 MHz, CDCl ₃) 156.0 (d, CH (Ar)), 149.4 (d, CH (Ar)), 136.5 (d, CH (Ar)), 132.4 (s, C (Ar)), 124.5 (d, CH (Ar)).
FT-IR	(CH ₂ Cl ₂ solution, cm ⁻¹) 3677 w, 3063 w, 1575 m, 1520 vs, 1470 m, 1421 s, 1399 s, 1329 w, 1202 s, 1107 w, 1022 s, 999 s.
HRMS	(ES +ve) [MH] ⁺ , C ₁₁ H ₅ O ₃ NF ₅ S Requires 325.9905. Found 325.9914.

Pentafluorophenyl methanesulfonate (11d)

Prepared as for **11a** to give the product as a colourless oil (364 mg, 1.39 mmol, 77%). Data consistent with literature values.⁸

¹H NMR	δ _H (300 MHz, CDCl ₃) 3.50 (3H, s, CH ₃ SO ₃).
¹³C NMR	δ _C (75 MHz, CDCl ₃) 47.9 (q, CH ₃ SO ₃).
FT-IR	(Thin film, cm ⁻¹) 2981 m, 2872 m, 1651 w, 1520 vs, 1472 m, 1460 m, 1318 w, 1293 w, 1239 w, 1182 s, 1141 s, 1116 m, 1022 s, 997 vs.

Thiophene-2-sulfonic acid pentafluorophenyl ester (11e)

Prepared as for **11a** to give the product as a white solid (452 mg, 1.37 mmol, 76%).

MP	84-86 °C (DCM / hexane)
¹H NMR	δ _H (300 MHz, CDCl ₃) 7.92 (1H, dd, <i>J</i> = 5.0, 1.4 Hz, ArH), 7.86 (1H, dd, <i>J</i> = 4.0, 1.3 Hz, ArH), 7.28 (1H, dd, <i>J</i> = 5.0, 4.0 Hz, ArH).
¹³C NMR	δ _C (75 MHz, CDCl ₃) 136.8 (d, CH (Ar)), 136.7 (d, CH (Ar)), 133.9 (s, C (Ar)), 128.5 (d, CH (Ar)).
FTIR	(CH ₂ Cl ₂ Solution, cm ⁻¹) 3100 w, 3056 m, 2987 m, 1651 w, 1520 vs, 1470 w, 1400 s, 1194 s, 1096 m, 1022 s.
LRMS	(EI +ve) 330 (M ⁺ , 26%), 147 ([M-OC ₆ F ₅] ⁺ , 100%).
HRMS	(ES +ve) [M+Na] ⁺ C ₁₀ H ₃ O ₃ S ₂ F ₅ Na Requires 352.9336. Found 352.9325.

5-Dimethylamino-naphthalene-1-sulfonic acid pentafluorophenyl ester (11f)

Prepared as for **11a** to give the product as a green solid (391 mg, 0.94 mmol, 52%).

¹H NMR	δ _H (300 MHz, CDCl ₃) 8.72 (1H, d, <i>J</i> = 8.7 Hz, ArH), 8.42 (1H, d, <i>J</i> = 8.7 Hz, ArH), 8.21 (1H, dd, <i>J</i> = 7.3, 1.3 Hz, ArH), 7.68 (1H, dd, <i>J</i> = 8.7, 7.7 Hz, ArH), 7.55 (1H, dd, <i>J</i> = 8.5, 7.4 Hz, ArH), 7.27 (1H, d, <i>J</i> = 7.3 Hz, ArH), 2.81 (6H, s, 2 x CH ₃).
¹³C NMR	δ _C (75 MHz, CDCl ₃) 152.3 (s, C (Ar)), 133.5 (d, CH (Ar)), 131.3 (d, CH (Ar)), 131.1 (s, C (Ar)), 130.6 (s, C (Ar)), 130.2 (s, C (Ar)), 129.7 (d, CH (Ar)), 123.3 (d, CH (Ar)), 119.5 (d, CH (Ar)), 116.3 (d, CH (Ar)), 45.8 (q, 2 x CH ₃).
FTIR	(Thin film, cm ⁻¹) 3433 br. w, 3056 w, 2948 m, 2793 w, 1650 w, 1614 w, 1587 m, 1571 m, 1520 vs, 1465 m, 1412 m, 1386 s, 1265 s, 1205 m, 1186 s, 1142 m.

Pentafluorophenyl 3-nitrobenzenesulfonate

Prepared as for **11a** to give the product as a white solid (617 mg, 1.69 mmol, 94%).

MP	64-67 °C (Ether / hexane)
¹H NMR	δ _H (300 MHz, CDCl ₃) 8.85 (1H, t, <i>J</i> = 2.0 Hz, ArH), 8.65 (1H, ddd, <i>J</i> = 8.2, 2.2, 1.0 Hz, ArH), 8.34 (1H, app. dt, <i>J</i> = 8.1, 1.4 Hz, ArH), 7.91 (1H, t, <i>J</i> = 8.1 Hz, ArH).
¹³C NMR	δ _C (75 MHz, CDCl ₃) 148.8 (s, C (Ar)), 137.4 (s, C (Ar)), 134.1 (d, CH (Ar)), 131.5 (d, CH (Ar)), 130.0 (d, CH (Ar)), 124.2 (d, CH (Ar)).
FTIR	(Thin film, cm ⁻¹) 3084 w, 2882 w, 1608 w, 1540 m, 1520 vs, 1470 w, 1404 m, 1353 s, 1200 s, 1022 w, 1000 vs.
LRMS	(EI +ve) 369 (M ⁺ , 59%), 186 ([M-C ₆ F ₅] ⁺ , 68%), 76 (100%).
HRMS	(ES +ve) [M+Na] ⁺ C ₁₂ H ₄ NO ₅ SF ₅ Na Requires 391.9628. Found 391.9654.

Pentafluorophenyl 4-methoxybenzenesulfonate

Prepared as for **11a** to give the product as a white solid (471 mg, 1.33 mmol, 74%).

MP	54-56 °C (Ether / hexane)
¹H NMR	δ _H (300 MHz, CDCl ₃) 7.90 (2H, d, <i>J</i> = 9.0 Hz, 2 x ArH), 7.06 (2H, d, <i>J</i> = 9.0 Hz, 2 x ArH), 3.93 (3H, s, OCH ₃).
¹³C NMR	δ _C (75 MHz, CDCl ₃) 165.4 (s, C (Ar)), 131.3 (d, 2 x CH (Ar)), 126.0 (s, C (Ar)), 115.1 (d, 2 x CH (Ar)), 56.3 (q, OCH ₃)
FTIR	(CH ₂ Cl ₂ Solution, cm ⁻¹) 3060 w, 2949 w, 2846 w, 1650 w, 1596 s, 1578 s, 1520 s, 1467 m, 1392 s, 1266 s, 1198 s, 1172 s, 1093 s.
LRMS	(EI +ve) 354 (M ⁺ , 18%), 171 ([M-OC ₆ F ₅] ⁺ , 100%).
HRMS	(ES +ve) [M+Na] ⁺ C ₁₃ H ₇ O ₄ SF ₅ Na Requires 376.9877. Found 376.9878

1-(3-Nitro-benzenesulfonyl)-piperidine

Prepared as for **10a** to give the product as an off-white solid (387 mg, 1.51 mmol, 84%). Data consistent with literature values.⁹

MP	107-110 °C (DCM / hexane). Lit. 108-110 (MeOH).
¹H NMR	δ _H (300 MHz, CDCl ₃) 8.58 (1H, t, <i>J</i> = 1.7 Hz, ArH), 8.45 (1H, ddd, <i>J</i> = 8.2, 2.3, 1.0 Hz, ArH), 8.09 (1H, ddd, <i>J</i> = 7.9, 1.7, 1.0 Hz, ArH), 7.78 (1H, dt, <i>J</i> = 8.2, 7.8 Hz, ArH), 3.05 (4H, br. t, <i>J</i> = 5.5 Hz, N(CH ₂) ₂), 1.73-1.60 (4H, m, (CH ₂) ₂), 1.51-1.40 (2H, m, CH ₂).
¹³C NMR	δ _C (75 MHz, CDCl ₃) 148.7 (s, C (Ar)), 139.3 (s, C (Ar)), 133.5 (d, CH (Ar)), 130.8 (d, CH (Ar)), 127.5 (d, CH (Ar)), 123.0 (d, CH (Ar)), 47.3 (t, 2 x CH ₂), 25.5 (t, 2 x CH ₂), 23.7 (t, CH ₂).
FTIR	(CH ₂ Cl ₂ Solution, cm ⁻¹) 3056 w, 2948 m, 2857 w, 1780 m, 1606 w, 1536 s, 1467 w, 1353 s, 1178 s, 1127 m.

2-(3-Nitro-benzenesulfonylamino)-benzoic acid ethyl ester

Prepared as for **10a** to give the product as an off-white solid (592 mg, 1.69 mmol, 94%).

MP	76-78 °C (CHCl ₃ / hexane)
¹H NMR	δ _H (300 MHz, CDCl ₃) 10.91 (1H, br. s, NH), 8.65 (1H, t, <i>J</i> = 1.9 Hz, ArH), 8.36 (1H, ddd, <i>J</i> = 8.3, 2.3, 1.1 Hz, ArH), 8.16 (1H, app. dt, <i>J</i> = 8.0, 1.3 Hz, ArH), 7.94 (1H, dd <i>J</i> = 8.0, 1.6 Hz, ArH), 7.73-7.63 (2H, m, 2 x ArH), 7.50 (1H, app. dt, <i>J</i> = 8.0, 1.5 Hz, ArH), 7.09 (1H, br. m, ArH), 4.32 (2H, q, <i>J</i> = 7.2 Hz, CO ₂ CH ₂ CH ₃), 1.35 (3H, t, <i>J</i> = 7.2 Hz, CO ₂ CH ₂ CH ₃).
¹³C NMR	δ _C (75 MHz, CDCl ₃) 168.3 (s, C=O), 148.5 (s, C (Ar)), 141.8 (s, C (Ar)), 139.9 (s, C (Ar)), 135.1 (d, CH (Ar)), 133.1 (d, CH (Ar)), 131.8 (d, CH (Ar)), 130.9 (d, CH (Ar)), 127.9 (d, CH (Ar)), 124.3 (d, CH (Ar)), 122.8 (d, CH (Ar)), 119.8 (d, CH (Ar)), 117.0 (s, C (Ar)), 62.4 (t, CO ₂ CH ₂ CH ₃), 14.5 (q, CO ₂ CH ₂ CH ₃).
FT-IR	(CH ₂ Cl ₂ solution, cm ⁻¹) 3151 br. w, 3090 w, 2985 w, 1685 s, 1605 m, 1585 m, 1537 s, 1492 s, 1410 m, 1353 s, 1310 m, 1255 s, 1213 w, 1178 s, 1087 s.
HRMS	(ES ⁺) [M+Na] ⁺ Requires 373.0465. Found 373.0459.

N-(4-Methyl-benzyl)-3-nitro-benzenesulfonamide

Prepared as for **10a** to give the product as a white solid (511 mg, 1.67 mmol, 93%).

MP	125-127 °C (CHCl ₃ / hexane)
¹H NMR	δ _H (300 MHz, CDCl ₃) 8.57 (1H, t, <i>J</i> = 1.9 Hz, ArH), 8.38 (1H, ddd, <i>J</i> = 8.3, 2.2, 1.0 Hz, ArH), 8.13 (1H, br. dt, <i>J</i> = 8.1, 1.4 Hz, ArH), 7.68 (1H, t, <i>J</i> = 8.0 Hz, ArH), 7.04 (4H, app. s, 4 x ArH), 5.09 (1H, t, <i>J</i> = 6.1 Hz, NH), 4.21 (2H, d, <i>J</i> = 6.0 Hz, CH ₂ NH), 2.28 (3H, s, ArCH ₃).
¹³C NMR	δ _C (75 MHz, CDCl ₃) 148.5 (s, C (Ar)), 142.9 (s, C (Ar)), 138.4 (s, C (Ar)), 132.9 (d, CH (Ar)), 132.7 (s, C (Ar)), 130.7 (d, CH (Ar)), 129.8 (d, 2 x CH (Ar)), 128.3 (d, 2 x CH (Ar)), 127.3 (d, CH (Ar)), 122.7 (d, CH (Ar)), 47.6 (t, CH ₂ NH), 21.4 (q, ArCH ₃).
FT-IR	(Nujol, cm ⁻¹) 3246 s, 2923 s, 1608 m, 1543 m, 1515 w, 1463 s, 1353 s, 1326 s, 1270 m, 1159 s, 1124 s, 1050 s.

(R)-2-(3-Nitro-benzenesulfonylamino)-3-phenyl-propionic acid ethyl ester

Prepared as for **10a** to give the product as a white solid (416 mg, 1.10 mmol, 61%).

MP	93-95 °C (CHCl ₃ / hexane)
¹H NMR	δ _H (300 MHz, CDCl ₃) 8.48 (1H, t, <i>J</i> = 1.9 Hz, ArH), 8.35 (1H, ddd, <i>J</i> = 8.3, 2.3, 1.0 Hz, ArH), 8.02 (1H, br. dt, <i>J</i> = 8.2, 1.3 Hz, ArH), 7.62 (1H, t, <i>J</i> = 8.0 Hz, ArH), 7.23-7.14 (3H, m, 3 x ArH), 7.09-7.03 (2H, m, 2 x ArH), 5.55 (2H, d, <i>J</i> = 9.4 Hz, NH), 4.25 (1H, ddd, <i>J</i> = 9.4, 7.5, 5.5 Hz, NHCHCO ₂ Et), 4.04 (2H, q, <i>J</i> = 7.2 Hz, CO ₂ CH ₂ CH ₃), 3.11 (1H, dd, <i>J</i> = 13.9, 5.5 Hz, NHCHCHH), 2.96 (1H, dd, <i>J</i> = 13.8, 7.5 Hz, NHCHCHH), 1.16 (3H, t, <i>J</i> = 7.2 Hz, CO ₂ CH ₂ CH ₃).
¹³C NMR	δ _C (75 MHz, CDCl ₃) 170.9 (s, C=O), 148.2 (s, C (Ar)), 142.1 (s, C (Ar)), 135.0 (s, C (Ar)), 132.6 (d, CH (Ar)), 130.4 (d, CH (Ar)), 129.4 (d, 2 x CH (Ar)), 128.7 (d, 2 x CH (Ar)), 127.4 (d, CH (Ar)), 122.4 (d, CH (Ar)), 62.2 (t, CO ₂ CH ₂ CH ₃), 57.3 (d, CHNH), 39.3 (t, NHCHCH ₂), 14.0 (q, CO ₂ CH ₂ CH ₃).
FTIR	(CH ₂ Cl ₂ solution, cm ⁻¹) 3340 w, 3265 br. w, 3055 m, 2986 m, 2305 w, 1734 s, 1607 w, 1536 s, 1512 s, 1394 m, 1358 s, 1265 s, 1172 s, 1127 w, 1029 w.

[4-(3-Nitro-benzenesulfonylamino)-phenyl]-acetic acid ethyl ester

Prepared as for **10a** to give the product as a white solid (575 mg, 1.58 mmol, 88%).

MP	99-102 °C (DCM / hexane)
¹H NMR	δ _H (300 MHz, CDCl ₃) 8.55 (1H, t, <i>J</i> = 1.9 Hz, ArH), 8.29 (1H, ddd, <i>J</i> = 8.2, 2.1, 1.1 Hz, ArH), 7.94 (1H, br. dt, <i>J</i> = 8.1, 1.3 Hz, ArH), 7.56 (1H, t, <i>J</i> = 8.0 Hz, ArH), 7.47 (1H, s, ArH), 7.05 (2H, d, <i>J</i> = 8.5 Hz, 2 x ArH), 6.88 (2H, d, <i>J</i> = 8.5 Hz, 2 x ArH), 4.11 (2H, q, <i>J</i> = 7.1 Hz, CO ₂ CH ₂ CH ₃), 3.49 (2H, s, ArCH ₂ CO ₂ Et), 1.19 (3H, t, <i>J</i> = 7.1 Hz, CO ₂ CH ₂ CH ₃).
¹³C NMR	δ _C (75 MHz, CDCl ₃) 172.2 (s, C=O), 148.3 (s, C (Ar)), 141.2 (s, C (Ar)), 134.6 (s, C (Ar)), 132.9 (d, CH (Ar)), 132.1 (s, C (Ar)), 130.6 (d, 2 x CH (Ar)), 130.5 (d, CH (Ar)), 127.5 (d, CH (Ar)), 122.7 (d, 2 x CH (Ar)), 122.5 (d, CH (Ar)), 61.4 (t, CO ₂ CH ₂ CH ₃), 40.6 t, ArCH ₂ CO ₂ Et), 14.2 (q, CO ₂ CH ₂ CH ₃).
FTIR	(CH ₂ Cl ₂ solution, cm ⁻¹) 3341 w, 3250 br. w, 3055 m, 2986 m, 2305 w, 1731 s, 1607 w, 1536 s, 1512 m, 1394 m, 1353 s, 1265 s, 1172 s, 1127 w, 1029 w.
HRMS	(ES +ve) [M+Na] ⁺ C ₁₆ H ₁₆ N ₂ SO ₆ Na Requires 387.0621. Found 387.0625.

References

1. Wedekind, E.; *Chem. Ber.* **1909**, *42*, 3939.
2. Whitehead, C. W.; Traverso, J. J.; *J. Am. Chem. Soc.* **1958**, *80*, 2182.
3. Tehrani, K. A.; NguyenVan, T.; Karikomi, M.; Rottiers, M.; De Kimpe, N.; *Tetrahedron* **2002**, *58*, 7145.
4. Foye *et al.* *J. Am. Pharm. Assoc.* **1952**, *41*, 273.
5. Patent: Sandoz Ltd., GB 908109, **1962**. *Chem. Abs.* **1963**, *58*, 11498.
6. Dang, H-S.; Roberts, B. P.; *J. Chem. Soc. Perkin Trans. I*, **1996**, *13*, 1493.
7. Forbes *et al.* *J. Chem. Soc.* **1959**, 2019.
8. Perec, V.; Bae, J-Y.; Zhao, M.; Hill, D. H.; *J. Org. Chem.* **1995**, *60*, 176.
9. Bell, C.; *J. Chem. Soc.* **1933**, 1290.