Divalent Activation in Temporary Phosphate Tethers: Highly Selective Cuprate Displacement Reactions

Alan Whitehead, James P. McParland, and Paul R. Hanson*

Department of Chemistry, University of Kansas, 1251 Wescoe Hall Drive, Lawrence, KS 66045-7582

phanson@ku.edu

Table of Contents

General Experimental Methods:	S 2
Experimental Section and Characterization Data	S 3- S 15
NMR Spectra for New Compounds	S 16- S 43

General Experimental Methods.

All air and moisture sensitive reactions were carried out in flame- or oven-dried glassware under argon atmosphere using standard gastight syringes, cannulae, and septa. Stirring was achieved with oven-dried magnetic stir bars. Et₂O, toluene, THF and CH₂Cl₂ were purified by passage through the Solv-Tek purification system employing activated Al₂O₃ (Grubbs, R.H.; Rosen, R.K.; Timmers, F.J.; *Organometallics* **1996**, *15*, 1518-1520). Et₃N was purified by passage over basic alumina and stored over KOH. Butyl Lithium was purchased from Aldrich and titrated prior to use. Dialkyl zinc and alkyl zinc halide reagents were purchased from Aldrich and were used at the reported molarities. Grubb's second-generation olefin metathesis catalyst was acquired from Materia and used without further purification. Flash column chromatography was performed with Merck silica gel (EM-9385-9, 230-400 mesh). Thin layer chromatography was performed on silica gel 60F₂₅₄ plates (EM-5717, Merck). Deuterated solvents were purchased from Cambridge Isotope laboratories. ¹H, ¹³C and ³¹P NMR spectra were recorded on a Bruker DRX-400 spectrometer operating at 400 MHz, 100 MHz, and 162 MHz respectively or a Bruker Avance operating at 500 MHz and 125 MHz respectively. High-resolution mass spectrometry (HRMS) and FAB spectra were obtained on a VG Instrument ZAB double-focusing mass spectrometer.

Experimental procedures and spectral data for compounds 1, 4-6, 8a, 9a-9h, 10a, 11, 12, 14, 15 and *syn*- and *anti*-ethyl-substituted acetonides.

(*S,S*)-acyclic diene phosphate triester: A solution of allylic alcohol **3** (2.05 g, 11.54 mmol) in THF (58 mL) was cooled to $-30\text{ }^{\circ}\text{C}$. BuLi (2.47M, 11.02 mmol) was slowly added, followed by ten minutes of stirring. A solution of dichloromethyl phosphate (781 mg, 5.25 mmol) in THF (10.5 mL) was slowly cannulated into the reaction vessel containing the alkoxide. The reaction continued to stir at $-30\text{ }^{\circ}\text{C}$ for 30 minutes, the cold bath was removed and the solution was warmed to room temperature and stirred for an additional two hours. Upon completion (monitored by TLC), the reaction was quenched with 20 mL of NH_4Cl (sat'd aq). The separated aqueous layer was extracted with Et_2O (3x) and the combined organic layers washed with brine then dried (MgSO_4). Flash chromatography (2:1 Hexanes/ EtOAc) provided 1.96 g (87% yield) of the diene product as a faint yellow oil. $[\alpha]_{\text{D}} +0.54$ ($c = 0.37$, CH_2Cl_2); IR (neat) 2930, 1452, 1266, 1016, 734, 698 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.4-7.3 (m, 10H), 5.91 (ddd, $J = 17.2$, 10.6, 6.4 Hz, 1H), 5.91 (ddd, $J = 17.2$, 10.6, 6.4 Hz, 1H), 5.45 (d, $J = 17.2$, 1H), 5.39 (d, $J = 17.2$, 1H), 5.30 (d, $J = 10.5$, 1H), 5.30 (d, $J = 10.5$, 1H), 4.98-5.09 (m, 2H), 4.59 (dd, $J = 12.2$, 5.1 Hz, 2H), 4.54 (dd, $J = 12$, 1.4 Hz, 2H), 3.72 (d, $J = 11.4$ Hz, 3H), 3.62 (dd, $J = 10.6$, 6.3 Hz, 2H), 3.57 (m, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.11, 134.05, 134.02, 133.99, 128.56, 127.89, 118.84, 118.65, 78.13, 78.10, 78.09, 78.05, 73.44, 73.41, 72.47(d, $J_{\text{CP}} = 0.8$ Hz), 72.42 (d, $J_{\text{CP}} = 1.5$ Hz), 54.48, 54.44; ^{31}P NMR (162 MHz, CDCl_3) δ 0.29; HRMS calculated for $\text{C}_{23}\text{H}_{29}\text{O}_6\text{P}$ ($\text{M}+\text{H}$) $^{+}$ 433.1780; found 433.1757 (FAB).

(*S,S*)-monocyclic phosphate 1: A flask containing the (*S,S*)-diene phosphate triester (1.45 g, 3.36 mmol) was charged with toluene (670 mL) that had been degassed for 20 minutes with argon. The solution was brought to reflux while continually sparging with argon, and $(\text{ImesH}_2)(\text{PCy}_3)(\text{Cl}_2)\text{Ru}=\text{CHPh}$ was added (200 mg, 0.235 mmol). The reaction was maintained at reflux under argon sparging for 30 minutes. Upon completion (monitored by TLC) the reaction was cooled to room temperature and concentrated under reduced pressure. Purification

via flash chromatography (1:1 Hexanes/EtOAc) supplied 927 mg (68% yield) of monocyclic phosphate **1**. $[\alpha]_D -52.6$ ($c = 1.6$, CH_2Cl_2); IR (neat) 3029, 1454, 1365, 1282, 1027 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.28-7.39 (m, 10H), 5.76 (ddd, $J = 12.0, 1.8, 1.6$ Hz, 1H), 5.72 (ddd, $J = 12.0, 1.8, 1.6$ Hz, 1H), 5.26-5.32 (m, 1H), 5.07-5.14 (m, 1H), 4.56-4.64 (m, 4H), 3.87 (d, $J_{\text{HP}} = 11.1$ Hz, 3H), 3.67 (dddd, $J = 6.4, 5.5, 5.1, 0.9$ Hz, 2H), 3.59 (dddd, $J = 5.3, 5.1, 2.1, 0.5$ Hz, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 137.56, 137.48, 129.49, 128.42, 128.09, 127.83, 127.81, 127.69, 127.63, 73.91, (d, $J_{\text{CP}} = 7.0$), 73.62 (d, $J_{\text{CP}} = 4.5$), 73.49, 73.33, 71.60 (d, $J_{\text{CP}} = 11.9$), 71.21 ($J_{\text{CP}} = 12.1$), 54.62 (d, $J_{\text{CP}} = 5.1$); ^{31}P NMR (162 MHz, CDCl_3) δ 3.61; HRMS calculated for $\text{C}_{21}\text{H}_{25}\text{O}_6\text{P}$ ($\text{M}+\text{H}$) $^+$ 405.1467; found 405.1437 (FAB).

(2S, 5S, 3Z)-1,6-bis(benzyloxy)hex-3-ene-2,5-diol 4: Phosphate **1** (30 mg, 0.074 mmol) was taken up in THF (0.20 mL) and cooled to 0 °C while stirring under argon. LiAlH_4 (0.010 mg, 0.24 mmol) was added slowly, and upon complete addition stirring continued for an additional 2 hours while warming to room temperature. The mixture was then cooled to 0 °C and slowly quenched with H_2O (0.1 mL) followed by 10% NaOH aq (0.2 mL). The layers were separated and the aqueous layer extracted with EtOAc (5x). The combined organic layers were concentrated under reduced pressure and dried (Na_2SO_4). Column chromatography (5:1 Heptane:EtOAc) provided **4** (18.3 mg, 75%) as a clear oil. $[\alpha]_D + 21.7$ ($c = 1.7$, CH_2Cl_2); IR (neat) 3417, 3025, 2902, 1493, 1465, 1101, 1075, 737, 698 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.28-7.37 (m, 10H), 5.58-5.64 (m, 2H), 4.63-4.68 (m, 2H), 4.57 (s, 2H), 3.53 (dd, $J = 9.4, 4.4$ Hz, 2H), 3.46 (dd, $J = 9.5, 7.4$ Hz, 1H), 2.65 (bs, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 137.62, 132.08, 128.47, 127.87, 127.84, 73.48, 73.42, 67.21. HRMS calculated for $\text{C}_{20}\text{H}_{24}\text{O}_4$ ($\text{M}+\text{Na}$) $^+$ 351.1572; found 351.1582 (FAB).

(S,S)-hydrogenated monocyclic phosphate 5: Post ring-closing metathesis (50 mg, 0.123 mmol) with $(\text{ImesH}_2)(\text{PCy}_3)(\text{Cl}_2)\text{Ru}=\text{CHPh}$ (33 mg, 0.040 mmol), the reaction was a concentrated under reduced pressure and the resulting brown oil was taken up in dry CH_2Cl_2 (10 mL). The reaction was cannulated to a hydrogen Parr bomb apparatus, followed by

implementing three quick cycles of evacuation and hydrogen purging. The bomb was capped and charged with H₂ (300 psi) while heating at 80 °C. After stirring for two hours, the reaction mixture was concentrated under reduced pressure and immediately subjected to column chromatography (1:1 EtOAc:Heptane) providing 45.4 mg of **5** (90 % yield) as a light brown oil (small ruthenium contamination estimated to be < 1%). [α]_D -2.15 (*c* = 0.33, CH₂Cl₂); IR (neat) 2917, 1279, 1098, 1000, 737, 698 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.25-7.37 (m, 10H), 4.52-4.61 (m, 5H), 4.31-4.35 (m, 1H), 3.82 (d, *J*_{HP} = 11.1 Hz, 3H), 3.43-3.65 (m, 4H), 2.06 (dd, *J* = 11.9, 4.3 Hz, 1H), 1.98 (dd, *J* = 10.6, 3.8 Hz, 1H), 1.73-1.82 (m, 2H); ¹³C NMR (100 MHz, CDCl₃) δ 137.67 (d, *J*_{CP} = 2.6 Hz), 128.35, 127.72, 127.65, 127.55, 77.94 (d, *J*_{CP} = 6.4 Hz), 76.38 (d, *J*_{CP} = 3.7 Hz), 73.32 (d, *J*_{CP} = 10.6 Hz), 72.59 (d, *J*_{CP} = 11.1 Hz), 72.08 (d, *J*_{CP} = 11.1 Hz), 54.26 (d, *J*_{CP} = 4.9 Hz), 30.78, 30.51; ³¹P NMR (162 MHz, CDCl₃) δ 3.02; HRMS calculated for C₂₁H₂₇O₆P (M+H)⁺ 407.1624; found 407.1649 (FAB).

(2S, 5S)-1,6-bis(benzyloxy)hexane-2,5-diol 6: Phosphate **5** (30 mg, 0.074 mmol) was dissolved in THF (1.2 mL) and cooled to 0 °C while stirring under argon. LiAlH₄ (0.0084 mg, 0.22 mmol) was added slowly, and upon complete addition stirring continued for an additional 2 hours while warming to room temperature. An additional amount of LiAlH₄ was added, and the solution stirred for an additional hour. The mixture was then cooled to 0 °C and slowly quenched with H₂O (0.1 mL) followed by 10% NaOH aq (0.2 mL), and then H₂O (0.1 mL). The layers were separated and the aqueous layer extracted with EtOAc (5x). The combined organic layers were concentrated under reduced pressure and dried (Na₂SO₄). Column chromatography (5:1 Heptane:EtOAc) provided **6** (21.9 mg, 90%) as a clear oil. [α]_D +7.06 (*c* = 0.09, CH₂Cl₂); IR (neat) 3436, 1096, 737, 698 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.28-7.37 (m, 10H), 4.55 (s, 2H), 3.82-3.88 (m, 2H), 3.49 (dd, *J* = 9.0, 3.5 Hz, 2H), 3.49 (dd, *J* = 9.0, 7.5 Hz, 2H), 2.25 (bs, 2H), 1.56-1.65 (m, 4H); ¹³C NMR (125 MHz, CDCl₃) δ 137.93, 128.46, 127.79, 127.75, 74.41, 73.34, 70.38, 29.53. HRMS calculated for C₂₀H₂₄O₄ (M+Na)⁺ 351.1572; found 351.1582 (FAB).

General Procedure A for Diakyl Zincate Derived Cuprates: To a solution of CuCN•2LiCl (1 mmol, 1M in THF) cooled to -30 °C was added dropwise the dialkyl zinc reagent as a solution in

THF (1 mmol). After stirring 30-45 min (Me_2Zn required longer stirring times) at $-25\text{ }^{\circ}\text{C}$ to $-30\text{ }^{\circ}\text{C}$, a solution of phosphate **1** (0.20 mmol) in THF (2 mL) was added. The reaction progress was monitored by TLC (products are baseline in EtOAc), and upon completion where quenched with 10% HCl aq. (1 mL) and stirred for 30 – 60 min. The aqueous layer was extracted was extracted with EtO_2 (4 x 1 mL) and the combined organic layers were subsequently washed with H_2O (1 x 1 mL) and filtered of the resulting white solids. The aqueous layer was removed and the organic layer was concentrated under reduced pressure. The crude acids were used without further purification.

General Procedure B for Alkyl Zinc halide Derived Cuprates: To a solution of $\text{CuCN}\cdot 2\text{LiCl}$ (1 mmol, 1M in THF) cooled to $-30\text{ }^{\circ}\text{C}$ was added dropwise the alkyl zinc halide reagent as a solution in THF (1 mmol, 1.0M). After stirring for one hour at $-25\text{ }^{\circ}\text{C}$ to $-30\text{ }^{\circ}\text{C}$, a solution of phosphate **1** (0.11 mmol) in THF (1.1 mL) was added. The reaction progress was monitored by TLC (products are baseline in EtOAc), and upon completion (~ 2 hours) were quenched with 10% HCl aq. (1 mL) and stirred for 30-60 min. The layers were separated and the aqueous layer extracted with CH_2Cl_2 (4 x 1 mL) and the combined organic layers were subsequently washed with H_2O (1 x 1mL) and the resulting white solids were filtered off. The aqueous layer was removed and the organic layer was concentrated under reduced pressure. The crude acids were used without further purification.

General Procedure for Making Alkyl Zinc Iodides (Cy-Zn-I):

To a heterogeneous mixture of activated Zn dust (288 mg, 4.4 mmol) in THF (330 μL) under argon was added dibromoethane (15 μL , 0.176 mmol), which was heated to $65\text{ }^{\circ}\text{C}$ for one minute and then cooled to room temperature. Trimethylsilyl chloride was then added (16 μL , 0.132 mmol) and the mixture was allowed to stir for 15 minutes, upon which time the appropriate alkyl iodide (4 mmol) in THF (1.6 mL) was added and the reaction heated to $40\text{ }^{\circ}\text{C}$ with stirring. Upon completion (monitored by GC or Zn disappearance), the reagent was used immediately.

General Procedure for Red-Al[®] of Phosphate Acids: A stirring solution of the phosphate acid (0.124 mmol) in toluene (1.24 mL) was cooled to $0\text{ }^{\circ}\text{C}$ and Red-Al (151 μL , 0.495 mmol) was added dropwise. The reaction was allowed to proceed for 30 minutes and monitored by TLC. If necessary, an additional 1 equivalent of Red-Al was added to ensure completion. Reaction was quenched with H_2O (1 mL) followed by slow addition of 10% HCl aq. (1 mL) at $0\text{ }^{\circ}\text{C}$. The layers

were separated and the aqueous layer extracted with CH₂Cl₂ (4 x) and the combined organic layers were subsequently washed with brine and dried (MgSO₄).

(2*S*,3*R*,*E*)-1,6-bis(benzyloxy)-3-methylhex-4-en-2-ol 9a: Following procedure A (100 mg of **1**) and phosphate acid removal, column chromatography (Heptane:EtOAc 4:1) afforded **9a** (67 mg, 83 %) as a clear oil over two steps. $[\alpha]_D^{+30.4}$ ($c = 1.0$, CH₂Cl₂); IR (neat) 3461, 2358, 1456, 1361, 1249, 1027 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.35-7.26 (m, 10H), 5.58-5.79 (m, 2H), 4.53 (d, $J = 1.8$ Hz, 2H), 4.48 (s, 2H), 3.96-3.97 (m, 2H), 3.64 (ddd, $J = 7.2, 7.2, 3.0$ Hz, 1H), 3.55 (dd, $J = 9.5, 3.1$, 1H), 3.39 (dd, $J = 9.7, 7.7$, 1H), 2.32-2.39 (m, 1H), 1.09 (d, $J = 6.8$ Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 138.26, 137.90, 135.62, 128.44, 127.77, 127.71, 127.59, 127.05, 73.53, 73.34, 72.66, 71.98, 70.60, 39.80, 15.9. HRMS calculated for C₂₁H₂₆O₃ (M+Na)⁺ 349.1780; found 349.1817 (FAB).

(2*S*,3*R*,*E*)-1,6-bis(benzyloxy)-3-ethylhex-4-en-2-ol 9b: Following procedure A (50 mg of **1**) and phosphate acid removal, column chromatography (Heptane:EtOAc 4:1) afforded **9b** (38 mg, 90%) as a clear oil over two steps; $[\alpha]_D^{+4.87}$ ($c = 0.58$, CH₂Cl₂); IR (neat) 3420, 2914, 1457, 1269, 1101, 1073, 737, 699 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.35-7.26 (m, 10H), 5.62 (dt, $J = 15.4, 6$ Hz, 1H), 5.44 (dd, $J = 15.4, 9.3$ Hz, 1H), 4.53 (d, $J = 4.4$ Hz, 2H), 4.49 (s, 2H), 3.99 (d, $J = 5.9$ Hz, 2H), 3.7-3.66 (m, 1H), 3.57 (d, $J = 9.5$ Hz, 1H), 3.38 (t, $J = 9.3$ Hz, 1H), 2.48 (s, 1H), 2.14-2.06 (m, 1H), 1.32-1.21 (m, 2H), 0.88 (t, $J = 7.5$ Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 138.26, 137.89, 133.60, 129.01, 128.51, 128.41, 128.36, 127.75, 127.72, 127.69, 127.57, 126.94, 73.32, 72.94, 72.49, 71.77, 70.50, 47.92, 23.32, 11.56. HRMS calculated for C₂₂H₂₈O₃ (M+NH₄)⁺ 358.2382; found 358.2381 (FAB).

(2*S*,3*R*,*E*)-1,6-bis(benzyloxy)-3-isopropylhex-4-en-2-ol 9c: Following procedure A (50 mg of **1**) and phosphate acid removal, column chromatography (Heptane:EtOAc 5:1) afforded **9c** (44.5 mg, 89 %) as a clear oil over two steps. $[\alpha]_D^{+17.5}$ (c , 2.1 CH₂Cl₂); IR (neat) 3446, 2348, 1101 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.38-7.27 (m, 10H), 5.59 (ddd, J = 15.6, 6.0, 6.0 Hz, 1H), 5.43 (dd, J = 15.4, 10.0 Hz, 1H), 4.53 (dd, J = 23.0, 11.7 Hz, 2H), 4.48 (dd, J = 12.2, 8.0 Hz, 2H), 3.92 (dd, J = 5.9, 1.0 Hz, 2H), 3.78-3.84 (m, 1H), 3.57 (dd, J = 9.5, 2.6 Hz, 1H), 3.33 (dd, J = 9.4, 7.6 Hz, 1H), 2.39 (bs, 1H), 2.13-2.21 (m, 1H), 2.07 (ddd, J = 9.6, 9.6, 3.1 Hz, 1H), 0.88 (d, J = 6.8 Hz, 3H), 0.86 (d, J = 6.9 Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 138.28, 137.90, 130.41, 130.11, 128.42, 128.37, 127.75, 127.72, 127.70, 127.58, 73.49, 73.34, 71.70, 70.46, 70.04, 51.79, 26.66, 21.38, 16.82. HRMS calculated for C₂₃H₃₀O₃ (M+Na)⁺ 377.2093; found 377.2075 (FAB).

(2*S*,3*R*,*E*)-1,6-bis(benzyloxy)-3-cyclohexylhex-4-en-2-ol 9d: Following procedure B (100 mg of **1**) and phosphate acid removal, column chromatography (Heptane:EtOAc 5:1) afforded **9d** (68 mg, 70 %) as a clear oil; $[\alpha]_D^{+9.84}$ (c = 1.58, CH₂Cl₂); IR (neat) 3453, 3031, 2922, 2850, 1496, 1450, 1098, 737, 698 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.37-7.28 (m, 10H), 5.57 (dt, J = 15.4, 5.9 Hz, 1H), 5.44 (dd, J = 15.4, 10 Hz, 1H), 4.53 (dd, J = 11.8, 10.6 Hz, 2H), 4.49 (s, 2H), 3.98 (d, J = 5.9 Hz, 2H), 3.86 (td, J = 8.4, 2.5 Hz, 1H), 3.57 (dd, J = 9.5, 2.6 Hz, 1H), 3.32 (dd, J = 9.4, 7.8 Hz, 1H), 2.42 (d, J = 4 Hz, 1H), 2.07 (td, J = 9.6, 3.9 Hz, 1H), 1.80-1.70 (m, 4H), 1.65 (d, J = 12.7 Hz, 1H), 1.50 (d, J = 12.8 Hz, 1H), 1.33-1.20 (m, 2H), 1.14-1.06 (m, 2H), 0.94 (ddd, J = 13, 12.8, 3.1 Hz, 1H); ¹³C NMR (125 MHz, CDCl₃) δ 138.57, 138.19, 131.42, 130.15, 128.66, 128.62, 127.99, 127.95, 127.93, 127.83, 73.30, 73.58, 72.01, 70.73, 69.68, 52.07, 37.37, 32.16, 28.09, 26.90, 26.84, 26.75; HRMS calculated for C₂₆H₃₄O₃ (M+Li)⁺ 401.2668; found 401.2678 (FAB).

(2*S*,3*R*,*E*)-3-benzyl-1,6-bis(benzyloxy)hex-4-en-2-ol 9e: Following procedure B (100 mg **1**) and phosphate acid removal, column chromatography (Heptane:EtOAc 4:1) afforded **9e** (81.0 mg, 82%) as a clear oil over two steps. $[\alpha]_D -6.2$ (c 1.4, CH_2Cl_2); IR (neat) 3444, 1494, 1452, 1097 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.38-7.15 (m, 15H), 5.48 (dd, $J = 15.7, 8.1\text{ Hz}$, 1H), 5.43 (ddd, $J = 15.7, 10.6, 5.3\text{ Hz}$, 1H), 4.56 (dd, $J = 11.9, 7.3\text{ Hz}$ 2H), 4.31 (dd, $J = 12.0, 0.1\text{ Hz}$, 2H), 3.91 (dd, $J = 12.8, 4.1\text{ Hz}$, 1H), 3.87 (dd, $J = 12.3, 5.0\text{ Hz}$, 1H), 3.78-3.73 (m, 1H), 3.65-3.57 (m, 1H), 3.50-3.43 (m, 1H), 3.12 (dd, $J = 13.4, 4.0\text{ Hz}$, 1H), 2.68 (dd, $J = 13.5, 9.3\text{ Hz}$, 1H), 2.62-2.52 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 139.79, 138.31, 137.00, 132.81, 129.52, 129.38, 128.45, 128.31, 128.07, 127.82, 127.76, 127.67, 127.51, 125.85, 73.37, 72.68, 71.81, 71.37, 70.27, 47.82, 37.27. HRMS calculated for $\text{C}_{37}\text{H}_{30}\text{O}_3$ ($\text{M}+\text{Na}$) $^+$ 425.2093; found 425.2083 (FAB).

(*R*,*E*)-8-(benzyloxy)-5-((*S*)-2-(benzyloxy)-1-hydroxyethyl)oct-6-enenitrile 9f:

Following procedure B (100 mg of **1**) and methylation/phosphate removal, column chromatography (Heptane:EtOAc 4:1) afforded **9f** (78.0 mg, 83%) as a clear oil. $[\alpha]_D = -12.0$ (c = 1.5, CH_2Cl_2); IR (neat) 3463, 3031, 2927, 2246, 742, 702 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.36-7.28 (m, 10H), 5.63 (ddd, $J = 15.5, 5.7, 5.7\text{ Hz}$, 1H), 5.43 (dd, $J = 15.5, 9.5\text{ Hz}$, 1H), 4.52 (dd, $J = 11.8, 7.7\text{ Hz}$, 2H), 4.48 (dd, $J = 9.1, 0.1\text{ Hz}$, 2H), 3.97 (d, $J = 5.6\text{ Hz}$, 2H), 3.62 (ddd, $J = 7.4, 5.4, 2.3\text{ Hz}$, 1H), 3.55 (dd, $J = 9.5, 9.4\text{ Hz}$, 1H), 3.36 (dd, $J = 9.4, 7.6\text{ Hz}$, 1H), 2.47 (bs, 1H), 2.32 (dddd, $J = 14.3, 7.3, 7.0, 2.6\text{ Hz}$ 2H), 2.18 (ddd, $J = 18.2, 10.1, 3.3\text{ Hz}$, 1H), 1.96-1.86 (m, 1H), 1.78-1.67 (m, 1H), 1.55 (ddd, $J = 6.9, 6.0, 4.8\text{ Hz}$, 1H), 1.34-1.45 (m, 1H); ^{13}C NMR (100 MHz, CDCl_3) δ 138.17, 137.78, 132.42, 129.90, 128.51, 128.46, 127.91, 127.80, 127.75, 127.72, 119.72, 73.44, 72.71, 72.49, 72.23, 70.30, 45.91, 29.76, 23.31, 17.27. HRMS calculated for $\text{C}_{24}\text{H}_{29}\text{NO}_3$ ($\text{M}+\text{NH}_4$) $^+$ 397.2491; found 397.2483 (FAB).

(2*S*,3*R*)-1-(benzyloxy)-3-((*E*)-3-(benzyloxy)prop-1-enyl)-7-chloroheptan-2-ol 9g:

Following procedure B (100mg of **1**) and methylation/phosphate removal, column chromatography (Heptane:EtOAc 5:1) afforded **9g** (84.7 mg, 85 %) as a clear oil. $[\alpha]_D = +5.8$ ($c = 0.625$, CH_2Cl_2); IR (neat) 3442, 3031, 2921, 2856, 1494, 1452, 1361, 1099, 736, 698 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.32-7.11 (m, 10H), 5.67-5.59 (m, 1H), 5.46 (dd, $J = 15.5, 9.4$ Hz, 1H), 4.54 (dd, $J = 17.4, 11.9$ Hz, 2H), 4.49 (app s, 2H), 3.99 (dd, $J = 5.9, 1.1$ Hz, 2H), 3.66 (ddd, $J = 7.8, 7.8, 2.9$ Hz, 1H), 3.57 (dd, $J = 9.5, 2.9$ Hz, 1H), 3.57-3.52 (m, 2H), 3.39 (dd, $J = 9.5, 7.5$ Hz, 1H), 2.22-1.88 (m, 3H), 1.84-1.69 (m, 3H), 1.48-1.24 (m, 4H); ^{13}C NMR (100 MHz, CDCl_3) δ 138.27, 137.86, 133.35, 129.24, 128.45, 128.38, 127.73, 127.70, 127.61, 73.38, 72.71, 72.85, 72.62, 71.90, 70.3420, 46.24, 45.00, 35.59, 29.70, 24.43. HRMS calculated for $\text{C}_{24}\text{H}_{31}\text{ClO}_3$ ($\text{M}+\text{NH}_4$) $^+$ 420.2305; found 420.2310 (FAB).

(2*S*,3*R*,*E*)-1-(benzyloxy)-3-(3-(benzyloxy)prop-1-enyl)oct-7-en-2-ol 9h: Following procedure B and phosphate acid removal, column chromatography (Heptane:EtOAc 4:1) afforded **9h** (61 mg, 65%) as a clear oil; $[\alpha]_D +3.53$ ($c = 0.17$, CH_2Cl_2); IR (neat) 3422, 1274, 750, 699 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ 7.36-7.30 (m, 10H), 5.80 (ddt, $J = 17, 10.2, 6.6$ Hz, 1H), 5.62 (dt, $J = 15.5, 6$ Hz, 1H), 5.46 (dd, $J = 15.5, 9.4$ Hz, 1H), 5.00 (dd, $J = 17.1, 1.4$ Hz, 1H), 4.94 (dd, $J = 10.2, 1$ Hz, 1H), 4.54 (dd, 18.3, 11.7, 2H), 4.49 (s, 2H), 3.99 (d, $J = 6$ Hz, 2H), 3.66 (td, $J = 7.7, 3$ Hz, 1H), 3.57 (dd, $J = 9.5, 3$ Hz, 1H), 3.38 (dd, $J_{\text{HH}} = 9.3, 7.7$ Hz, 1H), 2.40 (d, $J = 4.2$ Hz, 1H), 2.19 (ddd, $J = 9.4, 7.9, 3.6$ Hz, 1H), 2.09-2.00 (m, 2H), 1.77-1.73 (m, 1H), 1.49-1.45 (m, 1H), 1.31-1.27 (m, 2H); ^{13}C NMR (125 MHz, CDCl_3) δ 139.07, 138.53, 138.15, 133.98, 129.20, 128.67, 128.61, 128.01, 127.95, 127.94, 127.82, 114.65, 73.59, 73.10, 72.94, 72.06, 70.72, 46.50, 34.03, 30.19, 26.73. HRMS calculated for $\text{C}_{25}\text{H}_{32}\text{O}_3$ ($\text{M}+\text{NH}_4$) $^+$ 398.2695; found 398.2696 (FAB).

(2*S*,3*R*,*E*)-1,6-bis(benzyloxy)-3-methylhex-4-en-2-yl methyl hydrogen phosphate 8a: Using procedure A (50 mg of **1**) afforded **8a** (51.5 mg, 99%) as a clear viscous oil (compound was processed without further purification). $[\alpha]_D +15.6$ ($c = 4.5$, CH_2Cl_2); IR (neat) 2331, 1496, 1454, 1232 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 10.16 (bs, 1H), 7.37-7.26 (m, 10H), 5.71-5.62 (m, 2H), 4.58-4.49 (m, 2H), 4.48 (s, 2H), 4.36-4.32 (m, 1H), 3.96 (s, 2H), 3.71-3.53 (m, 5H), 2.73-2.61 (m, 1H), 1.11 (d, $J = 6.7$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 138.67, 138.30, 134.72, 128.81, 128.80, 128.25, 128.22, 128.11, 128.05, 81.45, 73.71, 72.44, 71.02, 70.45, 54.61, 38.80, 15.87; ^{31}P NMR (162 MHz, CDCl_3) δ 2.67. HRMS calculated for $\text{C}_{22}\text{H}_{29}\text{O}_6\text{P}$ ($\text{M}+\text{Na}$) $^+$ 443.1599; found 443.1600 (FAB).

General Methylation procedure:

To a stirring solution of phosphate acid (0.295 mmol) in MeOH (0.6 mL) was slowly added (trimethylsilyl)diazomethane (2.0M, 1.48 mmol). Upon addition, the reaction was allowed to run for 20 minutes at room temperature with stirring to ensure completion (monitored by TLC), then quenched by dropwise addition of acetic acid until effervescence ceases. The reaction was concentrated under reduced pressure and pushed through a small plug of silica (4:1 Hexanes/EtOAc) to supply the desired methyl phosphate ester.

(2*S*,3*R*,*E*)-1,6-bis(benzyloxy)-3-methylhex-4-en-2-yl dimethyl phosphate 10a: Using procedure A, methylation followed by column chromatography (EtOAc:Heptane 2:1) afforded **10a** (50 mg, 95%) as a clear oil over two steps. $[\alpha]_D -27.0$ ($c = 2.0$, CH_2Cl_2); IR (neat) 3062, 2952, 1454, 1363, 1278, 1041 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.36-7.24 (m, 10H), 5.69-5.62 (m, 2H), 4.58-4.40 (m, 5H), 3.97-3.94 (m, 2H), 3.73 (d, $J_{\text{HP}} = 11.2$, 2H), 3.69 (d, $J_{\text{HP}} = 11.2$, 2H), 3.64-3.56 (m, 2H), 2.73-2.64 (m, 1H), 1.10 (d, $J = 6.9$ Hz, 3H); ^{13}C NMR (125 MHz, CDCl_3) δ 138.19, 137.79, 134.00, 128.34, 128.31, 127.89, 127.71, 127.69, 127.65, 127.57, 80.95

(d, J_{CP} = 6.3), 73.16, 71.99, 70.55, 70.07 (d, J_{CP} = 1.3), 54.28 (d, J_{CP} = 6.3), 54.23 (d, J_{CP} = 6.3), 38.44 (d, J_{CP} = 5.0), 15.43; ^{31}P NMR (162 MHz, CDCl_3) δ 2.06; HRMS calculated for $\text{C}_{23}\text{H}_{32}\text{O}_6\text{P}$ ($\text{M}+\text{H}$) $^+$ 435.1937; found 435.1967 (FAB).

(S)-configured unsymmetrical monocyclic phosphate 11: A flask containing allylic alcohol **3** (0.680 g, 3.83 mmol) was diluted up in THF (0.2 M, 20 mL) and cooled to $-40\text{ }^\circ\text{C}$, followed by dropwise addition of BuLi (1.53, 2.5M). After 10 minutes of stirring, allyldiphenyl phosphate (1.45 g, 5 mmol) in THF (1M, 5 mL) was cannulated to the flask containing the alkoxide. The solution was stirred for eight hours while slowly warming to room temperature. The reaction was quenched with sat'd aq. NH_4Cl and the layers were separated and the aqueous layer extracted with Et_2O (2x). The combined organic layers were concentrated under reduced pressure and dried (Na_2SO_4). Column chromatography (1:1 Hex:EtOAc) afforded the acyclic phosphate ester (1.03 g, 71%) as an oil. To a flask containing the phosphate precursor was added degassed CH_2Cl_2 (0.05 M, 22.6 mL) and was subsequently fitted with a reflux condenser and needle, through which a slow steady stream of Ar flowed. Grubbs catalyst (ImesH_2)(PCy_3)(Cl_2) $\text{Ru}=\text{CHPh}$ (0.034 mg, 0.04 mmol) was added and the reaction lowered into an oil bath warmed to $50\text{ }^\circ\text{C}$. The reaction was monitored by TLC and after 1.0-1.5 hrs was deemed complete. Following cooling to room temperature, the reaction was concentrated and purified via column chromatography (2:1 Hex:EtOAc), to yield cyclic phosphate **11** as a 1:1 mixture of diastereomers at phosphorus (0.301 g, 76%); ^1H NMR (400 MHz, CDCl_3) δ 7.40-7.17 (m, 10H), 5.89-5.74 (m, 2H), 5.41-5.23 (m, 1H), 5.14-4.89 (m, 1H), 4.71-4.51 (m, 3H), 3.79-3.62 (m, 2H); ^{13}C NMR (100 MHz, CDCl_3) δ 150.38 (d, J_{CP} = 16.1), 150.22 (d, J_{CP} = 16.1), 137.44, 129.68 (d, J_{CP} = 4.0), 128.61, 128.38, 128.23, 127.78, 127.67 (d, J_{CP} = 7.0), 127.42, 127.08, 125.22, (d, J_{CP} = 6.0), 120.11 (d, J_{CP} = 5.0), 119.87 (d, J_{CP} = 4.0), 74.35 (d, J_{CP} = 7.0), 74.01 (d, J_{CP} = 4.0), 73.47, 73.28, 71.58 (d, J_{CP} = 9.1), 71.15 (d, J_{CP} = 11.1), 64.87 (d, J_{CP} = 7.0), 64.80 (d, J_{CP} = 8.0); ^{31}P NMR (162 MHz, CDCl_3) δ -3.61 and -2.87; HRMS calculated for $\text{C}_{18}\text{H}_{19}\text{O}_5\text{P}$ ($\text{M}+\text{Na}$) $^+$ 369.0868; found 369.0872 (FAB).

(*R,E*)-5-(benzyloxy)-2-ethylpent-3-en-1-ol 12: Following procedure A (90 mg of **11**) and methylation/phosphate removal, column chromatography (Heptane:EtOAc 3:1) afforded **12** (42.1 mg, 81%) as a clear oil. $[\alpha]_D + 6.2$ ($c = 1.6$, CH_2Cl_2); IR (neat) 3389, 2958, 1454, 1307, 1089, 1049 cm^{-1} ; ^1H NMR (400 MHz, CDCl_3) δ 7.22-7.19 (m, 5H), 5.63 (dt, $J = 15.4, 6.0$ Hz, 1H), 5.43 (dd, $J = 15.5, 8.7$ Hz, 1H), 4.45 (s, 2H), 3.49 (dd, $J = 10.6, 5.2$ Hz, 1H), 3.36 (dd, $J = 10.6, 8.0$ Hz, 1H), 2.12-2.03 (m, 1H), 1.71 (bs, 1H), 1.45-1.34 (m, 1H), 1.26-1.15 (m, 1H), 0.83 (t, $J = 7.5$ Hz, 3H); ^{13}C NMR (100 MHz, CDCl_3) δ 138.24, 135.15, 129.18, 128.43, 127.68, 127.58, 72.11, 70.67, 65.60, 47.26, 23.78, 11.67. HRMS calculated for $\text{C}_{14}\text{H}_{20}\text{O}_2$ ($\text{M}+\text{Na}$) $^+$ 243.1361; found 243.1361 (FAB).

(*S*)-configured unsymmetrical monocyclic phosphate 14: A flask containing allylic alcohol **3** (0.239 g, 1.34 mmol) was diluted up in THF (0.2 M, 6.7 mL) and cooled to $-30\text{ }^\circ\text{C}$, followed by dropwise addition of BuLi (0.537 mL, 2.5M). After 10 minutes of stirring, dichloromethyl phosphate (0.147, 1.47 mmol) was added to the flask containing the alkoxide. The solution was stirred for one hour while slowly warming to room temperature. The reaction was quenched with sat'd aq. NH_4Cl^- and the layers were separated and the aqueous layer extracted with Et_2O (2x). The combined organic layers were concentrated under reduced pressure and dried (Na_2SO_4). Column chromatography (4:1 Hex:EtOAc) afforded the monochloride phosphate ester (0.311 g, 80%) as an oil. The chloride was immediately used. A flask containing alcohol **13** (0.074 mg, 0.861 mmol) was charged with THF (0.2 M, 4.31 mL) and cooled to $-30\text{ }^\circ\text{C}$, followed by dropwise addition of BuLi (0.346 mL, 2.4 M). After 10 minutes of stirring, a solution of the previously generated monochloride phosphate ester (0.200 g, 0.688 mmol) in THF (0.7 M, 1 mL) was cannulated to the flask containing the alkoxide. The solution was stirred for 1.5 hrs while slowly warming to room temperature. The reaction was quenched with sat'd aq. NH_4^+Cl^- and the layers were separated and the aqueous layer extracted with Et_2O (2x). The combined organic layers were concentrated under reduced pressure and dried (Na_2SO_4). *Due to*

the instability of the acyclic phosphate, the reaction mixture was carried on without chromatography. To a flask containing the crude phosphate ester (assumed to be 0.688 mmol) was added degassed toluene (0.02 M, 34 mL) and was subsequently fitted with a reflux condenser, and needle through which a slow steady stream of Ar flowed. Grubbs catalyst (ImesH₂)(PCy₃)(Cl₂)Ru=CHPh (0.029 mg, 0.034 mmol) was added and the reaction lowered into an oil bath warmed to 90 °C. The reaction was monitored by TLC and after 1.0 hr was deemed complete. Following cooling to room temperature, the reaction was concentrated and purified via column chromatography (2:1 Hex:EtOAc), to yield cyclic phosphate **14** as a 1:1 mixture of diastereomers at phosphorus (0.100 g, 47 %). One diastereomer could be purified, and was used for characterization; IR (neat) 3029, 2966, 1101, 737, 698 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ 7.35-7.27 (m, 5H), 5.63 (dd, *J* = 12.1 Hz, *J*_{HP} = 2.3 Hz, 1H), 5.53 (dd, *J* = 12.1 Hz, *J*_{HP} = 3.1 Hz, 1H), 4.61 (dd, *J* = 12, 4.1 Hz, 2H), 3.81 (d, *J*_{HP} = 11.2 Hz, 3H), 3.67 (dd, *J* = 10.2, 6.9 Hz, 1H), 3.57 (ddd, *J* = 10.2, 5.2, 1.9 Hz, 1H), 1.75 (s, 3H), 1.48 (d, *J*_{HP} = 2.5 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃) δ 137.90, 137.66, 128.46, 127.68, 124.16, 82.10 (d, *J*_{CP} = 5.0 Hz), 73.55, 72.79 (d, *J*_{CP} = 6.3 Hz), 71.86 (d, *J*_{CP} = 10.1 Hz), 54.13 (d, *J*_{CP} = 5.0 Hz), 30.53 (d, *J*_{CP} = 11.3 Hz), 29.32; ³¹P NMR (162 MHz, CDCl₃) δ 0.07 and -0.90.

(2S,3S)-1-(benzyloxy)-3-ethyl-5-methylhex-4-en-2-ol 15: Following procedure A (90 mg of **14**) and methylation/phosphate removal, column chromatography (Heptane:EtOAc 3:1) afforded **15** (58.0 mg, 81%) as a clear oil. [α]_D = +13.1, (*c* = 1.5, CH₂Cl₂); IR (neat) 3431, 3029, 2966, 1101 cm⁻¹; ¹H NMR (400 MHz, CDCl₃) δ 7.31-7.22 (m, 5H), 5.02 (d, *J* = 8.9 Hz, 1H), 4.50 (s, 2H), 3.81-3.74 (m, 1H), 3.54 (dd, *J* = 9.6, 3.1 Hz, 1H), 3.45 (dd, *J* = 9.6, 7.8, 1H), 2.37-2.28 (m, 1H), 2.24 (bs, 1H), 1.78 (s, 3H), 1.65, (s, 3H), 1.61-1.51 (m, 1H), 1.34-1.24 (m, 1H), 0.86, (t, *J* = 7.3, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 138.15, 134.65, 129.61, 128.44, 127.76, 124.40, 115.34, 73.40, 73.21, 73.02, 42.61, 26.05, 24.70, 18.45.

4-(benzyloxymethyl)-5-ethyl-2,2-dimethyl-1,3-dioxane: ¹H NMR (400 MHz, CDCl₃) δ 7.37-7.28 (m, 5H), 4.62 (d, $J = 12.1$ Hz, 1H), 4.48 (d, $J = 12.1$ Hz, 1H), 4.26 (ddd, $J = 6.2, 6.2, 2.4$ Hz, 1H), 3.97 (ddd, $J = 11.9, 2.6, 1.11$ Hz, 1H), 3.82 (dd, $J = 11.9, 1.4$ Hz, 1H), 3.49 (dd, $J = 9.6, 6.6$ Hz, 1H), 3.43 (dd, $J = 9.7, 5.9$ Hz, 1H), 1.77-1.65 (m, 1H), 1.46 (s, 3H), 1.38 (s, 3H), 1.35-1.27 (m, 2H), 0.92 (t, $J = 7.2$ Hz).

(4*S*,5*S*)-4-(benzyloxymethyl)-5-ethyl-2,2-dimethyl-1,3-dioxane: ¹H NMR (400 MHz, CDCl₃) δ 7.36-7.22 (m, 5H), 4.63 (d, $J = 12.3$ Hz, 1H), 4.52 (d, $J = 12.3$ Hz, 1H), 3.87 (dd, $J = 11.7, 5.4$ Hz, 1H), 3.74 (ddd, $J = 10.3, 5.0, 2.8$ Hz, 1H), 85 (t, $J = 7.6$ Hz, 3H); ¹³C NMR (100 MHz, CDCl₃) δ 138.42, 128.33, 127.73, 127.56, 98.19, 73.58, 73.38, 71.50, 63.65, 37.02, 29.38, 21.08, 19.38, 10.05. HRMS calculated for C₁₆H₂₄O₃ (M+Na)⁺ 287.1623; found 287.1628 (FAB).

NMR Spectra for Compounds 1,4-6, 8a, 9a-9h, 10a, 11,12,14,15:

Bis((*S*)-1-(benzyloxy)but-3-en-2-yl) methyl phosphate

(*S,S*)-Monocyclic phosphate- **Compound 1**

(2*S*,5*S*,*Z*)-1,6-bis(benzyloxy)hex-3-ene-2,5-diol- **Compound 4**

(*S,S*)-Hydrogenated monocyclic phosphate- **Compound 5**

(2*S*,5*S*)-1,6-bis(benzyloxy)hexane-2,5-diol- **Compound 6**

(2*S*,3*R*,*E*)-1,6-bis(benzyloxy)-3-methylhex-4-en-2-ol-Compound 9a

(2*S*,3*R*,*E*)-1,6-bis(benzyloxy)-3-ethylhex-4-en-2-ol- **Compound 9b**

(2*S*,3*R*,*E*)-1,6-bis(benzyloxy)-3-isopropylhex-4-en-2-ol- **Compound 9c**

(2*S*,3*R*,*E*)-1,6-bis(benzyloxy)-3-cyclohexylhex-4-en-2-ol- **Compound 9d**

(2*S*,3*R*,*E*)-3-benzyl-1,6-bis(benzyloxy)hex-4-en-2-ol- **Compound 9e**

(*R,E*)-8-(benzyloxy)-5-((*S*)-2-(benzyloxy)-1-hydroxyethyl)oct-6-enitrile- Compound 9f

(2*S*,3*R*)-1-(benzyloxy)-3-((*E*)-3-(benzyloxy)prop-1-enyl)-7-chloroheptan-2-ol- **Compound 9g**

(2*S*,3*R*)-1-(benzyloxy)-3-((*E*)-3-(benzyloxy)prop-1-enyl)oct-7-en-2-ol- **Compound 9h**

(2*S*,3*R*,*E*)-1,6-bis(benzyloxy)-3-methylhex-4-en-2-yl methyl hydrogen phosphate-**Compound 8a**

(2S,3R,E)-1,6-bis(benzyloxy)-3-methylhex-4-en-2-yl dimethyl phosphate-**Compound 10a**

Unsymmetric Monophosphate- **Compound 11**

(*R,E*)-5-(benzyloxy)-2-ethylpent-3-en-1-ol- **Compound 12**

Unsymmetrical dimethyl phosphate – **Compound 14**

(2*S*,3*S*)-1-(benzyloxy)-3-ethyl-5-methylhex-4-en-2-ol – **Compound 15**

4-(benzyloxymethyl)-5-ethyl-2,2-dimethyl-1,3-dioxane

(4*S*,5*S*)-4-(benzyloxymethyl)-5-ethyl-2,2-dimethyl-1,3-dioxane