A Divergent Synthetic Strategy Leading to Structurally Diverse Pyrrolidines and Piperidines from Common γ -Aminoalkyl Butenolide and Aldehyde Precursors

José M. Aurrecoechea, Rubén Suero and Esther de Torres*

Departamento de Química Orgánica II, Facultad de Ciencia y Tecnología, Universidad del País
Vasco, Apartado 644, 48080 Bilbao, Spain

SUPPORTING INFORMATION

Table of Contents

1. General Experimental Methods.	S2
2. Preparation of Precursors.	S2
3. Preparation of Pyrrolidines 13 , 16 , 19 .	S8
4. Preparation of Tetrahydropyridines 17 .	S16
5. Preparation of Piperidines 20 .	S19
6. Stereochemical Elucidation	S25
7. References	S28
8. Copies of ^1H NMR Spectra.	S29

1. General Experimental Methods.

All reactions involving air- and moisture-sensitive materials were performed under an atmosphere of dry Ar. Dichloromethane, diethyl ether, benzene and triethylamine were freshly distilled from CaH₂. Methanol, ethanol and *t*-butanol were made anhydrous with Mg-I₂, distilled and stored over molecular sieves. Tetrahydrofuran (THF) was freshly distilled from sodium/benzophenone and, for reactions with SmI₂, it was deoxygenated prior to use. SmI₂ (*ca* 0.1 M in THF) was prepared from Sm and diiodomethane using a literature procedure.¹ Flash column chromatography was performed on silica gel (230-400 mesh). HPLC purifications were carried out with a μ -Bondapak column (10 μ , 19 mm x 15 cm) using a refraction index detector. Routine NMR spectra were obtained at 250, 300 or 500 MHz for ¹H and 62.9, 75.4 or 125.8 MHz for ¹³C. In most cases, CDCl₃ was used as solvent and internal reference (δ 7.26 for ¹H and δ 77.0 for ¹³C). Alternatively, DMSO-d₆ was used instead (δ = 2.50 for ¹H and δ = 39.5 for ¹³C). The DEPT sequence was routinely used for ¹³C multiplicity assignment. NMR (¹H and ¹³C) peak assignments were established using a combination of selective homodecoupling, COSY, and HSQC experiments. Stereochemical elucidation details are given in Section 6 of this Supporting Information. IR data include only characteristic absorptions. Mass spectra were obtained at 70 eV. GC/MS analysis were performed at 70-280 °C (20 °C/min) with a stationary phase of methylphenylsilicone (0.25 μ m, 30 m x 0.25 mm).

2. Preparation of Precursors

***N*-(*t*-Butoxycarbonyl)-*N*-methoxymethylbenzylamine (2).** To *N*-(Boc)benzylamine (1)² (10.4 g, 50.0 mmol) in THF (150 mL) at -78 °C was added *n*-BuLi (1.60 M, 34.0 mL, 55.0 mmol) dropwise and the solution was stirred for 30 min. Chloromethyl methyl ether (4.60 mL, 60.0

mmol) was added, the mixture was stirred 30 min at -78 °C, allowed to reach room temperature, stirred further 20 h, and then poured over a saturated NaHCO₃ solution (100 mL). The aqueous layer was extracted with CH₂Cl₂ (3 x 75 mL), and the combined organic layers were washed with brine (50 mL) and dried (Na₂SO₄). The residue after evaporation was purified by flash column chromatography (96:4 hexanes/EtOAc) to afford **2** (11.3 g, 90%) as a colorless oil: ¹H NMR (500 MHz, DMSO-d₆, 70 °C) δ 1.41 (s, 9H), 3.22 (s, 3H), 4.42 (s, 2H), 4.65 (s, 2H), 7.24-7.34 (m, 5H); ¹³C NMR (125.8 MHz, DMSO-d₆, 70 °C) δ 27.5 (CH₃), 48.6 (CH₂), 54.6 (CH₃), 78.0 (C), 79.3 (CH₂), 126.4 (CH), 126.9 (CH), 127.8 (CH), 138.1 (C), 154.5 (C); IR (neat) ν 1700 (C=O) cm⁻¹; MS (FAB) m/z (%) 251 (M, 3), 250 (17), 220 (92), 194 (10), 164 (base); HRMS (FAB) calculated for C₁₄H₂₀NO₃ (M-1) 250.1443, observed 250.1443.

5-[N-Benzyl-N-(*t*-butoxycarbonyl)aminomethyl]furan-2(5*H*)-one (3) and 3-[N-Benzyl-N-(*t*-butoxycarbonyl)aminomethyl]furan-2(5*H*)-one (5). To a stirred solution of **2** (7.38 g, 29.4 mmol) in CH₂Cl₂ (140 mL) at -80 °C was added dropwise trimethylsilyloxyfuran (TMSOF) (5.00 mL, 29.4 mmol), followed by trimethylsilyl triflate (TMSOTf) (1.06 mL, 5.88 mmol). The reaction mixture was stirred 14 h at the same temperature and poured over a saturated NaHCO₃ solution (125 mL). The aqueous layer was extracted with CH₂Cl₂ (3 x 100 mL), and the combined organic layers were washed with brine (50 mL) and dried (Na₂SO₄). The residue after evaporation (9.72 g) contained a 5:1 mixture of **3** and **5** that was subjected to successive precipitations from 4:1 hexanes/diethyl ether at 4 °C to yield **3** (5.95 g, 67%) as a white solid. A pure sample of the regioisomer **5** was obtained for characterization purposes by repeated flash column chromatography of the oily residue obtained by evaporation of the supernatant liquids. Data for **3** (rotamer mixture): mp 88-90 °C (dryness); ¹H NMR (250 MHz, CDCl₃) δ 1.43 and 1.52 (2 s, 9H), 3.47-3.62 (m, 2H), 4.37-4.63 (m, 2H), 5.13-5.22 (m, 1H), 6.07-6.09 (m, 1H), 7.18-

7.35 (m, 5H), 7.42-7.54 (m, 1H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 27.8 (CH_3), 47.7 (CH_2), 50.7 (CH_2), 51.8 (CH_2), 80.1 (C), 80.3 (C), 82.0 (CH), 82.8 (CH), 121.0 (CH), 121.6 (CH), 126.8 (CH), 127.0 (CH), 127.4 (CH), 128.2 (CH), 137.2 (C), 137.5 (C), 154.0 (CH), 154.5 (CH), 154.9 (C), 155.2 (C), 172.0 (C), 172.4 (C); IR (neat) ν 1760 (C=O), 1690 (C=O), 1600 (C=C) cm^{-1} ; MS (FAB) m/z (%) 304 (10), 248 (base); HRMS (FAB) calculated for $\text{C}_{17}\text{H}_{22}\text{NO}_4$ (M+1) 304.1549, observed 304.1558. Data for **5** (rotamer mixture): ^1H NMR (250 MHz, CDCl_3) δ 1.47 (s, 9H), 4.02-4.04 (br, 2H), 4.50 (s, 2H), 4.73-4.76 (m, 2H), 7.06-7.35 (m, 6H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 28.3 (CH_3), 41.9 (CH_2), 50.5 (CH_2), 51.4 (CH_2), 70.4 (CH_2), 80.5 (C), 127.5 (CH), 128.0 (CH), 128.6 (CH), 130.8 (C), 131.4 (C), 137.8 (C), 145.8 (CH), 147.2 (CH), 155.4 (C), 173.0 (C), 173.4 (C); IR (neat) ν 1750 (C=O), 1690 (C=O) cm^{-1} ; MS (EI) m/z (%) 247 (27), 230 (5), 202 (55), 150 (23), 106 (base), 91 (69); HRMS calculated for $\text{C}_{17}\text{H}_{21}\text{NO}_4$ 303.1471, observed 303.1474.

5-[(Benzylamino)methyl]furan-2(5H)-one (4). To a stirred solution of **3** (0.303 g, 1.00 mmol) in CH_2Cl_2 (12 mL) at 0 °C was added dropwise TFA (1.31 mL, 17.0 mmol). The solution was stirred 10 min at 0 °C, was allowed to reach room temperature, stirred further 1 h and poured over a stirred saturated NaHCO_3 solution (20 mL). The aqueous layer was extracted with CH_2Cl_2 (3 x 15 mL). The combined organic layers were washed with brine (5 mL) and dried (Na_2SO_4). Solvent evaporation afforded the free amine **4**³ (203 mg, 100%) as an oil that was used without further purification: ^1H NMR (250 MHz, CDCl_3) δ 1.76 (s, 1H), 2.80 (dd, J = 12.9, 6.9 Hz, 1H), 2.99 (dd, J = 12.9, 4.4 Hz, 1H), 3.81* (s, 2H), 5.11-5.17 (m, 1H), 6.12 (dd, J = 5.8, 2.0 Hz, 1H),

* Reported at δ 3.47 in ref. 3.

7.20-7.35 (m, 5H), 7.44 (dd, $J = 5.8, 1.6$ Hz, 1H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 50.7 (CH_2), 53.5 (CH_2), 83.0 (CH), 122.0 (CH), 127.1 (CH), 127.9 (CH), 128.4 (CH), 139.4 (C), 154.8 (CH), 172.8 (C); IR (neat) ν 3320 (N-H), 1755 (C=O), 1665 (C=C) cm^{-1} ; MS (EI) m/z (%) 121 (5), 120 (51), 92 (8), 91 (base); HRMS calculated for $\text{C}_{12}\text{H}_{13}\text{NO}_2$ 203.0946, observed 203.0941.

(5*R, 2'*R**)-5-[*N*-(Boc)Pyrrolidin-2-yl]furan-2(5*H*)-one (7a).** To a stirred solution of **6a** ($n = 1$)^{4,5} (3.69 g, 17.15 mmol) in CH_2Cl_2 (68 mL) at -80°C under Ar was added dropwise *t*-butyldimethylsilyloxyfuran⁶ (3.74 g, 18.86 mmol) in CH_2Cl_2 (2 mL) and then TMSOTf (0.31 mL, 1.72 mmol). After stirring the mixture 12 h at the same temperature, a saturated NaHCO_3 solution (50 mL) was added and the whole was allowed to reach room temperature. After separation, the aqueous layer was extracted with CH_2Cl_2 (3 x 100 mL). The combined organic layers were washed with brine (40 mL), and dried (MgSO_4). The residue after evaporation was purified by flash column chromatography (hexanes/EtOAc, 80:20) to yield **7a**⁷ (3.50 g, 81%, mixture of rotamers at 25°C) as a white solid: ^1H NMR (250 MHz, DMSO-d_6 , 70°C) δ 1.42 (s, 9H), 1.64-2.07 (m, 4H), 3.14-3.23 (m, 1H), 3.36 (dt, $J = 10.7, 7.6$ Hz, 1H), 4.17-4.23 (m, 1H), 5.26-5.30 (m, 1H), 6.16 (dd, $J = 5.7, 1.9$ Hz, 1H), 7.70 (dd, $J = 5.7, 1.5$ Hz, 1H); ^{13}C NMR (62.9 MHz, DMSO-d_6 , 60°C) δ 23.0 (CH_2), 26.4 (CH_2), 27.8 (CH_3), 46.4 (CH_2), 57.0 (CH), 78.8 (C), 84.3 (CH), 120.2 (CH), 153.6 (C), 155.7 (CH), 172.2 (C); IR (neat) ν 1750 (C=O), 1690 (C=O), 1590 (C=C) cm^{-1} ; MS (EI) m/z (%) 254 ($M+1$, 38), 219 (7), 199 (11), 198 (base), 180 (3), 170 (20), 154 (52); HRMS calculated for $\text{C}_{13}\text{H}_{20}\text{NO}_4$ ($M+1$) 254.1392, observed 254.1392.

(5*R, 2'*R**)-5-(Pyrrolidin-2-yl)furan-2(5*H*)-one (9a).** To a stirred solution of **7a** (0.200 g, 0.79 mmol) in CH_2Cl_2 (4 mL) at -25°C was added dropwise TMSOTf (0.14 mL, 0.79 mmol) and stirring was continued 15 min at the same temperature. The solution was allowed to reach -5°C over 2.5 h and was extracted with an aqueous ammonia solution (25%, 10 mL). The aqueous

phase was back-extracted with CH₂Cl₂ (3 x 10 mL), and the combined organic layers were washed with brine (5 mL) and dried (Na₂SO₄). Partial removal of solvent afforded an oil that contained **9a** as a single isomer, together with remaining solvent and silylated residues. Evaporation to dryness removed residual solvent and silyl groups, and provided a 8:1 mixture of **9a** and its epimer (0.129 mg, 84%). Upon further standing at room temperature the sample underwent complete epimerization and eventually degradation: ¹H NMR (250 MHz, CDCl₃) δ 1.46-1.91 (m, 4H), 2.14 (br s, 1H), 2.83-2.97 (m, 2H), 3.21-3.29 (m, 1H), 4.91 (dt, *J* = 6.3, 1.7 Hz, 1H), 6.13 (dd, *J* = 5.7, 1.7 Hz, 1H), 7.42 (dd, *J* = 5.7, 1.7 Hz, 1H); ¹³C NMR (62.9 MHz, CDCl₃, data from a 8:1 epimeric mixture) δ 25.2 (CH₂), 26.7 (CH₂), 46.6 (CH₂), 59.7 (CH), 86.6 (CH), 122.6 (CH), 154.4 (CH), 173.0 (C); IR (neat) ν 3320 (br, N-H), 1755 (C=O), 1640 (C=C) cm⁻¹; MS (EI) *m/z* (%) 153 (M, 6), 152 (10), 130 (13), 97 (6), 83 (6), 70 (base); HRMS calculated for C₈H₁₁NO₂ 153.0790, observed 153.0781.

(5*R, 2'*R**)- and (5*S**, 2'*R**)-5-[*N*-(Boc)Piperidin-2-yl]furan-2(5*H*)-one (7b and 8b).** The procedure described above for the preparation of **7a** was followed starting from **6b** (*n* = 2)^{4,5,8} to afford, after flash column chromatography (75:25 hexanes/EtOAc), a 5:1 mixture of **7b**⁷ and its epimer **8b** (1.40 g, 82%) as a white solid mass. HPLC separation (85:15 hexanes/EtOAc, 9 mL/min) afforded the individual isomers. The major isomer **7b** (*t_R* = 18 min) has been previously described.⁷ Data for **8b** (mixture of rotamers): *t_R* = 12 min; ¹H NMR (250 MHz, CDCl₃) δ 1.39 (s, 9H), 1.46-1.64 (m, 5H), 1.80-2.02 (m, 1H), 2.66-3.06 (br, 1H), 3.86-4.28 (br, 2H), 5.22 (dt, *J* = 8.5, 1.7 Hz, 1H), 6.08 (dd, *J* = 5.8, 2.0 Hz, 1H), 7.39 (br d, *J* = 5.4 Hz, 1H); ¹³C NMR (62.9 MHz, CDCl₃) δ 19.2 (CH₂), 24.7 (CH₂), 25.2 (CH₂), 28.2 (CH₃), 39.8 (CH₂), 41.1 (CH₂), 52.1 (CH), 53.8 (CH), 80.2 (C), 80.5 (CH), 81.9 (CH), 121.2 (CH), 154.4 (C), 155.6 (CH), 172.7 (C); IR (neat) ν 1790 (C=O), 1760 (C=O), 1690 (C=C) cm⁻¹; MS (FAB) *m/z* (%) 268 (M+1, 9), 267

(M, 2), 259 (5), 212 (base), 184 (32), 168 (25), 166 (22); HRMS (FAB) calculated for C₁₄H₂₂NO₄ (M+1) 268.1549, observed 268.1547.

***N*-Boc-2-Ethoxyazepine (6c, n= 3).** Prepared in 83% yield as a colorless oil following the literature procedure (no spectroscopic data given).⁴ Data for the rotamer mixture: ¹H NMR (250 MHz, CDCl₃) δ 0.99-1.75 (m, 19H), 1.09 (t, *J* = 7.0 Hz, included in m at 0.99-1.75), 1.41 (s, included in m at 0.99-1.75), 2.09-2.24 (m, 1H), 2.78-2.95 (m, 1H), 3.22-3.50 (m) and 3.61 (dt, *J* = 14.2, 3.2 Hz) (total 3H), 5.22 and 5.38 (2 t, *J* = 7.9 Hz, 1H); ¹³C NMR (62.9 MHz, CDCl₃) δ 15.0 (CH₃), 15.1 (CH₃), 22.4 (CH₂), 22.6 (CH₂), 27.4 (CH₂), 28.0 (CH₂), 28.2 (CH₃), 28.3 (CH₃), 29.8 (CH₂), 29.9 (CH₂), 34.6 (CH₂), 34.9 (CH₂), 40.2 (CH₂), 40.6 (CH₂), 61.9 (CH₂), 62.4 (CH₂), 79.3 (C), 79.6 (C), 83.5 (CH), 84.1 (CH), 155.1 (C), 156.2 (C); IR (neat) ν 1700 (C=O) cm⁻¹; MS (EI) *m/z* (%) 198 (14), 170 (12), 158 (29), 143 (13), 142 (base), 114 (29), 98 (33); HRMS calculated for C₁₃H₂₅NO₃ 243.1834, observed 243.1841.

(5*R, 2'*R*')-5-[*N*-(Boc)Azepin-2-yl]furan-2(5*H*)-one (7c).** The procedure described above for the preparation of **7a** was followed starting from **6c** to afford, after flash column chromatography (80:20 hexanes/EtOAc), **7c**⁷ (79%) as a white solid (rotamer mixture): mp 114-115 °C (dryness); ¹H NMR (250 MHz, CDCl₃) δ 1.12-1.45 (m, 13H, that includes 2 s at δ 1.38 and 1.45), 1.60-2.16 (m, 4H), 2.72-2.90 (m, 1H), 3.59-3.67 (m) and 3.82 (dt, *J* = 14.5, 2.1 Hz) (total 1H), 4.28 (ddd, *J* = 12.4, 5.9, 3.0 Hz) and 4.34 (ddd, *J* = 12.7, 6.0, 1.6 Hz) (total 1H), 5.01-5.05 (m, 1H), 5.96 (dd, *J* = 5.8, 1.8 Hz) and 6.08 (dd, *J* = 5.8, 2.0 Hz) (total 1H), 7.38 and 7.53 (2 dd, *J* = 5.8, 1.6 Hz, 1H); ¹³C NMR (62.9 MHz, CDCl₃) δ 25.1 (CH₂), 27.9 (CH₃), 28.0 (CH₃), 28.6 (CH₂), 29.0 (CH₂), 29.1 (CH₂), 29.3 (CH₂), 30.4 (CH₂), 30.8 (CH₂), 43.0 (CH₂), 43.6 (CH₂), 55.6 (CH), 56.5 (CH), 79.2 (C), 79.8 (C), 85.2 (CH), 86.1 (CH), 119.7 (CH), 121.2 (CH), 153.9 (CH), 154.4 (C), 155.0 (CH), 155.5 (C), 172.2 (C), 172.9 (C); IR (KBr) 1750 (C=O), 1680 (C=O), 1600 (C=C)

cm⁻¹; MS (FAB) m/z (%) 282 (M+1, 16), 227 (13), 226 (base), 198 (20), 182 (11), 154 (13); HRMS (FAB) calculated for C₁₅H₂₄NO₄ (M+1) 282.1705, observed 282.1703.

3. Preparation of Pyrrolidines 13, 16, 19

(1*R, 5*R**)-7-Benzyl-2-oxa-7-azabicyclo[3.3.0]octan-3-one (13a).** Prepared in 67% yield from **4** and hydroxymethylbenzotriazole using the General Reductive Cyclization Procedure. Eluted with 78:20:2 hexanes/EtOAc/Et₃N. Oil: ¹H NMR (250 MHz, CDCl₃) δ 2.25-2.34 (m, 2H, H-6, H-8), 2.40 (dd, *J* = 18.3, 3.9 Hz, 1H, H-4), 2.67-2.82 (m, 2H, H-4, H-6), 2.91-3.03 (m, 1H, H-5), 3.09 (d, *J* = 11.1 Hz, 1H, H-8), 3.55 and 3.63 (AB q, *J* = 13.1 Hz, 2H), 4.96 (dd, *J* = 7.5, 4.9 Hz, 1H, H-1), 7.19-7.33 (m, 5H); ¹³C NMR (62.9 MHz, CDCl₃) δ 34.9 (CH₂), 36.6 (CH), 58.4 (CH₂), 59.8 (CH₂), 60.0 (CH₂), 83.0 (CH), 126.9 (CH), 128.2 (CH), 128.2 (CH), 138.1 (C), 177.4 (C); IR (neat) ν 1770 (C=O) cm⁻¹; MS (EI) m/z (%) 217 (29), 216 (19), 140 (29), 133 (13), 132 (11), 126 (30), 92 (38), 91 (base); HRMS calculated for C₁₃H₁₅NO₂ 217.1103, observed 217.1106.

(1*R, 2*R**, 6*R**)-3-Oxa-8-azatricyclo[6.4.0.0^{2,6}]dodecan-4-one (13c).** Prepared from **7b** and hydroxymethylbenzotriazole using the General Reductive Cyclization Procedure. Also isolated in this reaction was **13d**, resulting from epimerization of the intermediate amine **9b**. The extent of epimerization was estimated in at least 10% from the crude NMR spectrum. The isolated yield of **13c** corrected for epimerization was 61% (three steps from **7b**). Eluted with 78:20:2 hexanes/EtOAc/Et₃N. Data for **13c**: Oil; ¹H NMR (250 MHz, CDCl₃) δ 1.09-1.28 (m, 1H), 1.39-1.63 (m, 3H), 1.71-1.95 (m, 4H), 2.23 (dd, *J* = 9.3, 6.5 Hz, 1H, H-7), 2.38 (dd, *J* = 17.1, 2.7 Hz, 1H, H-5), 2.74-2.94 (m, 3H, H-5, H-6, H-7), 3.04 (dt, *J* = 10.7, 2.7 Hz, 1H, H-9), 4.80 (dd, *J* = 7.1, 4.2 Hz, 1H, H-2); ¹³C NMR (62.9 MHz, CDCl₃) δ 23.8 (CH₂), 24.8 (CH₂), 25.1 (CH₂), 34.9

(CH), 35.6 (CH₂), 52.7 (CH₂), 61.7 (CH₂), 67.8 (CH), 84.7 (CH), 177.7 (C); IR (neat) ν 1780 (C=O) cm⁻¹; MS (EI) m/z (%) 181 (M, 14), 180 (11), 121 (24), 120 (39), 117 (19), 97 (base), 93 (14); HRMS calculated for C₁₀H₁₅NO₂ 181.1103, observed 181.1102.

(1*S, 2*R**, 6*R**)-3-Oxa-8-azatricyclo[6.4.0.0^{2,6}]dodecan-4-one (13d)**. Prepared from **8b** and hydroxymethylbenzotriazole using the General Reductive Cyclization Procedure. Also isolated in this reaction was **13c**, resulting from epimerization of the intermediate amine **10b**. The extent of epimerization was estimated in at least 25% from the crude NMR spectrum. The isolated yield of **13d** corrected for epimerization was 66% (three steps from **8b**). Eluted with 78:20:2 hexanes/EtOAc/Et₃N. Data for **13d**: Oil; ¹H NMR (250 MHz, CDCl₃) δ 1.10-2.13 (m, 9H), 2.34 (dd, J = 18.6, 4.2 Hz, 1H, H-5), 2.73 (dd, J = 18.6, 10.9 Hz, 1H, H-5), 2.89-3.06 (m, 2H, H-6, H-9), 3.40 (t, J = 8.5 Hz, 1H, H-7), 4.52 (dd, J = 8.9, 6.1 Hz, 1H, H-2); ¹³C NMR (62.9 MHz, CDCl₃) δ 23.9 (CH₂), 24.6 (CH₂), 28.8 (CH₂), 32.9 (CH₂), 34.7 (CH), 52.4 (CH₂), 61.1 (CH₂), 70.0 (CH), 87.0 (CH), 177.3 (C); IR (neat) ν 1780 (C=O) cm⁻¹; MS (EI) m/z (%) 181 (M, 79), 180 (52), 124 (27), 98 (35), 97 (base), 96 (64); HRMS calculated for C₁₀H₁₅NO₂ 181.1103, observed 181.1101.

(1*R, 2*R**, 6*R**)-3-Oxa-8-azatricyclo[6.5.0.0^{2,6}]tridecan-4-one (13e)**. Prepared in 54% overall yield (three steps) from **7c** and hydroxymethylbenzotriazole using the General Reductive Cyclization Procedure. Eluted with 88:10:2-78:20:2 hexanes/EtOAc/Et₃N gradient. Oil: ¹H NMR (250 MHz, CDCl₃) δ 1.45-1.87 (m, 8H), 2.08-2.18 (m, 2H), 2.38-2.47 (m, 2H), 2.72-3.00 (m, 4H), 4.79 (dd, J = 7.1, 4.8 Hz, 1H, H-2); ¹³C NMR (62.9 MHz, CDCl₃) δ 23.9 (CH₂), 26.5 (CH₂), 26.5 (CH₂), 27.8 (CH₂), 35.3 (CH), 35.4 (CH₂), 54.0 (CH₂), 62.7 (CH₂), 70.1 (CH), 86.8 (CH), 177.5 (C); IR (neat) ν 1750 (C=O) cm⁻¹; MS (EI) m/z (%) 195 (M, 52), 194 (32), 180 (7), 167

(6), 166 (62), 153 (63), 152 (28), 134 (42), 122 (78), 111 (base), 96 (77); HRMS calculated for $C_{11}H_{17}NO_2$ 195.1259, observed 195.1258.

7-Benzyl-6-propyl-2-oxa-7-azabicyclo[3.3.0]octan-3-one (16a). Prepared in 71% overall yield from **4** and butanal using the General Reductive Cyclization Procedure. Also obtained in this reaction was **17a** (6%, characterization data given in tetrahydropyridines section). Eluted with 88:10:2 hexanes/EtOAc/Et₃N. Data for the major isomer (**1R***, **5R***, **6S***)-**16a**: Oil; ¹H NMR (250 MHz, CDCl₃) δ 0.94-1.00 (distorted t, $J \approx 6.9$ Hz, 3H), 1.22-1.41 (m, 3H), 1.69-1.84 (m, 1H), 2.11 (dd, $J = 11.5, 5.6$ Hz, 1H, H-8), 2.34 (dd, $J = 9.7, 5.8$ Hz, 1H, H-6), 2.49 (dd, $J = 18.3, 10.8$ Hz, 1H, H-4), 2.69 (dd, $J = 18.3, 5.7$ Hz, 1H, H-4), 2.95-3.09 (m, 3H, H-5, PhCH, H-8), 3.02 (d, $J = 13.3$ Hz, PhCH, included in m at 2.95-3.09), 3.06 (d, $J = 11.5$ Hz, H-8, included in m at 2.95-3.09), 4.06 (d, $J = 13.3$ Hz, 1H, PhCH), 4.88 (dd, $J = 7.9, 5.4$ Hz, 1H, H-1), 7.19-7.33 (m, 5H); ¹³C NMR (62.9 MHz, CDCl₃) δ 14.3 (CH₃), 19.4 (CH₂), 28.2 (CH₂), 30.7 (CH₂), 40.1 (CH), 56.1 (CH₂), 59.9 (CH₂), 65.9 (CH), 80.9 (CH), 126.9 (CH), 128.2 (CH), 128.3 (CH), 138.2 (C), 177.7 (C); IR (neat) ν 1770 (C=O) cm⁻¹; MS (EI) m/z (%) 259 (M, 3), 217 (14), 216 (94), 91 (base); HRMS calculated for $C_{16}H_{21}NO_2$ 259.1572, observed 259.1564. Data for the minor isomer (**1R***, **5R***, **6R***)-**16a** (from a 3:1 mixture with the major diastereoisomer): Oil; ¹H NMR (250 MHz, CDCl₃) δ 0.94-1.00 (apparent t, 3H), 1.25-1.49 (m, 3H), 1.70-1.81 (m, 1H), 2.31-2.56 (m, 3H), 2.64-2.82 (m, 2H), 3.25-3.30 (m, 2H), 3.99 (d, $J = 13.1$ Hz, 1H), 4.87-4.94 (m, 1H), 7.22-7.36 (m, 5H); ¹³C NMR (62.9 MHz, CDCl₃) δ 14.5 (CH₃), 18.6 (CH₂), 32.7 (CH₂), 33.5 (CH₂), 42.9 (CH), 56.2 (CH₂), 58.8 (CH₂), 69.1 (CH), 81.6 (CH), 127.2 (CH), 128.4 (CH), 128.6 (CH), 138.3 (C), 177.1 (C); IR (neat) ν 1760 (C=O) cm⁻¹; GC/MS t_R = 16.7 min, MS (EI) m/z (%) 217 (59), 216 (81), 170 (20), 92 (38), 91 (base), HRMS calculated for $C_{16}H_{21}NO_2$

259.1572, observed 259.1577; t_R = 17.0 min, MS (EI) m/z (%) 217 (79), 216 (base), 160 (10), 92 (54), 91 (base), HRMS calculated for $C_{16}H_{21}NO_2$ 259.1572, observed 259.1576.

7-Benzyl-6-[(3,4-dimethoxyphenyl)ethyl]-2-oxa-7-azabicyclo[3.3.0]octan-3-one (16b).

Prepared in 84% overall yield from **4** and 3-(3,4-dimethoxyphenyl)propanal⁹ following the General Reductive Cyclization Procedure, where benzene was used as substitute for CH_2Cl_2 in the first step. The crude product was purified by flash column chromatography (the major isomer eluted with 65:35 hexanes/EtOAc and the minor isomer with 60:40 hexanes/EtOAc). Data for the major isomer (**1R***, **5R***, **6S***)-**16b**: Oil; 1H NMR (500 MHz, $CDCl_3$) δ 1.61-1.69 (m, 1H), 2.11-2.18 (m, 2H), 2.13 (dd, J = 11.4, 5.6 Hz, H-8, included in m at 2.11-2.18), 2.36-2.40 (m, 1H, H-6), 2.48-2.63 (m, 3H, 2 H-2', H-4), 2.53 (dd, J = 18.3, 11.2 Hz, H-4, included in m at 2.48-2.63), 2.75 (dd, J = 18.3, 5.5 Hz, 1H, H-4), 3.02-3.08 (m, 3H, H-5, H-8, PhCH), 3.85 and 3.88 (2 s, 6H, CH_3O), 4.04 (d, J = 13.2 Hz, 1H, PhCH), 4.88 (dd, J = 7.6, 5.6 Hz, 1H, H-1), 6.69-6.81 (m, 3H), 7.20-7.28 (m, 5H); ^{13}C NMR (125.8 MHz, $CDCl_3$) δ 28.3 (CH_2 , C-4), 30.4 (CH_2), 32.0 (CH_2), 40.1 (CH, C-5), 55.8 (CH_3), 55.8 (CH_3), 56.0 (CH_2), 59.8 (CH_2 , C-8), 65.6 (CH, C-6), 80.8 (CH, C-1), 111.2 (CH), 111.3 (CH), 119.8 (CH), 126.9 (CH), 128.2 (CH), 128.2 (CH), 133.6 (C), 138.0 (C), 147.3 (C), 148.8 (C), 177.4 (C); IR (neat) ν 1780 (C=O) cm^{-1} ; MS (EI) m/z (%) 381 (M, 6), 243 (93), 230 (60), 228 (16), 217 (base), 152 (100), 151 (100), 146 (16), 137 (38), 126 (87), 91 (100); HRMS calculated for $C_{23}H_{27}NO_4$ 381.1940, observed 381.1951. Data for the minor isomer (**1R***, **5R***, **6R***)-**16b**: Oil; 1H NMR (500 MHz, $CDCl_3$) δ 1.75-1.77 (m, 1H), 2.04-2.10 (m, 1H), 2.41-2.58 (m, 4H, H-4, H-6, H-8, H-2'), 2.66-2.72 (m, 1H), 2.75-2.82 (m, 2H, H-4, H-5), 3.29 (dd, J = 10.6, 6.5 Hz, 1H, H-8), 3.34 (d, J = 13.0 Hz, 1H, PhCH), 3.86 and 3.87 (2 s, 6H, CH_3O), 4.00 (d, J = 13.0 Hz, 1H, PhCH), 4.92-4.95 (m, 1H, H-1), 6.68-6.81 (m, 3H), 7.24-7.33 (m, 5H); ^{13}C NMR (125.8 MHz, $CDCl_3$) δ 31.1 (CH_2), 32.0 (CH_2), 33.6 (CH_2 , C-4), 42.8

(CH, C-5), 55.8 (CH₃), 55.9 (CH₃), 56.2 (CH₂), 58.7 (CH₂, C-8), 68.9 (CH, C-6), 81.5 (CH, C-1), 111.3 (CH), 111.4 (CH), 119.9 (CH), 127.2 (CH), 128.4 (CH), 128.6 (CH), 134.0 (C), 138.1 (C), 147.4 (C), 149.0 (C), 176.8 (C); IR (neat) ν 1775 (C=O) cm⁻¹; MS (EI) m/z (%) 217 (2), 216 (94), 164 (8), 152 (3), 149 (2), 91 (base); HRMS calculated for C₂₃H₂₇NO₄ 381.1940, observed 381.1929.

7-Benzyl-6-(4'-methoxycarbonylbutyl)-2-oxa-7-azabicyclo[3.3.0]non-3-one (16c). Prepared in 66% overall yield from **4** and methyl 6-oxohexanoate using the General Reductive Cyclization Procedure. Also obtained in this reaction was **17c** (6%, characterization data given in tetrahydropyridines section). Eluted with 73:25:2 hexanes/EtOAc/Et₃N. Data for the major isomer (**1R***, **5R***, **6S***)-**16c**: Oil; ¹H NMR (500 MHz, CDCl₃) δ 1.22-1.40 (m, 3H), 1.60-1.77 (m, 2H), 1.80-1.85 (m, 1H), 2.12 (dd, J = 11.6, 5.5 Hz, 1H, H-8), 2.28-2.39 (m, 3H, H-4', H-6), 2.50 (dd, J = 18.3, 10.9 Hz, 1H, H-4), 2.67 (dd, J = 18.3, 5.7 Hz, 1H, H-4), 2.99-3.09 (m, 3H, H-5, PhCH, H-8), 3.03 (d, J = 12.9 Hz, PhCH, included in m at 2.99-3.09), 3.08 (d, J = 11.6 Hz, H-8, included in m at 2.99-3.09), 3.68 (s, 3H, CH₃O), 4.04 (d, J = 12.9 Hz, 1H, PhCH), 4.89 (dd, J = 7.8, 5.6 Hz, 1H, H-1), 7.22-7.31 (m, 5H); ¹³C NMR (125.8 MHz, CDCl₃) δ 25.0 (CH₂), 25.7 (CH₂), 28.3 (CH₂), 28.3 (CH₂, C-4), 33.7 (CH₂), 40.2 (CH, C-5), 51.6 (CH₃), 56.2 (CH₂), 60.1 (CH₂, C-8), 66.0 (CH, C-6), 80.8 (CH, C-1), 127.0 (CH), 128.3 (CH), 128.4 (CH), 138.2 (C), 173.8 (C), 177.5 (C); IR (neat) ν 1770 (C=O), 1730 (C=O) cm⁻¹; MS (EI) m/z (%) 331 (M, 3), 300 (2), 217 (15), 216 (base), 92 (8), 91 (97); HRMS calculated for C₁₉H₂₅NO₄ 331.1784, observed 331.1776. Data for the minor isomer (**1R***, **5R***, **6R***)-**16c** (from diastereomeric mixtures enriched in the minor isomer): Oil; ¹H NMR (250 MHz, CDCl₃) δ 1.25-1.76 (m, 6H), 2.31-2.48 (m, 5H), 2.63-2.82 (m, 2H), 3.24-3.31 (m, 2H), 3.67 (s, 3H), 3.96 (d, J = 13.1 Hz, 1H), 4.90 (dd, J = 11.1, 6.7 Hz, 1H), 7.26-7.31 (m, 5H); ¹³C NMR (62.9 MHz, CDCl₃) δ 24.6 (CH₂),

25.2 (CH₂), 29.8 (CH₂), 33.4 (CH₂), 33.8 (CH₂), 42.7 (CH), 51.6 (CH₃), 56.2 (CH₂), 58.8 (CH₂), 68.9 (CH), 81.5 (CH), 127.2 (CH), 128.4 (CH), 128.6 (CH), 138.0 (C), 173.9 (C), 177.0 (C); IR (neat) ν 1770 (C=O), 1730 (C=O) cm⁻¹; GC/MS t_R = 17.3 min, MS (EI) m/z (%) 217 (5), 216 (37), 92 (7), 91 (base); HRMS calculated for C₁₉H₂₆NO₄ (M + 1) 332.1862, observed 332.1871; t_R = 18.0 min, MS (EI) m/z (%) 217 (4), 216 (36), 96 (1), 92 (6), 91 (base); HRMS calculated for C₁₉H₂₆NO₄ (M + 1) 332.1862, observed 332.1871.

7-Benzyl-6-(1'-methylethyl)-2-oxa-7-azabicyclo[3.3.0]octan-3-one (16d). Prepared in 84% overall yield from **4** and isobutanal using the General Reductive Cyclization Procedure. Eluted with 83:15:2 hexanes/EtOAc/Et₃N. Data for the major isomer (**1R***, **5R***, **6S***)-**16d**: Oil; ¹H NMR (250 MHz, CDCl₃) δ 1.09 (apparent t, $J \approx 6.8$ Hz, 6H), 2.07-2.21 (m, 2H), 2.16 (dd, $J = 11.7, 6.1$ Hz, H-8, included in m at 2.07-2.21), 2.30 (t, $J = 5.6$ Hz, 1H, H-6), 2.52 (dd, $J = 17.1, 9.4$ Hz, 1H, H-4), 2.90-3.09 (m, 4H, PhCH, H-4, H-5, H-8), 4.15 (d, $J = 12.9$ Hz, 1H, PhCH), 4.87 (t, $J = 6.8$ Hz, 1H, H-1), 7.20-7.32 (m, 5H); ¹³C NMR (62.9 MHz, CDCl₃) δ 18.3 (CH₃), 20.4 (CH₃), 26.9 (CH), 30.1 (CH₂), 40.6 (CH), 56.8 (CH₂), 59.9 (CH₂), 70.5 (CH), 80.6 (CH), 126.9 (CH), 128.1 (CH), 128.2 (CH), 138.6 (C), 177.8 (C); IR (neat) ν 1770 (C=O) cm⁻¹; MS (EI) m/z (%) 259 (4), 258 (6), 217 (97), 216 (94), 174 (3), 160 (5), 108 (7), 91 (base); HRMS calculated for C₁₆H₂₁NO₂ 259.1572, observed 259.1562. Data for the minor isomer (**1R***, **5R***, **6R***)-**16d** (from a 36:64 mixture with the major isomer): Oil; ¹H NMR (250 MHz, CDCl₃) δ 0.97 (d, $J = 6.7$ Hz, 6H), 2.07-2.58 (m, 4H), 2.67-2.81 (m, 2H), 2.90-3.17 (m, 1H), 3.38 (dd, $J = 10.5, 6.7$ Hz, 1H), 4.03 (d, $J = 12.9$ Hz, 1H), 4.79-4.89 (m, 1H), 7.22-7.31 (m, 5H); ¹³C NMR (62.9 MHz, CDCl₃) δ 15.2 (CH₃), 19.9 (CH₃), 27.1 (CH), 34.2 (CH₂), 37.4 (CH), 56.9 (CH₂), 59.4 (CH₂), 74.7 (CH), 81.0 (CH), 127.0 (CH), 128.2 (CH), 128.4 (CH), 128.6 (CH), 138.3 (C), 177.1 (C); IR (neat) ν 1770 (C=O) cm⁻¹; GC-MS t_R = 13.5 min, MS (EI) m/z (%) 216 (92), 199 (4), 184

(10), 105 (7), 91 (base), HRMS calculated for $C_{13}H_{14}NO_2$ (M-*i*-Pr) 216.1025, observed 216.1016; t_R = 13.8 min, MS (EI) m/z (%) 217 (2), 216 (base), 199 (2), 184 (3), 105 (4), 91 (base), HRMS calculated for $C_{13}H_{14}NO_2$ (M-*i*-Pr) 216.1025, observed 216.1031; HRMS (mixture) calculated for $C_{16}H_{21}NO_2$ 259.1572, observed 259.1576.

7-Benzyl-6-phenyl-2-oxa-7-azabicyclo[3.3.0]octan-3-one (16e). Prepared in 44% overall yield from **4** and benzaldehyde using the General Reductive Cyclization Procedure with the following modification. The first step was carried out in THF. After filtration of the sieves and evaporation, the crude product was redissolved in THF, one equivalent each of benzaldehyde and benzotriazole were added, together with the appropriate amount of 4Å MS, and the mixture was stirred 22 h at room temperature. The General Reductive Cyclization Procedure was then followed as described above. The isomers were separated by flash column chromatography (silica gel saturated with Et_3N , 88:10:2 hexanes/EtOAc/ Et_3N). The minor isomer was further purified on silica gel with 80:20 hexanes/EtOAc as eluent. The pure samples were finally obtained by precipitation with hexanes. Data for the major isomer (**1R***, **5R***, **6S***)-**16e**: mp 116-118 °C (dryness); 1H NMR (250 MHz, $CDCl_3$) δ 2.42-2.50 (m, 2H, H-4, H-8), 2.63 (ddd, J = 18.2, 8.9, 1.0 Hz, 1H, H-4), 2.77-2.88 (m, 1H, H-5), 3.09 (d, J = 13.0 Hz, 1H, PhCH), 3.22 (d, J = 8.7 Hz, 1H, H-6), 3.60 (dd, J = 11.1, 6.7 Hz, 1H, H-8), 3.78 (d, J = 13.0 Hz, 1H, PhCH), 5.02 (td, J = 7.0, 4.0 Hz, 1H, H-1), 7.21-7.46 (m, 10H); ^{13}C NMR (62.9 MHz, $CDCl_3$) δ 31.6 (CH_2), 48.1 (CH), 56.6 (CH_2), 58.9 (CH_2), 74.1 (CH), 81.4 (CH), 127.2 (CH), 127.7 (CH), 128.3 (CH), 128.6 (CH), 129.0 (CH), 137.9 (C), 139.5 (C), 176.5 (C); IR (neat) ν 1770 (C=O) cm^{-1} ; MS (EI) m/z (%) 293 (M, 26), 292 (15), 216 (26), 202 (38), 194 (11), 118 (44), 117 (15), 91 (base); HRMS calculated for $C_{19}H_{19}NO_2$ 293.1416, observed 293.1414. Data for the minor isomer (**1R***, **5R***, **6R***)-**16e**: mp 120-122 °C (dryness); 1H NMR (250 MHz, $CDCl_3$) δ 2.25-2.37 (m, 3H, 2 H-4, H-8), 2.34

(dd, $J = 11.5, 4.8$ Hz, H-8, included in m at 2.25-2.37), 3.04 (d, $J = 13.7$ Hz, 1H, PhCH), 3.19 (quint, $J = 7.3$ Hz, 1H, H-5), 3.37 (d, $J = 11.5$ Hz, 1H, H-8), 3.70 (d, $J = 7.5$ Hz, 1H, H-6), 4.02 (d, $J = 13.7$ Hz, 1H, PhCH), 5.03 (dd, $J = 7.3, 4.8$ Hz, 1H, H-1), 7.22-7.44 (m, 10H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 30.2 (CH_2), 42.7 (CH), 56.7 (CH_2), 58.9 (CH_2), 71.0 (CH), 81.6 (CH), 127.1 (CH), 127.9 (CH), 128.2 (CH), 128.3 (CH), 128.4 (CH), 128.9 (CH), 137.5 (C), 137.7 (C), 177.3 (C); IR (neat) ν 1770 ($\text{C}=\text{O}$) cm^{-1} ; MS (EI) m/z (%) 293 (M, 32), 292 (15), 216 (14), 208 (10), 202 (27), 194 (14), 118 (55), 117 (10), 92 (16), 91 (base); HRMS calculated for $\text{C}_{19}\text{H}_{19}\text{NO}_2$ 293.1416, observed 293.1408.

(1*R, 2*R**, 6*R**, 7*S**)-7-Propyl-3-oxa-8-azatricyclo[6.3.0.0^{2,6}]undecan-4-one (19).** Prepared in 40% overall yield from **9a** and butanal using the General Reductive Cyclization Procedure with the following modification. After treatment with SmI_2 , the reaction mixture was poured over an aqueous ammonia solution (25%, 25 mL). After separation, the aqueous layer was extracted with EtOAc (4 x 50 mL). The combined organic layers were washed with brine (10 mL) and aqueous ammonia solution (25%, 5 mL), and were dried (Na_2SO_4). Flash chromatography (silica gel saturated with NH_4OH , 98:1:1 $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$) afforded **19** as an oil: ^1H NMR (500 MHz, CDCl_3) δ 0.94 (t, $J = 7.2$ Hz, 3H, CH_3), 1.31-1.45 (m, 3H), 1.62-1.70 (m, 1H), 1.79-2.03 (m, 4H, H-10, H-11), 2.40 (td, $J = 8.3, 4.7$ Hz, 1H, H-9), 2.52-2.60 (m, 3H, 2 H-5, H-9), 2.79-2.84 (m, 1H, H-7), 2.92 (ddd, $J = 7.4, 5.8, 4.5$ Hz, 1H, H-1), 3.07 (td, $J = 15.5, 7.2$ Hz, 1H, H-6), 4.80 (dd, $J = 7.2, 4.5$ Hz, 1H, H-2); ^{13}C NMR (62.9 MHz, CDCl_3) δ 14.3 (CH_3), 20.3 (CH_2), 21.6 (CH_2), 25.9 (CH_2), 29.6 (CH_2 , C-5), 31.0 (CH_2), 43.7 (CH, C-6), 46.4 (CH_2 , C-9), 62.0 (CH, C-7), 70.8 (CH, C-1), 83.8 (CH, C-2), 177.4 (C); IR (neat) ν 1770 ($\text{C}=\text{O}$) cm^{-1} ; MS (EI) m/z (%) 209 (M, 11), 167 (9), 166 (81), 125 (43), 110 (14), 96 (base); HRMS calculated for $\text{C}_{12}\text{H}_{19}\text{NO}_2$ 209.1416, observed 209.1416.

4. Preparation of Tetrahydropyridines 17

(1*R, 5*R**)-8-Benzyl-6-ethyl-2-oxa-8-azabicyclo[3.4.0]non-6-en-3-one (17a).** Prepared from **4** and butanal using the General Procedure. The crude product was purified by flash column chromatography (85:15 hexanes/EtOAc) to yield **17a** (73%) as a solid: mp 80-81 °C (dryness); ¹H NMR (250 MHz, CDCl₃) δ 1.00 (t, *J* = 7.4 Hz, 3H), 1.85-2.04 (m, 2H, CH₂CH₃), 2.36 (dd, *J* = 17.4, 6.2 Hz, 1H, H-4), 2.74 (dd, *J* = 17.4, 8.5 Hz, 1H, H-4), 2.83-3.04 (m, 3H), 3.99 (s, 2H), 4.71-4.78 (m, 1H, H-1), 5.93 (s, 1H, H-7), 7.24-7.37 (m, 5H); ¹³C NMR (62.9 MHz, CDCl₃) δ 12.8 (CH₃), 25.2 (CH₂), 34.3 (CH), 34.6 (CH₂), 47.8 (CH₂), 59.1 (CH₂), 77.4 (CH), 109.9 (C), 127.2 (CH), 128.0 (CH), 128.4 (CH), 131.9 (CH), 137.4 (C), 176.6 (C); IR (neat) ν 1780 (C=O), 1660 (C=C) cm⁻¹; MS (EI) *m/z* (%) 257 (M, 16), 243 (2), 242 (68), 106 (2), 92 (4), 91 (base); HRMS calculated for C₁₆H₁₉NO₂ 257.1416, observed 257.1425.

(1*R, 5*R**)-8-Benzyl-6-[(3,4-dimethoxyphenyl)methyl]-2-oxa-8-azabicyclo[3.4.0]non-6-en-3-one (17b).** Prepared from **4** and 3-(3,4-dimethoxyphenyl)propanal⁹ following the General Procedure. The crude product **17b** (~ 100% yield) was used without further purification for subsequent reactions: Oil; ¹H NMR (250 MHz, CDCl₃) δ 2.30 (dd, *J* = 17.4, 6.1 Hz, 1H, H-4), 2.56 (dd, *J* = 17.4, 8.5 Hz, 1H, H-4), 2.76-2.85 (m, 1H, H-5), 2.86-2.98 (m, 2H, H-9), 3.17 and 3.22 (AB q, *J* = 15.3 Hz, 2H), 3.84 and 3.86 (2 s, 6H, CH₃O), 4.01 and 4.08 (AB q, *J* = 14.7 Hz, 2H), 4.66-4.73 (m, 1H, H-1), 6.05 (s, 1H, H-7), 6.67-6.80 (m, 3H), 7.25-7.38 (m, 5H); ¹³C NMR (62.9 MHz, CDCl₃) δ 34.1 (CH), 34.6 (CH₂), 38.9 (CH₂), 47.7 (CH₂), 55.8 (CH₃), 55.8 (CH₃), 59.1 (CH₂), 77.3 (CH), 107.9 (C), 111.0 (CH), 111.6 (CH), 120.5 (CH), 127.5 (CH), 128.0 (CH), 128.5 (CH), 132.3 (C), 134.0 (CH), 137.2 (C), 147.5 (C), 148.9 (C), 176.6 (C); IR (neat) ν 1770

(C=O), 1650 (C=C) cm^{-1} ; MS (EI) m/z (%) 379 (M, 89), 294 (8), 229 (10), 151 (10), 91 (base); HRMS calculated for $\text{C}_{23}\text{H}_{25}\text{NO}_4$ 379.1784, observed 379.1776.

(1*R, 5*R**)-8-Benzyl-6-(3-methoxycarbonylpropyl)-2-oxa-8-azabicyclo[3.4.0]non-6-en-3-one (17c).** Prepared from **4** and methyl 6-oxohexanoate following the General Procedure. The crude product was purified by flash column chromatography (silica gel saturated with Et_3N , 73:25:2 hexanes/ EtOAc / Et_3N) to yield **17c** (78%) as an oil: ^1H NMR (250 MHz, CDCl_3) δ 1.55-2.08 (m, 4H), 2.27-2.41 (m, 3H, 2 H-3', H-4), 2.74 (dd, $J = 17.3, 8.5$ Hz, 1H, H-4), 2.92 (d, $J = 4.9$ Hz, 2H, H-9), 2.94-3.03 (m, 1H, H-5, overlapped with signals of H-9), 3.67 (s, 3H, CH_3O), 4.00 (s, 2H), 4.74 (dt, $J = 6.7, 4.9$ Hz, 1H, H-1), 5.95 (s, 1H, H-7), 7.22-7.37 (m, 5H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 23.2 (CH_2), 31.8 (CH_2), 33.2 (CH_2), 34.0 (CH), 34.6 (CH_2), 47.7 (CH_2), 51.4 (CH_3), 59.0 (CH_2), 77.3 (CH), 106.3 (C), 127.4 (CH), 128.0 (CH), 128.4 (CH), 133.5 (C), 137.3 (CH), 173.9 (C), 176.5 (C); IR (neat) ν 1770 (C=O), 1730 (C=O), 1650 (C=C) cm^{-1} ; MS (EI) m/z (%) 329 (M, 29), 298 (6), 243 (16), 242 (86), 92 (10), 91 (base); HRMS calculated for $\text{C}_{19}\text{H}_{23}\text{NO}_4$ 329.1627, observed 329.1628.

(1*R, 5*R**)-(*E*)-8-Benzyl-6-(4-ethoxycarbonylbut-3-en-1-yl)-2-oxa-8-azabicyclo[3.4.0]non-6-en-3-one (17d).** Prepared from **4** and ethyl 7-oxohept-2-enoate¹⁰ following the General Procedure at 50 °C (oil bath temperature) for 14 h. The crude product was purified by flash column chromatography (silica gel saturated with Et_3N , 70:28:2 hexanes/ EtOAc / Et_3N) to yield **17d** (81%) as an oil: ^1H NMR (250 MHz, CDCl_3) δ 1.28 (t, $J = 7.1$ Hz, 3H), 2.02–2.39 (m, 5H), 2.35 (dd, $J = 17.4, 5.4$ Hz, H-4, included in m at 2.02-2.39), 2.73 (dd, $J = 17.4, 8.3$ Hz, 1H, H-4), 2.90-2.99 (m, 3H, H-5, 2 H-9), 3.97 and 4.02 (AB q, $J = 15.1$ Hz, 2H, CH_2Ph), 4.18 (q, $J = 7.1$ Hz, 2H), 4.73 (dt, $J = 6.7, 4.9$ Hz, 1H, H-1), 5.81 (dd, $J = 14.9, 1.6$ Hz, 1H), 5.91 (s, 1H, H-7), 6.93 (dt, $J = 14.9, 6.7$ Hz, 1H), 7.21–7.36 (m, 5H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 14.1 (CH_3),

30.8 (CH₂), 34.2 (CH), 34.5 (CH₂), 47.5 (CH₂), 59.0 (CH₂), 60.1 (CH₂), 77.3 (CH), 105.7 (C), 121.7 (CH), 127.4 (CH), 127.9 (CH), 128.4 (CH), 133.6 (CH), 137.2 (C), 148.1 (CH), 166.3 (C), 176.3 (C); IR (neat) ν 1720 (C=O), 1710(C=O), 1660 (C=C) cm⁻¹; MS (EI) m/z (%) 355 (M, 1), 243 (17), 242 (base), 91 (81); HRMS calculated for C₂₁H₂₅NO₄ 355.1784, found 355.1779.

(1*R, 2*R**, 6*R**)-7-Ethyl-3-Oxa-9-azatricyclo[7.3.0.0^{2,6}]dodec-7-en-4-one (17e).** Prepared from **9a** and butanal following the General Procedure. The crude product was purified by flash column chromatography (silica gel saturated with Et₃N, 83:15:2 hexanes/EtOAc/Et₃N) to yield **17e** (42%) as an oil: ¹H NMR (250 MHz, CDCl₃) δ 1.00 (t, *J* = 7.4 Hz, 3H, CH₃), 1.72-2.19 (m, 6H), 2.50 (d, *J* = 17.1 Hz, 1H, H-5), 2.77 (dd, *J* = 17.1, 7.7 Hz, 1H, H-5), 3.03-3.19 (m, 4H), 4.72 (dd, *J* = 5.9, 1.6 Hz, 1H, H-2), 6.05 (s, 1H, H-8); ¹³C NMR (62.9 MHz, CDCl₃) δ 13.0 (CH₃), 23.5 (CH₂), 24.1 (CH₂), 26.9 (CH₂), 34.7 (CH₂), 36.0 (CH), 50.1 (CH₂), 56.7 (CH), 81.0 (CH), 108.0 (C), 129.0 (CH), 177.0 (C); IR (neat) ν 1770 (C=O), 1650 (C=C) cm⁻¹; MS (EI) m/z (%) 237 (M, 20), 206 (4), 193 (13), 192 (base); HRMS calculated for C₁₂H₁₇NO₂ 207.1259, observed 207.1253.

(1*R, 2*R**, 6*R**)-7-Ethyl-3-oxa-9-azatricyclo[7.4.0.0^{2,6}]tridec-7-en-4-one (17f).** Prepared from **7b** and butanal following the General Procedure. Also observed in this reaction was a product derived from epimerization of the intermediate amine **9b**. The extent of epimerization was estimated in at least 10% from the crude NMR spectrum. The crude product was purified by flash column chromatography (silica gel saturated with Et₃N, 88:10:2 hexanes/EtOAc/Et₃N) to yield **17f** (42%, corrected for epimerization) as an oil: ¹H NMR (250 MHz, CDCl₃) δ 0.97 (t, *J* = 7.4 Hz, 3H, CH₃), 1.18-1.37 (m, 1H), 1.48-2.01 (m, 7H), 2.41-2.53 (m, 3H), 2.67 (dd, *J* = 17.3, 8.0 Hz, 1H, H-5), 3.03-3.11 (m, 2H), 4.54 (dd, *J* = 5.7, 1.6 Hz, 1H, H-2), 5.63 (s, 1H, H-8); ¹³C

NMR (62.9 MHz, CDCl₃) δ 12.7 (CH₃), 24.1 (CH₂), 24.3 (CH₂), 25.3 (CH₂), 28.1 (CH₂), 33.6 (CH₂), 35.0 (CH), 53.6 (CH₂), 57.3 (CH), 81.8 (CH), 109.3 (C), 133.6 (CH), 176.6 (C); IR (neat) ν 1760 (C=O), 1660 (C=C) cm⁻¹; MS (EI) m/z (%) 221 (M, 21), 220 (3), 207 (13), 206 (base), 162 (3), 134 (2), 122 (3), 106 (2), 69 (3); HRMS calculated for C₁₃H₁₉NO₂ 221.1416, observed 221.1415.

(1*R, 2*R**, 6*R**)-7-Ethyl-3-oxa-9-azatricyclo[7.5.0.0^{2,6}]tetradec-7-en-4-one (17g).** Prepared from **7c** and butanal following the General Procedure under refluxing conditions. The crude product was purified by flash column chromatography (silica gel saturated with Et₃N, 88:10:2 hexanes/EtOAc/Et₃N) to yield **17g** (68%) as an oil: ¹H NMR (250 MHz, CDCl₃) δ 0.98 (t, J = 7.4 Hz, 3H), 1.24-2.02 (m, 10H), 2.39 (dd, J = 17.2, 1.8 Hz, 1H, H-5), 2.68 (dd, J = 17.2, 7.9 Hz, 1H, H-5), 3.00-3.22 (m, 4H), 4.63 (dd, J = 6.5, 1.8 Hz, 1H, H-2), 5.84 (s, 1H, H-8); ¹³C NMR (62.9 MHz, CDCl₃) δ 13.1 (CH₃), 24.2 (CH₂), 27.4 (CH₂), 29.2 (CH₂), 30.2 (CH₂), 31.2 (CH₂), 34.0 (CH₂), 35.7 (CH), 53.6 (CH₂), 58.2 (CH), 85.0 (CH), 105.6 (C), 134.1 (CH), 177.5 (C); IR (neat) ν 1775 (C=O), 1660 (C=C) cm⁻¹; MS (EI) m/z (%) 235 (M, 21), 234 (3), 233 (2), 221 (19), 220 (base), 192 (7), 176 (7), 164 (7), 148 (8), 136 (8); HRMS calculated for C₁₄H₂₁NO₂ 235.1572, observed 235.1570.

5. Preparation of Piperidines 20

(1*R, 5*R**, 6*S**)-8-Benzyl-6-ethyl-2-oxa-8-azabicyclo[3.4.0]nonan-3-one (20a).** Prepared from crude tetrahydropyridine **17a** following the General Hydrogenation Procedure. The crude hydrogenation product was purified by flash column chromatography (silica gel, 75:25 hexanes/EtOAc) to yield **20a** and its C-6-epimer (9:1 dr, 77% for two steps from amine **4**) as oils. Data for **20a**: ¹H NMR (500 MHz, CDCl₃) δ 0.93 (t, J = 7.4 Hz, 3H, CH₃), 1.19-1.31 (m, 2H),

1.86-2.01 (m, 3H, H-6, H-7, H-9), 1.99 (dd, $J = 11.6, 9.4$ Hz, H-9), 2.28 (dd, $J = 17.3, 8.6$ Hz, 1H, H-4), 2.46 (dd, $J = 17.3, 13.1$ Hz, 1H, H-4), 2.70 (dd, $J = 6.9, 1.8$ Hz, 1H, H-7), 2.81-2.86 (m, 1H, H-5), 3.09 (ddd, $J = 11.6, 6.0, 1.8$ Hz, 1H, H-9), 3.52 and 3.60 (AB q, $J = 13.2$ Hz, 2H), 4.66 (ddd, $J = 9.4, 7.3, 6.1$ Hz, 1H, H-1), 7.27-7.36 (m, 5H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 11.4 (CH_3), 24.5 (CH_2), 26.9 (CH_2), 36.7 (CH, C-5), 36.7 (CH, C-6), 53.4 (CH_2), 54.1 (CH_2), 62.4 (CH_2), 76.7 (CH, C-1), 127.2 (CH), 128.2 (CH), 128.8 (CH), 137.5 (C), 176.5 (C); IR (neat) ν 1780 (C=O) cm^{-1} ; MS (EI) m/z (%) 259 (M, 10), 258 (10), 182 (17), 168 (89), 120 (33), 92 (5), 91 (base); HRMS calculated for $\text{C}_{16}\text{H}_{21}\text{NO}_2$ 259.1572, observed 259.1561. Data for the C-6-epimer (from a 1:2 mixture with **20a**): ^1H NMR (250 MHz, CDCl_3) δ 0.90 (distorted t, 3H), 1.10-1.60 (m, 3H), 1.68-2.03 (m, 2H), 2.65 (dd, $J = 13.9, 7.1$ Hz, 1H), 2.75-2.88 (m, 1H), 3.24 (br d, $J = 13.5$ Hz, 1H) 3.51 and 3.62 (AB q, $J = 13.5$ Hz, 2H), 4.44-4.48 (m, 1H), 7.24-7.36 (m, 5H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 11.3 (CH_3), 24.8 (CH_2), 36.0 (CH_2), 38.2 (CH), 38.9 (CH), 53.9 (CH_2), 55.4 (CH_2), 62.3 (CH_2), 78.2 (CH), 127.0 (CH), 128.2 (CH), 128.8 (CH), 137.5 (C), 176.9 (C); IR (neat) ν 1780 (C=O) cm^{-1} ; GC/MS $t_R = 13.8$ min, MS (EI) m/z (%) 259 (M, 3), 258 (2), 182 (5), 168 (51), 120 (11), 91 (base), HRMS calculated for $\text{C}_{16}\text{H}_{21}\text{NO}_2$ 259.1572, observed 259.1572; $t_R = 14.0$ min, MS (EI) m/z (%) 259 (M, 4), 258 (4), 182 (8), 168 (54), 120 (19), 91 (base), HRMS calculated for $\text{C}_{16}\text{H}_{21}\text{NO}_2$ 259.1572, observed 259.1565.

(1*R, 5*R**, 6*S**)-8-Benzyl-6-[3,4-dimethoxyphenylmethyl]-2-oxa-8-azabicyclo[3.4.0]octan-3-one (20b)**. Prepared from crude tetrahydropyridine **17b** following the General Hydrogenation Procedure. The crude hydrogenation product (**20b** and epimer at C-6, 9:1 dr) was purified by flash column chromatography (silica gel, 70:30 hexanes/EtOAc) to yield **20b** (75%, two steps from amine **4**). Data for **20b**: Oil; ^1H NMR (250 MHz, CDCl_3) δ 1.93-2.04 (m, 2H), 2.27-2.40 (m, 3H), 2.47-2.69 (m, 4H), 3.06 (dd, $J = 11.6, 6.0$ Hz, 1H, H-9), 3.42 and 3.64 (AB q, $J = 13.1$

Hz, 2H), 3.86 (s, 6H, CH₃O), 4.54-4.63 (m, 1H, H-1), 6.64 (s, 1H), 6.66 (d, $J = 8.7$ Hz, 1H), 6.78 (d, $J = 8.7$ Hz, 1H), 7.23-7.36 (m, 5H); ¹³C NMR (62.9 MHz, CDCl₃) δ 27.2 (CH₂), 36.4 (CH), 37.1 (CH), 37.5 (CH₂), 53.5 (CH₂), 54.0 (CH₂), 55.8 (CH₃), 62.4 (CH₂), 76.7 (CH), 111.1 (CH), 111.4 (CH), 120.5 (CH), 127.3 (CH), 128.3 (CH), 128.9 (CH), 131.1 (C), 137.3 (C), 147.5 (C), 148.9 (C), 176.4 (C); MS (EI) m/z (%) 381 (M, 50), 290 (10), 243 (11), 229 (28), 178 (10), 177 (10), 152 (base); HRMS calculated for C₂₃H₂₇NO₄ 381.1940, observed 381.1950.

(1*R, 5*R**, 6*S**)-8-Benzyl-6-(3-methoxycarbonylpropyl)-2-oxa-8-azabicyclo[3.4.0]nonan-3-one (20c).** Prepared from crude tetrahydropyridine **17c** following the General Hydrogenation Procedure. The crude hydrogenation product was purified by flash column chromatography (silica gel saturated with Et₃N, 73:25:2 hexanes/EtOAc/Et₃N) to yield **20c** and its C-6-epimer (95%, 94:6 dr) as oils. Data for **20c**: ¹H NMR (250 MHz, CDCl₃) δ 1.16-1.29 (m, 2H), 1.51-1.73 (m, 2H), 1.82-2.07 (m, 3H, H-6, H-7, H-9), 1.86 (t, $J = 11.1$ Hz, H-7, included in m at 1.82-2.07), 1.95 (dd, $J = 11.7, 9.4$ Hz, H-9, included in m at 1.82-2.07), 2.22-2.32 (m, 3H, 1 H-4, 2 H-3'), 2.44 (dd, $J = 17.1, 12.8$ Hz, 1H, H-4), 2.63-2.69 (m, 1H, H-7), 2.74-2.92 (m, 1H, H-5), 3.06 (ddd, $J = 11.7, 6.0, 1.6$ Hz, 1H, H-9), 3.49 and 3.57 (AB q, $J = 13.2$ Hz, 2H), 3.66 (s, 3H, CH₃O), 4.63 (ddd, $J = 9.4, 7.3, 6.0$ Hz, 1H, H-1), 7.23-7.36 (m, 5H); ¹³C NMR (62.9 MHz, CDCl₃) δ 22.1 (CH₂), 27.0 (CH₂), 31.1 (CH₂), 33.7 (CH₂), 34.9 (CH), 36.9 (CH), 51.6 (CH₃), 53.4 (CH₂), 54.2 (CH₂), 62.4 (CH₂), 76.7 (CH, C-1), 127.3 (CH), 128.4 (CH), 128.8 (CH), 137.4 (C), 173.6 (C), 176.4 (C); IR (neat) ν 1770 (C=O), 1730 (C=O) cm⁻¹; MS (EI) m/z (%) 331 (M, 18), 330 (10), 258 (8), 254 (11), 240 (38), 180 (9), 120 (37), 91 (base); HRMS calculated for C₁₉H₂₅NO₄ 331.1784, observed 331.1781. Data for the minor isomer are given from a 8:5 mixture with **20c**: ¹H NMR (250 MHz, CDCl₃) δ 1.11-2.03 (m, 7H), 2.21-2.50 (m, 4H), 2.61-2.67 (m, 1H), 2.74-2.92 (m, 1H), 3.03-3.08 and 3.19-3.25 (2 m, total 1H), 3.46-3.60 (m, 2H), 3.66 (s, 3H), 4.44-4.48

and 4.59-4.68 (2 m, 1H), 7.21-7.35 (m, 5H); ^{13}C NMR (62.9 MHz, CDCl_3) δ 22.1 (CH_2), 27.1 (CH_2), 31.2 (CH_2), 31.6 (CH_2), 33.8 (CH_2), 33.9 (CH_2), 34.9 (CH), 35.9 (CH_2), 36.7 (CH), 36.9 (CH), 39.0 (CH), 51.6 (CH_3), 51.6 (CH_3), 53.5 (CH_2), 53.8 (CH_2), 54.2 (CH_2), 55.7 (CH_2), 62.3 (CH_2), 62.5 (CH_2), 76.7 (CH , C-1), 78.0 (CH , C-1), 127.2 (CH), 127.3 (CH), 128.3 (CH), 128.4 (CH), 128.8 (CH), 128.9 (CH), 137.4 (C), 137.4 (C), 173.6 (C), 173.7 (C), 176.4 (C), 176.7 (C); IR (neat) ν 1770 (C=O), 1730 (C=O) cm^{-1} ; GC/MS t_{R} = 20.2 min, MS (EI) m/z (%) 331 (M, 1), 330 (3), 258 (2), 254 (5), 240 (64), 180 (4), 120 (19), 91 (base), HRMS calculated for $\text{C}_{19}\text{H}_{25}\text{NO}_4$ 331.1784, observed 331.1793; t_{R} = 20.6 min, MS (EI) m/z (%) 331 (M, 2), 330 (2), 258 (2), 254 (5), 240 (61), 180 (4), 120 (20), 91 (base), HRMS calculated for $\text{C}_{19}\text{H}_{25}\text{NO}_4$ 331.1784, observed 331.1778.

(1*R, 2*R**, 6*R**, 7*S**)-7-Ethyl-3-oxa-9-azatricyclo[7.3.0.0^{2,6}]dodecan-4-one (20e).** Prepared from crude tetrahydropyridine **17e** following the General Hydrogenation Procedure. The crude hydrogenation product was purified by flash column chromatography (silica gel saturated with Et_3N , 78:20:2 hexanes/EtOAc/ Et_3N) to yield **20e** and its C-7-epimer (11:1 dr, 62% for two steps starting from **9a**). Data for **20e**: ^1H NMR (500 MHz, CDCl_3) δ 0.89 (t, J = 7.3 Hz, 3H, CH_3), 1.23-1.36 (m, 2H), 1.68-1.77 (m, 3H, 2 H-11, H-12), 1.82-1.92 (m, 2H, H-7, H-12), 2.14-2.18 (m, 1H, H-10), 2.36 (dd, J = 17.8, 10.5 Hz, 1H, H-5), 2.58 (dd, J = 17.8, 8.4 Hz, 1H, H-5), 2.61 (dd, J = 10.6, 7.6 Hz, 1H, H-8), 2.68 (t, J = 10.6 Hz, 1H, H-8), 2.72-2.83 (m, 2H, H-1, H-6), 2.89-2.92 (m, 1H, H-10), 4.62 (dd, J = 8.8, 2.8 Hz, 1H, H-2); ^{13}C NMR (125.8 MHz, CDCl_3) δ 11.6 (CH_3), 23.3 (CH_2 , C-11), 23.7 (CH_2 , C-12), 26.1 (CH_2), 29.2 (CH_2 , C-5), 34.8 (CH , C-7), 36.0 (CH , C-6), 52.8 (CH_2 , C-8), 54.3 (CH_2 , C-10), 59.7 (CH , C-1), 80.1 (CH , C-2), 177.9 (C); IR (neat) ν 1770 (C=O) cm^{-1} ; MS (EI) m/z (%) 209 (M, 53), 208 (37), 180 (9), 152 (8), 124 (66), 110 (base); HRMS calculated for $\text{C}_{12}\text{H}_{19}\text{NO}_2$ 209.1416, observed 209.1406. Data for the C-7-epimer (from a

2:7 mixture with **20e**): ^1H NMR (250 MHz, CDCl_3) δ 0.86-0.95 (m, 3H), 1.24-2.09 (m, 9H), 2.12-3.03 (m, 6H), 4.32 (dd, $J = 8.7, 6.5$ Hz, H-2); ^{13}C NMR (62.9 MHz, CDCl_3) δ 12.2 (CH_3), 21.2 (CH_2), 23.6 (CH_2), 26.6 (CH_2), 31.9 (CH_2), 37.9 (CH), 38.8 (CH), 51.4 (CH_2), 54.3 (CH_2), 64.1 (CH, C-1), 82.0 (CH, C-2), 176.6 (C); IR (neat) ν 1770 ($\text{C}=\text{O}$) cm^{-1} ; GC-MS $t_{\text{R}} = 13.8$ min, MS (EI) m/z (%) 209 (M, 29), 208 (34), 180 (6), 152 (4), 124 (60), 110 (base), HRMS calculated for $\text{C}_{12}\text{H}_{19}\text{NO}_2$ 209.1416, observed 209.1409; $t_{\text{R}} = 14.2$ min, MS (EI) m/z (%) 209 (M, 43), 208 (31), 180 (6), 152 (4), 124 (60), 110 (base), HRMS calculated for $\text{C}_{12}\text{H}_{19}\text{NO}_2$ 209.1416, observed 209.1415.

(1*R, 2*R**, 6*R**, 7*S**)-7-Ethyl-3-oxa-9-azatricyclo[7.4.0.0^{2,6}]tridecan-4-one (20f)** and **(1*R**, 2*R**, 3*R**, 4*S**)-4-Ethyl-3-(ethoxycarbonylmethyl)-2-hydroxy-6-azabicyclo[4.4.0]decane (33)**. Prepared from **17f** following the General Hydrogenation Procedure using a 60 psi H_2 pressure for 20 h. The crude product was purified by flash column chromatography (silica gel saturated with Et_3N , 88:10:2 hexanes/ EtOAc / Et_3N) to yield **33** (3%) and **20f** (88%) as oils, together with a 5% recovered **17f**. Data for **20f**: ^1H NMR (500 MHz, CDCl_3) δ 0.88 (t, $J = 7.3$ Hz, 3H, CH_3), 1.21 (qt, $J = 12.7, 4.0$ Hz, 1H, H-12, overlapped with signals of H-1'), 1.22-1.32 (m, 1H, H-1'), 1.46-1.77 (m, 6H, H-7, 2 H-11, 2 H-13, 1 H-1'), 1.78 (d, $J = 12.7$ Hz, 1H, H-12), 1.98 (td, $J = 11.5, 3.3$ Hz, 1H, H-10), 2.15 (dd, $J = 11.8, 5.1$ Hz, 1H, H-8), 2.19 (d, $J = 11.2$ Hz, 1H, H-1), 2.44 (dd, $J = 17.3, 8.7$ Hz, 1H, H-5), 2.54 (dd, $J = 17.3, 5.2$ Hz, 1H, H-5), 2.66-2.74 (m, 2H, H-6, H-8), 2.72 (dd, $J = 11.8, 6.5$ Hz, H-8), 2.82 (d, $J = 11.2$ Hz, 1H, H-10), 4.33 (dd, $J = 6.8, 3.3$ Hz, 1H, H-2); ^{13}C NMR (125.8 MHz, CDCl_3) δ 12.0 (CH_3), 22.6 (CH_2), 24.4 (CH_2 , C-12), 25.4 (CH_2 , C-11), 27.2 (CH_2 , C-13), 31.9 (CH_2 , C-5), 35.1 (CH, C-7), 36.8 (CH, C-6), 55.8 (CH_2 , C-8), 56.6 (CH_2 , C-10), 60.7 (CH, C-1), 81.7 (CH, C-2), 177.5 (C); IR (neat) ν 1770 ($\text{C}=\text{O}$) cm^{-1} ; MS (EI) m/z (%) 223 (M, 35), 222 (34), 194 (12), 164 (11), 138 (76), 124 (30), 97

(18), 83 (base); HRMS calculated for $C_{13}H_{21}NO_2$ 223.1572, observed 223.1573. Data for **33**: 1H NMR (250 MHz, $CDCl_3$) δ 0.87 (t, $J = 7.0$ Hz, 3H), 1.22-2.11 (m, 17H), 1.25 (t, $J = 7.1$ Hz, included in m at 1.22-2.11), 2.43 (dd, $J = 15.6, 6.7$ Hz, 1H), 2.64 (dd, $J = 15.6, 8.3$ Hz, 1H), 2.76-2.81 (m, 2H), 3.34 (m, 1H), 4.12 and 4.13 (2 q, $J = 7.1$ Hz, 2H); ^{13}C NMR (62.9 MHz, $CDCl_3$) δ 14.2 (CH_3), 14.3 (CH_3), 21.6 (CH_2), 24.0 (CH_2), 25.6 (CH_2), 28.2 (CH_2), 35.0 (CH_2), 39.2 (CH), 41.0 (CH), 56.3 (CH_2), 59.4 (CH_2), 60.2 (CH_2), 66.6 (CH), 72.2 (CH), 173.4 (C); IR (neat) ν 3530 (O-H), 1730 (C=O) cm^{-1} ; MS (EI) m/z (%) 269 (M, 4), 223 (36), 222 (42), 194 (10), 182 (34), 164 (51), 138 (base), 124 (84), 110 (15), 97 (88), 83 (90); HRMS calculated for $C_{15}H_{27}NO_3$ 269.1991, observed 269.2000.

(1*R, 2*R**, 6*R**, 7*S**)-7-Ethyl-3-oxa-9-azatricyclo[7.5.0.0^{2,6}]tetradecan-4-one (20g).**

Prepared in 93% yield from tetrahydropyridine **17g** following the General Hydrogenation Procedure. The crude product was purified by flash column chromatography (silica gel saturated with Et_3N , 93:5:2 hexanes/ $EtOAc$ / Et_3N) to yield **20g** as an oil: 1H NMR (250 MHz, $CDCl_3$) δ 0.89 (t, $J = 7.3$ Hz, 3H, CH_3), 1.17-2.01 (m, 11H), 2.31 (dd, $J = 17.2, 10.3$ Hz, 1H), 2.43 (t, $J = 10.3$ Hz, 1H), 2.55-2.96 (m, 6H), 4.52 (dd, $J = 8.7, 2.4$ Hz, 1H, H-2); ^{13}C NMR (62.9 MHz, $CDCl_3$) δ 11.5 (CH_3), 24.3 (CH_2), 26.4 (CH_2), 28.2 (CH_2), 28.9 (CH_2), 29.2 (CH_2), 29.5 (CH_2), 33.8 (CH), 35.7 (CH), 53.5 (CH_2), 54.9 (CH_2), 61.5 (CH, C-1), 84.3 (CH, C-2), 178.1 (C); IR (neat) ν 1770 (C=O) cm^{-1} ; MS (EI) m/z (%) 237 (M, 22), 208 (41), 195 (13), 194 (25), 180 (12), 164 (34), 152 (38), 150 (41), 138 (37), 124 (32), 97 (82), 68 (base); HRMS calculated for $C_{14}H_{23}NO_2$ 237.1729, observed 237.1737.

6. Stereochemical Elucidation

Stereochemical assignments for pyrrolidine- (**13**, **16**, **35**) and piperidine-type (**17**, **19**, **20**, **26**) products were made based on data obtained from NOE experiments. Thus, a *syn*-relationship between ring methine hydrogens was generally revealed by the observation of relatively strong NOE's, that were either absent or significantly weaker in the isomers with a corresponding *anti*-relationship. The most significant NOE data are collected in Chart 1. In the case of **20b,c,g** and **26a** unambiguous use of NOE's was difficult due to signal overlap, and further support for the proposed stereochemistry was gained by the observation of *J* values for H-1 and H-6 (in **20c**) or for H-6 and H-7 (in the case of **26a**) that indicated axial dispositions for all of those hydrogens. Additionally, where appropriate comparisons with another isomer were possible (e.g. **20a-g**), the all-*syn* isomers displayed a significant shielding (1.6-5.4 ppm) in the ¹³C resonances corresponding to both the methine carbon bearing the R³ substituent and the adjacent bridgehead position, and this was interpreted in terms of the increased steric congestion associated to those isomers.¹¹

Chart 1. Summary of NOE Data

Chart 1. Summary of NOE Data (cont.)

7. References

- (1) Molander, G. A. In *Organic Reactions*; Paquette, L. A., Ed.; John Wiley & Sons: New York, 1994; Vol. 46, pp 211-367.
 - (2) Kanazawa, A. M.; Correa, A.; Denis, J. N.; Luche, M. J.; Greene, A. E. *J. Org. Chem.* **1993**, *58*, 255-257.
 - (3) Ha, H.-J.; Kang, K.-H.; Ahn, Y.-G.; Oh, S.-J. *Heterocycles* **1997**, *45*, 277-286.
 - (4) Nagasaka, T.; Tamano, H.; Hamaguchi, F. *Heterocycles* **1986**, *24*, 1231-2.
 - (5) Siu, T.; Li, W.; Yudin, A. K. *J. Comb. Chem.* **2000**, *2*, 545-549.
 - (6) Rassu, G.; Zanardi, F.; Battistini, L.; Gaetani, E.; Casiraghi, G. *J. Med. Chem.* **1997**, *40*, 168-180.
 - (7) de Oliveira, M. C. F.; Silva-Santos, L.; Pilli, L. A. *Tetrahedron Lett.* **2001**, *42*, 6995-6997.
- This paper reports the preparation of **7a-c** and **8b**. Spectroscopic data (NMR, IR, MS) are provided for **7b** only.
- (8) Pilli, R. A.; Dias, L. C. *Synth. Commun.* **1991**, *21*, 2213-2229.
 - (9) Padwa, A.; Brodney, M. A.; Marino, J., J. P.; Sheehan, S. M. *J. Org. Chem.* **1997**, *62*, 78-87.
 - (10) (a) Aurrecoechea, J. M.; López, B.; Fernández, A.; Arrieta, A.; Cossío, F. P. *J. Org. Chem.* **1997**, *62*, 1125-1135. (b) Nishida, A.; Kawahara, N.; Nishida, M.; Yonemitsu, O. *Tetrahedron* **1996**, *52*, 9713-9734.
 - (11) Aurrecoechea, J. M.; Fernández, A.; Gorgojo, J. M.; Saornil, C. *Tetrahedron* **1999**, *55*, 7345-7362.

¹H NMR Spectrum (500 MHz, DMSO-d₆, 70 °C) of 2

^1H NMR Spectrum (250 MHz, CDCl_3) of **3**

^1H NMR Spectrum (250 MHz, CDCl_3) of **5**

^1H NMR Spectrum (250 MHz, CDCl_3) of **4**

^1H NMR Spectrum (250 MHz, DMSO- d_6 , 70 °C) of 7a

^1H NMR Spectrum (250 MHz, CDCl_3) of 8b

^1H NMR Spectrum (250 MHz, CDCl_3) of **7c**

^1H NMR Spectrum (250 MHz, CDCl_3) of crude 9a including remaining solvent (CH_2Cl_2 , δ 5.28) and silyl residues ($\delta \sim 0.1$)

^1H NMR Spectrum (250 MHz, CDCl_3) of 13a

^1H NMR Spectrum (250 MHz, CDCl_3) of 13c

^1H NMR Spectrum (250 MHz, CDCl_3) of 13d

^1H NMR Spectrum (250 MHz, CDCl_3) of 13e

^1H NMR Spectrum (250 MHz, CDCl_3) of 16a (major isomer)

^1H NMR Spectrum (250 MHz, CDCl_3) of 16a (minor isomer; 3:1 mixture with major isomer)

16a (minor isomer; mixture with major isomer)

^1H NMR Spectrum (500 MHz, CDCl_3) of **16b (major isomer)**

^1H NMR Spectrum (500 MHz, CDCl_3) of **16b (minor isomer)**

^1H NMR Spectrum (500 MHz, CDCl_3) of **16c (major isomer)**

^1H NMR Spectrum (250 MHz, CDCl_3) of 16c (minor isomer; 1:2 mixture with major isomer)

^1H NMR Spectrum (250 MHz, CDCl_3) of 16d (major isomer)

^1H NMR Spectrum (250 MHz, CDCl_3) of 16d (minor isomer; 36:64 mixture with major isomer)

16d (minor isomer; mixture with major isomer)

^1H NMR Spectrum (250 MHz, CDCl_3) of 16e (major isomer)

^1H NMR Spectrum (250 MHz, CDCl_3) of 16e (minor isomer)

^1H NMR Spectrum (500 MHz, CDCl_3) of 19

^1H NMR Spectrum (250 MHz, CDCl_3) of 17a

^1H NMR Spectrum (250 MHz, CDCl_3) of crude 17b

^1H NMR Spectrum (250 MHz, CDCl_3) of 17c

¹H NMR Spectrum (250 MHz, CDCl₃) of 17d

^1H NMR Spectrum (250 MHz, CDCl_3) of 17e

17e

^1H NMR Spectrum (250 MHz, CDCl_3) of 17f

17f

^1H NMR Spectrum (250 MHz, CDCl_3) of 17g

17g

^1H NMR Spectrum (300 MHz, CDCl_3) of 26b

^1H NMR Spectrum (250 MHz, CDCl_3) of 26a

^1H NMR Spectrum (250 MHz, CDCl_3) of 26c (93:7 mixture with 26a)

^1H NMR Spectrum (250 MHz, CDCl_3) of 32a

32a

^1H NMR Spectrum (250 MHz, CDCl_3) of 32b (9:1 mixture with 32a)

^1H NMR Spectrum (500 MHz, CDCl_3) of 32c/32d (2:3 mixture)

^1H NMR Spectrum (300 MHz, CDCl_3) of 32e

32e

^1H NMR Spectrum (500 MHz, CDCl_3) of 20a

^1H NMR Spectrum (250 MHz, CDCl_3) of a 2:1 mixture of 20a with minor isomer

^1H NMR Spectrum (250 MHz, CDCl_3) of 20b

20c

¹H NMR Spectrum (250 MHz, CDCl₃) of a 5:8 mixture of 20c with minor isomer

^1H NMR Spectrum (500 MHz, CDCl_3) of 20e

^1H NMR Spectrum (250 MHz, CDCl_3) of a 7:2 mixture of 20e with minor isomer

^1H NMR Spectrum (500 MHz, CDCl_3) of 20f

20f

^1H NMR Spectrum (250 MHz, CDCl_3) of 33

^1H NMR Spectrum (250 MHz, CDCl_3) of 20g

^1H NMR Spectrum (250 MHz, CDCl_3) of 34a

^1H NMR Spectrum (250 MHz, CDCl_3 , 50 $^\circ\text{C}$) of 34b

^1H NMR Spectrum (500 MHz, CDCl_3) of 35

^1H NMR Spectrum (250 MHz, CDCl_3) of 36 (3:1 Diastereomeric Mixture)

^1H NMR Spectrum (250 MHz, CDCl_3) of 37

¹H NMR Spectrum (250 MHz, CDCl₃, 50 °C) of 38a

^1H NMR Spectrum (250 MHz, CDCl_3 , 50 $^\circ\text{C}$) of 38b