Diverse functionalization of thiazolo ring-fused 2-pyridones

*Erik Chorell, Pralay Das and Fredrik Almqvist**

Department of Chemistry, Umeå University, SE-90187 Umeå Sweden

Fredrik.almqvist@chem.umu.se

Supporting information

Table of contents

General synthesis	3
¹ H NMR spectra of compound 9a	4
¹³ C NMR spectra of compound 9a	5
¹ H NMR spectra of compound 9b	6
¹³ C NMR spectra of compound 9b	7
¹ H NMR spectra of compound 10a	8
¹³ C NMR spectra of compound 10a	9
¹ H NMR spectra of compound 10b	10
¹³ C NMR spectra of compound 10b	11
¹ H NMR spectra of compound 11a	12
¹³ C NMR spectra of compound 11a	13
¹ H NMR spectra of compound 11b	14
¹³ C NMR spectra of compound 11b	15
¹ H NMR spectra of compound 13a	16
¹³ C NMR spectra of compound 13a	17
¹ H NMR spectra of compound 13b	18
¹³ C NMR spectra of compound 13b	19
¹ H NMR spectra of compound 14a	20
¹³ C NMR spectra of compound 14a	21
¹ H NMR spectra of compound 14b	22
¹³ C NMR spectra of compound 14b	23
¹ H NMR spectra of compound 15a	24
¹³ C NMR spectra of compound 15a	25
¹ H NMR spectra of compound 15b	26

¹³ C NMR spectra of compound 15b	27
¹ H NMR spectra of compound 16a	28
¹³ C NMR spectra of compound 16a	29
¹ H NMR spectra of compound 16b	30
¹³ C NMR spectra of compound 16b	31
¹ H NMR spectra of compound 17a	32
¹³ C NMR spectra of compound 17a	33
¹ H NMR spectra of compound 17b	34
¹³ C NMR spectra of compound 17b	35
¹ H NMR spectra of compound 18a	36
¹³ C NMR spectra of compound 18a	37
¹ H NMR spectra of compound 18b	38
¹³ C NMR spectra of compound 18b	39
¹ H NMR spectra of compound 19a	40
¹³ C NMR spectra of compound 19a	41
¹ H NMR spectra of compound 19b	42
¹³ C NMR spectra of compound 19b	43
¹ H NMR spectra of compound 20a	44
¹³ C NMR spectra of compound 20a	45
¹ H NMR spectra of compound 20b	46
¹³ C NMR spectra of compound 20b	47
Temp, pressure and power graphs for microwave heating of compound 14a	48
Temp, pressure and power graphs for microwave heating of compound 14b	49
Temp, pressure and power graphs for microwave heating of compound 15a	50
Temp, pressure and power graphs for microwave heating of compound 15b	51

General synthesis

All reactions were carried out under an inert atmosphere with dry solvents under anhydrous conditions, unless otherwise stated. MeCN was freshly distilled from calcium hydride, THF was freshly distilled from potassium, DMF was dried using P₂O₅, distilled and stored over 3 Å molecular sieves, MeOH was dried using calcium hydride, distilled and stored over 3 Å molecular sieves.

All microwave reactions were carried out in a monomode reactor (Smith Synthesizer, Biotage AB) using process vials sealed with Teflon septa and an aluminum crimp top. Reaction times refer to irradiation time at the target temperature, not the total irradiation time. The temperature was measured with an IR sensor. Heck reactions described in the experimental procedure were carried out in process vials with a filling volume of 0.5-2.0 mL and with a solvent volume of 1.2 mL.

TLC was performed on Silica Gel 60 F₂₅₄ aluminium backed plates using UV light detection (254 nm). Flash column chromatography (eluents given in brackets) employed normal phase silica gel (60 Å, 35-70 µm). The ¹H and ¹³C NMR spectra were recorded at 400 or 100 MHz respectively at 298 K in CDCl₃ [residual CHCl₃ (δ_H 7.26 ppm) or CDCl₃ (δ_C 77.0 ppm) as internal standard] MeOH [residual CD₂HOD (δ_H 3.30 ppm) or CD₃OD (δ_C 49.0 ppm) as internal standard]. It should be noted that the presence of fluorine substituents leads to reduced peak intensities. IR spectra were recorded on an FTIR spectrometer. Melting points are uncorrected.

^1H NMR spectra of compound **9a** in CDCl_3

^{13}C NMR spectra of compound **9a** in CDCl_3

^1H NMR spectra of compound **9b** in CDCl_3

^{13}C NMR spectra of compound **9b** in CDCl_3

^1H NMR spectra of compound **10a** in CDCl_3

^{13}C NMR spectra of compound **10a** in CDCl_3

^1H NMR spectra of compound **10b** in CDCl_3

^{13}C NMR spectra of compound **10b** in CDCl_3

^1H NMR spectra of compound **11a** in CDCl_3

^{13}C NMR spectra of compound **11a** in CDCl_3

^1H NMR spectra of compound **11b** in CDCl_3

^{13}C NMR spectra of compound **11b** in CDCl_3

^1H NMR spectra of compound **13a** in CD_3OD

^{13}C NMR spectra of compound **13a** in CD_3OD

^1H NMR spectra of compound **13b** in CD_3OD

^{13}C NMR spectra of compound **13b** in CD_3OD

^1H NMR spectra of compound **14a** in CDCl_3

^{13}C NMR spectra of compound **14a** in CDCl_3

^1H NMR spectra of compound **14b** in CDCl_3

^{13}C NMR spectra of compound **14b** in CDCl_3

^1H NMR spectra of compound **15a** in CDCl_3

^{13}C NMR spectra of compound **15a** in CDCl_3

^1H NMR spectra of compound **15b** in CDCl_3

^{13}C NMR spectra of compound **15b** in CDCl_3

^1H NMR spectra of compound **16a** in CDCl_3

^{13}C NMR spectra of compound **16a** in CDCl_3

^1H NMR spectra of compound **16b** in CDCl_3

^{13}C NMR spectra of compound **16b** in CDCl_3

^1H NMR spectra of compound **17a** in CDCl_3

^{13}C NMR spectra of compound **17a** in CDCl_3

^1H NMR spectra of compound **17b** in CDCl_3

^{13}C NMR spectra of compound **17b** in CDCl_3

^1H NMR spectra of compound **18a** in CDCl_3

^{13}C NMR spectra of compound **18a** in CDCl_3

^1H NMR spectra of compound **18b** in CDCl_3

^{13}C NMR spectra of compound **18b** in CDCl_3

^1H NMR spectra of compound **19a** in CDCl_3

^{13}C NMR spectra of compound **19a** in CDCl_3

^1H NMR spectra of compound **19b** in CDCl_3

^{13}C NMR spectra of compound **19b** in CDCl_3

^1H NMR spectra of compound **20a** in CDCl_3

^{13}C NMR spectra of compound **20a** in CDCl_3

^1H NMR spectra of compound **20b** in CDCl_3

^{13}C NMR spectra of compound **20b** in CDCl_3

Temp, pressure and power graphs for microwave heating of compound **14a**

Process parameters

Temperature: 105°C
Time: 1500 s
Fixed Hold Time: On
Absorption Level: High
Pre-stirring: 30 s

Graphs

Temp, pressure and power graphs for microwave heating of compound **14b**

Process parameters

Temperature: 105°C
Time: 1500 s
Fixed Hold Time: On
Absorption Level: High
Pre-stirring: 30 s

Graphs

Temp, pressure and power graphs for microwave heating of compound **15a**

Process parameters

Temperature: 105°C
Time: 1500 s
Fixed Hold Time: On
Absorption Level: High
Pre-stirring: 30 s

Graphs

Temp, pressure and power graphs for microwave heating of compound **15b**

Process parameters

Temperature: 105°C
Time: 1500 s
Fixed Hold Time: On
Absorption Level: High
Pre-stirring: 30 s

Graphs