Dual Catalyst Control in the Enantioselective Intramolecular Morita-Baylis-Hillman Reaction

Carrie E. Aroyan, Melissa M. Vasbinder and Scott J. Miller*

Department of Chemistry, Merkert Chemistry Center, Boston College, Chestnut Hill, Massachusetts 02467-3860 Email: scott.miller.1@bc.edu

Supporting Information

General Procedures. Proton NMR spectra were recorded on a Varian 400 MHz spectrometer. Proton chemical shifts are reported in ppm (δ) relative to internal tetramethylsilane (TMS, δ 0.0 ppm) or with the solvent reference relative to TMS employed as the internal standard (CDCl_3 , δ 7.26 ppm; d_6 -DMSO, δ 2.50 ppm; CD_3OD , δ 3.31 ppm; D_2O , δ 4.79 ppm). Data are reported as follows: chemical shift (multiplicity [singlet (s), doublet (d), triplet (t), quartet (q), and multiplet (m)], coupling constants [Hz], integration). Carbon NMR spectra were recorded on a Varian 400 (100 MHz) spectrometer with complete proton decoupling. Carbon chemical shifts are reported in ppm (δ) relative to TMS with the respective solvent resonance as the internal standard (CDCl_3 , δ 77.0 ppm). Infrared spectra were obtained on a Perkin-Elmer Spectrum 1000 spectrometer. Analytical thin-layer chromatography (TLC) was performed using Silica Gel 60 Å F254 precoated plates (0.25 mm thickness). TLC R_f values are reported. Visualization was accomplished by irradiation with a UV lamp and/or staining with KMnO_4 or cerium ammonium molybdenate (CAM) solutions. Flash column chromatography was performed using Silica Gel 60A (32-63 μm).¹ Optical rotations were recorded on a Rudolf Research Analytical Autopol IV Automatic polarimeter at the sodium D line (path length 50 mm). High resolution mass spectra were obtained at the Mass Spectrometry Facility of either Boston College (Chestnut Hill, MA) or the University of Illinois (Urbana-Champaign, IL). The method of ionization is given in parentheses.

Analytical normal phase HPLC was performed on a Hewlett-Packard 1100 Series chromatograph equipped with a diode array detector (214 nm and 254 nm). All reactions were carried out under an argon or nitrogen atmosphere employing oven- and flame-dried glassware. All solvents were distilled from appropriate drying agents prior to use. *N*-methyl imidazole and acetic anhydride were distilled and stored in a Schlenk flask for no more than two weeks prior to use. Intramolecular Baylis-Hillman substrate **1** was prepared as reported previously in the literature² as well as by the general procedure described.

General Procedure for the Preparation of Intramolecular Morita-Baylis-Hillman Substrates (1, 4, 6, 8, 10).

(*E*)-7-oxo-7-phenylhept-5-enal (1). Phosphorane³ **14** (9.1 g, 24 mmol) was added to a solution of 5,5-dimethoxypentanal⁴ (**15**, 2.5 g, 17 mmol) and CH₂Cl₂ (43 mL). The solution was stirred for 24 h and then concentrated. Flash chromatography of the crude material (hexane-EtOAc, 5:1) yielded the (*E*) Wittig product in 90% purity as determined by ¹H NMR. This material was then dissolved in THF (60 mL) and 1N HCl (20 mL) and stirred at rt. After 2 h, THF was removed in vacuo from the yellow solution and the resultant residue was diluted with Et₂O (100 mL) and washed with water (50 mL), saturated NaHCO₃ (50 mL), and brine (50 mL). The aqueous layers were back extracted with ether (2 x 50 mL). The combined organic layers were then dried over Na₂SO₄ and concentrated to give a yellow oil. Flash chromatography of the crude material (toluene-EtOAc, 5:1) gave **1** as a pale yellow oil (960 mg, 4.7 mmol, 28%). Compound **1**, prepared in this manner, is identical to that prepared previously.² Compounds **4**, **6**, **8**, and **10** were prepared by analogous methodology.

(*E*)-7-(4-chlorophenyl)-7-oxohept-5-enal (4).

¹H NMR (CDCl₃, 400 MHz) δ 9.83 (s, 1H), 7.92 (d, *J* = 8.8 Hz, 2H), 7.49 (d, *J* = 8.8 Hz, 2H), 7.08 (m, 1H), 6.92 (d, *J* = 15.6 Hz, 1H), 2.58 (t, *J* = 7.2 Hz, 2H), 2.42 (q, *J* = 7.2 Hz, 2H), 1.92 (quintet, *J* = 7.6 Hz, 2H); **¹³C NMR** (CDCl₃, 100 MHz) δ 201.6, 189.3, 148.7, 139.3, 136.2, 130.1, 129.0, 126.2, 43.1, 32.0, 20.6; **IR** (film, cm⁻¹) 2942, 2823, 2722, 1727, 1671, 1621, 1583, 1401, 1293, 1224, 1092, 1016; **TLC** R_f 0.31 (3:1 toluene:EtOAc); **Exact mass** calcd for [C₁₃H₁₃O₂Cl + H]⁺ requires *m/z* 237.0682. Found 237.0693 (ESI⁺).

(*E*)-7-(4-bromophenyl)-7-oxohept-5-enal (6).

¹H NMR (CDCl₃, 400 MHz) δ 9.83 (s, 1H), 7.83 (d, *J* = 8.8 Hz, 2H), 7.64 (d, *J* = 8.8 Hz, 2H), 7.06 (dt, *J* = 15.2, 6.8 Hz, 1H), 6.89 (d, *J* = 15.2 Hz, 1H), 2.56 (dt, *J* = 7.2 Hz, 1.2 Hz, 2H), 2.40 (q, *J* = 6.8 Hz, 2H), 1.91 (quintet, *J* = 7.2 Hz, 2H); **¹³C NMR** (CDCl₃, 100 MHz) δ 200.8, 188.7, 148.1, 136.0, 131.4, 129.6, 127.4, 125.6, 42.8, 31.7, 20.2; **IR** (film, cm⁻¹) 2949, 2886, 2829, 2722, 1721, 1671, 1620, 1583, 1400, 1073, 1010; **TLC** R_f 0.36

(3:1 toluene:EtOAc); **Exact mass** calcd for $[C_{13}H_{13}O_2Br + H]^+$ requires m/z 281.0177. Found 281.0188 (ESI⁺).

(E)-7-oxo-7-*o*-tolylhept-5-enal (8).

¹H NMR (CDCl₃, 400 MHz) δ 9.80 (s, 1H), 7.42-7.35 (m, 2H), 7.30-7.24 (m, 2H), 6.71 (dt, J = 15.6, 6.8 Hz, 1H), 6.53 (d, J = 16.0 Hz, 1H), 2.55-2.53 (dt, J = 7.2, 1.2 Hz, 2H), 2.42 (bs, 3H), 2.34 (q, J = 7.2 Hz, 2H), 1.85 (quintet, J = 7.6 Hz, 2H); **¹³C NMR** (CDCl₃, 100 MHz) δ 201.1, 196.1, 149.2, 138.5, 136.6, 131.0, 130.2, 127.8, 125.1, 43.0, 31.8, 20.4, 20.2; **IR** (film, cm⁻¹) 2930, 2829, 2728, 1728, 1652, 1627; **TLC** R_f 0.53 (3:1 toluene:EtOAc); **Exact mass** calcd for $[C_{14}H_{16}O_2 + H]^+$ requires m/z 217.1229. Found 217.1238 (ESI⁺).

(E)-7-oxo-7-(thiophen-2-yl)hept-5-enal (10).

¹H NMR (CDCl₃, 400 MHz) δ 9.82 (s, 1H), 7.77 (dd, J = 3.6, 0.8 Hz, 1H), 7.66 (dd, J = 4.8, 0.8 Hz, 1H), 7.16 (dd, J = 4.8, 4.0 Hz, 1H), 7.07 (dt, J = 15.2, 7.2 Hz, 1H), 6.83 (d, J = 15.2 Hz, 1H), 2.53 (dt, J = 7.2, 1.2 Hz, 2H), 2.37 (q, J = 6.8 Hz, 2H), 1.88 (quintet, J = 7.2 Hz, 2H); **¹³C NMR** (CDCl₃, 100 MHz) δ 201.2, 181.7, 146.9, 144.7, 133.6, 131.7, 128.0, 126.0, 43.0, 31.7, 20.5; **IR** (film, cm⁻¹) 3094, 2936, 2829, 2722, 1728, 1659, 1614, 1520, 1420; **TLC** R_f 0.46 (3:1 toluene:EtOAc); **Exact mass** calcd for $[C_{11}H_{12}O_2S + H]^+$ requires m/z 209.0636. Found 209.0634 (ESI⁺).

General Procedure for the Catalytic Intramolecular Morita-Baylis-Hillman Reactions.

Pipecolinic acid (14.2 mg, 0.11 mmol), water (233 μ L), THF (659 μ L), and NMI (41 μ L of a 1.37 M solution in THF) were added sequentially to a vessel containing aldehyde 4 (132 mg, 0.56 mmol). The reaction was stirred at room temperature for 48 hours, whereupon the yellow solution was added to 30 mL of EtOAc and washed with saturated NH₄Cl (20 mL), NaHCO₃ (20 mL), and brine (20 mL). The organic layer was dried over Na₂SO₄, filtered, and concentrated to afford a pale yellow oil (121 mg, 92% yield of 5). The product isolated in this manner appears as a single compound with slight impurities manifested by low intensity peaks in the baseline of the ¹H NMR spectrum (see figure 1). This material may be purified further by flash chromatography using a gradient eluent

(toluene:ethyl acetate 100:0 to 0:100) affording 5 (74 mg) as a clear oil in 56% isolated yield.

2-benzoyl-1-hydroxycyclohex-2-ene (2).

The characterization data for this compound matched that which has previously been reported.²

[α]_D + 30 (*c* = 1.0, CHCl₃, 80% ee); **Assay of enantiomeric purity:** Enantiomers of product were separated by chiral HPLC employing a Chiralcel OD column (Daicel). Conditions: 98:2 hexanes: ethanol; Flow rate 0.75 mL/min; 18.8 min (major ent), 22.2 min (minor ent).

The absolute stereochemistry was assigned by Mosher analysis.⁵

Compound 5.

¹H NMR (CDCl₃, 400 MHz) δ 7.62 (d, *J* = 8.4 Hz, 2H), 7.42 (d, *J* = 8.8 Hz, 2H), 6.70 (t, *J* = 4.4 Hz, 1H), 4.73 (br s, 1H), 3.38 (s, 1H), 2.39-2.27 (m, 2H), 1.97-1.85 (m, 3H), 1.68-1.64 (m, 1H); **¹³C NMR** (CDCl₃, 100 MHz) δ 197.6, 146.7, 139.9, 138.2, 135.9, 130.6, 128.4, 63.9, 29.9, 26.5, 17.5; **IR** (film, cm⁻¹) 3440, 2936, 2867, 1646, 1583; **TLC R_f** 0.37 (3:1 toluene:EtOAc); **[α]_D** + 18 (*c* = 1.0, CHCl₃, 79% ee); **Exact mass** calcd for [C₁₃H₁₃O₂Cl + Na] + requires *m/z* 259.0502. Found 259.0503 (ESI⁺). **Assay of enantiomeric purity:** Enantiomers of product were separated by chiral HPLC employing a Chiralcel OJ column (Daicel). Conditions: 98:2 hexanes: ethanol; Flow rate 0.60 mL/min; 28.3 min (major ent), 32.7 min (minor ent).

Compound 7.

¹H NMR (CDCl₃, 400 MHz) δ 7.59 (d, *J* = 8.8 Hz, 2H), 7.53 (d, *J* = 8.8 Hz, 2H), 6.70 (t, *J* = 4.4 Hz, 1H), 4.73 (br s, 1H), 3.38 (s, 1H), 2.39-2.22 (m, 2H), 1.96-1.83 (m, 3H), 1.69-1.62 (m, 1H); **¹³C NMR** (CDCl₃, 100 MHz) δ 197.7, 146.8, 139.9, 136.4, 131.4, 130.7, 126.8, 63.9, 29.8, 26.5, 17.5; **IR** (film, cm⁻¹) 3414, 2936, 2867, 1634; **TLC R_f** 0.36 (3:1

toluene:EtOAc); $[\alpha]_D + 15.4$ ($c = 1.0$, CHCl_3 , 79% ee); **Exact mass** calcd for $[\text{C}_{13}\text{H}_{13}\text{O}_2\text{Br} + \text{Na}] +$ requires m/z 302.9997. Found 302.9996 (ESI^+). **Assay of enantiomeric purity:** Enantiomers of product were separated by chiral HPLC employing a Chiralcel OJ column (Daicel). Conditions: 98:2 hexanes: ethanol; Flow rate 0.60 mL/min; 30.8 min (major ent), 35.4 min (minor ent).

Compound 9.

^1H NMR (CDCl_3 , 400 MHz) δ 7.32 (m, 1H), 7.21 (m, 3H), 6.6 (t, $J = 4.0$ Hz, 1H), 4.78 (br s, 1H), 3.52 (br s, 1H), 2.30 (s, 3H), 2.34-2.15 (m, 2H), 1.93-1.78 (m, 3H), 1.66-1.58 (m, 1H); **^{13}C NMR** (CDCl_3 , 100 MHz) δ 201.3, 148.8, 141.4, 138.2, 135.8, 130.6, 129.6, 127.6, 124.9, 63.5, 29.9, 26.7, 19.7, 17.8; **IR** (film, cm^{-1}) 3478, 2943, 2867, 1633; **TLC R_f** 0.38 (3:1 toluene:EtOAc); $[\alpha]_D + 4.2$ ($c = 1.0$, CHCl_3 , 51% ee); **Exact mass** calcd for $[\text{C}_{14}\text{H}_{16}\text{O}_2 + \text{Na}] +$ requires m/z 239.1048. Found 239.1045 (ESI^+); **Assay of enantiomeric purity:** Enantiomers of product were separated by chiral HPLC employing a Chiralcel OD column (Daicel). Conditions: 98:2 hexanes: ethanol; Flow rate 0.75 mL/min; 14.2 min (major ent), 27.6 min (minor ent).

Compound 11.

^1H NMR (CDCl_3 , 400 MHz) δ 7.66 (dd, $J = 4.8$ Hz, 1.1 Hz, 1H), 7.62 (dd, $J = 4.0$ Hz, 1.1 Hz, 1H), 7.12 (m, 1H), 6.95 (m, 1H), 4.68 (br s, 1H), 3.46 (br s, 1H), 2.42-2.35 (m, 1H), 2.32-2.23 (m, 1H), 1.96-1.81 (m, 3H), 1.70-1.65 (m, 1H); **^{13}C NMR** (CDCl_3 , 100 MHz) δ 189.8, 143.2, 142.6, 139.9, 133.4, 133.3, 127.6, 64.1, 29.9, 26.2, 17.4; **IR** (film, cm^{-1}) 3440, 2936, 2867, 1621; **TLC R_f** 0.32 (3:1 Toluene:EtOAc); $[\alpha]_D + 37$ ($c = 1.0$, CHCl_3 , 74% ee); **Exact mass** calcd for $[\text{C}_{11}\text{H}_{12}\text{O}_2\text{S} + \text{Na}] +$ requires m/z 231.0456. Found 231.0459 (ESI^+). **Assay of enantiomeric purity:** Enantiomers of product were separated by chiral HPLC employing a Chiralcel OD column (Daicel). Conditions: 95:5 hexanes: ethanol; Flow rate 0.60 mL/min; 25.7 min (major ent), 23.2 min (minor ent).

Procedure for the Tandem Enantioselective Intramolecular MBH Reaction – Asymmetric Acylation Reaction

Pipecolic acid (2.6 mg, 0.02 mmol), water (41 μ L), THF (94 μ L), and NMI (30 μ L of a 0.33 M solution in THF) were added sequentially to a vessel containing aldehyde 1 (20 mg, 0.1 mmol). The reaction was stirred at room temperature for 48 hours, whereupon the yellow solution was concentrated, transferred to a round bottom flask, and azeotroped with toluene (5 x 10 mL). The residue was then dissolved in toluene (16.8 mL) and peptide 12 (7 mg, 0.005 mmol) and acetic anhydride (37.4 μ L, 0.4 mmol) were added. The solution was maintained at rt for 24 h and then quenched with methanol (2 mL). After concentration, the residue was dissolved in EtOAc (30 mL), washed with saturated NH_4Cl (20 mL), NaHCO_3 (20 mL), and brine (20 mL), and dried (Na_2SO_4). Concentration gave a yellow oil which was purified by flash chromatography (toluene:ethyl acetate 7:1) affording 2 (9.9 mg, 50% yield, >98% ee) as a clear oil.

ent-2-Ac

^1H NMR (CDCl_3 , 400 MHz) δ 7.70 (d, J = 7.0 Hz, 2H), 7.53 (t, J = 7.3 Hz, 1H), 7.43 (t, J = 7.3 Hz, 2H), 6.72 (m, 1H), 5.99 (m, 1H), 2.46-2.39 (m, 1H), 2.24-2.15 (m, 1H), 2.05-1.98 (overlapping s and m, 4H), 1.86-1.70 (m, 3H); **^{13}C NMR** (CDCl_3 , 100 MHz) δ 195.3, 169.8, 146.0, 137.8, 136.7, 131.7, 129.0, 127.9, 65.0, 28.0, 25.8, 21.3, 17.3; **IR** (film, cm^{-1}) 2942, 2867, 1733, 1652; **TLC** R_f 0.51 (30% EtOAc:hexanes); **Exact mass** calcd for $[\text{C}_{15}\text{H}_{16}\text{O}_3 + \text{Na}]$ + requires m/z 267.0997. Found 267.0988 (ESI^+). **Assay of enantiomeric purity:** Enantiomers of product were separated by chiral HPLC employing a Chiralcel OD column (Daicel). Conditions: 98:2 hexanes: ethanol; Flow rate 0.75 mL/min; 15.3 min (major ent), 16.6 min (minor ent).

Aldol Dimerization By-Product (3).

^1H NMR (CDCl_3 , 400 MHz) δ 9.42 (s, 1H), 7.92 (t, J = 7.6 Hz, 4H), 7.58-7.52 (m, 2H), 7.49-7.43 (m, 4H), 7.08-6.84 (m, 4H), 6.54 (t, J = 7.6 Hz, 1H), 2.51-2.34 (m, 8H), 1.75 (quintet, J = 7.6 Hz, 2H); **^{13}C NMR** (CDCl_3 , 100 MHz) δ 194.3, 190.2, 154.4, 147.8, 147.6, 142.2, 137.6, 132.6, 128.4, 126.4, 32.4, 31.5, 28.7, 27.3, 23.2; **IR** (film, cm^{-1}) 3056, 2936, 2861, 2722, 1671, 1647, 1620, 1596, 1451, 1287, 1224; **TLC** R_f 0.59 (3:1 toluene:EtOAc); **Exact mass** calcd for $[\text{C}_{26}\text{H}_{26}\text{O}_3 + \text{H}]$ + requires m/z 387.1960. Found 387.1943 (ESI^+).

4

6

8

10

[illegible]

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	18.637	MM	0.7152	4981.65039	116.09653	92.9642
2	22.165	MM	0.7643	377.02399	8.22196	7.0358

Totals : 5358.67438 124.31850

Results obtained with enhanced integrator!

Signal 2: DAD1 B, Sig=254,8 Ref=360,100

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	18.633	MM	0.7159	6719.94678	156.44354	92.1521
2	22.258	MM	0.8070	572.29016	11.81874	7.8479

Signal 1: DAD1 A, Sig=215,10 Ref=360,100

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	28.339	IM	0.9551	5366.59717	93.64401	90.4835
2	32.650	IM	0.9106	564.42731	10.33063	9.5165

Totals :

Results obtained with enhanced integrator!

Signal 2: DAD1 B, Sig=254,8 Ref=360,100

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	28.339	MM	0.9440	7522.39697	132.80659	89.7551
2	32.654	MM	0.9459	858.62701	15.12957	10.2449

```

NAME: 7
EXPNO: 1
PROCNO: 1
PROCNAME: 7
F2: 1
F3: 1
F4: 1
F5: 1
F6: 1
F7: 1
F8: 1
F9: 1
F10: 1
F11: 1
F12: 1
F13: 1
F14: 1
F15: 1
F16: 1
F17: 1
F18: 1
F19: 1
F20: 1
F21: 1
F22: 1
F23: 1
F24: 1
F25: 1
F26: 1
F27: 1
F28: 1
F29: 1
F30: 1
F31: 1
F32: 1
F33: 1
F34: 1
F35: 1
F36: 1
F37: 1
F38: 1
F39: 1
F40: 1
F41: 1
F42: 1
F43: 1
F44: 1
F45: 1
F46: 1
F47: 1
F48: 1
F49: 1
F50: 1
F51: 1
F52: 1
F53: 1
F54: 1
F55: 1
F56: 1
F57: 1
F58: 1
F59: 1
F60: 1
F61: 1
F62: 1
F63: 1
F64: 1
F65: 1
F66: 1
F67: 1
F68: 1
F69: 1
F70: 1
F71: 1
F72: 1
F73: 1
F74: 1
F75: 1
F76: 1
F77: 1
F78: 1
F79: 1
F80: 1
F81: 1
F82: 1
F83: 1
F84: 1
F85: 1
F86: 1
F87: 1
F88: 1
F89: 1
F90: 1
F91: 1
F92: 1
F93: 1
F94: 1
F95: 1
F96: 1
F97: 1
F98: 1
F99: 1
F100: 1

```

Signal 1: DAD1 A, Sig=215,10 Ref=360,100

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	30.834	MM	1.1348	2.84151e4	417.31009	89.2532
2	35.438	MM	1.1117	3421.38232	51.29292	10.7468

Totals : 3.18364e4 468.60301

Results obtained with enhanced integrator!

Signal 2: DAD1 B, Sig=254,8 Ref=360,100

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	30.834	MM	1.1400	4.22750e4	618.08105	88.9597
2	35.445	MM	1.1284	5246.50146	77.49103	11.0403

01 0
 02 0
 03 0
 04 0
 05 0
 06 0
 07 0
 08 0
 09 0
 10 0
 11 0
 12 0
 13 0
 14 0
 15 0
 16 0
 17 0
 18 0
 19 0
 20 0
 21 0
 22 0
 23 0
 24 0
 25 0
 26 0
 27 0
 28 0
 29 0
 30 0
 31 0
 32 0
 33 0
 34 0
 35 0
 36 0
 37 0
 38 0
 39 0
 40 0
 41 0
 42 0
 43 0
 44 0
 45 0
 46 0
 47 0
 48 0
 49 0
 50 0
 51 0
 52 0
 53 0
 54 0
 55 0
 56 0
 57 0
 58 0
 59 0
 60 0
 61 0
 62 0
 63 0
 64 0
 65 0
 66 0
 67 0
 68 0
 69 0
 70 0
 71 0
 72 0
 73 0
 74 0
 75 0
 76 0
 77 0
 78 0
 79 0
 80 0
 81 0
 82 0
 83 0
 84 0
 85 0
 86 0
 87 0
 88 0
 89 0
 90 0
 91 0
 92 0
 93 0
 94 0
 95 0
 96 0
 97 0
 98 0
 99 0
 100 0
 101 0
 102 0
 103 0
 104 0
 105 0
 106 0
 107 0
 108 0
 109 0
 110 0
 111 0
 112 0
 113 0
 114 0
 115 0
 116 0
 117 0
 118 0
 119 0
 120 0
 121 0
 122 0
 123 0
 124 0
 125 0
 126 0
 127 0
 128 0
 129 0
 130 0
 131 0
 132 0
 133 0
 134 0
 135 0
 136 0
 137 0
 138 0
 139 0
 140 0
 141 0
 142 0
 143 0
 144 0
 145 0
 146 0
 147 0
 148 0
 149 0
 150 0
 151 0
 152 0
 153 0
 154 0
 155 0
 156 0
 157 0
 158 0
 159 0
 160 0
 161 0
 162 0
 163 0
 164 0
 165 0
 166 0
 167 0
 168 0
 169 0
 170 0
 171 0
 172 0
 173 0
 174 0
 175 0
 176 0
 177 0
 178 0
 179 0
 180 0
 181 0
 182 0
 183 0
 184 0
 185 0
 186 0
 187 0
 188 0
 189 0
 190 0
 191 0
 192 0
 193 0
 194 0
 195 0
 196 0
 197 0
 198 0
 199 0
 200 0
 201 0
 202 0
 203 0
 204 0
 205 0
 206 0
 207 0
 208 0
 209 0
 210 0
 211 0
 212 0
 213 0
 214 0
 215 0
 216 0
 217 0
 218 0
 219 0
 220 0
 221 0
 222 0
 223 0
 224 0
 225 0
 226 0
 227 0
 228 0
 229 0
 230 0
 231 0
 232 0
 233 0
 234 0
 235 0
 236 0
 237 0
 238 0
 239 0
 240 0
 241 0
 242 0
 243 0
 244 0
 245 0
 246 0
 247 0
 248 0
 249 0
 250 0
 251 0
 252 0
 253 0
 254 0
 255 0
 256 0
 257 0
 258 0
 259 0
 260 0
 261 0
 262 0
 263 0
 264 0
 265 0
 266 0
 267 0
 268 0
 269 0
 270 0
 271 0
 272 0
 273 0
 274 0
 275 0
 276 0
 277 0
 278 0
 279 0
 280 0
 281 0
 282 0
 283 0
 284 0
 285 0
 286 0
 287 0
 288 0
 289 0
 290 0
 291 0
 292 0
 293 0
 294 0
 295 0
 296 0
 297 0
 298 0
 299 0
 300 0
 301 0
 302 0
 303 0
 304 0
 305 0
 306 0
 307 0
 308 0
 309 0
 310 0
 311 0
 312 0
 313 0
 314 0
 315 0
 316 0
 317 0
 318 0
 319 0
 320 0
 321 0
 322 0
 323 0
 324 0
 325 0
 326 0
 327 0
 328 0
 329 0
 330 0
 331 0
 332 0
 333 0
 334 0
 335 0
 336 0
 337 0
 338 0
 339 0
 340 0
 341 0
 342 0
 343 0
 344 0
 345 0
 346 0
 347 0
 348 0
 349 0
 350 0
 351 0
 352 0
 353 0
 354 0
 355 0
 356 0
 357 0
 358 0
 359 0
 360 0
 361 0
 362 0
 363 0
 364 0
 365 0
 366 0
 367 0
 368 0
 369 0
 370 0
 371 0
 372 0
 373 0
 374 0
 375 0
 376 0
 377 0
 378 0
 379 0
 380 0
 381 0
 382 0
 383 0
 384 0
 385 0
 386 0
 387 0
 388 0
 389 0
 390 0
 391 0
 392 0
 393 0
 394 0
 395 0
 396 0
 397 0
 398 0
 399 0
 400 0
 401 0
 402 0
 403 0
 404 0
 405 0
 406 0
 407 0
 408 0
 409 0
 410 0
 411 0
 412 0
 413 0
 414 0
 415 0
 416 0
 417 0
 418 0
 419 0

Signal 1: DND1 A, Sig=215,10 Ref=360,100

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	14.226	MM	0.5523	1.23251e4	371.93393	75.6058
2	27.571	MM	0.9235	3976.69873	71.76760	24.3942
Totals:				1.63018e4	443.70153	

Results obtained with enhanced integrator!

Signal 2: DAD1 B, Sig=254,8 Ref=360,100

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	14.227	IMP	0.5698	6423.48047	187.90344	75.1236
2	27.573	IMP	0.9629	2127.06934	36.81874	24.8764

[illegible]

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	23.247	MS	0.7706	3229.65259	69.85147	12.7838
2	25.737	MS	1.1360	2.20340e4	323.28015	87.2162
Totals:				2.52636e4	393.13162	

Signal 2: MWD1 B, Sig=254,16 Ref=360,100

Peak #	RetTime [min]	Type	Width [min]	Area [mAU*s]	Height [mAU]	Area %
1	23.247	MF	0.7676	4433.10254	96.25753	12.8845
2	25.736	MF	1.1380	2.99733e4	438.95941	87.1155

Figure 1. (Compound purity after extractive work-up)

14

13

14

-
- ¹ Still, W.C.; Kahn, M.; Mitra, J. *J. Org. Chem.* **1978**, *43*, 2923.
- ² (a) Montgomery, J.; Savchenko, A.V.; Zhao, Y. *J. Org. Chem.* **1995**, *60*, 5699-5701. (b) Richards, E.L.; Murphy, P.J.; Dinon, F.; Fratucello, S.; Brown, P.M.; Gelbrich, T.; Hursthouse, M.B. *Tetrahedron* **2001**, *57*, 7771-7784.
- ³ (a) Ramirez, F.; Dershowitz, S. *J. Org. Chem.* **1957**, *22*, 41-45. (b) Denney, D.B.; Smith, L.C.; Song, J.; Rossi, C.J.; *J. Org. Chem.* **1963**, *28*, 778-780.
- ⁴ Schreiber, S.L.; Claus, R.E.; Reagan, J. *Tetrahedron Lett.*, **1982**, *23*, 3867-3870.
- ⁵ (a) Dale, J.A.; Dull, D.L.; Mosher, H.S. *J. Org. Chem.* **1969**, *34*, 2543-2549. (b) Dale, J.A.; Mosher, H.S. *J. Am. Chem. Soc.* **1973**, *95*, 512-519. (c) Ward, D.E.; Rhee, C.K. *Tetrahedron Lett.*, **1991**, *32*, 7165-7166. (d) Ohtani, I.; Kusumi, T.; Kashman, Y.; Kakisawa, H. *J. Org. Chem.* **1991**, *56*, 1296-1298.