This item was submitted to [Loughborough's Research Repository](#) by the author.
Items in Figshare are protected by copyright, with all rights reserved, unless otherwise indicated.

Dust source identification using MODIS: a comparison of techniques applied to the Lake Eyre Basin, Australia

PLEASE CITE THE PUBLISHED VERSION

PUBLISHER

© Elsevier

VERSION

AM (Accepted Manuscript)

LICENCE

CC BY-NC-ND 4.0

REPOSITORY RECORD

Baddock, Matthew C., Joanna E. Bullard, and Robert G. Bryant. 2019. "Dust Source Identification Using MODIS: A Comparison of Techniques Applied to the Lake Eyre Basin, Australia". figshare.
<https://hdl.handle.net/2134/4523>.

This item was submitted to Loughborough's Institutional Repository (<https://dspace.lboro.ac.uk/>) by the author and is made available under the following Creative Commons Licence conditions.

For the full text of this licence, please go to:
<http://creativecommons.org/licenses/by-nc-nd/2.5/>

To be cited as: Baddock, M.C., Bullard, J.E., Bryant, R.G. 2009. Dust source identification using MODIS: a comparison of techniques applied to the Lake Eyre Basin, Australia, *Remote Sensing of Environment*, doi:10.1016/j.rse.2009.03.002

Dust source identification using MODIS: a comparison of techniques applied to the Lake Eyre Basin, Australia

^aMatthew C. Baddock, ^{a*}Joanna E. Bullard, and ^bRobert G. Bryant.

^aDepartment of Geography, Loughborough University, Loughborough, Leicestershire, LE11 3TU UK.

^bDepartment of Geography, The University of Sheffield, Western Bank, Sheffield, S10 2TN UK

* Corresponding author

Abstract

The impact of mineral aerosol (dust) in the Earth's system depends on particle characteristics which are initially determined by the terrestrial sources from which the sediments are entrained. Remote sensing is an established method for the detection and mapping of dust events, and has recently been used to identify dust source locations with varying degrees of success. This paper compares and evaluates five principal methods, using MODIS Level 1B and MODIS Level 2 aerosol data, to: (a) differentiate dust (mineral aerosol) from non-dust, and (2) determine the extent to which they enable the source of the dust to be discerned. The five MODIS L1B methods used here are: (1) un-processed false colour composite (FCC), (2) brightness temperature difference, (3) Ackerman's (1997: J.Geophys. Res., 102, 17069-17080) procedure, (4) Miller's (2003:Geophys. Res. Lett. 30, 20, art.no.2071) dust enhancement algorithm and (5) Roskovensky and Liou's (2005: Geophys. Res. Lett. 32, L12809) dust differentiation algorithm; the aerosol product is MODIS Deep Blue (Hsu et al., 2004: IEEE Trans. Geosci. Rem. Sensing, 42, 557-569), which is optimised for use over bright surfaces (i.e. deserts). These

are applied to four significant dust events from the Lake Eyre Basin, Australia. OMI AI was also examined for each event to provide an independent assessment of dust presence and plume location. All of the techniques were successful in detecting dust when compared to FCCs, but the most effective technique for source determination varied from event to event depending on factors such as cloud cover, dust plume mineralogy and surface reflectance. Significantly, to optimise dust detection using the MODIS L1B approaches, the recommended dust/non-dust thresholds had to be considerably adjusted on an event by event basis. MODIS L2 aerosol data retrievals were also found to vary in quality significantly between events; being affected in particular by cloud masking difficulties. In general, we find that OMI AI and MODIS AQUA L1B and L2 data are complementary; the former are ideal for initial dust detection, the latter can be used to both identify plumes and sources at high spatial resolution. Overall, approaches using brightness temperature difference (BT10-11) are the most consistently reliable technique for dust source identification in the Lake Eyre Basin. One reason for this is that this enclosed basin contains multiple dust sources with contrasting geochemical signatures. In this instance, BTD data are not affected significantly by perturbations in dust mineralogy. However, the other algorithms tested (including MODIS Deep Blue) were all influenced by ground surface reflectance or dust mineralogy; making it impossible to use one single MODIS L1B or L2 data type for all events (or even for a single multiple-plume event). There is, however, considerable potential to exploit this anomaly, and to use dust detection algorithms to obtain information about dust mineralogy.

Dust source identification using MODIS: a comparison of techniques applied to the Lake Eyre Basin, Australia

1. Introduction

Atmospheric mineral aerosols (termed here dust) play an important role in the land-atmosphere-ocean system (Ridgwell, 2002; Jickells et al., 2005; Waeles et al., 2007). For example, they affect soil nutrients at source and sink (McTainsh & Strong, 2007; Muhs et al., 2007; Li et al., 2007; Reynolds et al., 2006; Soderberg & Compton, 2007; Swap et al., 1992; Wang et al., 2006), the radiative forcing of the atmosphere (Haywood & Boucher, 2000; Hsu et al., 2000; Satheesh & Moorthy, 2005; Yoshioka et al., 2007) and may regulate phytoplankton activity of oceans (de Baar et al., 2005; Erickson et al., 2003; Mackie et al., 2008; Piketh et al., 2000; Wolff et al., 2006). The impact of dust in the Earth's system depends on characteristics such as particle size, shape and mineralogy (in particular iron content: Jickells et al., 2005; Mahowald et al., 2005). Whilst these characteristics can change during dust transport (Desboeufs, 2005; Mackie et al., 2005) they are initially determined by the terrestrial sources from which the particles are entrained.

The detection and mapping of dust events and dust transport pathways has benefited greatly from the use of remote sensing, and at the global scale major dust source regions have been identified using satellite data, such as from the Total Ozone Mapping Spectrometer (TOMS; Prospero et al., 2002; Washington et al., 2003). The passage of dust along specific regional transport pathways over land and ocean and the behavior of individual dust events have also been tracked using TOMS and OMI (Ozone Monitoring Instrument; e.g. Alpert et al. 2004) and at higher temporal and spatial resolutions using data from, amongst others, AVHRR (Advanced Very High Resolution Radiometer; e.g., Evan et al., 2006; Zhu et al., 2007), GOES-VISSR (Geostationary Operational Environmental Satellite, Visible Infra-Red Spin-Scan Radiometer, e.g., MacKinnon et al., 1996), METEOSAT (e.g., Moorthy et al., 2007), MODIS (Moderate Resolution Imaging Spectroradiometer, e.g., Badarinath et al., 2007; Gassó & Stein, 2007;

Kaskaoutis et al. 2008; McGowan & Clark, 2008; Zha & Li, 2007), MSG-SEVIRI (Meteosat Second Generation-Spinning Enhanced Visible and InfraRed Imager; e.g., Schepanski et al., 2007) and SeaWIFS (Sea-viewing Wide Field-of-View Sensor; e.g., Eckardt & Kuring, 2005). Sensor-retrieved parameters (such as MODIS aerosol size parameters; Dubovik et al., 2008; Jones & Christopher, 2007; Kaufman et al. 2005) or complex statistical analyses (such as Principal Component Analysis; e.g. Argarwal et al. 2007; Jones & Christopher, 2008; Zubko et al. 2007) have also been used to differentiate dust and non-dust with some success.

Systematic determination of both the geomorphological and geochemical variability of dust sources, and hence the variability of the sediments which are entrained and transported, requires as accurate and precise an identification of the upwind (source) end of the dust plume as possible. Researchers have recently started to use remote sensing data to achieve this (e.g., Bullard et al., 2008; Lee et al., 2008; Zhang et al., 2008), but with varying levels of success. The ability to use remotely-sensed data both to detect a dust plume and identify the location from which it has originated is affected by several factors including the radiative transfer properties of the material emitted, the radiative properties of the ground/ocean surface over which the plume is transported, the size and density of the dust plume, the time of satellite overpass relative to dust emission, the presence or absence of cloud, the horizontal and vertical plume trajectory, and the sensor characteristics and radiative transfer model used to detect dust. In many respects, the relative impacts of these factors on dust source determination are hard to determine without close reference to surface meteorological data (e.g. wind speed and visibility records) and ground-based aerosol determination records (e.g. AERONET – Aerosol Robotic Network) which can allow comparative characterisation of individual dust events (e.g. Bullard et al., 2008; Mahowald et al., 2007). Even where these records exist, the direct comparison of ground and remote sensing data retrievals to determine dust sources can be problematic, with some remote sensing data products being unable consistently to detect dust events due to the factors listed above; particularly the presence of cloud, and the existence of low contrast between

dust plume and ground/ocean surface (e.g. Gassó & Stein, 2007; Bullard et al., 2008). The principal aim of this paper is to evaluate in detail the use of MODIS data, one of the most widely and successfully-used sensors, for improved identification of dust source locations. This paper varies in emphasis from many previous studies because the focus is on the *precision* with which the upwind (source) location of the plume can be discerned, rather than on the simple determination of plume location, density and trajectory. Specifically, we compare five methods of using MODIS Level 1 band data and one MODIS Level 2 aerosol product and evaluate them in terms of: (a) how well they enable the differentiation of dust and non-dust (cloud, smoke, volcanic aerosols) and, (b) the extent to which it is possible to discern the location of the dust source (i.e. the upwind part of the dust plume - or 'dust head') and how much this varies from method to method. The influence of environmental factors such as plume density and mineralogy on source detection by MODIS will also be evaluated.

2. Data and Methods

2.1 Data

Mineral aerosol (dust) can be detected and mapped through remote sensing via inversion of radiative transfer models which operate in the following wavelengths: (a) ultraviolet (UV 0.315-0.4 μm) via absorption (e.g. TOMS AI; Torres *et al.*, 1998), (b) visible (VIS 0.38-79 μm) via scattering (e.g. Tanré and Legrand, 1991), and (c) thermal infrared (TIR 8-15 μm) via contrasting land/aerosol emissivity and/or temperature (e.g. Ackerman, 1997). Due to constraints of sensor design, observations by remote sensing systems operating in VIS wavelengths can be determined at higher resolution (pixel size = x) than those made in the TIR (pixel size = $x*2-4$) and UV (pixel size = $x*100-200$), and this has implications for both plume and source detection using these approaches. Radiative transfer model inversion of aerosol observations made within (or via combinations of) each of the three wavelength ranges often provides either a relative indication of aerosol concentration (e.g. via TOMS AI), or a calibrated (e.g. through comparison

with AERONET observations) measure of wavelength-dependent total aerosol optical thickness/depth (AOT/D). The success of the radiative transfer model inversion in each case is often complicated by factors such as the non-spherical nature of the mineral aerosol, changes in the chemical/physical nature of the material, and location within the atmosphere during transport. In addition, over very bright surfaces (e.g. desert regions and urban areas), in the presence of cloud, and at night, mineral aerosol detection using UV/VIS/TIR wavelengths can become increasingly uncertain (e.g. Kaufman et al., 2000). The short-term nature of some mineral aerosol events (often <1 day) also means that an understanding of any bias associated with mineral aerosol detection at the time of satellite over-passes and temporal sampling (i.e. either am or pm data collection time) is needed in order to characterize fully the emission and transport process. In order to evaluate, compare and contrast mineral aerosol detection approaches, a range of remote sensing data are used here (see Table 1).

<Insert Table 1>

2.1.1 MODIS Data

Data from the Moderate Resolution Imaging Spectroradiometer (MODIS) were used to make comparisons of retrievals using VIS and TIR (often combined) approaches. MODIS makes observations using 36 spectral bands with wavelengths from 0.41 to 14.4 μm and nadir spatial resolutions of 0.25 km, 0.5 km, and 1 km. It is currently operating onboard the NASA Earth Observing System (EOS) Terra and Aqua satellites, launched in December 1999 and May 2002, respectively. Daily MODIS Level 1B (L1B) 1 km data (MOD021KM = Terra, and MYD021KM = Aqua) used in this work have been processed to convert the sensor's on-orbit responses in digital numbers to radiometrically calibrated and geo-located data products (v5.06 processing for Terra and v5.07 for Aqua). Data were obtained from the Level 1 and Atmosphere Archive and Distribution System (LAADS; <http://ladsweb.nascom.nasa.gov/>). Details of images dates and subsequent processing of MODIS L1B data are outlined below.

Daily MODIS Level 2 Aerosol data are produced at the spatial resolution of a 10 x 10 km (at nadir) pixel array. There are two MODIS Aerosol data product file types: MOD04_L2, containing data collected from the Terra platform and MYD04_L2, containing data collected from the Aqua platform. Here we only use the MYD04 Aqua product because to date Deep Blue (see below) retrievals are not yet available for MOD04 Terra data. Aerosol properties within MYD04_L2 are derived by the inversion of MODIS observed reflectances at 500 m resolution using pre-computed radiative transfer look-up tables based on dynamical aerosol models (Kaufman et al., 1997; Remer et al., 2005). Derivation of aerosol from these data is far from straightforward and, in initial versions of the MODIS aerosol product, the ability to retrieve aerosol optical thickness (AOT) and single scattering albedo over bright-reflecting surfaces has been problematic because the algorithm relies in part on the initial detection of dark surfaces or targets (Kaufman et al., 2000). In addition, the cloud screening has been shown to have problems where mis-identification of some dust plumes as cloud has led to artifacts in the final data (e.g. as noted by Brindley and Ignatov, 2006). These products have been under continued and careful evaluation and development, and product MYD04 (see http://modis-atmos.gsfc.nasa.gov/C005_Changes/C005_Aerosol_5.2.pdf) has recently received an improved aerosol determination (via reprocessing to collection 5.1/2; Levy et al., 2006, 2007; Remer et al., 2006) over bright surfaces through the integration of a revised determination of AOT over land (Levy et al., 2007), and inclusion of the Deep Blue algorithm (Hsu et al., 2004; Hsu et al., 2006). Here we evaluate the Deep Blue algorithm, which relies on the blue wavelengths and libraries of surface reflectance to make retrievals over bright surfaces (Hsu et al., 2004).

The Deep Blue processing approach involves the following processing elements: (1) Rayleigh Correction for Terrain Elevation in the following MODIS channels: R8 (0.405-0.42 μm), R3 (0.459-0.479 μm) and R1 (0.62-0.67 μm); (2) Cloud Screening using: R8 (3 x 3 pixel spatial variance) and R3/R8 AI; (3) the surface reflectance for a given pixel is determined from a clear-scene database based upon its geo-location; (4) R8, R3 and R1

reflectances are then compared to radiances contained in a lookup table with dimensions consisting of solar zenith, satellite zenith, and relative azimuth angles, surface reflectance, AOT, and single scattering albedo; (5) a maximum likelihood method is used to compute a mixing ratio between dust and smoke models until the calculated spectral reflectances make the best match with those that are measured; and (6) for mixed aerosol conditions, once the aerosol models and the mixing ratio that produce the best match are determined, the values of AOT and Ångström exponent are reported. For dust-dominant cases, the values of single scattering albedo are retrieved in addition to these parameters. MODIS Deep Blue data within MYD04_L2 includes AOT (τ) determination at 0.412, 0.47, 0.55 and 0.66 μm , although only the 0.412 μm data are used here. MYD04_L2 data were obtained from the Level 1 and Atmosphere Archive and Distribution System (LAADS; <http://ladsweb.nascom.nasa.gov/>). The typical aerosol optical thickness for visible light in clear air is 0.1, very hazy skies have AOTs of ≥ 0.3 . During initial processing, typical scale (0.001) and offset (0) values were applied to MYD04_L2 AOT data prior to display and subsequent data processing.

2.1.2 AURA OMI

This paper focuses on an evaluation of MODIS data but for each case study, in addition to MODIS L1B and L2 aerosol data, co-incident data from an independent sensor, the Ozone Monitoring Instrument (OMI) were also acquired. OMI is on the Aura satellite (launch date: July 2004) which flies as part of the NASA A-Train constellation (http://aqua.nasa.gov/doc/pubs/A-Train_Fact_sheet.pdf) a few minutes behind the Aqua satellite. OMI is designed to continue the Total Ozone Mapping Spectrometer (TOMS) record for total ozone and other atmospheric parameters related to ozone chemistry and climate. OMI measurements are sensitive to aerosol absorption in UV wavelengths, thus providing an independent source of information relating to mineral aerosol detection in the scene under observation. In addition, and unlike MODIS, OMI AI (Absorbing Aerosol Index: e.g. Torres et al., 2007) is sensitive to aerosol absorption even when the particles are above cloud and AAI is therefore derived successfully in both cloudless and cloudy conditions (although see Ahn et al., 2008). OMI has a ground resolution of 13 x 24 km

(nadir) and uses a retrieval algorithm similar to the one used by TOMS (Torres *et al.*, 1998). The OMI AI is defined as follows:

$$\text{OMI AI} = 100 \log_{10} (I_{360}^{\text{Meas}} / I_{360}^{\text{Calc}}) \quad (\text{Eq.1})$$

where I_{360}^{Meas} is the measured 360 nm OMI radiance and I_{360}^{Calc} is the calculated 360 nm OMI radiance for a Rayleigh atmosphere. Under most conditions, the AI (Eq.1) is positive for absorbing aerosols and negative for non-absorbing aerosols (pure scattering). An AI >1 is typical of absorbing aerosols such as smoke or dust (Gassó & Stein, 2007; Kubilay *et al.*, 2005; Washington *et al.*, 2003). In this instance, we have chosen to use the OMI-Aura_OMTO3E data, which is a daily Level 3 global gridded product which is generated by binning the original pixels from the Level 2 data products (15 orbits per day; 13 x 24 km spatial resolution at nadir) into a 0.25 x 0.25 degree global grid.

2.2 Methods

2.2.1 Study region and event selection

The performance of different MODIS dust detection methods in identifying source locations involved the analysis of four dust events which all originate in the same drainage basin. The Lake Eyre Basin (LEB), Australia was chosen for several reasons. First, it has been identified as a persistent and significant southern hemisphere dust source on the basis of surface observations (Middleton, 1986) and using TOMS AI (Washington *et al.*, 2003). Second, it is the only inland basin dust source region in Australia, a geographically-isolated continent distant from other dust sources. Consequently, within the LEB there is less potential for interaction with other major dust sources than would be the case, for example, in the Sahara (Prospero *et al.* 2002) or China (Shao & Wang, 2003). Third, the basin is large enough to give rise to several major dust events each year, but not such an intense dust source as to make it difficult to discern individual plumes.

The LEB covers 1.14 million km², with mean annual rainfall of less than 125 mm and annual potential evaporation in excess of 2500 mm. There are several different sedimentary environments in the LEB, all of which emit dust. The most significant of these are: (1) aeolian deposits covering 33% of the basin area and accounting for 37% of the dust plumes, (2) alluvial deposits and floodplains (11.55% area, 30% dust plumes), and (3) ephemeral lakes and playas which cover only 2.26% of the basin area but from which originate 29% of the dust plumes making these the most intense dust sources (figures averaged over 2003-6: Bullard et al., 2008). Inter- and intra-annual variability of dust storm frequency in the LEB is high, responding to changes in synoptic pressure distributions across the continent (Ekstrom et al., 2004). The Sprigg Model, which characterizes dust transporting wind systems in Australia (Sprigg, 1982), suggests that as frontal systems pass over the LEB, pre-frontal northerly and post-frontal southerly winds can entrain dusts which travel southeast or northwest respectively. It is important to note, however, that the estimated total annual number of dust events in the LEB varies not only in response to climate but also as a result of differences in how events are defined. In a previous study (Bullard et al., 2008), we examined MODIS imagery for all days (between July 2003 and June 2006) where at least one meteorological station in the LEB (or within 250 km of the catchment boundary) recorded a dust-induced reduction in visibility to ≤ 1 km (which corresponds to the WMO definition of a dust storm). Whilst there are some inconsistencies in the relationship between visibility records from meteorological stations and other indicators of dust emissions, (including AERONET, TOMS AI and TOMS AOD: Mahowald et al., 2007) and the spatial distribution of meteorological stations across the arid LEB is sparse which means a number of events will be missed, visibility remains a useful criteria for identifying days on which significant dust events have occurred. From the 43 days on which dust events were identified four case studies were chosen to illustrate key types of event that occur in the LEB, and also to include factors which can significantly affect dust plume and source identification (i.e. single /multiple dust plumes and varying amounts of cloud; see Figure 1; Table 2). Although there are versions of some dust detection algorithms designed to work at night (e.g. Wald et al., 1998), we focus on daytime events

here so that the influence of surface reflectance on dust source identification can be explored.

<Insert Figure 1> -

<Insert Table 2>

2.2.2 MODIS Level 1B Processing Algorithms

As outlined earlier, mineral aerosol is sometimes detectable on unadjusted VIS satellite images (particularly over the ocean), but because mineral aerosol can have similar reflectivity to the desert surfaces from which it is entrained it can be difficult to detect over land. In addition, mineral aerosol is often hard to differentiate from cloud, sea salt and anthropogenic pollution. As a result of this, and also due to problems with the performance of MODIS L2 aerosol products (section 2.1.1), a number of studies have used changes in brightness temperature (TIR) to detect mineral aerosol over land surfaces. Initial attempts using single TIR channel data, such as that by Shenk & Curran (1974) using Nimbus-THIR (Temperature Humidity Infrared Radiometer) 11 μm data, had limited success because changes in surface emissivity at this wavelength can be misinterpreted as dust (Roskovensky & Liou, 2003; 2005). As a result of observed variability in the emissive and transmissive nature of mineral aerosols within multiple TIR wavelength ranges, other researchers have used methods based on brightness temperature difference (BTD) in either two or three wavelength ranges, typically 11-12 μm bands (bi-spectral split window technique) or near 8, 11 and 12 μm bands (tri-spectral) (e.g. Ackerman, 1997). BTD values from this method reveal temperature differences that exist between the ground surface and cooler mineral aerosol while at the same time are largely unaffected by absorption from other atmospheric gases (Darmenov & Sokolik, 2005). In addition to detecting dust over land, these approaches may also allow discrimination between cloud and dust when both exist in the vicinity of each other.

Here we initially apply the simple BTD approach detailed by Ackerman (1997) to MODIS L1B data (Table 3). Using this methodology it has been inferred that (BTD; 11.03-12.02 μm or MODIS BT₃₁-BT₃₂) values <0 K signify

the presence of mineral aerosol (dimensionless) and BT_D values ≥ 0 K indicate no mineral aerosol. While developing the MODIS cloud mask, Ackerman et al. (2002) have also placed the mineral aerosol detection threshold at < -1 K. Although Ackerman's (1997) analysis implied that the 0 K threshold could be widely used over a range of land surfaces, it is likely that this will vary slightly according to variability in the emissive/transmissive nature of the mineral aerosol. This in turn is determined by factors such as mineralogy as well as processes acting upon the aerosol as it is transported in the atmosphere. Mineralogical composition is an important control on the TIR radiative properties of mineral aerosol and can vary significantly from region to region (e.g. Claquin et al., 1999; Caquineau et al., 2002; Satheesh & Moorthy, 2005). Darmenov and Sokolik (2005) investigated the TIR radiative signature of dust transported over oceans from 7 different regions and located the BT_D (11.03 -12.02 μm) aerosol detection threshold at 0.5, -0.2, -1.0 and -0.4 K for the Nubian, Thar, Gobi/Taklimakan and Australian deserts respectively; but could not locate a clear threshold to distinguish mineral aerosol from cloud for dust over oceans sourced from NW Africa, Libya or the Iranian desert. It may also be the case that the threshold varies for a single geographical region, the precise value being dependent on factors such as the density of the dust plume (Darmenov and Sokolik, 2005) or local variation in dust source mineralogy (e.g. iron-rich sources versus illite-rich sources). This simple bi-spectral split window approach will be applied here to identify appropriate aerosol detection thresholds over land for the Lake Eyre Basin.

Using BT_D as a basis, a range of more complex algorithms has been developed that combine BT_D and VIS wavelengths to detect mineral aerosol over land and remove the effects of dense cloud cover, which can obscure dust, and cirrus clouds which have similar reflectance and BT_D properties to fine dust particles. In this paper we evaluate two of these cloud-removal approaches applying them to MODIS L1B data for the LEB. The first is the multispectral dust enhancement algorithm of Miller (2003) which exploits the fact that dust particles can have contrasting VIS reflective properties when compared to cloud (Table 3). In this model, an inverse brightness temperature difference is used (BT_D; 12.02-11.03 μm or MODIS BT₃₂ – BT₃₁)

which is rescaled/normalized to lie within the -2 to $+2$ K range. Based on Miller's (2003) algorithm the mineral aerosol output (D) has values constrained between 1.3 and 2.7 (dimensionless). In addition this approach, through manipulation of the red (R), green (G) and blue (B) display, enables mineral aerosol to be visually differentiated from cloud using colour (D is loaded on the red color gun). The second approach is that of Roskovensky and Liou (2005) and Hansell et al., (2007) which focuses on the differentiation of mineral aerosol from cirrus clouds by combining BTD ($11.03\text{--}12.02\text{ }\mu\text{m}$) and VIS wavelengths (reflectance ratio of $0.54\text{ }\mu\text{m}/0.86\text{ }\mu\text{m}$). In the final output image, values of $D > 1$ (dimensionless) indicate mineral aerosol is present and values ≤ 1 indicate cirrus cloud or non-mineral aerosol in the scene (Table 3). Inclusion of the reflectance ratio in this case reduces the amount of false detection of dust over land observed by Ackerman et al. (2002).

Although the majority of studies cited above have used data from MODIS it is worth noting that similar approaches have been explored using data from other sensors such as AVHRR, HIRS/2, GOES-8 and MSG-SEVIRI with varying degrees of success (e.g. Legrand et al. 1989, Sokolik, 2002, Schepanski et al., 2007).

<Insert Table 3>

Table 3 includes a summary of the default threshold values used to differentiate dust from non-dust in each of the original algorithms used here. The threshold values used in these algorithms are sensitive to varying atmospheric conditions, surface reflectance, dust density and dust mineralogy, but are formulated in a manner such that they allow a certain degree of tuning to adjust for specific conditions such as regional variability (Darmonov & Sokolik, 2005), or for dust blowing over land or ocean (Roskovensky & Liou (2005). In this study, we verified the published models by using the authors' original data and study-events both to check the set up of the algorithms and to ensure we could reproduce the initial values of dust/non-dust threshold and coefficients used. For the four case studies presented here, we therefore established event-specific thresholds using the

approach suggested by each author (Table 4). The definition of thresholds therefore involved the interrogation of pixel histograms for each scene-algorithm combination (see below). In each case, peaks were found to be attributable to specific scene components (e.g. densities and types of cloud and aerosol), and thresholds were chosen to represent the value which was best able to identify dust in the scene, as judged by the user (Figure 2a). In each case, peaks representing dust were relatively easy to identify and disinter from other scene components, and the effects of the choice of dust threshold in each case is outlined below. Given the rather inflexible nature of this approach, data from other study regions, where atmospheric conditions and water vapor concentrations vary more significantly, may pose a challenge to the straightforward identification and threshold determination for dust peaks outlined here.

Figure 2 shows the histograms used to derive BTD thresholds and Miller's D for each event. For the Roskovensky and Liou (2005) output the dust/non-dust threshold remained at 1, but the D-parameter scaling factor 'a' and BTD offset 'b' were adjusted for each event by using the midpoints between the clear sky and dust histograms of the reflectance ratio and BTD respectively (Figure 3).

<Insert Figure 2>

<Insert Figure 3>

<Insert Table 4>

2.2.3 Evaluation of output images

To evaluate the different MODIS dust detection algorithms it is necessary to have a common reference against which to compare the output data. For each of the four dust events examined, an eight panel figure was produced. In each case, panel (a) represents the MODIS VIS image (where red = band 1, green = band 4, blue = band 3). Panel (b) represents the bi-spectral brightness temperature difference ($BTD = BT_{31} - BT_{32}$) with no dust threshold applied. This is the principal image against which the outputs from the different algorithms outlined in panels (c) Ackerman (1997), (d) Roskovensky & Liou (2005) and (e) Miller (2003) were compared, because a

bi-spectral BTM is a common component of each of these Level 1B MODIS algorithms. What is evaluated therefore is the extent to which the additional components of the algorithms actually led to improved dust source detection. In addition, a simple objective comparison of the outputs from each of the Ackerman (1997), Roskovensky & Liou (2005) and Miller (2003) algorithms is shown in panel (f). To produce this, boolean outputs from panels (c), (d) and (e), where pixels were categorized as dust (=1) or non-dust (=0) were colored red, green and blue respectively and combined to create a color composite output image. For example, if a pixel was categorized as dust following Ackerman's (1997) procedure it will appear red; if categorized as dust by both Ackerman (1997) and Miller (2003) it will appear pink; if all three algorithms categorize it as dust it will appear white; if all three categorize it as non-dust it will appear black (Figure 4).

<Insert Figure 4>

Panels (g) and (h) represent the two dust products, MODIS L2 aerosol (MYD04) Deep Blue AOT and OMI AI respectively. Although the spatial resolution of the data is lower than the MODIS, OMI AI provides an independent check on the spatial extent and intensity of aerosol retrieval for all panels because it is not derived from MODIS.

Whilst a comparison of the different approaches to dust detection will help to understand how MODIS can best be used to identify the presence or absence of dust, the main aim of this paper is to evaluate the use of MODIS for identifying dust sources. This means that the way in which the upwind edge of the dust plume is depicted is of most interest. In these comparison figures, the areas highlighting the active sources for each event (denoted by coloured squares; e.g. figure 5), which were used to compare the outcome of the techniques, were determined by an informed approach. Since the BTM principle is a component of all of the evaluated algorithms (panels c-e), sources were determined from a combination of the scene BTM plus other readily available information, including the use of wind direction data to ascertain the upwind side of plumes. From the companion dust source

inventory work of Bullard et al. (2008), certain sources in the LEB region could be identified as relatively recurrent points of emission; locations which had a record of acting as source areas for several different dust events across the three year study period. The case study events here were therefore chosen to ensure that several of the plume origins used for comparison were from 'proven' dust sources. Consequently, it is worth noting that the persistence of certain key sources allowed their identification with a further confidence when flagged as active in each BTM scene. One such example is the point-source located at the south east margin of Lake Callabonna, South Australia (centred on 140°15'0E, 30°S.). This well-studied source location was seen to be active in both the third and fourth case studies, where BTM data are able accurately to indicate it as an area where a plume has originated (Figures 7 and 8). The relative performance of the dust enhancement models was evaluated on this basis. It is worth noting that extensive background knowledge of dust events in the LEB was used to help verify source locations determined from BTM data and this may not be possible in areas where comparable auxiliary data are unavailable. Nevertheless, in this study, we were able to carefully and thoroughly assess model performance for detection of dust emanating from known source locations.

As noted earlier, there are some additional caveats to the use of remote sensing to determine (or infer) dust source locations; these include the relative timings of the satellite overpass and the onset of dust emissions (which might affect not only the location of the plume head, but also the density of the dust), and the fact that only the upwind dust source can be located with any additional contributing sources lying under the dust plume possibly going undetected. To test the likely impact of some of these issues, HYSPLIT (<http://www.arl.noaa.gov/ready>) was used to calculate possible trajectories and plume concentrations for each event. These data are not presented here but confirm that the dust emitted during the events was close to, or at, source at the time of data capture and rarely reached an altitude of more than 500 m. This suggests that overpass timings for remote sensing data capture were likely to have provided data suitable for source identification.

3. Results

3.1 Event 1: 7th October 2005

For this event, raised dust is quite difficult to observe in the visible scene (Figure 5a) and sources are not at all apparent regardless of the level of contrast enhancement applied. However, BTD analysis (Figure 5b) reveals the presence of dust plumes, which appear as dark streaks at the centre of the image. Much of the cloud in the lower left of Figure 5b also appears dark, indicating some overlap in the thermal signature of cloud and dust in this scene. This is further highlighted in Figure 5c where a dust/non-dust threshold of 0 (dust <0 , cloud >0) has been applied. Here, not only are parts of the dust plumes categorized as dust, but so too are some of the patches of cloud. Although this simple threshold effectively separates cirrus cloud (white/light grey in Figure 5b) from the dust plumes, the thicker areas of cloud (which have similar BTD values to the dust) are mis-identified. Adjustment of the dust/non-dust threshold for this scene highlighted that there was no single BTD value that could differentiate these two components. In terms of identification of the sources of dust in this scene, there was an observed offset between the upwind dust heads shown in Figure 5c and the dust sources identified using Figure 5b. This is most likely to be because the dust at source is still close to the ground surface and therefore has a less pronounced thermal contrast with the ground surface than airborne dust further downwind of the source which will have risen to a higher atmospheric level. Given the dust/non-dust threshold applied to the scene is 0, this also indicates that some of the dust from this location can have a BTD value of >0 (Figure 5b).

<Figure 5>

Application of the Roskovensky and Liou (2005) algorithm (Figure 5d) effectively removes the cirrus cloud from the scene, and only a small area of the remaining cloud is included when a dust threshold (using event-specific 'a'

and 'b' coefficients; Table 4) is applied. The upwind ends of the main dust plumes map on to the same source locations inferred from BTM (Figure 5b), with the exception of the most northwestern plume which is not detected. Miller's (2003) algorithm clearly differentiates the cloud from the main dust plume (Figure 5e) which is picked out in red. With the exception of the northernmost plume, source detection is comparable to those in Figure 5b. However, there are some parts of this scene where dust is likely to have been mis-identified. These areas (marked 'FS' in Figure 5e) are patches on the ground surface where fires have changed the ground surface reflectance characteristics significantly (Jacobberger-Jellison, 1994), and suggest that the Miller algorithm is sensitive to ground reflectance variability. Given that the fire scar is clearly discernible in the visible image (Figure 5a), but not when the other techniques (which rely more heavily on BTM to detect dust) are used, this implies that the component within the Miller (2003) algorithm that uses VNIR wavelengths is slightly over weighted in this application. Figure 5f shows the extent to which the three MODIS L1B algorithms agree and highlight co-incident pixels containing dust. Whilst all three pick out the main central plumes of dust, there are considerable differences elsewhere in the scene. In particular, the Miller (2003) algorithm suggests a much more extensive plume of dust than the other two approaches, especially in the northeast. All of the techniques misidentify some of the most dense cloud as dust, with Ackerman (1997) and Miller (2003) performing particularly poorly.

The MODIS Deep Blue AOT image (Figure 5g) shows that the cloud mask applied in this instance is effective in separating the clear or dusty sky from the clouds, and some dust is detected (AOT values close to 0). The dust plumes in the far right of the scene that are highlighted in previous panels (Figure 5 b, c, and d) are clearly defined, but the main central plume is less obvious. In comparison, despite the relatively coarse resolution, the OMI AI image (Figure 5h) depicts clearly the central and far right plumes, and with similar AI values. This suggests that the inability of Deep Blue to detect both of these plumes is not likely to be due to vastly different aerosol densities in each plume. Instead, one possibility (explored further in section 4.6) is that MODIS Deep Blue data are actually more sensitive to variations in dust colour

(mineralogy) than OMI. There is also an area in Figure 5g that is excluded by the Deep Blue cloud mask (marked 'FP' in Figure 5g) where cloud is not apparent. On the ground, this is a floodplain and the high surface reflectance characteristics also cause confusion for the cloud mask when applied to event 3 (Figure 7g). Overall, the location of the dust plumes outlined by OMI AI data correspond very closely to the position of the plumes in the MODIS 1B algorithm outputs. In addition, the gridded AI data capture the extent and variation of aerosol density apparent in the other panels. Although plume identification is acceptable, the coarse resolution of the aggregated OMI AI data ($0.25^\circ \times 0.25^\circ$) mean these data are less able to define the dust source location or the nature of the surface sedimentary environments with the same precision as can be achieved using the combination of outputs from the MODIS L1B algorithms (e.g., Bullard et al., 2008; Lee et al., 2008).

3.2 Event 2: 24th September 2006

In the second event, the downwind (northerly) limit of the advancing dust is very distinct in the MODIS VIS (Figure 6a). The upwind edges of the plume are, however, not distinct and are in places difficult to differentiate from the underlying bright desert surface. For Event 1, the dust/non-dust thresholds or coefficients chosen are the same as those recommended in the published techniques. However, if these values are applied to Event 2 some problems become evident. Figure 6 shows the visible MODIS (panel a) and BTD values (panel b) for Event 2 and the results of applying the Event 1 thresholds (Table 4; Figure 6 panels c-f). There is no possibility of identifying dust sources using the Roskovensky and Liou (2005) or Miller (2003) algorithms with these thresholds. The Roskovensky and Liou (2005) output suggests that the dust plume fills most of the panel, whilst no dust is highlighted using the Miller (2003) approach. This is emphasized in Figure 6f which shows there are no areas of the image that all the different approaches identify as containing dust.

<Figure 6>

For this reason, thresholds and parameters appropriate to this event were determined using the histogram approach (see Table 4 for values). The results of applying these event-specific thresholds are shown in Figure 7. The dust source areas for this event and extent of the plume are reasonably well discerned using BTD (Figure 7b), which also reveals several other minor plumes that are not evident in the VIS. When an event-specific threshold is applied to the BTD (Figure 7c) most of the plume is highlighted but some of the thin, discrete plumes are not identified as dust or are foreshortened. This again suggests the use of the dust/non-dust threshold can affect the accurate identification of dust sources. The main plume is successfully identified using Roskovensky and Liou's (2005) algorithm (Figure 7d) but the source areas are poorly represented. Despite extensive experimentation with the 'a' and 'b' coefficients to improve dust detection, it was not possible to pick out the westernmost dust plumes without introducing a significant component of the ground surface reflectivity to the dust determination.

<Figure 7>

Using Miller's (2003) algorithm (Figure 7e), the maximum D value for this event falls below the published dust >1.3 threshold (see Table 3) necessitating an adjustment of this threshold such that dust >-0.55 . Although this adjustment enhances the dust visualization significantly, it does not do so without introducing further artifacts. First, some areas of the plume evident in Figure 7b were not highlighted, for example the thin streaks to the left of the main plume that were also not identified using the default Ackerman (1997) or Roskovensky and Liou (2005) thresholds (Figures 7c and d). Second, whilst detection of airborne dust is improved, some areas of the ground surface are also mis-identified as dust. Some of these are the same areas (marked FS in Figure 7e) that caused difficulties in event 1 due to changes in surface reflectivity caused by fire scars, and can clearly be seen in other panels (Figure 7 a-d).

Figure 7f shows that the agreement between the MODIS L1B enhancement methods for dust (white) is restricted to the most dense part of

the plume. Ackerman (1997) performs best at highlighting the more subtle (perhaps less dense) dust plumes in the west of the scene. The Roskovensky and Liou (2005) threshold mis-identifies dust not only to the west but also to the south and so under-represents its spatial extent; with obvious implications for upwind source detection.

The central dust plume is shown clearly in the MODIS Deep Blue aerosol product for this event (Figure 7g). Not only is the sharp advancing dust front apparent, but these data also indicate higher dust concentrations in the southerly, upwind source area of the plume. At best, however, these data are only able to provide a broad regional indication of the plume's source because the thin, discrete plumes in the west are not detected. The MODIS Deep Blue product also suggests that the highest AOT values in the scene are associated with the plume at the extreme eastern edge of the scene (marked 'X'). This contrasts with the MODIS BTM analyses where dust appears to have a higher concentration at the furthest downwind edge of the plume (marked 'Y'). Similarly, OMI AI data clearly outline the main plume, and also suggest higher aerosol density in the east (Figure 7h). Comparison of the Deep Blue and OMI data highlight the performance of the cloud masks used in these products. The Deep Blue cloud mask only removes the cloud in the top left of the scene, which is clearly present in MODIS VIS, whilst the OMI cloud-mask obscures as much as 15% of the scene.

3.3 Event 3: 2nd February 2005

Event 3 was an extensive dust event during which 35 meteorological stations recorded a reduction in visibility to ≤ 1 km. Here we concentrate on two areas in the central LEB: the first is where two parallel dust plumes can clearly be seen blowing northwards out of Lake Eyre North and the second is in the lower right corner of the image and is difficult to see in the VIS (Figure 8a), but is clearly shown on the BTM (Figure 8b). The previous two events used a BTM threshold of < 0 with varying success. For this event (Figure 8c), a significantly lower threshold for dust detection (dust < -1.2) was required to identify dust plumes (Table 4). Using this value, most of the pixels highlighted

contain dust although the area marked 'GS' on Figure 8c is not dust, but is the ground surface. Crucially, the <-1.2 threshold in this instance does allow the observed dust plumes to be traced all the way back to the source areas.

<Figure 8>

The Roskovensky and Liou (2005) algorithm (Figure 8d) is effective at picking out the two main dust areas, but the origin of the twin plumes is situated north of the known dust source (the bed of Lake Eyre North). The parallel plumes are also visible when Miller's (2003) algorithm (Figure 8e) is applied with an adjusted dust threshold, but this dense dust is only enhanced by the model (coloured red) at the downwind end of the plume, and not in the source locations. Furthermore, in comparison to BTD (Figure 8b), the origin of the dust in Figure 8e (in white) would be placed approximately 70km north of the actual lake bed source. The 'best' dust threshold that could be determined for the Miller (2003) algorithm in this instance also seems to divorce the apparent upwind boundary of the plume from the source area marked 'X' in the right of the image (Figure 8e). However, the entire aerosol outbreak to the lower right is highlighted in red using this approach, and the plume clearly extends back to the assumed source. This demonstrates that there can be significant variability in the performance of this enhancement approach within a single scene. The composite image (Figure 8f) for the MODIS L1B enhancement techniques highlights the problem of surface reflectance evident when applying the Miller algorithm, as to the northwest of the twin plumes the fire scar (FS) is clearly shown in blue.

In Figure 8g, the distinctive parallel dust plumes are mainly excluded by the Deep Blue cloud mask, and only the downwind end of the plume is detected. Other parts of the image, where no cloud is present (cf. Figure 8a), are also excluded by the cloud mask. For example, the dry bed of Lake Eyre and small patches across the whole area of the scene are flagged as no-data. In Figure 8h, the OMI AI data show dust over the majority of the scene, suggesting a widespread dust haze. The AI maximum of 5.4 is very high for Australia but, whilst all the main areas of dust are identified in a manner

broadly consistent with the MODIS L1B algorithms, the spatial resolution is insufficient to illustrate the detail of the parallel plumes or the specific source locations. Indeed, although these data clearly have limited utility for determining the specific point-sources in this scene, the OMI data do suggest the presence of diffuse raised dust across the scene which would be expected given the number of meteorological stations recording the event.

3.4 *Event 4: 30th August 2005*

The weather systems that promote LEB dust storms (thunderstorms, pre- and post-frontal winds; Sprigg, 1982) mean that dust events are often associated with cloud cover. Event 4 was selected to explore further the extent to which dust and cloud can be distinguished. Dust is visible in the centre of the VIS scene between the bands of cloud (Figure 9a) and can also be identified using BTD (Figure 9b); although large areas of thicker cloud can be seen which exhibit a similar BTD as the dust, making initial interpretation of this scene using BTD alone problematic. A dust threshold of <-0.35 was applied to BTD in this instance (Figure 9c), and was effective in isolating the major dust plumes that exist between the clouds, but at the expense of the thinner plumes which are removed when this particular threshold is applied.

<Figure 9>

Most of the cloud is removed from the image by application of Roskovensky and Liou's (2005) algorithm (Figure 9d) and the inferred sources for the two main dust plumes clearly map on to those identified from BTD (Figure 9b). For this event, the Miller (2003) algorithm (Figure 9e) was scaled using $\text{dust} > 0.45$, and can be seen to be very effective as the principal dust plumes are easily discernable and source determination is possible. The enhancements discussed here are unable to ameliorate the blanketing effect of thick cloud when it obscures active dust sources or plumes; they can only enhance the dust that can be 'seen' between the cloud banks. Interestingly, none of the BTD-based methods pick up the thin plume which is best seen on the MODIS visible panel (marked 'X', Figure 9a). The most notable feature,

other than the general agreement of spatial extent and source location for the two plumes shown in Figure 9h, is the appearance of the blue areas where the Miller (2003) output has confused the thickest cloud with dust.

The cloud masking of the Deep Blue (Figure 9g) scene seems to work well for much of the cloud coverage in the image, but does also remove much of the northernmost dust evident in the other panels (Figures 9b-f). The shape and downwind extent of the central plume is indicated by raised AOT values, but the MODIS Deep Blue data also suggest an origin for the dust that is some distance removed downwind from the source when compared with the BTD-based approaches. The OMI data (Figure 9h) again show more of the dust plume than the MODIS Deep Blue product, and less of the image is affected by cloud masking.

4. Discussion

The main aim of this paper is to evaluate the use of MODIS for detecting dust sources. In some instances dust plumes may be discernible on the MODIS VIS (e.g. Figure 6); but this certainly is not always the case (e.g. Figure 4). From the results presented above, we can confirm that all the dust enhancement techniques used here make it easier to detect dust. However, with respect to source determination, the results suggest that, of the MODIS L1 processing techniques, the 'best' approach varies from event to event. For events 1 and 2 arguably the best source detection came from the simple brightness temperature difference calculation (BTD), often with no dust threshold applied. Of the more complex processing techniques, that of Roskovensky & Liou (2005) works well for event 1 (Figure 5) as it is very effective at removing cloud cover, whereas the Ackerman (1997) is better for events 2 and 3, where cloud cover is less of an issue. The cloud cover in event 4 (Figure 8) makes it much harder to determine sources from the BTD alone, but both Roskovensky & Liou (2005) and Ackerman (1997) work well. For these four events, the Miller (2003) algorithm is extremely useful for visualizing dust, but there are significant problems with precise source identification and determination of dust plume extent in all cases except event

4. For the majority of events and algorithms the published, or indicative, thresholds under-perform and the values vary from event to event. This makes it difficult to suggest appropriate regional scale thresholds. Whilst some of this variation is due to factors specific to the algorithms or individual events, other causes such as diurnal and seasonal variations in surface temperature/dust contrast (which affect BTD) will affect all the methods. The advantages and disadvantages of each of the approaches from an operational perspective are discussed in detail below.

4.1 *Brightness Temperature Difference (bispectral split window)*

Calculating BTD is straightforward, and keeping the full range of values (rather than applying a dust threshold) is often preferable for both dust plume and source detection. The procedure does not appear to be very sensitive to observed mineralogical variability either within or between plumes, and so all dust, regardless of source, is enhanced provided it can be differentiated thermally from the ground surface. The main disadvantages are that because no definition of dust/non-dust is applied the interpretation of the BTD data becomes subjective and data retrieval can suffer through lack of cloud cover elimination. With the exception of event 4, this is not a major problem in the case studies presented here, but it is likely to be important for anyone interpreting the data, to have a good understanding of how and why ground surface characteristics may vary.

4.2 *Ackerman (1997)*

Although Ackerman (1997) did not explicitly present a dust/non-dust threshold of zero, he observed negative differences in $BT_{11}-BT_{12}$ for dust storms and a universal threshold of $dust < 0$ could be implied. Darmenov & Sokolik (2005) demonstrated that this dust threshold was in fact variable when applied to dust over oceans and suggested that dust sourced from the Lake Eyre Basin and travelling southeast over the Tasman Sea had a value of < -0.4 K. All the dust plumes examined here are over land, and whilst the threshold of zero worked effectively for events 1 and 2, adjustments had to be

made for events 3 and 4. For event 3, in order to eliminate interference from the ground surface, the threshold had to be lowered to <-1.2 K; for event 4 the threshold was <-0.35 to eliminate cloud. Interestingly, whilst it was possible to find a dust/non-dust threshold for event 4 where all cloud could be removed this was not possible for event 1. Here (Figure 5c), a dust threshold lower than zero removed more dust in the scene so the threshold was left at 0. Another factor affecting the BTD threshold is likely to be the thickness of the dust plume. Where there is low AOT (as confirmed by comparison with MODIS Deep Blue) we have determined BTD differences of >0 for pixels populated by dust. One possibility is that where the dust plume thickness/density is low, BTD becomes increasingly affected by the ground surface temperature signal. Using this approach, therefore, may involve a compromise between dust detection and the elimination of cloud. If both dust and cloud are dense/opaque then it is straightforward to identify and implement a dust threshold. If the dust is thin and cloud cover is dense (as in event 1), then it can be hard to identify an appropriate dust/non-dust threshold. Where the cloud cover is sparse and the dust plume is dense/opaque (as in event 4) then their differentiation through threshold adjustment is straightforward. From this study it is also apparent that the thickness/density of the dust plume also affects the degree to which the dust-head can be pinpointed, and an inappropriate threshold value may foreshorten plumes.

4.3 *Roskovensky & Liou (2005)*

This approach was designed explicitly as a simple method for the differentiation of dust from cirrus cloud, and is very effective at doing so in both of the cloudy scenes examined here (events 1 and 4). For all events it was necessary to adjust the scaling factor 'a' and BTD offset value 'b'. Roskovensky and Liou (2005) calculated these to be 1.1 and 0 over ocean (around the Korean peninsula) and 3 and 0 over land (the Gobi desert). All of the events examined here occurred over land and the coefficients determined were variable (values of 'a' ranged from 0.25 to 1.2 and values of 'b' ranged from -0.5 to $+1$) and made a significant difference to both the number of

pixels classified as dust and the inferred location of the dust sources. Although the algorithm is slightly more computationally complex to calculate than simple BTM, it is easy to tune it for specific events, and certainly worth the extra effort. Overall this model worked best on dense dust; there was little observed confusion with ground surface reflectance, and the inferred upwind plume source locations compared well with those suggested by BTM alone.

4.4 *Miller (2003)*

The Miller (2003) algorithm is designed to provide improved differentiation of dust from water/ice clouds over bright desert surfaces and was found to be visually very effective for all events observed in this study. In particular, there is generally a clear distinction between dust and cloud. In a similar manner to optimizing the Ackerman (1997) data, the Miller (2003) dust threshold also had to be tuned for each event to be effective (Table 4) and in most cases it was necessary to decrease the lower threshold value to well below Miller's suggested +1.3 (as low as -0.55 in one instance). Unfortunately tuning this algorithm was not straightforward, and although it worked very well for some events, this was not always the case.

4.5 *The Aerosol Products (Deep Blue and OMI)*

The main utility of the aerosol products is in the detection of dust because for dust source identification the coarse spatial resolution of the products is a limitation. Average AI values over a long time series have been used to detect persistent, regional scale dust sources (e.g. Washington et al., 2003) but accurate and event-specific source identification requires the clear delineation of the upwind margin of the plumes. We have presented values of Deep Blue AOT and OMI AI which, whilst not comparable in terms of absolute values can be compared relatively. There are occasions where OMI agrees well with the much higher resolution MODIS, but often the upwind margin of a dust plume is difficult to detect using AI.

In some cases, it was difficult to determine not only the upwind dust sources but also the extent of the dust plumes because the in-built cloud masks of the products eliminated them. For example, in Figure 8g, the origin of the parallel twin plumes was not discernible because MYD04 returned no data from the bright dry lake surface, which was classified as cloud. This is a recognised limitation and is probably due to the colour and density of the dust or reflectance of the surface. The twin parallel plumes comprise white coloured dense dust sourced from Lake Eyre and are sufficiently bright to saturate the pixels causing misidentification as cloud (a known problem with Level 2 aerosol product http://modis-atmos.gsfc.nasa.gov/MOD04_L2/ga.htm). There are obvious implications for dust source identification – ephemeral lakes are often very bright surfaces and have been seen in this study to be routinely masked out as cloud even in cloud-free and dust-free scenes, yet are common dust sources not only in the LEB (Bullard et al. 2008), but also in the USA (e.g. Reynolds et al. 2007), southern Africa (Mahowald et al. 2003) and other dryland regions. The authors are also evaluating the Aura-OMI Aerosol Data Product; OMAERUV (V003) which provides aerosol extinction and optical depth via swath data (rather than global gridded) at the native 13 x 24 km pixel (see: <http://daac.gsfc.nasa.gov/Aura/OMI/omaeruv.shtml>). This may offer further potential for dust source identification, but requires further validation.

4.6 Impacts of Dust Mineralogy and Surface Reflectance on Data Retrieval

One issue that can be explored briefly here, and will be developed further as a future project, is the impact of dust mineralogy on both the Miller (2003) dust thresholds and MODIS Deep Blue retrievals. A range of different sedimentary environments emit dust within the Lake Eyre Basin (Bullard et al., 2008) and these have different mineralogical compositions which in turn control the infrared radiative properties of the dust (Claquin et al., 1999; Sokolik, 2002). An example can be seen in event 3 (Figure 8) where a dust plume was observed originating from the bed of Lake Eyre (which is illite-rich), in the west of the scene and a dust plume originating from dune sands with iron-rich coatings in the southeast of the scene. Using threshold values

942 where $0.6 < \text{dust} < 1.98$ it was possible to pick out the dust sourced from the
943 dunes in red, but not the near-source dust from the lake (although part of the
944 distal end of the plume is highlighted). When a threshold was selected to
945 highlight both plumes large areas of ground surface were also included.

946
947 The effect of dust mineralogy on detection is also apparent in the
948 MYD04 Deep Blue data presented here. In the Deep Blue algorithm, surface
949 reflectivity over desert regions is assumed to be low in blue wavelengths and,
950 dust aerosol slightly more reflective (Hsu et al., 2004). However, dust
951 reflectivity is variable depending on its chemistry and can decrease
952 significantly with increased iron concentration (Dubovic et al., 2002; Arimoto
953 et al., 2002). Consequently red, or iron-rich, dust will have relatively low blue
954 reflectivity and therefore potentially lower contrast relative to background
955 reflectance, whereas white dust (composed of carbonates, bleached quartz or
956 evaporite minerals) will have higher reflectance in the VIS and significantly
957 higher contrast with respect to underlying soils and vegetation. This is
958 illustrated in Figures 7g and 8g where the white, dry lake bed sources of dust
959 are more distinct than the iron-oxide rich dust from dune sources (noting
960 however that the OMI product suggests these red plumes are also less dense
961 than the white dust plumes which will affect AOT).

962
963 For the Lake Eyre Basin events there are some problems with the
964 differentiation of dust from the underlying desert surface. Specifically, large
965 areas of the LEB dunefields comprise red brown to orange, highly reflective
966 sands with a partial vegetation cover; although sand color is variable
967 especially between the redder Simpson dunefield and the less red Strzelecki
968 dunefield (Pell et al., 2000; Bullard and White, 2002). Where the vegetation
969 cover has not been disturbed by fire, the Miller (2003) algorithm works well.
970 However, where vegetation has been removed by fire and large areas of
971 bright sand are exposed the algorithm cannot distinguish dust from the
972 surface. Fire changes both surface reflectance characteristics, through the
973 removal of absorbing vegetation, and sediment reflectance characteristics,
974 through fire-induced reddening or changes in mineralogy (Jacobberger-
975 Jellison, 1994). The way in which the performance of the Miller (2003)

algorithm is affected by these changes has important implications for dust provenance because areas which have been de-vegetated through fire may be mis-identified as a dust source. A complicating issue is that fire scars can act as sources of dust under certain conditions (Bullard et al. 2008; McGowan & Clark, 2008). This highlights two issues. First, it is important to deploy event-specific dust/non-dust thresholds to limit the influence of surface reflectance as far as possible. Second, familiarity with the underlying surface characteristics is essential, and examination of more than one scene, including known dust-free images is desirable. The latter means that permanent or semi-permanent ground characteristics can be discerned. For example, the firescars in the LEB have distinctive shapes that can be identified on most of the VIS images taken post-2001 when widespread burning occurred. Cross-referencing with other MODIS-derived output can also be useful. For example, in event 1 the fire scar is only clearly discernible on the VIS and Miller (2003) panels which suggests that it was not a dust source. In contrast, in event 2 the fire scars are discernable in all panels (Figure 7 a-e) which may suggest that they acted as dust sources at this time.

4.7. *Tools for dust source detection*

Figure 10 presents a summary of the how the different data sources and analyses used here complement one another. At the global-regional scale, dust events can be detected using visibility criteria, as in this paper, or OMI AI. An alternative that we have not discussed in detail here is the MOD/MYD08 Global Gridded Atmospheric Product (1 x 1 deg) which could also be used (Bryant et al., 2007), although it has some limitations over bright desert surfaces (Chu et al., 2002). If none of these three indicates dust it does not necessarily mean that dust is not present as the relative timing of satellite overpass or visibility observation may result in no dust being recorded (see quality control indicators), but the events missed are likely to be minor. If any one of these indicates the presence of dust then there is the potential for determining dust sources at higher resolution. The choice of higher resolution technique depends on the precise research question to be answered. The MOD/MYD04 aerosol products give data processed to a common standard

that enables comparison from one region to another. The versatile MOD/MYD02 data can be processed simply using brightness temperature difference to enhance the dust signal. Where cloud is present, or if it is necessary to highlight the dust plume then one of the methods for employing a dust/non-dust threshold can be used, but it is recommended that event-specific thresholds are calculated where possible (as opposed to using global or regional thresholds).

5. Conclusions

This paper set out to evaluate the use of MODIS data for identifying dust sources. Through objective and subjective comparison of several different approaches to dust detection several conclusions can be drawn. Whilst these conclusions have implications for regional and global scale studies of dust, the clear outcomes are that even within a single drainage basin, dust events should be examined on an event by event basis, and that the 'best' algorithm for identifying dust sources varies considerably.

(1) For the region examined here (the Lake Eyre Basin), no single MODIS technique was found to be ideal for source determination.

(2) MODIS VIS full colour composite data can be useful but are often insufficient to discern dust plumes over reflective desert surfaces. In particular, for event detection MODIS VIS (particularly via quicklooks) should be used with caution, or in combination with other data sources, because not all dust plumes will be visible over the bright surfaces. An ideal combination for rapid detection of dust activity is OMI AI (or equivalents) and MODIS but the former can not be used to derive source location due to its low spatial resolution.

(3) BTd data are simple to calculate and very effective at highlighting dust that is not seen in VIS data. If the user is familiar with the underlying ground surface characteristics and has the VIS image available to discern cloud cover, this can be the most simple and possibly most accurate method of source determination. From the four example events presented here, BTd is the approach that is least sensitive to dust mineralogy.

- (4) For scenes where cloud is present, there is potential for confusion between cloud and dust using just BTM and the algorithms designed to differentiate these have been demonstrated as effective for screening out cirrus cloud (for which they were designed) although thick cloud can remain.
- (5) Whilst there are published dust/non-dust thresholds for each method compared here, for each method thresholds may need adjusting on a regional and/or event scale. Although there are clear guidelines for positioning thresholds, a considerably amount of informed (but subjective) judgement can be required. It is important therefore for users of these techniques to consider the effects of not adjusting thresholds for each event. This paper suggests that for an event-based study it is essential to derive event-specific thresholds. However it seems likely that for global or longer-term studies of this nature to be effective, it may be pragmatic to use regional thresholds.
- (6) In the Lake Eyre Basin, it is not possible to use a single dust/non-dust threshold for all events for any of the algorithms tested here. One possible reason is that it is a basin with multiple potential dust sources with different mineralogies; where different sources (e.g. iron-rich dunes, illite-rich lake beds) can emit dust simultaneously it is necessary to use event-specific, or even plume-specific thresholds. In regions with a single definable source (for example a large playa such as the Magkadigadki, Botswana) it may be possible to discern a single dust/non-dust threshold, however any use of a regional (or global) threshold is likely to result in errors or inconsistencies.
- (7) Some sedimentary environments are more intense dust sources than others. In particular, previous studies have noted ephemeral lakes and areas devegetated by fire as prominent dust sources. Significantly, these are two of the ground surface types that have proved most problematic for establishing dust/non-dust thresholds due to confusion caused by their bright surfaces which cause false positive dust signals or may be falsely identified as cloud.
- (8) Whilst the findings of this research may be challenging in some respects, what is clear is that there is considerable potential for using

1078 MODIS data to obtain information about dust mineralogy by interpreting
1079 the shifts to the thresholds (or coefficients) that need to be made.
1080 Possible mineral aerosol information that could be gained can not only
1081 assist in identifying the dust source, but also its radiative properties.

1082

1083 Using data from sensors such as MODIS will inevitably mean that some
1084 dust activity is missed due to the relative timing of overpass and dust
1085 emissions or cloud cover. Other sources of data such as MSG can minimize
1086 the problem of overpass timings but cloud cover is still a problem and
1087 coverage is not yet global. This paper has focused on daytime dust
1088 emissions only, but analysis of night time dust emissions and source
1089 identification are the subject of future research. It is worth noting that the
1090 products and data used here are subject to continual development and
1091 improvement and consequently some of the issues raised here may be
1092 resolved or change.

1093

1094

1095

Acknowledgements

This research was part of a project funded by The Leverhulme Trust, (JEB), for which MCB was the post-doctoral research fellow. The authors would like to thank Steve Miller, Clare Salustro and Christina Hsu for their technical assistance and useful discussions. The highly valuable comments of three anonymous reviewers also helped to improve this paper.

References

- Ackerman, S.A. (1997). Remote sensing aerosols using satellite infrared observations, *Journal of Geophysical Research*, 102, 17069–17080
- Ackerman, S.A., Strabala, K.I., Menzel, W.P., Frey, R., Moeller, C., Gumley, L., Baum, B.A., Seeman, S.W., & Zhang, H. (2002). Discriminating clear-sky from cloud with MODIS algorithm theoretical basis document (MOD35). In ATBD Ref.ATBD-MOD-06, version 4.0, 115 pp. Greenbelt, Maryland: NASA Goddard Space Flight Center
- Ahn, C., Torres, O., & Bhartia, P.K. (2008). Comparison of ozone monitoring instrument UV aerosol products with Aqua/Moderate Resolution Imaging Spectroradiometer and Multiangle Imaging Spectroradiometer observations in 2006. *Journal of Geophysical Research*, 113, D16S27
- Agarwal, A., El-Askary, J.M., El-Ghazawi, T., Kafatos, M. and Le-Moine, J. (2007). Hierarchical PCA techniques for fusing spatial and spectral observations with application to MISR and monitoring dust storms. *IEEE Geoscience and Remote Sensing Letters*, 4, 4, 678-682.
- Alpert, P., Kishcha, P., Shtivelman, A., Krichak, S.O. & Joseph, J.H. (2004). Vertical distribution of Saharan dust based on 2.5-year model predictions. *Atmospheric Research*, 70, 109-130.
- Arimoto, R., Balsam, W., Schloesslin, C. 2002. Visible spectroscopy of aerosol particles collected on filters: iron-oxide minerals. *Atmospheric Environment*, 36, 89-96.
- Badarinath, K.V.S., Kumar Kharol, S., Kaskaoutis, D.G. & Kambezidis, H.D. (2007). Dust storm over Indian region and its impact on the ground reaching solar radiation – a case study using multi-satellite data and ground measurements. *Science of the Total Environment*, 384, 316-332.
- Brindley, H. E., & Ignatov, A. (2006). Retrieval of mineral aerosol optical depth and size information from Meteosat Second Generation SEVIRI solar reflectance bands. *Remote Sensing of Environment*, 102, 344-363, doi:10.1016/j.rse.2006.02.024
- Bryant, R. G., Bigg, G. R., Mahowald, N. M., Eckardt, F. D., & Ross, S. G., (2007). Dust emission response to climate in southern Africa. *Journal of*

1135 *Geophysical Research*, 112, D09207, doi:10.1029/2005JD007025.

1136 Bullard, J. E., Baddock, M. C., McTainsh, G. H., & Leys, J. F., (2008). Sub-
 1137 basin scale dust source geomorphology detected using MODIS.
 1138 *Geophysical Research Letters* 35, 1, art.no.L15404,
 1139 doi:2008GL033928.

1140 Bullard, J. E., & White, K. H., (2002). Quantifying iron oxide coatings on dune
 1141 sands using spectrometric measurements: an example from the
 1142 Simpson-Strzelecki Desert, Australia. *Journal of Geophysical*
 1143 *Research*, 107, doi:10.1029/2001JB000454.

1144 Caquineau, S., Gaudichet, A., Gomes, L., & Legrand, M. (2002). Mineralogy
 1145 of Saharan dust transported over northwestern tropical Atlantic Ocean
 1146 in relation to source regions. *Journal of Geophysical Research*,
 1147 107(D15), 4251, doi:10.1029/2000JD000247

1148 Chu, D.A., Kaufman, Y.J., Ichoku, C., Remer, L.A., Tanré, D. & Holben, B.N.
 1149 (2002). Validation of MODIS aerosol optical depth retrieval over land.
 1150 *Geophysical Research Letters*, 29, 12, 8007,
 1151 doi:10.1029/2001GL013205.

1152 Claquin, T., Schulz, M., & Balkanski, Y.J. (1999). Modeling the mineralogy of
 1153 atmospheric dust sources. *Journal of Geophysical Research*, 104,
 1154 22243-22256

1155 Darmenov, A. & I.N. Sokolik (2005). Identifying the regional thermal-IR
 1156 radiative signature of mineral dust with MODIS. *Geophysical Research*
 1157 *Letters*, 32, L16803, doi:10.1029/2005GL023092

1158 de Baar, H.J.W. and 33 others (2005). Synthesis of iron fertilization
 1159 experiments: from the iron age in the age of enlightenment. *Journal of*
 1160 *Geophysical Research-Oceans*, 110, C9, C09S16.

1161 Desboeufs, K.V., Sofikitis, A., Losno, R., Colin, J.L., & Ausset, P. (2005).
 1162 Dissolution and solubility of trace metals from natural and
 1163 anthropogenic particulate matter. *Chemosphere*, 58, 2, 195-203.

1164 Dubovik, O., Holben, B., Eck, T.F., Snirnov, A., Kaufman, Y.J., King, M.D.,
 1165 Tanré, D. & Slutsker, I. 2002. Variability of absorption and optical
 1166 properties of key aerosol types observed in worldwide locations.
 1167 *Journal of the Atmospheric Sciences*, 59, 590-608.

1168 Dubovik, O., Lapyonok, T., Kaufman, Y.J., Chin, M., Ginoux, P., Kahn, R.A. &

1169 Sinyuk, A. (2008). Retrieving global aerosol sources from satellites
 1170 using inverse modeling. *Atmospheric Chemistry and Physics*, 8, 2,
 1171 209-250.

1172 Eckardt, F.D., & Kuring, N. (2005). SeaWIFS identified dust sources in the
 1173 Namib Desert. *International Journal of Remote Sensing*, 26, 19, 4159-
 1174 4167

1175 Ekstrom, M., McTainsh, G.H., & Chappell, A. (2004). Australian dust storms:
 1176 temporal trends and relationships with synoptic pressure distributions
 1177 (1960-1999). *International Journal of Climatology*, 24, 12, 1581-1599.

1178 Erickson, D.J., Hernandez, J.L., Ginoux, P., Gregg, W.W., McClain, C., &
 1179 Christian, J. (2003). Atmospheric iron delivery and surface ocean
 1180 biological activity in the Southern Ocean and Patagonian region.
 1181 *Geophysical Research Letters*, 30, 12, doi:10.1029/2003GL017241

1182 Evan, A. T., A. K. Heidinger, and M. J. Pavolonis (2006) Development of a
 1183 new over water Advanced Very High Resolution Radiometer dust
 1184 detection algorithm. *International Journal of Remote Sensing*, 27(18),
 1185 pp. 3903-3924. doi: 10.1080/01431160600646359

1186 Gassó, S. & A.F. Stein (2007). Does dust from Patagonia reach the sub-
 1187 Antarctic Atlantic Ocean. *Geophysical Research Letters*, 34, L01801,
 1188 doi:10.1029/2006GL027693

1189 Hansell, R.S., Ou, S.C., Liou, K.N., Roskovensky, J.K., Tsay, S.C., Hsu, C., &
 1190 Ji, Q. (2007). Simultaneous detection/separation of mineral dust and
 1191 cirrus clouds using MODIS thermal infrared window data. *Geophysical*
 1192 *Research Letters*, 34, L11808, doi:10.1029/2007GL029388

1193 Haywood, J., & Boucher, O. (2000). Estimates of the direct and indirect
 1194 radiative forcing due to tropospheric aerosols: a review. *Reviews of*
 1195 *Geophysics*, 38, 4, 513-543.

1196 Hsu, N.C., Herman, J. R. & Weaver, C. (2000). Determination of radiative
 1197 forcing of Saharan dust using combined TOMS and ERBE data.
 1198 *Journal of Geophysical Research*, 105, 20649–20661

1199 Hsu, N. C., Tsay, S. C. King, M. D. & Herman J. R. (2004). Aerosol properties
 1200 over bright-reflecting source regions, *IEEE Transactions on Geoscience*
 1201 *and Remote Sensing*, 42, 557-569

1202 Hsu, N.C., Tsay, S-C, King, M.D., & Herman, J.R. (2006). Deep Blue retrievals

1203 of Asian aerosol properties during ACE-Asia. *IEEE Transactions on*
1204 *Geoscience and Remote Sensing*. 44, 11, 3180-3195.
1205 doi:10.1109/TGRS.2006.879540

1206 Jacobberger-Jellison, P.A. (1994). Fire-induced reddening of dune sands.
1207 *Journal of Arid Environments*, 26, 305-313.

1208 Jickells, T.D. and 18 others, (2005). Global iron connections between desert
1209 dust, ocean biogeochemistry and climate. *Science*, 308, 67-71

1210 Jones, T.A. & Christopher, S.A. (2007). MODIS-derived fine mode fraction
1211 characteristics of marine dust and anthropogenic aerosols over the
1212 ocean constrained by GOCART, MOPITT and TOMS. *Journal of*
1213 *Geophysical Research – Atmospheres*, 112, D22, D22204.

1214 Jones, T.A. & Christopher, S.A. (2008). Multispectral analysis of aerosols over
1215 oceans using principal components. *IEEE Transactions on*
1216 *Geoscience and Remote Sensing*, 46, 9, 2659-2665.

1217 Kaskaoutis, D.G., Kambezidis, H.D., Nastos, P.T. & Kosmopoulos, P.G.
1218 (2008). Study on an intense dust storm over Greece. *Atmospheric*
1219 *Environment*, 42, 9, 6884-6896.

1220 Kaufman, Y.J., Tanré, D., Remer, L., Vermote, E.F., Chu, A., & Holben, B.N.
1221 (1997). Operational remote sensing of tropospheric aerosol over the
1222 land from EOS-MODIS. *Journal of Geophysical Research*, 102, 17051-
1223 17068

1224 Kaufman, Y.J., Karnieli, A., & Tanré, D. (2000). Detection of dust over deserts
1225 using satellite data in the solar wavelengths. *IEEE Transactions on*
1226 *Geoscience and Remote Sensing*, 38, 1, 525-531

1227 Kaufman, Y.J., Boucher, O., Tanré, D., Chin, M., Remer, L.A. & Takemura, T.
1228 (2005). Aerosol anthropogenic component estimated from satellite
1229 data. *Geophysical Research Letters*, 32, 17, L17804.

1230 Kubilay, N, Oguz, T., Kocak, M., & Torres, O. (2005). Ground-based
1231 assessment of Total Ozone Mapping Spectrometer (TOMS) data for
1232 dust transport over the northeastern Mediterranean. *Global*
1233 *Biogeochemical Cycles*, 19, GB1022.1

1234 Lee, J.A., Gill, T.E., Mulligan, K.R., Acosta, M.D., & Perez, A.E. (2008). Land
1235 use/land cover and point sources of the 15 December 2003 dust storm

1236 in southwestern North America. *Geomorphology*, 105, 1-2. 18-27,
 1237 doi:10.1016/j.geomorph.2007.12.016.

1238 Legrand, M., Nertrand, J.J., Desbois, M., Menenger, L., & Fouquart, Y. (1989).
 1239 The potential of infrared satellite data for the retrieval of Saharan dust
 1240 optical depth over Africa. *Journal of Climatology and Applied*
 1241 *Meteorology*, 28, 309-318.

1242 Levy, R.C., Remer, L.A., Kaufman, Y.J., Tanré, D., Mattoo, S., Vermote, E. &
 1243 O. Dubovik, (2006). Revised Algorithm Theoretical Basis Document:
 1244 MODIS Aerosol Products MOD/MYD04. Available at [http://modis-](http://modis-atmos.gsfc.nasa.gov/reference_atbd.html)
 1245 [atmos.gsfc.nasa.gov/reference_atbd.html](http://modis-atmos.gsfc.nasa.gov/reference_atbd.html)

1246 Levy, R.C., L.A. Remer, S. Mattoo, E.F. Vermote, & Y.J. Kaufman (2007).
 1247 Second-generation operational algorithm: Retrieval of aerosol
 1248 properties over land from inversion of Moderate Resolution Imaging
 1249 Spectroradiometer spectral reflectance, *Journal of Geophysical*
 1250 *Research*, 112, D13211, doi:10.1029/2006JD007811

1251 Li, J., Okin, G.S., Alvarez, L., & Epstein, H. (2007). Quantitative effects of
 1252 vegetation cover on wind erosion and soil nutrient loss in a desert
 1253 grassland of southern New Mexico, USA. *Biogeochemistry*, 85, 3, 317-
 1254 332

1255 Mackie, D.S., Boyd, P.W., Hunter, K.A., & McTainsh, G.H. (2005). Simulating
 1256 the cloud processing of iron in Australian dust: pH and dust
 1257 concentration. *Geophysical Research Letters*, 32, 6, L06809

1258 Mackie, D.S., Boyd, P.W., McTainsh, G.H., Tindale, N.W., Westberry, T.K., &
 1259 Hunter, K.A. (2008). Biogeochemistry of iron in Australian dust: from
 1260 eolian uplift to marine uptake. *Geochemistry, Geophysics,*
 1261 *Geosystems*. 9, doi:10.1029/2007GC001813

1262 MacKinnon, D. J., Chavez, P. S., Jr., Fraser, R. S., Niemeyer, T. C., and
 1263 Gillette, D. A., (1996), Calibration of GOES-VISSR, visible-band
 1264 satellite data and its application to the analysis of a dust storm at
 1265 Owens Lake, California, *Geomorphology* 17, pp. 229 - 250

1266 McGowan, H.A., & Clark, A. (2008). A vertical profile of PM10 dust
 1267 concentrations measured during a regional dust event identified by
 1268 MODIS Terra, western Queensland, Australia. *Journal of Geophysical*
 1269 *Research*, 113, F02S03, doi:10.1029/2007JF000765.

- 1270 McTainsh, G. H. & Strong, C.S. (2007). The role of aeolian dust in
1271 ecosystems. *Geomorphology*, 89, 1-2, 39-54
- 1272 Mahowald, N.; Baker, A.; Bergametti, G.; Brooks, N.; Duce, R.; Jickells, T.;
1273 Kubilay, N.; Prospero, J.; Tegen, I.(2005) Atmospheric global dust
1274 cycle and iron inputs to the ocean *Global Biogeochemical Cycles*, Vol.
1275 19, No. 4, GB4025,doi: 10.1029/2004GB002402, 2005
- 1276 Mahowald, N., Ballentine, J.A., Feddema J., & Ramankutty N. (2007). Global
1277 trends in visibility: implications for dust sources, *Atmospheric Chemistry*
1278 *and Physics*, 7, 3309-3339.
- 1279 Mahowald, N.M., Bryant, R.G., del Corral, J. & Steinberger, L. (2003).
1280 Ephemeral lakes and dust sources. *Geophysical Research Letters*, 30,
1281 2, art.no.1074.
- 1282 Middleton, N.J., Goudie, A.S. & Wells, G.L. (1986). The frequency and source
1283 areas of dust storms. In Nickling, W.G. (ed) *Aeolian Geomorphology*.
1284 Allen & Unwin, Boston, 237-259.
- 1285 Miller, S.D. (2003). A consolidated technique for enhancing desert dust
1286 storms with MODIS. *Geophysical Research Letters*, 30, 20,
1287 art.no.2071.
- 1288 Moorthy, K.K., Babu, S.S., Satheesh, S.K., Srinivasan, J., & Dutt, C.B.S.
1289 (2007). Dust absorption over the “Great Indian Desert” inferred using
1290 ground-based and satellite remote sensing. *Journal of Geophysical*
1291 *Research – Atmospheres*, 112, D9, D09206.
- 1292 Muhs, D.R., Budahn, J.R., Prospero, J.M., & Carey, S.N. (2007).
1293 Geochemical evidence for African dust inputs to soils of western
1294 Atlantic islands: Barbados, the Bahamas and Florida. *Journal of*
1295 *Geophysical Research – Earth Surface*. 112,
1296 doi:10.1029/2005JF000445.
- 1297 Pell, S.D., Chivas, A.R., & Williams, I.S. (2000). The Simpson, Strzelecki and
1298 Tirari Deserts: development and sand provenance. *Sedimentary*
1299 *Geology*, 130, 1-2, 107-130.
- 1300 Piketh, S.J., Tyson, P.D., & Steffen, W. (2000). Aeolian transport from
1301 southern Africa and iron fertilization of marine biota in the South India
1302 Ocean. *South African Journal of Science*, 96, 244-246.

1303 Prospero, J.M., Ginoux, P., Torres, O., Nicholson, S.E. & Gill, T.E. (2002).
 1304 Environmental characterization of global sources of atmospheric soil
 1305 dust identified with the Nimbus 7 Total Ozone Mapping Spectrometer
 1306 (TOMS) absorbing aerosol product. *Reviews of Geophysics*, 40, 1,
 1307 art.no.1002.

1308 Remer, L.A., Tanré, D., Kaufman, Y.J., Levy, R. & Mattoo, S. (2006).
 1309 Algorithm for Remote Sensing of Tropospheric Aerosol from MODIS for
 1310 Collection 005 Products: 04_L2, 08_D3, 08_E3, 08_M3. Available from:
 1311 [http://modis-](http://modis-atmos.gsfc.nasa.gov/_docs/MOD04:MYD04_ATBD_C005_rev1.pdf)
 1312 [atmos.gsfc.nasa.gov/_docs/MOD04:MYD04_ATBD_C005_rev1.pdf](http://modis-atmos.gsfc.nasa.gov/_docs/MOD04:MYD04_ATBD_C005_rev1.pdf)

1313 Remer, L.A., Kaufman, Y.J., Tanré, D., Mattoo, S., Chu, D. A., Martins, J. V.,
 1314 Li, R.-R., Ichoku,, Levy, R. C., Kleidman, R. G., Eck, T. F., Vermote, E.,
 1315 & Holben, B. N. (2005). The MODIS aerosol algorithm, products and
 1316 validation, *Journal of Atmospheric Sciences*, 62, 4, 947–973. Reynolds,
 1317 R., Neff, J., Reheis, M., & Lamothe, P. (2006). Atmospheric dust in
 1318 modern soil on Aeolian sandstone, Colorado Plateau (USA): variation
 1319 with landscape position and contribution to potential plant nutrients.
 1320 *Geoderma*, 130, 1-2, 108-123.

- 1321 Reynolds, R., Neff, J., Reheis, M. & Lamothe, P. (2006). Atmospheric dust in
1322 modern soil on aeolian sandstone, Colodaro Plateau (USA): variation
1323 with landscape position and contribution to potential plant nutrients.
1324 *Geoderma*, 130, 1-2, 108-123.
- 1325 Reynolds, R., Yount, J.C., Reheis, M., Goldstein, H., Chavez, P., Fulton, R.,
1326 Whitney, J., Fuller, C. & Forester, R.M. 2007. Dust emission from wet
1327 and dry playas in the Mojave desert, USA. *Earth Surface Processes*
1328 *and Landforms*, 32, 12, 1811-1827.
- 1329 Ridgwell, A.J. (2002). Dust in the Earth system: the biogeochemical linking of
1330 land, air and sea. *Philosophical Transactions of the Royal Society of*
1331 *London*, A. 360, 1801, 2905-2924.
- 1332 Roskovensky, J.K. & K.N. Liou (2003). Detection of thin cirrus from 1.38
1333 mm/0.65 mm reflectance ratio combined with 8.6– 11 mm brightness
1334 temperature difference. *Geophysical Research Letters*, 30, 19,
1335 doi:10.1029/2003GL018135
- 1336 Roskovensky, J.K. & K.N. Liou (2005). Differentiating airborne dust from cirrus
1337 clouds using MODIS data. *Geophysical Research Letters*, 32, L12809,
1338 doi:10.1029/2005GL022798.
- 1339 Satheesh, S.K. & Moorthy, K.K. (2005). Radiative effects of natural aerosols,
1340 *Atmospheric Environment*, 39, 2089-2110.
- 1341 Shao, Y.P., & Wang, J.J. (2003). A climatology of northeast Asian dust events.
1342 *Meteorologische Zeitschrift*, 12, 4, 187-196.
- 1343 Shenk, W.E. & Curran, R.J. (1974). The detection of dust storms over land
1344 and water with satellite visible and infrared measurements, *Monthly*
1345 *Weather Review*, 102, 830-837.
- 1346 Schepanski, K., Tegen, I., Laurent, B., Heinold, B. & Macke, A. (2007). A new
1347 Saharan dust source activation frequency map derived from MSG-
1348 SEVIRI IR-channels. *Geophysical Research Letters*, 34, L18803,
1349 doi:10.1029/2007GL030168.
- 1350 Soderberg, K. & Compton, J.S. (2007). Dust as a nutrient source for fynbos
1351 ecosystems, South Africa. *Ecosystems*, 10, 550-561.
- 1352 Sokolik, I. (2002). The spectral radiative signature of wind-blown mineral dust:
1353 implications for remote sensing in the thermal IR region. *Geophysical*
1354 *Research Letters*, 29, 2154, doi:10.1029/2002GL015910.

- Sprigg, R.C. (1982). Alternating wind cycles of the Quaternary era, and their influence on Aeolian sedimentation in and around the dune deserts of southeastern Australia. In Wasson, R.J. (ed). *Quaternary Dust Mantles of China, New Zealand and Australia*. Proceedings International Union for Quaternary Research (INQUA) Loess Commission Workshop, Canberra, 211-240.
- Swap, R., Garstang, M., Greco, S., Talbot, R. & Kallberg, P. (1992). Saharan dust in the Amazon Basin. *Tellus Series B- Chemical and Physical Meteorology*, 44, 2, 133-149.
- Tanré, D. & Legrand, M., (1991). On the satellite retrieval of Saharan dust optical thickness over land: Two different approaches, *Journal of Geophysical Research*, 96, 5221–5227.
- Torres, O., Bhartia, P.K., Herman, J. R., Ahmad, Z. & Gleason J. (1998). Derivation of aerosol properties from a satellite measurement of backscattered ultraviolet radiation: theoretical basis, *Journal of Geophysical Research*, 103, 17,099-17,110.
- Torres, O., Tanskanen, A., Veihelmann, B., Ahn, C., Braak, R., Bhartia, P. K., Veefkind, P. & Levelt P. (2007). Aerosols and surface UV products from Ozone Monitoring Instrument observations: an overview, *Journal of Geophysical Research*, 112, D24S47, doi:10.1029/2007JD008809.
- Waeles, M., Baker, A.R., Jickells, T., & Hoogewerff, J. (2007). Global dust teleconnections: aerosol iron solubility and stable isotope composition. *Environmental Chemistry*, 4, 4, 233-237 doi:10.1071/EN07013.
- Wald, A.E., Kaufman, Y.J., Tanré, D., Gao, B.C. (1998). Daytime and nighttime detection of mineral dust over desert using infrared spectral contrast. *Journal of Geophysical Research-Atmospheres*, 103, D24, 32307-32313.
- Wang, X.B., Enema, O., Hoogmed, W.B., Perdok, U.D., & Cai, D.X. (2006). Dust storm erosion and its impact on soil carbon and nitrogen losses in northern China. *Catena*, 66, 3, 221-227.
- Washington, R., Todd, M., Middleton, N.J., & Goudie, A.S. (2003). Dust-storm source areas determined by the Total Ozone Monitoring Spectrometer and surface observations. *Annals of the Association of American Geographers*, 93, 2, 297-313.

1389 Wolff, E.W. and 28 others. (2006). Southern Ocean sea-ice extent,
 1390 productivity and iron flux over the past eight glacial cycles. *Nature*, 440,
 1391 7083, 491-496.

1392 Yoshioka, M., Mahowald, N.M., Conley, A.J., Collins, W.D., Fillmore, D.W.,
 1393 Zender, C.S., & Coleman, D.B. (2007). Impact of desert dust radiative
 1394 forcing on Sahel precipitation: relative importance of dust compared to
 1395 sea surface temperature variations, vegetation changes and
 1396 greenhouse gas warming. *Journal of Climate*, 20, 8, 1445-1467,
 1397 doi:10.1175/JCL14056.

1398 Zha, Y., & Li, L. (2007). Influence of the 17 April 2006 dust storm on Moderate
 1399 Resolution Imaging Spectroradiometer data for land cover
 1400 identification. *Journal of Geophysical Research*, 112, D14205,
 1401 doi:10.1029/2006JD007869.

1402 Zhang, B., Tsunekawa, A. & Tsubo, M. (2008). Contributions of sandy lands
 1403 and stony deserts to long-distance dust emission in China and
 1404 Mongolia during 2000-2006. *Global and Planetary Change*, 60, 3-4,
 1405 487-504.

1406 Zhu, A., Ramanathan, V., Li, F. & Kim, D. (2007). Dust plumes over the pacific,
 1407 Indian and Atlantic oceans: climatology and radiative impact. *Journal of*
 1408 *Geophysical Research – Atmospheres*, 112, D16, D16208.

1409 Zubko, V., Kaufman, Y.J., Burg, R.I. & Martins, J.V. (2007). Principal
 1410 components analysis of remote sensing of aerosols over oceans. *IEEE*
 1411 *Transactions on Geoscience and Remote Sensing*, 45, 3, 730-745.

1412

Figure Captions

Figure 1. The Lake Eyre Basin of Australia, and the coverage extents of the four dust event case studies by event number.

Figure 2. Histograms illustrating (a) the basic method for determining dust/non-dust threshold; (b) dust/non-dust thresholds derived for BTD (Ackerman, 1997) (dust < threshold; and (c) dust/non-dust threshold derived for Miller (2003) algorithm outputs (dust > threshold) for events 1-4.

Figure 3. Histograms showing the derivation of scaling factor 'a', based on reflectance ratio, for events 1-4 (graphs a-d), and BTD offset 'b' for events 1-4 (graphs e-h) necessary for applying Roskovensky & Liou's (2005) algorithm. For event 4 the scaling factor = 0.25 (not shown). Note: vertical axes vary.

Figure 4. Key to interpreting the colour composite combining dust/non-dust detection using Ackerman (1997), Miller (2003) and Roskovensky & Liou (2005) and shown in Figures 5-8, panel (f).

Figure 5. Event 1: 7th October 2005. Across all panels, squares highlight the prominent active dust source areas, as identified from the BTD split window product. The dust/non-dust thresholds used are detailed in Table 4. Panels (a) to (e) are derived from MODIS (Aqua) data. (a) MODIS visible (wind direction marked with arrow); (b) BTD ($BT_{11}-BT_{12}$); (c) threshold applied to BTD (after Ackerman, 1997); (d) Roskovensky & Liou (2005) dust enhancement algorithm; (e) Miller's (2003) dust enhancement algorithm (dust is red); (f) Composite image of Ackerman, Miller and Roskovensky and Liou output for key see Fig.4. Panels (g) and (h) are aerosol products for the same event where (g) MYD04 Deep Blue AOT $0.1^{\circ} \times 0.1^{\circ}$ spatial resolution; (h) OMI AAI $0.25^{\circ} \times 0.25^{\circ}$ spatial resolution.

Figure 6. Event 2: 24th September 2006. Dust/non-dust thresholds used are those recommended in the published literature and the same as those used for Event 1.

Figure 7. Event 2: 24th September 2006. Dust/non-dust thresholds used are specific to Event 2 and adjusted using the histogram approach (Figure 3).

Figure 8. Event 3: 2nd February 2005. For panel explanations see Figure 5. Dust/non-dust thresholds used are specific to Event 3 and adjusted using the histogram approach (Figure 3)

Figure 9. Event 4: 30th August 2005. For panel explanations see Figure 5. Dust/non-dust thresholds used are specific to Event 4 and adjusted using the histogram approach (Figure 3)

Figure 10. Summary of the relationships between the methods and approaches discussed in this paper.

Table 1. Spatial and temporal characteristics of remote sensing data used in this study.

Data Type	Spatial Resolution at Nadir (km)	Scenes used per day	Archive Length	Typical Overpass: (am/pm)
MOD02 [Terra]	0.25 x 0.25 (VIS ¹) 0.5 x 0.5 (VIS + NIR ²) 1 x 1 (TIR ³ + all bands)	1	1999 - date	am (10:30) ect [†]
MYD02 [Aqua]	0.25 x 0.25 (VIS) 0.5 x 0.5 (VIS + NIR) 1 x 1 (TIR + all bands)	1	2002 - date	pm (13:30) ect
MYD04 [Deep Blue]	10 x 10	1	2002 - date	pm (13:30) ect
OMI	13 x 24	1	2004 - date	pm (13:38) ect

¹Visible; ²Near-infrared; ³Thermal-infrared; [†]Equatorial crossing time

Table 2. Summary of dust event case studies used in this study to evaluate
MODIS dust detection algorithms

Event	Date DD/MM/YY (Julian)	Satellite overpass time (UTC)	BoM [†] stations recording visibility ≤1km	Selection Criteria	Synoptic Conditions
1	07/10/05 (2005:280)	AQUA (01:35) AURA (01:46)	1	Dust and 30% cloud cover	Pre-frontal northerly
2	24/09/06 (2006:267)	AQUA (04:20) AURA (04:31)	2	Dust - Single frontal plume	Post-frontal southerly
3	02/02/05 (2005:033)	AQUA (04:10) AURA (04:21)	35	Dust - Multiple plumes and source types	Post-frontal southerly
4	30/08/05 (2005:242)	AQUA (04:50) AURA (05:01)	1	Dust and 50% cloud cover	Pre-frontal northerly

[†]Bureau of Meteorology

1473 Table 3. Summary of dust detection algorithms applied to MODIS L1B
1474 data in this study
1475

Algorithm	Display parameters	Source
$D = (BT_{31} - BT_{32})$ where: $BT_{31} = BT_{10.780-11.280 \text{ } \mu\text{m}}$ $BT_{32} = BT_{11.770-12.27 \text{ } \mu\text{m}}$	Dust < 0.0	Ackerman (1997) Sokolik (2002)
$D = \exp \left\{ - \left[\frac{R_4}{R_{16}} * a + ((BT_{31} - BT_{32}) - b) \right] \right\}$ where: a=Scaling Factor (0.8) $R_{16} = R_{0.862-0.877 \text{ } \mu\text{m}}$ b=btd offset (2.0) $BT_{31} = BT_{10.780-11.280 \text{ } \mu\text{m}}$ $R_4 = R_{0.545-0.565 \text{ } \mu\text{m}}$ $BT_{32} = BT_{11.770-12.27 \text{ } \mu\text{m}}$	Dust > 1.0	Roskovensky & Liou (2003) Roskovensky & Liou (2005) Hansell et al., (2007)
$D = \left[(BT_{32} - BT_{31})^a + (2R_1 - R_3 - R_4 - BT_{31})^b - (R_{26})^c + (1 - BT_{31})^d \right]$ where: $R_1 = R_{0.620-0.670 \text{ } \mu\text{m}}$ $R_3 = R_{0.459-0.479 \text{ } \mu\text{m}}$ $R_4 = R_{0.545-0.565 \text{ } \mu\text{m}}$ $R_{26} = R_{1.360 - 1.390 \text{ } \mu\text{m}}$ $BT_{31} = BT_{10.780-11.280 \text{ } \mu\text{m}}$ $BT_{32} = BT_{11.770-12.27 \text{ } \mu\text{m}}$	Normalization: $a = (-2 \text{ to } +2)$ $b = (-5 \text{ to } +0.25)$ $c = (\text{if } M_{26} > 0.05, c=0, \text{ otherwise } c=1)$ $d = \text{occurrences of } M_{31} \text{ are normalized in this equation as } (M_{31max} - 21K \text{ if } M_{31max} < 301K) \text{ or } ((M_{31max} - 273)/4 + 273) \text{ otherwise.}$	1.3 < Dust < 2.7 Miller (2003)

1476
1477
1478

Table 4. Comparison of recommended dust/non dust thresholds and thresholds identified for events described in this study.

Technique	Suggested threshold values	Dust/non-dust thresholds			
		Event 1 7 Oct 2005	Event 2 24 Sept 2006	Event 3 2 Feb 2005	Event 4 30 Aug 2004
Ackerman (1997)	Dust < 0.0	Dust < 0.0	Dust < 0.0	Dust < -1.2	Dust < -0.35
Miller (2003)	1.3 < Dust < 2.7	1.3 < Dust < 2.27*	-0.55 < Dust < 1.11*	0.3 < Dust < 2.05*	0.45 < Dust < 1.08*
Roskovensky & Liou (2005)	Dust > 1.0 a = 0.8, b = 2	Dust > 1.0 a = 0.8, b = 1	Dust > 1.0 a = 0.9, b = -0.5	Dust > 1.0 a=1.2, b=-0.5	Dust > 1.0 a=0.25, b=0

*upper value is scene maximum

1485 Figure 1

1486
1487
1488

Figure 2

1492 Figure 3

1493

1494

1495

1495

Figure 4

		Dust pixels		
		A	M	R & L
Dust pixels	A	<div></div>		
	M	<div></div>	<div></div>	
	R & L	<div></div>	<div></div>	<div></div>
ALL indicate dust pixels		<div></div>		
ALL indicate non-dust pixels		<div></div>		

1496

Figure 5

1497
1498

1499
1500

Figure 8

Figure 9

1502
1503

Figure 10