Supporting Material

Efficient Isomer-Pure Synthesis of a Benzo[*b*]thiophene Analog of Pentacene

Brigitte Wex,[†] Bilal R. Kaafarani,[‡] and Douglas C. Neckers^{†,*}

neckers@photo.bgsu.edu

[†] Center for Photochemical Science, Bowling Green State University, Bowling Green, OH 43403

[‡] School of Chemistry and Biochemistry, Georgia Institute of Technology, Atlanta, GA 30332

Table of Contents.

Spectroscopic data for 5	S2
NMR Spectra for 7 - 8	S3 – S4
NMR Spectra of 2 (n=2)	S5

2,5-Thiophenedicarboxyaldehyde (5). The title compound was synthesized according to a literature procedure.¹ mp 113-114 °C (109-112 °C); ¹H NMR (CDCl₃, 400 MHz): δ 7.84 (2H, s), 10.04 (2H, s, CO-H); ¹³C NMR (CDCl₃, 100 MHz) δ 135.7, 149.8, 184.0; GC-MS (EI, 70 eV) *m/z* (%): 140 (100)[M⁺], 139 (72) [M⁺-1], 111 (25) [M⁺-29].

¹ Feringa, B. L.; Hulst, R.; Rikers, R.; Brandsma, L. *Synthesis* **1988**, (4), 316-318.

Figure S1. ^1H (a) and APT (b) NMR spectra of 2,5-bis(3-bromo-[2]thienylmethyl)thiophene (**7**).

Figure S2. ¹H (a) and ¹³C (b) NMR spectra of 2,5-bis[(3-formyl-[2]thienylmethyl)thiophene (**8**).

Figure S3. ^1H (a) and ^{13}C (b) NMR spectra of thieno[3,2-f:4,5-f']bis[1]benzothiophene (2, n=2).