SUPPORTING INFORMATION

**Efficient method for the synthesis of hetarenoindanones on the basis of 3-
arylheterenes and their conversion into hetarenoindenes**

Igor A. Kashulin and Ilya E. Nifant'ev.*

Department of Chemistry, Moscow State University, Moscow 119992, Russia

Fax: +7(095) 939 4523.

inif@org.chem.msu.ru

TABLE OF CONTENTS

1. Spectral and analytical data – S2
2. References – S6

Spectral and analytical data for **1a**, **1c-1e**, **2a**, **2c-2e**, **2h**, **3a**, **3c-3e**, **3h**:

3-phenyl-1-benzofuran¹ (1a)

¹H NMR (CDCl₃, 25 °C) δ: 7.93 (d, 1H *J*=7.4 Hz); 7.85 (s, 1H); 7.72 (d, 2H *J*=6.9 Hz); 7.63 (d, 1H *J*=7.4 Hz); 7.55 (t, 2H *J*=6.9 Hz); 7.42-7.35 (m, 3H). Anal Calcd for C₁₄H₁₀O (%): 86.57C; 5.19H. Found (%): 86.52C; 5.23H.

3-phenyl-1-benzothiophene² (1c)

¹H NMR (CDCl₃, 30°C) δ: 7.37-7.42 (4H, m); 7.48 (2H, t, *J* = 7.5 Hz); 7.58 (2H, d, *J* = 7.8 Hz); 7.90-7.92 (2H, m). ¹³C NMR (CDCl₃, 30°C) δ: 140.7, 138.1, 137.9, 136.0, 128.7, 127.5, 124.4, 124.3, 123.4, 122.9. Anal Calcd for C₁₄H₁₀S (%): 79.69C; 4.79H. Found (%): 79.66C; 4.81H.

2-methyl-4-phenylthiophene (1d)

¹H NMR (CDCl₃, 30°C) δ: 7.66 (d, 2H, *J* = 7.5 Hz), 7.47 (t, 2H, *J* = 7.5 Hz), 7.37 (t, 1H, *J* = 7.5 Hz), 7.28 (d, 1H *J* = 7.5 Hz), 7.16 (s, 1H), 2.61 (s, 3H, CH₃). ¹³C NMR (CDCl₃, 30°C) δ: 141.9, 140.3, 136.0, 128.6, 126.8, 126.1, 124.5, 117.9, 15.3. Anal Calcd for C₁₁H₁₀S (%): 75.82C; 5.78H. Found (%): 75.84C; 5.79H. mp 75-77 °C.

4-(4-tert-butylphenyl)-2-methylthiophene (1e)

¹H NMR (CDCl₃, 30°C) δ: 7.55 (d, 2H, *J* = 8.1 Hz), 7.46 (d, 2H, *J* = 8.1 Hz), 7.21 (s, 1H), 7.10 (s, 1H), 2.57 (s, 3H), 1.40 (s, 9H). ¹³C NMR (CDCl₃, 30°C) δ: 149.8, 141.9, 140.1, 133.3, 125.8, 125.5, 124.6, 117.4, 34.4, 31.2, 15.3. Anal Calcd for C₁₅H₁₈S (%): 78.21C; 7.88H. Found (%): 78.29C; 7.92H. mp 58-59 °C.

6H-indeno[2,1-b][1]benzofuran-6-one (2a)

^1H NMR (CDCl_3 , 25°C) δ : 7.73 (d, 1H, $J = 7.8$ Hz); 7.55 (d, 1H, $J = 7.8$ Hz); 7.46 (t, 1H, $J = 7.8$ Hz); 7.41-4.29 (m, 3H); 7.21-7.14 (m, 2H). ^{13}C NMR (CDCl_3 , 25°C) δ : 180.4, 161.4, 154.5, 141.2, 135.9, 134.8, 133.7, 128.5, 128.5, 124.6, 123.9, 121.9, 121.9, 120.1, 113.64. Anal Calcd for $\text{C}_{15}\text{H}_8\text{O}_2$ (%): 81.81C; 3.66H. Found (%): 81.78C; 3.55H. mp $105\text{-}106^\circ\text{C}$ (lit.³ mp $109\text{-}110^\circ\text{C}$).

6H-indeno[2,1-b][1]benzothiophen-6-one (2c)

^1H NMR (CDCl_3 , 25°C) δ : 7.87 (m, 1H); 7.81 (m, 1H); 7.76-7.38 (m, 3H); 7.34 (m, 2H); 7.16 (m, 1H). ^{13}C NMR (CDCl_3 , 25°C) δ : 186.9, 152.7, 148.2, 140.1, 137.0, 136.9, 133.6, 131.8, 127.98, 127.4, 125.7, 124.4, 123.8, 123.7, 119.5. Anal Calcd for $\text{C}_{15}\text{H}_8\text{OS}$ (%): 76.25C; 3.41H. Found (%): 76.11C; 3.49H. mp $198\text{-}199^\circ\text{C}$ (lit.⁴ mp $194\text{-}196^\circ\text{C}$).

2-methyl-8H-indeno[2,1-b]thiophen-8-one (2d)

^1H NMR (CDCl_3 , 30°C) δ : 7.44 (d, 1H, $J = 7.5$ Hz), 7.29 (t, 1H, $J = 7.5$ Hz), 7.14 (t, 1H, $J = 7.5$ Hz), 7.07 (d, 1H, $J = 7.5$ Hz), 6.79 (d, 1H, $J = 1.0$ Hz), 2.55 (d, 3H, $J = 1.0$ Hz). ^{13}C NMR (CDCl_3 , 30°C) δ : 185.2, 159.0, 156.1, 139.6, 137.4, 134.1, 133.1, 127.8, 123.3, 119.1, 118.9, 16.4. Anal Calcd for $\text{C}_{12}\text{H}_8\text{OS}$ (%): 71.97C; 4.03H. Found (%): 72.88C; 4.00H. mp $125\text{-}126^\circ\text{C}$.

6-tert-butyl-2-methyl-8H-indeno[2,1-b]thiophen-8-one (2e)

^1H NMR (CDCl_3 , 30°C) δ : 7.53 (d, 1H, $J = 1.9$ Hz), 7.29 (dd, 1H, $J = 7.5$ Hz and 1.9 Hz), 7.01 (d, 1H, $J = 7.5$ Hz), 6.80 (d, 1H, $J = 0.9$ Hz), 2.55 (d, 3H, $J = 0.9$ Hz), 1.33 (s, 9H). ^{13}C NMR (CDCl_3 , 30°C) δ : 185.8, 159.1, 156.0, 151.5, 137.6, 136.8, 134.0, 129.2, 121.3, 118.9, 118.8, 34.8, 30.9, 16.5. Anal Calcd for $\text{C}_{16}\text{H}_{16}\text{OS}$ (%): 74.96C; 6.29H. Found (%): 75.02C; 6.39H.

9H-pyrrolo[1,2-a]indol-9-one (2h)

¹H NMR (CDCl₃, 30°C) δ: 7.58 (d, 1H, *J* = 6.8 Hz), 7.44 (dt, 1H, *J* = 6.8 Hz, and 1.3 Hz), 7.16-7.08 (m, 3H), 6.78 (d, 1H, *J* = 3.7 Hz), 6.32 (dd, 1H, *J* = 3.7 Hz and 2.6 Hz). ¹³C NMR (CDCl₃, 30°C) δ: 179.4, 143.6, 133.9, 131.8, 130.1, 125.2, 124.3, 119.3, 115.7, 113.7, 110.1. Anal Calcd for C₁₁H₇NO (%): 78.09C; 4.17H; 8.28N. Found (%): 78.18C; 4.06H; 8.21N. mp 121-122 °C (lit.³ mp 121-122 °C).

6H-indeno[2,1-b][1]benzofuran (3a)

¹H NMR (CDCl₃, 25°C) δ: 7.83 (d, 1H, *J* = 7.4 Hz); 7.67 (d, 1H, *J* = 7.4 Hz); 7.60 (d, 1H, *J* = 7.4 Hz); 7.50 (d, 1H, *J* = 7.4 Hz); 7.39 (m, 3H); 7.24 (t, 1H, *J* = 7.4 Hz); 3.81 (s, 2H). ¹³C NMR (CDCl₃, 25°C) δ: 165.4, 160.2, 141.6, 137.3, 127.0, 124.9, 124.1, 124.0, 123.6, 123.3, 123.1, 119.5, 119.3, 112.1, 31.2. Anal Calcd for C₁₅H₁₀O (%): 87.36C; 4.89H. Found (%): 87.25C; 4.80H. mp 95-98 °C.

6H-indeno[2,1-b][1]benzothiophene (3c)

¹H (CDCl₃, 25°C) δ: 8.20 (d, 1H, *J* = 7.8 Hz); 7.92 (d, 1H, *J* = 7.8 Hz); 7.88 (d, 1H, *J* = 7.8 Hz); 7.56 (d, 1H, *J* = 7.8 Hz); 7.52 (t, 1H, *J* = 7.8 Hz); 7.44 (t, 1H, *J* = 7.8 Hz); 7.39 (t, 1H, *J* = 7.8 Hz); 7.27 (t, 1H, *J* = 7.8 Hz), 3.97 (s, 2H). ¹³C NMR (CDCl₃, 25°) δ: 146.2, 145.6, 144.8, 140.9, 139.5, 133.0, 126.7, 124.5, 124.5, 124.2, 123.7, 123.5, 121.7, 118.9, 35.2. Anal Calcd for C₁₅H₁₀S (%): 81.04C; 4.53H. Found (%): 80.91C; 4.58H. mp 112-113 °C (lit.⁵ mp 111-112 °C).

2-methyl-8H-indeno[2,1-b]thiophene (3d).

¹H NMR (CDCl₃, 30°C) δ: 7.51 (m, 2H), 7.37 (t, 1H, *J* = 7.9 Hz), 7.23 (t, 1H, *J* = 7.9 Hz), 7.01 (m, 1H), 3.81 (s, 2H), 2.61 (d, 3H, *J* = 0.8 Hz). ¹³C NMR (CDCl₃, 30°C) δ: 146.8, 146.2, 143.4,

141.3, 139.5, 126.5, 124.5, 123.9, 118.7, 116.6, 34.6, 16.0. Anal Calcd for C₁₂H₁₀S (%): 77.37C; 5.41H. Found (%): 77.26C; 5.36H. mp 77-78 °C.

6-tert-butyl-2-methyl-8H-indeno[2,1-b]thiophene (3e)

¹H NMR (CDCl₃, 30°C) δ: 7.58 (s, 1H), 7.46-7.37 (m, 2H), 6.98 (s, 1H), 3.82 (s, 2H), 2.62 (s, 3H), 1.44 (s, 9H). Anal Calcd for C₁₆H₁₈S (%): 79.29C; 7.49H. Found (%): 79.21C; 7.37H. mp 139-141 °C.

9H-pyrrolo[1,2-a]indole (3h)

¹H NMR (CDCl₃, 30°C) δ: 7.47 (d, 1H, *J* = 6.8 Hz), 7.41- 7.32 (m, 2H), 7.22-7.12 (m, 2H), 6.54 (t, 1H, *J* = 3.2 Hz), 6.25 (m, 1H), 3.91 (s, 2H). ¹³C NMR (CDCl₃, 30°C) δ: 141.0, 135.2, 134.8, 128.2, 127.2, 125.6, 122.8, 113.0, 109.5, 101.5, 28.8. Anal Calcd for C₁₁H₉N (%): 85.13C; 5.85H; 9.03N. Found (%): 86.02C; 5.80H; 8.94N. mp 91-92 °C (lit.⁶ mp 90-91 °C).

References

1. Davies, W; Middleton, S. *J.Chem. Soc.*, **1958**, 822.
2. Katritzky, A. R.; Kirichenko, K.; Yu Ji; Prakash; I. *Chemistry of Heterocyclic Compounds*, **2002**, 38, 156.
3. Campo, M. A.; Larock, R. C. *J. Org. Chem.*, **2002**, 67, 5616.
4. Krubsack, A. J; Higa, T. *Tetrahedron Lett.*, **1972**, 4823.
5. Jeffries, A. T.; Moore, K. C.; Ondeyka, D.M.; Springsteen, A.W.; MacDowell, D. W. H. *J. Org. Chem.*, **1981**, 46, 2885.
6. Caddick, S.; Aboutayab, K.; Jenkins, K.; West, R. I. *J. Chem. Soc., Perkin Trans.1*, **1996**, 675.