Supporting Information

Efficient Oxidative Radical Cyclizations of Ester Enolates with Carbocation Desilylation as Termination - Synthesis of Cyclopentanoid Monoterpenes and Analogs

Ullrich Jahn*, Philip Hartmann, Emmi Kaasalainen

Institut für Organische Chemie, Technische Universität Braunschweig, Hagenring 30, D-38106

Braunschweig; FAX: +49-531-391-5388, e-mail: u.jahn@tu-bs.de

General: All reactions were conducted in flame dried glassware under nitrogen atmosphere. THF, DME, DMF, DMSO, HMPA, diisopropylamine, and 2,2,6,6-tetramethylpiperidine were dried after standard methods under nitrogen atmosphere. TLC plates POLYGRAM SIL G/UV₂₅₄ (Macherey-Nagel) were used for monitoring reactions. Flash column chromatographic separations were performed on silica gel 60 (Fluka, 230 - 400 mesh) with *n*-hexane/ethyl acetate as eluent in the given ratio. IR spectra were taken on a Nicolet DX-320 FT-IR spectrometer. ¹H and ¹³C NMR spectra were recorded, unless otherwise noted, in CDCl₃ on Bruker DRX 400 or AC 200 spectrometers at 400 or 200, and 100 or 50 MHz, respectively. Connectivity was determined by ¹H-¹H COSY experiments. ¹³C NMR assignments were obtained from DEPT and C,H-correlation experiments. EI-mass spectra were recorded on a Finnigan MAT 8430 and MAT 8400 spectrometer at 70 eV. Combustion analyses were performed at the Microanalytical Laboratories of the Technical University of Braunschweig.

Diethyl 2-(6-Trimethylsilylhex-4-enyl)malonates or Diethyl 2-(7-Trimethylsilylhept-4-enyl)malonates 9a-d,f (General Procedure):

At 0°C under N₂, 3.0 mL (30 mmol) diethyl malonate were added to a suspension of 450 mg (15 mmol) NaH (80% in mineral oil) in 10 mL dry DMF and 35 mL dry THF (for the preparation of **9c** in 50 mL neat DMF). The mixture was stirred for one hour and the resulting clear solution was transferred via canula to a solution of 10 mmol of the corresponding mesylate or bromide **8a-d,f**¹ in 70 mL dry THF at -78°C (DMF for **8c**). The temperature was raised to room temperature, 149 mg (1.0 mmol) of anhydrous NaI was added and the mixture was stirred under reflux until the reaction was complete by tlc (ca. 72 h). The reaction mixture was hydrolyzed by 10 mL sat'd NaHCO₃ solution, concentrated under reduced pressure and partitioned between water and diethyl ether. After separation of the layers, the aqueous layer was extracted

¹ **8a,b**: Paquette, L. A.; Mendez-Andino, J. L. *J. Org. Chem.* **1998**, 63, 9061-9068. **8c**: Moret, E.; Franzini, L.; Schlosser, M. *Chem. Ber./Recueil* **1997**, 130, 335-339. **8d**: Hudson, C. M.; Marzabadi, M. R.; Moeller, K. D.; New, D. G. *J. Am. Chem. Soc.* **1991**, 113, 7372-7385. **8f**: Lee, T. V.; Channon, J. A.; Cregg, C.; Porter, J. R.; Roden, F. S.; Yeoh, H. T.-L. *Tetrahedron* **1989**, 45, 5877-5886.

three times with ether. The combined organic layers were dried over Na_2SO_4 . The solvent was removed *in vacuo*, and excess diethyl malonate was removed by distillation at $55^\circ\text{C}/0.6$ mbar. The residue was further purified by flash column chromatography. Explicite conditions are provided at the individual compounds.

(E/Z)-Diethyl 2-[5-Methyl-6-(trimethylsilyl)hex-4-enyl]malonate (9a): Flash column chromatography (25/1 gradient to 10/1) yielded 2.95 g (90%) **9a** (*E/Z* 1.8:1) [R_f (10/1) = 0.43] as a colorless oil. - IR (neat): $\tilde{\nu}$ = 2957 cm^{-1} , 1754, 1736, 1248, 1153, 858. - (*E*)-**9a**: - ^1H NMR (200 MHz): δ = 0.00 (s, 9 H, SiCH_3), 1.27 (t, J = 7.1 Hz, 6 H, OCH_2CH_3), 1.41 (m, 2 H, $=\text{CHCH}_2\text{CH}_2$), 1.46 (s, 2 H, CH_2Si), 1.59 (s, 3 H, $=\text{CCH}_3$), 1.84 – 2.07 (m, 4 H, $=\text{CHCH}_2\text{CH}_2\text{CH}_2$), 3.32 (t, J = 7.6 Hz, 1 H, CHCO_2), 4.20 (q, J = 7.1 Hz, 4 H, OCH_2), 4.94 (m, 1 H, $=\text{CH}$). - ^{13}C NMR (50 MHz): δ = -1.3 (q, SiCH_3), 14.0 (q, OCH_2CH_3), 18.5 (q, $=\text{CCH}_3$), 27.6 (t), 27.7 (t), 28.3 (t), 29.8 (t, CH_2Si), 51.9 (d, CHCO_2), 61.1 (t, OCH_2), 121.6 (d, $=\text{CH}$), 133.3 (s, $=\text{C}$), 169.5 (s, CO_2). - (*Z*)-**9a**: - ^1H NMR (200 MHz): δ = 0.03 (s, 9 H, SiCH_3), 1.27 (t, J = 7.1 Hz, 6 H, OCH_2CH_3), 1.41 (m, 2 H, $=\text{CHCH}_2\text{CH}_2$), 1.50 (s, 2 H, CH_2Si), 1.67 (d, J = 1.1 Hz, 3 H, $=\text{CCH}_3$), 1.84 – 2.07 (m, 4 H, $=\text{CHCH}_2\text{CH}_2\text{CH}_2$), 3.32 (t, J = 7.6 Hz, 1 H, CHCO_2), 4.19 (q, J = 7.1 Hz, 4 H, OCH_2), 4.94 (m, 1 H, $=\text{CH}$). - ^{13}C NMR (50 MHz): δ = -0.8 (q, SiCH_3), 14.0 (q, OCH_2CH_3), 23.2 (t, CH_2Si), 26.21 (q, $=\text{CCH}_3$), 27.6 (t), 28.0 (t), 28.5 (t), 52.0 (d, CHCO_2), 61.1 (t, OCH_2), 121.3 (d, $=\text{CH}$), 133.5 (s, $=\text{C}$), 169.5 (s, CO_2). - MS m/z (%): 328 (15) [M^+], 283 (5), 255 (4), 245 (24), 217 (17), 209 (6), 127 (25), 73 (100). - HRMS: $\text{C}_{17}\text{H}_{32}\text{O}_4\text{Si}$: calcd. 328.2070; found 328.2060 ± 2 ppm. - $\text{C}_{17}\text{H}_{32}\text{O}_4\text{Si}$ (328.5): calcd. C 62.15, H 9.82; found C 62.26, H 9.92.

(Z)-Diethyl 2-[6-(Trimethylsilyl)hex-4-enyl]malonate (9b): Flash column chromatography (25/1 gradient to 10/1) yielded 2.39 g (76%) of **9b** (*E/Z* 1:10) [R_f (15/1) = 0.32] as a colorless oil. - IR (neat): $\tilde{\nu}$ = 2956 cm^{-1} , 1754, 1736, 1370, 1248, 1153, 858. - ^1H NMR (200 MHz): δ = 0.00 (s, 9 H, SiCH_3), 1.27 (t, J = 7.1 Hz, 6 H, OCH_2CH_3), 1.39 (m, 2 H, $=\text{CHCH}_2\text{CH}_2$), 1.46 (d, J = 8.5 Hz, 2 H, CH_2Si), 1.87 (m, 2 H, CH_2CHCO_2), 2.02 (q, J = 7.1 Hz, 2 H, $=\text{CHCH}_2\text{CH}_2$), 3.33 (t, J = 7.5 Hz, 1 H, CHCO_2), 4.20 (q, J = 7.1 Hz, 4 H, OCH_2), 5.23 (dt, J = 10.8, 6.9 Hz, 1 H, $=\text{CHCH}_2\text{CH}_2$), 5.39 (m, 1 H, $=\text{CHCH}_2\text{Si}$). - ^{13}C NMR (50 MHz): δ = -1.9 (q, SiCH_3), 14.0 (q, OCH_2CH_3), 18.4 (t, CH_2Si), 26.6 (t), 27.4 (t), 28.4 (t), 51.94 (d, CHCO_2), 61.2 (t, OCH_2), 126.1 (d, $=\text{CH}$), 126.4 (d, $=\text{CH}$), 169.4 (s, CO_2). - MS m/z (%): 314 (27) [M^+], 299 (10), 269 (7), 245 (41), 217 (18), 195 (18), 127 (21), 80 (25), 73 (100). - HRMS: $\text{C}_{16}\text{H}_{30}\text{O}_4\text{Si}$: calcd. 314.1913; found 314.1904 ± 2 ppm. - $\text{C}_{16}\text{H}_{30}\text{O}_4\text{Si}$ (314.5): calcd. C 61.11, H 9.62; found C 61.18, H 9.58.

Resonances, assigned to (*E*)-**9b**: - ^{13}C NMR (50 MHz): δ = -2.1 (q, SiCH_3), 22.5 (t), 27.5 (t), 28.1 (t), 32.2 (t), 51.88 (d, CHCO_2), 126.9 (d, $=\text{CH}$), 127.7 (d, $=\text{CH}$).

(E/Z)-Diethyl 2-[6-Methyl-7-(trimethylsilyl)hept-5-en-2-yl]malonate (9c): Flash column chromatography (25/1 gradient to 10/1) yielded 0.96 g (50%) of **9c** (*Z/E* 5:1) [R_f (10/1) = 0.36] as a

colorless oil. - IR (neat): $\tilde{\nu}$ = 2961 cm^{-1} , 1756, 1735, 1249, 1156, 858. - (Z)-**9c**: - ^1H NMR (400 MHz, C_6D_6): δ = 0.00 (s, 9 H, SiCH_3), 0.91 (t, J = 7.1 Hz, 6 H, OCH_2CH_3), 1.06 (d, J = 6.8 Hz, 3 H, CHCH_3), 1.31 (m, 1 H, $=\text{CHCH}_2\text{CH}_2$), 1.46 (d, J_{AB} = 13.5 Hz, 1 H, CH_2Si), 1.50 (d, J_{AB} = 13.5 Hz, 1 H, CH_2Si), 1.62 (m, 1 H, $=\text{CHCH}_2\text{CH}_2$), 1.64 (s, 3 H, $=\text{CCH}_3$), 1.90 – 2.06 (m, 2 H, $=\text{CHCH}_2$), 2.46 (m, 1 H, CHCH_3), 3.35 (d, J = 7.6 Hz, 1 H, CHCO_2), 3.93 (q, J = 7.1 Hz, 4 H, OCH_2), 5.05 (t, J = 6.8 Hz, 1 H, $=\text{CH}$). - ^{13}C NMR (100 MHz, C_6D_6): δ = -0.6 (q, SiCH_3), 14.07 (q, OCH_2CH_3), 14.09 (q, OCH_2CH_3), 17.2 (q, CHCH_3), 23.3 (t, CH_2Si), 26.4 (q, $=\text{CCH}_3$), 26.4 (t, $=\text{CHCH}_2$), 33.4 (d, CHCH_3), 35.0 (t, $=\text{CHCH}_2\text{CH}_2$), 58.0 (d, CHCO_2), 60.8 (t, OCH_2), 60.9 (t, OCH_2), 123.3 (d, $=\text{CH}$), 133.2 (s, $=\text{C}$), 168.6 (s, CO_2), 168.7 (s, CO_2). - (E)-**9c**: - ^1H NMR (400 MHz, C_6D_6): δ = -0.01 (s, 9 H, SiCH_3), 0.91 (t, J = 7.1 Hz, 6 H, OCH_2CH_3), 1.06 (d, J = 6.8 Hz, 3 H, CHCH_3), 1.31 (m, 1 H, $=\text{CHCH}_2\text{CH}_2$), 1.43 (d, J = 5.8 Hz, 2 H, CH_2Si), 1.55 (s, 3 H, $=\text{CCH}_3$), 1.62 (m, 1 H, $=\text{CHCH}_2\text{CH}_2$), 1.90 – 2.06 (m, 2 H, $=\text{CHCH}_2$), 2.46 (m, 1 H, CHCH_3), 3.35 (d, J = 7.6 Hz, 1 H, CHCO_2), 3.93 (q, J = 7.1 Hz, 4 H, OCH_2), 4.98 (t, J = 7.0 Hz, 1 H, $=\text{CH}$). - ^{13}C NMR (100 MHz, C_6D_6): δ = -1.2 (q, SiCH_3), 14.07 (q, OCH_2CH_3), 14.09 (q, OCH_2CH_3), 17.1 (q, CHCH_3), 18.7 (q, $=\text{CCH}_3$), 25.9 (t, $=\text{CHCH}_2$), 30.0 (t, CH_2Si), 33.2 (d, CHCH_3), 35.1 (t, $=\text{CHCH}_2\text{CH}_2$), 57.8 (d, CHCO_2), 60.8 (t, OCH_2), 60.9 (t, OCH_2), 122.4 (d, $=\text{CH}$), 133.1 (s, $=\text{C}$), 168.6 (s, CO_2), 168.7 (s, CO_2). - MS m/z (%): 342 (17) [M^+], 269 (6), 259 (39), 141 (49), 73 (100). - HRMS: $\text{C}_{18}\text{H}_{34}\text{O}_4\text{Si}$: calcd. 342.2226; found 342.2226.

(E/Z)-Diethyl 2-[6-Methyl-7-(trimethylsilyl)hept-5-enyl]malonate (**9d**): Flash column chromatography (30/1 gradient to 10/1) yielded 2.43 g (71%) of **9d** (Z/E 1.2:1) [R_f (10/1) = 0.42] as a colorless oil. - IR (neat): $\tilde{\nu}$ = 2956 cm^{-1} , 1754, 1736, 1249, 1155, 859. - (E)-**9d**: - ^1H NMR (400 MHz): δ = -0.03 (s, 9 H, SiCH_3), 1.25 (t, J = 7.1 Hz, 6 H, OCH_2CH_3), 1.32 (m, 4 H, $=\text{CHCH}_2\text{CH}_2\text{CH}_2$), 1.43 (s, 2 H, CH_2Si), 1.56 (s, 3 H, $=\text{CCH}_3$), 1.87 (m, 2 H, CH_2CHCO_2), 1.95 (m, 2 H, $=\text{CHCH}_2$), 3.29 (t, J = 7.6 Hz, 1 H, CHCO_2), 4.17 (q, J = 7.1 Hz, 4 H, OCH_2), 4.89 (dt, J = 7.1, 0.9 Hz, 1 H, $=\text{CH}$). - ^{13}C NMR (100 MHz): δ = -1.3 (q, SiCH_3), 14.0 (q, OCH_2CH_3), 18.5 (q, $=\text{CCH}_3$), 27.0 (t), 27.8 (t), 28.6 (t), 29.7 (t), 29.8 (t, CH_2Si), 52.04 (d, CHCO_2), 61.2 (t, OCH_2), 122.2 (d, $=\text{CH}$), 132.8 (s, $=\text{C}$), 169.5 (s, CO_2). - (Z)-**9d**: - ^1H NMR (400 MHz): δ = 0.00 (s, 9 H, SiCH_3), 1.25 (t, J = 7.1 Hz, 6 H, OCH_2CH_3), 1.32 (m, 4 H, $=\text{CHCH}_2\text{CH}_2\text{CH}_2$), 1.47 (s, 2 H, CH_2Si), 1.64 (d, J = 1.2 Hz, 3 H, $=\text{CCH}_3$), 1.87 (m, 4 H, $=\text{CHCH}_2\text{CH}_2\text{CH}_2\text{CH}_2$), 3.30 (t, J = 7.6 Hz, 1 H, CHCO_2), 4.17 (q, J = 7.1 Hz, 4 H, OCH_2), 4.95 (m, 1 H, $=\text{CH}$). - ^{13}C NMR (100 MHz): δ = -0.8 (q, SiCH_3), 14.0 (q, OCH_2CH_3), 23.2 (t, CH_2Si), 26.2 (q, $=\text{CCH}_3$), 27.1 (t), 28.2 (t), 28.69 (t), 29.5 (t), 52.01 (d, CHCO_2), 61.2 (t, OCH_2), 121.9 (d, $=\text{CH}$), 133.1 (s, $=\text{C}$), 169.5 (s, CO_2). - MS m/z (%): 342 (23) [M^+], 245 (39), 217 (16), 173 (24), 73 (100). - HRMS: $\text{C}_{18}\text{H}_{34}\text{O}_4\text{Si}$: calcd. 342.2227; found 342.2216 $\pm$ 2 ppm. - $\text{C}_{18}\text{H}_{34}\text{O}_4\text{Si}$ (342.5): calcd. C 63.11, H 10.00; found C 63.01, H 10.07.

Diethyl 2-[(4-Trimethylsilylmethyl)pent-4-enyl]malonate (**9f**): Flash column chromatography (30/1 gradient to 10/1) yielded 1.51 g (48%) of **9f** [R_f (10/1) = 0.40] as a colorless oil. - IR (neat): $\tilde{\nu}$ = 2957

cm⁻¹, 1754, 1736, 1249, 115, 858. - ¹H NMR (200 MHz): δ = 0.00 (s, 9 H, SiCH₃), 1.26 (t, J = 7.1 Hz, 6 H, OCH₂CH₃), 1.46 (m, 2 H, =CCH₂CH₂), 1.49 (d, J = 0.6 Hz, 2 H, CH₂Si), 1.83 – 2.04 (m, 4 H, CH₂CH₂CH₂CHCO₂), 3.32 (t, J = 7.5 Hz, 1 H, CHCO₂), 4.19 (q, J = 7.1 Hz, 4 H, OCH₂), 4.51 (br. s, 1 H, =CH₂), 4.57 (d, J = 1.3 Hz, 1 H, =CH₂). - ¹³C NMR (50 MHz): δ = -1.4 (q, SiCH₃), 14.0 (q, OCH₂CH₃), 25.4 (t), 26.6 (t, CH₂Si), 28.4 (t), 37.6 (t, =CCH₂CH₂), 51.9 (d, CHCO₂), 61.2 (t, OCH₂), 107.3 (t, =CH₂), 146.7 (s, =C), 169.4 (s, CO₂). - MS m/z (%): 314 (3) [M⁺], 269 (8), 245 (19), 217 (17), 154 (56), 127 (23), 73 (100). - HRMS: C₁₆H₃₀O₄Si: calcd. 314.1913; found 314.1904 $\pm$ 2 ppm. - C₁₆H₃₀O₄Si (314.5): calcd. C 61.11, H 9.62; found C 61.31, H 9.88.

Ethyl 3,7-Dimethyl-8-(trimethylsilyl)oct-6-enoate (9e) via Krapcho Reaction: A solution of 684 mg (2.0 mmol) **9c**, 170 mg (4.0 mmol) anhydrous LiCl, and 0.072 mL (4.0 mmol) water in 7 mL dry DMSO was heated to 185°C until the reaction was finished by tlc. After cooling to room temperature, the solution was diluted with 5 mL sat'd NaHCO₃ solution and 50 mL of water and extracted four times with ether. The combined ethereal layers were washed with brine and dried over Na₂SO₄. The solvent was removed *in vacuo* and the residue was purified by flash column chromatography (50/1 gradient to 10/1) to give 433 mg (80%) of **9e** (Z/E 7:1) [R_f (10/1) = 0.47] as a colorless oil. - IR (neat): $\tilde{\nu}$ = 2959 cm⁻¹, 1738, 1249, 1159, 855. - (Z)-**9e**: - ¹H NMR (400 MHz): δ = 0.00 (s, 9 H, SiCH₃), 0.92 (d, J = 6.6 Hz, 3 H, CHCH₃), 1.20 (m, 1 H, =CHCH₂CH₂), 1.234 (t, J = 7.1 Hz, 3 H, OCH₂CH₃), 1.32 (m, 1 H, =CHCH₂CH₂), 1.47 (s, 2 H, CH₂Si), 1.63 (d, J = 1.2 Hz, 3 H, =CCH₃), 1.80 - 2.00 (m, 3 H, =CHCH₂CH₂CH), 2.08 (dd, J = 14.6, 8.2 Hz, 1 H, CH₂CO₂), 2.28 (ddd, J = 14.5, 5.8, 2.9 Hz, 1 H, CH₂CO₂), 4.10 (q, J = 7.1 Hz, 2 H, OCH₂), 4.94 (t, J = 6.8 Hz, 1 H, =CH). - ¹³C NMR (100 MHz): δ = -0.7 (q, SiCH₃), 14.3 (q, OCH₂CH₃), 19.64 (q, CHCH₃), 23.2 (t, CH₂Si), 25.9 (t, =CHCH₂), 26.2 (q, =CCH₃), 30.2 (d, CHCH₃), 36.9 (t, =CHCH₂CH₂), 41.9 (t, CH₂CO₂), 60.1 (t, OCH₂), 121.9 (d, =CH), 133.1 (s, =C), 173.3 (s, CO₂). - (E)-**9e**: - ¹H NMR (400 MHz): δ = -0.03 (s, 9 H, SiCH₃), 0.92 (d, J = 6.6 Hz, 3 H, CHCH₃), 1.20 (m, 1 H, =CHCH₂CH₂), 1.230 (t, J = 7.0 Hz, 3 H, OCH₂CH₃), 1.32 (m, 1 H, =CHCH₂CH₂), 1.43 (s, 2 H, CH₂Si), 1.56 (br. s, 3 H, =CCH₃), 1.80 - 2.00 (m, 3 H, =CHCH₂CH₂CH), 2.08 (m, 1 H, CH₂CO₂), 2.28 (m, 1 H, CH₂CO₂), 4.10 (m, 2 H, OCH₂), 4.89 (t, J = 7.2 Hz, 1 H, =CH). - ¹³C NMR (100 MHz): δ = -1.3 (q, SiCH₃), 14.3 (q, OCH₂CH₃), 18.6 (q, =CCH₃), 19.6 (q, CHCH₃), 25.6 (t, =CHCH₂), 29.8 (t, CH₂Si), 30.0 (d, CHCH₃), 37.1 (t, =CHCH₂CH₂), 41.9 (t, CH₂CO₂), 60.1 (t, OCH₂), 122.2 (d, =CH), 132.8 (s, =C), 173.3 (s, CO₂). - MS m/z (%): 270 (17) [M⁺], 187 (58), 73 (100), 69 (50). - HRMS: C₁₅H₃₀O₂Si: calcd. 270.2015; found 270.2020.

Oxidative Cyclization of the Lithium Enolates of Silylated Malonates 9a-d,f (General Procedure): At -70°C under N₂, 1.0 mmol of malonate **9a-d,f** dissolved in 1.0 mL dry DME was added to a solution of 1.3 mmol LDA prepared from 0.19 mL *i*-Pr₂NH and 0.81 mL *n*-BuLi (1.6 M in hexanes) in 19 mL dry DME. The mixture was stirred for 30 min with rising temperature. At 0°C, 2.3-3 equiv. of **2** was added in

portions to the solution until no more oxidant was consumed and the reaction mixture remained green-blue. The inhomogeneous mixture was stirred at 0°C for additional 90 min and quenched by a few drops of sat'd NH₄Cl solution. The inhomogeneous mixture was diluted with 20 mL of ether and filtered through a pad of silica gel. After removing the solvent *in vacuo*, the inhomogeneous residue was preadsorbed on silica gel, transferred to a silica gel flash column and purified (50/1 gradient to 1/1). As the first fraction pure ferrocene was eluted, followed by the cyclic compounds **10**. If necessary, the products can be further purified as indicated under the individual derivatives.

Diethyl 2-Isopropenylcyclopentane-1,1-dicarboxylate (10a): Flash chromatography (50/1) gave 220 mg (86%) of **10a** [*R_f* (10/1) = 0.61] as a colorless oil. - IR (neat): $\tilde{\nu}$ = 2980 cm⁻¹, 1729, 1448, 1261, 1211, 1176, 1158, 1031. - ¹H NMR (400 MHz): δ = 1.16 (t, *J* = 7.1 Hz, 3 H, OCH₂CH₃), 1.21 (t, *J* = 7.0 Hz, 3 H, OCH₂CH₃), 1.52 (m, 1 H, CHCH₂CH₂), 1.72 (s, 3 H, =CCH₃), 1.74 (m, 1 H, CHCH₂), 1.84 (m, 2 H, CHCH₂CH₂), 1.98 (ddd, *J* = 14.1, 8.4, 3.1 Hz, 1 H, CH₂CH₂C), 2.58 (ddd, *J* = 13.8, 9.6, 8.1 Hz, 1 H, CH₂CH₂C), 3.30 (dd, *J* = 9.5, 7.9 Hz, 1 H, CH), 4.06 (q, *J* = 7.1 Hz, 2 H, OCH₂), 4.12 (q, *J* = 7.1 Hz, 2 H, OCH₂), 4.73 (d, *J* = 1.0 Hz, 1 H, =CH₂), 4.74 (d, *J* = 1.4 Hz, 1 H, =CH₂). - ¹³C NMR (100 MHz): δ = 13.9 (q, OCH₂CH₃), 14.0 (q, OCH₂CH₃), 23.2 (q, CCH₃), 23.4 (t), 30.8 (t), 35.2 (t), 51.7 (d, CH), 60.9 (t, OCH₂), 61.1 (t, OCH₂), 64.0 (s, CCH₂), 112.2 (t, =CH₂), 145.2 (s, =C), 170.9 (s, CO₂), 172.8 (s, CO₂). - MS *m/z* (%): 254 (22) [M⁺], 180 (100), 163 (26), 135 (55), 107 (32). - HRMS: C₁₄H₂₂O₄: calcd. 254.1518; found 254.1518 ± 2 ppm.

Diethyl 2-Vinylcyclopentane-1,1-dicarboxylate (10b): Flash chromatography (50/1) gave 235 mg (98%) of **10b** [*R_f* (10/1) = 0.42] as a colorless oil. - IR (neat): $\tilde{\nu}$ = 2982 cm⁻¹, 1729, 1263, 1223, 1183, 1098. - ¹H NMR (200 MHz): δ = 1.22 (t, *J* = 7.1 Hz, 3 H, OCH₂CH₃), 1.25 (t, *J* = 7.1 Hz, 3 H, OCH₂CH₃), 1.65 (m, 2 H), 1.70 - 2.15 (m, 3 H), 2.46 (m, 1 H), 3.50 (m, 1 H, =CHCH), 4.08 (dq, *J* = 10.7, 7.1 Hz, 1 H, OCH₂), 4.16 (dq, *J* = 10.8, 7.1 Hz, 2 H, OCH₂), 4.24 (dq, *J* = 10.7, 7.1 Hz, 1 H, OCH₂), 4.98 (dd, *J* = 10.3, 1.9 Hz, 1 H, =CH₂), 5.13 (dd, *J* = 17.2, 1.9 Hz, 1 H, =CH₂), 5.81 (ddd, *J* = 17.2, 10.2, 7.9 Hz, 1 H, =CH). - ¹³C NMR (50 MHz): δ = 13.9 (q, OCH₂CH₃), 22.8 (t), 30.7 (t), 34.4 (t), 47.9 (d, =CHCH), 60.7 (t, OCH₂), 61.0 (t, OCH₂), 64.0 (s, CCO₂), 115.6 (t, =CH₂), 137.7 (s, =C), 171.5 (s, CO₂), 172.6 (s, CO₂). - MS *m/z* (%): 240 (11) [M⁺], 195 (10), 167 (20), 166 (100), 149 (17), 138 (22), 120 (18), 93 (28). - HRMS: C₁₃H₂₀O₄: calcd. 240.1362; found 240.1356 ± 3 ppm.

Diethyl 2-Isopropenyl-5-methylcyclopentane-1,1-dicarboxylate (10c): Flash column chromatography (50/1 gradient to 30/1) yielded 158 mg (59%) of **10c** [*R_f* (10/1) = 0.39] as an oily, inseparable *trans/cis*-diastereomeric mixture (2.5:1). - IR (neat): $\tilde{\nu}$ = 2964 cm⁻¹, 1725, 1448, 1255, 1209, 1181. - *trans*-**10c**: - ¹H NMR (400 MHz): δ = 0.90 (d, *J* = 7.0 Hz, 3 H, CHCH₃), 1.13 (t, *J* = 7.2 Hz, 3 H,

OCH₂CH₃), 1.19 (t, $J = 7.1$ Hz, 3 H, OCH₂CH₃), 1.22 (m, 1 H, CH₂CHCH₃), 1.62 (m, 1 H, =CCHCH₂), 1.65 (d, $J = 0.7$ Hz, 3 H, =CCH₃), 1.85 – 2.01 (m, 2 H, CHCH₂CH₂), 2.85 (dq, $J = 10.3, 7.0$ Hz, 1 H, CH₂CHCH₃), 3.33 (dd, $J = 8.7, 8.1$ Hz, 1 H, =CCHCH₂), 3.93 (dq, $J = 10.7, 7.1$ Hz, 1 H, OCH₂), 4.06 (dq, $J = 10.8, 7.1$ Hz, 1 H, OCH₂), 4.07 (dq, $J = 10.7, 7.1$ Hz, 1 H, OCH₂), 4.17 (dq, $J = 10.8, 7.1$ Hz, 1 H, OCH₂), 4.67 (d, $J = 0.7$ Hz, 1 H, =CH₂), 4.69 (d, $J = 1.5$ Hz, 1 H, =CH₂). - ¹³C NMR (100 MHz): $\delta = 13.8$ (q, OCH₂CH₃), 14.1 (q, OCH₂CH₃), 17.2 (q, CHCH₃), 23.0 (q, CCH₃), 29.8 (t, =CCHCH₂), 32.7 (t, =CCHCH₂CH₂), 40.8 (d, CHCH₃), 52.1 (d, =CCH), 60.6 (t, OCH₂), 60.7 (t, OCH₂), 67.5 (s, CCO₂), 112.3 (t, =CH₂), 146.1 (s, =C), 171.0 (s, CO₂), 171.4 (s, CO₂). - *cis*-**10c**: - ¹H NMR (400 MHz): $\delta = 1.09$ (d, $J = 6.9$ Hz, 3 H, CHCH₃), 1.16 (t, $J = 7.1$ Hz, 3 H, OCH₂CH₃), 1.20 (t, $J = 7.1$ Hz, 3 H, OCH₂CH₃), 1.67 (m, 1 H, CH₂CHCH₃), 1.69 (s, 3 H, =CCH₃), 1.80 (m, 1 H, =CCHCH₂), 1.87 (m, 1 H, CH₂CHCH₃), 1.95 (m, 1 H, =CCHCH₂), 2.30 (m, 1 H, CH₂CHCH₃), 3.18 (dd, $J = 10.7, 8.7$ Hz, 1 H, =CCHCH₂), 3.98 – 4.21 (m, 4 H, OCH₂), 4.73 (s, 1 H, =CH₂), 4.74 (s, 1 H, =CH₂). - ¹³C NMR (100 MHz): $\delta = 13.98$ (q, OCH₂CH₃), 14.01 (q, OCH₂CH₃), 15.4 (q, CHCH₃), 23.1 (q, CCH₃), 27.7 (t, =CCHCH₂), 30.7 (t, =CCHCH₂CH₂), 44.5 (d, CHCH₃), 53.4 (d, =CCH), 60.2 (t, OCH₂), 60.8 (t, OCH₂), 66.7 (s, CCO₂), 112.2 (t, =CH₂), 144.7 (s, =C), 169.2 (s, CO₂), 172.1 (s, CO₂). - MS m/z (%): 268 (12) [M⁺], 222 (12), 194 (88), 177 (18), 149 (27), 148 (32), 121 (100). - HRMS: C₁₅H₂₄O₄: calcd. 268.1675; found 268.1676.

Figure 1: NOE investigations of **10c** gave the following key results:

Diethyl 2-Isopropenylcyclohexane-1,1-dicarboxylate (10d): Flash column chromatography (50/1 gradient to 30/1) yielded 183 mg (68%) of **10d** [R_f (10/1) = 0.47] as a colorless oil. - IR (neat): $\tilde{\nu} = 2938$ cm⁻¹, 1733, 1263, 1244, 1213, 1143. - ¹H NMR (200 MHz): $\delta = 1.15$ (t, $J = 7.1$ Hz, 3 H, OCH₂CH₃), 1.17 (t, $J = 7.0$ Hz, 3 H, OCH₂CH₃), 1.23 – 1.53 (m, 4 H), 1.73 (s, 3 H, =CCH₃), 1.76 (m, 2 H), 1.94 – 2.23 (m, 2 H), 2.79 (t, $J = 5.9$ Hz, 1 H, =CCH), 4.06 (q, $J = 7.1$ Hz, 2 H, OCH₂), 4.09 (q, $J = 7.1$ Hz, 2 H, OCH₂), 4.71 (d, $J = 2.0$ Hz, 1 H, =CH₂), 4.73 (d, $J = 1.5$ Hz, 1 H, =CH₂). - ¹³C NMR (50 MHz): $\delta = 13.89$ (q, OCH₂CH₃), 13.94 (q, OCH₂CH₃), 22.3 (t), 23.1 (t), 23.5 (q, CCH₃), 26.8 (t), 31.2 (t), 46.7 (d, =CCH), 58.5 (s, CCO₂), 60.6 (t, OCH₂), 61.0 (t, OCH₂), 113.1 (t, =CH₂), 146.5 (s, =C), 170.7 (s, CO₂), 171.8 (s, CO₂). - MS m/z (%): 268 (12) [M⁺], 222 (21), 195 (100), 194 (27), 167 (38), 149 (50), 148 (30), 121 (42), 93 (21), 79 (28). - HRMS: C₁₅H₂₄O₄: calcd. 268.1675; found 268.1668 ± 2 ppm.

Diethyl 3-Methylenecyclohexane-1,1-dicarboxylate (10f): Flash column chromatography (50/1 gradient to 30/1) yielded 200 mg (83%) of **10f** [R_f (10/1) = 0.45] as a colorless oil. - IR (neat): $\tilde{\nu}$ = 2940 cm^{-1} , 1733, 1254, 1244, 1189, 1066. - ^1H NMR (200 MHz): δ = 1.18 (t, J = 7.1 Hz, 6 H, OCH_2CH_3), 1.62 (m, 2 H), 2.02 (m, 4 H), 2.60 (s, 2 H, CCH_2CCO_2), 4.10 (q, J = 7.1 Hz, 2 H, OCH_2), 4.11 (q, J = 7.1 Hz, 2 H, OCH_2), 4.67 (s, 2 H, $=\text{CH}_2$). - ^{13}C NMR (50 MHz): δ = 14.0 (q, OCH_2CH_3), 24.1 (t), 31.0 (t), 33.9 (t), 39.5 (t, $=\text{CCH}_2\text{CCO}_2$), 56.5 (s, CCO_2), 61.1 (t, OCH_2), 110.4 (t, $=\text{CH}_2$), 144.1 (s, $=\text{C}$), 171.1 (s, CO_2). - MS m/z (%): 240 (26) [M^+], 195 (20), 194 (27), 167 (46), 166 (100), 139 (26), 138 (61), 137 (37), 121 (17), 93 (84). - HRMS: $\text{C}_{13}\text{H}_{20}\text{O}_4$: calcd. 240.1362; found 240.1356 $\pm$ 2 ppm. - $\text{C}_{13}\text{H}_{20}\text{O}_4$ (240.3): calcd. C 64.98, H 8.39; found C 65.15, H 8.50.

Oxidative Cyclization of Ester 9e: At -78°C under N_2 , 270 mg (1.0 mmol) of ester **9e** dissolved in 0.5 mL dry THF was added to a solution of 1.3 mmol LDA prepared from 0.19 mL $i\text{-Pr}_2\text{NH}$ and 0.81 mL $n\text{-BuLi}$ (1.6 M in hexanes) in 19 mL dry THF. The mixture was stirred for 30 min between -78 and -70°C . At this temperature, **2** was added in portions to the solution until no more oxidant was consumed and the reaction mixture remained green-blue (2.7 equiv.). The inhomogeneous mixture was stirred at -78°C for additional 90 min, and quenched by a few drops of sat'd NH_4Cl solution. The inhomogeneous mixture was diluted with 20 mL of ether and filtered through a pad of silica gel. After removing the solvent *in vacuo*, the inhomogeneous residue was preadsorbed on silica gel, transferred to a silica gel flash column and purified (50/1 gradient to 1/1). As the first fraction, pure ferrocene eluted, followed by a mixture of the cyclic compounds **10e** and unreacted **9e**. Through further flash column chromatography, the mixture was enriched in **10e**, but a complete removal of **9e** was impossible. The yield and distereomeric ratio of **10e** is based on ^1H NMR. The assignment of the diastereomers is in agreement with published data.²

1,5-trans-2,5-trans-10e: - ^1H NMR (400 MHz): δ = 1.16 (d, J = 7.1 Hz, 3 H, CH_3CH), 1.18 - 1.31 (m, 1 H, CH_3CHCH_2), 1.21 (t, J = 7.1 Hz, 3 H, $\text{CH}_3\text{CH}_2\text{O}$), 1.72 (s, 3 H, $\text{CH}_3\text{C}=\text{CH}_2$), 1.73-1.95 (m, 2 H, $\text{CH}_2\text{CHC}=\text{CH}_2$), 2.02 (m, 1 H, CH_3CHCH_2), 2.40 (m, 1 H, CH_3CH), 2.53 (m, 1 H, $\text{CHC}=\text{O}$), 2.75 (m, 1 H, $\text{CHC}=\text{CH}_2$), 3.98 (dq, J = 10.8, 7.1 Hz, 1 H, CH_2O), 4.03 (dq, J = 10.8, 7.1 Hz, 1 H, CH_2O), 4.67 (br. s, 1 H, $\text{CH}_2=\text{CH}_2$), 4.71 (br. s, 1 H, $\text{CH}_2=\text{CH}_2$). - ^{13}C NMR (100 MHz): δ = 14.3 (q, $\text{CH}_3\text{CH}_2\text{O}$), 21.3 (q, CH_3CH), 22.8 (q, CH_3CH), 29.8 (t, $\text{CH}_2\text{CHC}=\text{CH}_2$), 33.8 (t, CH_3CHCH_2), 36.9 (d, CHCH_3), 49.8 (d, $\text{CHC}=\text{CH}_2$), 55.4 (d, $\text{CHC}=\text{O}$), 59.8 (t, $\text{CH}_3\text{CH}_2\text{O}$), 110.8 (t, $\text{C}=\text{CH}_2$), 145.4 (s, $\text{C}=\text{CH}_2$), 174.7 (s, $\text{C}=\text{O}$). - MS m/z (%): 196 (9), 167 (3), 155 (20), 153 (17), 150 (13), 141 (28), 128 (24), 123 (100), 115 (28), 107 (39), 87 (33), 81 (100), 69 (39), 67 (48), 41 (67). - **1,5-trans-2,5-cis-10e:** - ^1H NMR (400 MHz): δ = 0.98 (d, J = 6.4 Hz, 3 H, CH_3CH), 1.17 (t, J = 7.1 Hz, 3 H, $\text{CH}_3\text{CH}_2\text{O}$), 1.23 (m, 1 H, CH_3CHCH_2), 1.52 (m, 1 H, $\text{CH}_2\text{CHC}=\text{CH}_2$), 1.64 (s, 3 H, $\text{CH}_3\text{C}=\text{CH}_2$), 1.86 (m, 2 H, CH_3CHCH_2 , $\text{CH}_2\text{CHC}=\text{CH}_2$), 2.14 (m, 2 H, $\text{CHC}=\text{O}$, CH_3CH), 2.79 (m, 1 H, $\text{CHC}=\text{CH}_2$), 4.06 (q, J = 7.1 Hz, 2 H, CH_2O), 4.61 (br. s, 1 H, $\text{CH}_2=\text{CH}_2$), 4.66 (br. s, 1 H, $\text{CH}_2=\text{CH}_2$). - ^{13}C NMR (100MHz): δ = 14.3 (q,

CH₃CH₂O), 19.5 (q, CH₃CH), 20.1 (q, CH₃-C=), 29.9 (t, CH₂CHC=), 33.1 (t, CH₃CHCH₂), 39.7 (d, CHCH₃), 51.8 (d, CHC=), 56.7 (d, CHC=O), 60.1 (t, CH₃CH₂O), 109.9 (t, C=CH₂), 146.5 (s, C=CH₂), 175.6 (s, C=O).

Figure 2: Characteristic NOE enhancements of *1,5-trans-2,5-cis-10e*.

One-Pot Synthesis of (*E/Z*)-Ethyl 8-(Trimethylsilyl)citronellate (9e**):** Ozone was bubbled through a mixture of 2.95 g (14.9 mmol) ethyl citronellate (**12**) in 27 mL dry CH₂Cl₂ at -78°C until a light blue colour persisted. Nitrogen was bubbled through the solution for 15 min to remove excess O₃, then 3 mL of Me₂S was added. After 30 min, the mixture was warmed to room temperature. The volatile components were evaporated and the residue was partitioned between water and ether. The aqueous layer was extracted three times with ether and dried over Na₂SO₄. The solvent was evaporated to give 2.56 g of aldehyde **15** that was used without purification in the following Wittig reaction.

In a second flask, 14.9 mL (23.8 mmol) *n*-BuLi (1.6 *M* in hexanes) was added slowly to a suspension of 8.3 g (22.3 mmol) ethyltriphenylphosphonium bromide in 100 mL dry THF at 0°C. After 15 min, the icebath was removed and the red homogeneous mixture was stirred for 30 min at room temperature. Trimethylsilylmethyl iodide (23.8 mmol, 3.54 mL) was added slowly at room temperature and the colour changed to brown-red. The mixture was stirred overnight during which a precipitate was formed. At -78°C, 14.75 mL (23.4 mmol) *n*-BuLi (1.6 *M* in hexanes) was added and the red mixture was stirred at room temperature for 90 min. A solution of 2.56 g (14.9 mmol) **15** in 5 mL dry THF was added at -78°C. After 30 min at -78°C, the mixture was stirred at room temperature for 4 hours during which a precipitate was formed. Sat'd NH₄Cl solution (40 mL) was added, the mixture was filtered. The aqueous layer was extracted with ether, the combined organic layers were washed with 5 mL of brine and dried over Na₂SO₄. A short flash column chromatography gave a mixture consisting of **9e**, PPh₃, and **16** as indicated by ¹H NMR. Triphenylphosphine was removed by refluxing the mixture with excess MeI for 3 h, followed by filtration of the phosphonium salt. The remaining mixture of **9e** and **16** was then separated by flash column chromatography yielding 30% of a 1:1 diastereomeric mixture of *E*- and *Z*-**9e** [*R_f*(10/1) = 0.47] and 16% of a 1:1 mixture of *anti/syn*-**16** [*R_f*(10/1) = 0.40] as colorless oils. The analytical data of **9e** are identical to the compound obtained by Krapcho dealkoxycarbonylation of **9c** (*vide supra*). - *syn/anti*-**16**: -

² Sakai, T.; Morita, K.; Matsumura, C.; Sudo, S.; Tsuboi, S.; Takeda, A. *J. Org. Chem.* **1981**, *46*, 4774-4779.

IR (neat): $\tilde{\nu}$ = 2958 cm⁻¹, 2875, 1738, 1459, 1450, 1372, 1299, 1287, 1251, 1199, 1159, 1096, 1074, 1035, 897, 842, 750. - ¹H NMR (400 MHz, C₆D₆): δ = -0.004 (s, 9 H, SiCH₃), 0.000 (s, 9 H, SiCH₃), 0.757 (d, J = 6.4 Hz, 3 H, CH₃CH), 0.761 (d, J = 6.4 Hz, 3 H, CH₃CH), 0.85 (t, J = 7.1 Hz, 2 $\times$ 3 H, CH₃CH₂O), 1.15 (m, 2 $\times$ 1 H, CH₂CH₂CHOTMS), 1.22-1.44 (m, 2 $\times$ 3 H, CH₂CHOTMS, CH₂CH₂COTMS), 1.52 (br. s, 2 $\times$ 3 H, CH₃C=), 1.88 (m, 2 $\times$ 2 H, CH₂C=O, CHCH₃), 2.05 (m, 2 $\times$ 1 H, CH₂C=O), 3.83 (m, 2 $\times$ 3 H, CH₃CH₂O, CHOTMS), 4.63 (s, 2 $\times$ 1 H, CH₂=), 4.79 (s, 2 $\times$ 1 H, CH₂=). - ¹³C NMR (100MHz, C₆D₆): δ = 0.2 (q, SiCH₃), 14.3 (q, CH₃CH₂O), 17.16 (q, CH₃C=), 17.23 (q, CH₃C=), 19.8 (q, CH₃CH), 19.9 (q, CH₃CH), 30.5 (d, CHCH₃), 32.8 (t, CH₃CHCH₂), 33.69 (t, CH₂COTMS), 33.75 (t, CH₂COTMS), 41.83 (t, CH₂C=O), 41.84 (t, CH₂C=O), 59.8 (t, CH₂O), 77.0 (d, CHOTMS), 77.1 (d, CHOTMS), 110.9 (t, CH₂=), 111.0 (t, CH₂=), 147.9 (s, C=), 148.0 (s, C=), 172.3 (s, C=O). - GC/MS m/z (%): 286 (2), 271 (4), 241 (3), 216 (6), 198 (4), 187 (7), 173 (3), 158 (4), 144 (13), 143 (100), 130 (3), 109 (8), 75 (5), 73 (22).

Tandem Alkoxy carbonylation/Oxidative Radical Cyclization/Carbocation Desilylation of 9e: At -78°C under N₂, 1.63 mL (2.6 mmol) of *n*-BuLi (1.6 M in hexanes) was added to a mixture of 367 mg (0.44 mL, 2.6 mmol) of 2,2,6,6-tetramethylpiperidine in 19.5 mL dry DME. After 30 min, 270 mg (1.0 mmol) of **9e** in 0.5 mL DME was added. After stirring for 30 min at -78°C, 0.114 mL (1.2 mmol) of ethyl chloroformate was added. After completion of the carboxylation, **2** was added in portions at 0°C until a blue-green color of the mixture persisted (2.3 equiv.). The reaction mixture was stirred for additional 30 min, and quenched by a few drops of sat'd NH₄Cl solution. The inhomogeneous mixture was diluted with 20 mL of ether and filtered through a pad of silica gel. After removing the solvent *in vacuo*, the inhomogeneous residue was preadsorbed on silica gel, transferred to a silica gel flash column and purified (50/1 gradient to 30/1). As the first fraction, pure ferrocene was eluted, followed by 250 mg (93%) of a 2:1 diastereomeric mixture of *trans/cis*-**10c** (*vide supra*).

Cyclic Alcohols 19 by Hydroboration/Oxidation (General Procedure): At 0°C under N₂, 0.94 mL (0.47 mmol) of 9-BBN (0.5 M in THF) was added dropwise to a mixture of 0.43 mmol **10a-c** in 0.5 mL dry THF and stirred for 1.5 h at room temperature. During a time of four hours, 4 $\times$ 0.28 mL (0.14 mmol) 9-BBN solution was added at 0°C. After stirring the mixture for 12 h at room temperature, the starting material disappeared by tlc. At 0°C, 1.5 mL 10% NaOH solution was added slowly. After stirring for 15 min, 1.5 mL of 30% H₂O₂ solution was added dropwise. The inhomogeneous mixture was stirred for 40 min. After addition of 10 mL of sat'd NH₄Cl solution, the aqueous layer was extracted with ethyl acetate and the combined organic phases were washed with 3 mL of 10% Na₂S₂O₃ solution. The organic layer was dried over Na₂SO₄, and the solvent was evaporated *in vacuo*. Purification by column chromatography (30/1 gradient to 1/1) gave alcohols **19a-c** as colorless oils.

(2*R)-Diethyl 2-[(*S**)-2-Hydroxy-1-methylethyl]cyclopentane-1,1-dicarboxylate (19a):** Yield: 73 mg (62%) [R_f (1/1) = 0.48]. - IR (neat): $\tilde{\nu}$ = 3531 cm^{-1} , 2979, 1725, 1260, 1178, 1113, 1036. - ^1H NMR (200 MHz): δ = 0.96 (d, J = 6.8 Hz, 3 H, CHCH_3), 1.19 (t, J = 7.1 Hz, 3 H, OCH_2CH_3), 1.21 (t, J = 7.1 Hz, 3 H, OCH_2CH_3), 1.31 - 1.66 (m, 2 H), 1.66 - 1.80 (m, 2 H), 1.84 - 1.98 (m, 2 H), 2.26 (br. s, 1 H, OH), 2.33 - 2.54 (m, 2 H), 3.34 (dd, J = 10.7, 6.5 Hz, 1 H, CH_2OH), 3.55 (dd, J = 10.7, 4.1 Hz, 1 H, CH_2OH), 4.08 (qd, J = 10.8, 7.1 Hz, 1 H, OCH_2), 4.12 (m, 1 H, OCH_2), 4.16 (qd, J = 10.8, 7.1 Hz, 1 H, OCH_2), 4.17 (qd, J = 10.8, 7.0 Hz, 1 H, OCH_2). - ^{13}C NMR (50 MHz): δ = 13.9 (q, OCH_2CH_3), 16.5 (q, CHCH_3), 22.8 (t, $\text{CH}_2\text{CH}_2\text{CCO}_2$), 29.3 (t, CHCH_2CH_2), 36.7 (t, CH_2CCO_2), 37.5 (d, CHCH_3), 49.7 (d, CHCH_2CH_2), 60.9 (t, OCH_2), 61.1 (t, OCH_2), 62.4 (s, CCO_2), 66.7 (t, CH_2OH), 171.7 (s, CO_2), 173.2 (s, CO_2). - MS m/z (%): 272 (0.6) [M^+], 254 (3), 241 (23), 227 (20), 213 (41), 181 (40), 180 (46), 173 (100), 167 (80), 166 (55), 153 (42), 141 (58), 140 (29), 139 (54), 138 (21), 135 (33), 127 (47), 109 (33), 95 (81), 93 (47), 81 (30), 79 (32), 67 (72), 55 (27). - HRMS: $\text{C}_{14}\text{H}_{24}\text{O}_5$ [$\text{M}^+ - \text{H}_2\text{O}$]: calcd. 254.1518; found 254.1513 $\pm$ 2 ppm.

Diethyl 2-(2-Hydroxyethyl)cyclopentane-1,1-dicarboxylate (19b): Yield: 82 mg (74%) [R_f (1/1) = 0.35]. - IR (neat): $\tilde{\nu}$ = 3542 cm^{-1} , 2960, 1727, 1258, 1194, 1181, 1158, 1064. - ^1H NMR (200 MHz): δ = 1.219 (t, J = 7.1 Hz, 3 H, OCH_2CH_3), 1.221 (t, J = 7.1 Hz, 3 H, OCH_2CH_3), 1.34 - 1.66 (m, 3 H, $\text{CH}_2\text{CH}_2\text{CHCH}_2\text{CH}_2\text{OH}$), 1.69 - 1.83 (m, 2 H, $\text{CH}_2\text{CH}_2\text{CHCH}_2\text{CH}_2\text{OH}$), 1.84 - 2.03 (m, 2 H, $\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}$), 2.39 (m, 1 H, CH_2CCO_2), 2.60 (m, 1 H, CHCH_2), 2.92 (br. s, 1 H, OH), 3.61 (m, 2 H, CH_2OH), 4.09 (qd, J = 10.8, 7.1 Hz, 1 H, OCH_2), 4.11 (qd, J = 10.8, 7.0 Hz, 1 H, OCH_2), 4.18 (qd, J = 10.8, 7.0 Hz, 1 H, OCH_2), 4.20 (qd, J = 10.6, 7.0 Hz, 1 H, OCH_2). - ^{13}C NMR (50 MHz): δ = 13.9 (q, OCH_2CH_3), 14.0 (q, OCH_2CH_3), 22.7 (t, $\text{CH}_2\text{CH}_2\text{CCO}_2$), 31.2 (t), 34.3 (t), 34.7 (t), 42.6 (d, CHCH_2), 60.9 (t), 61.06 (t), 61.13 (t), 63.0 (s, CCO_2), 171.4 (s, CO_2), 172.9 (s, CO_2). - MS; m/z (%): 240 (5), 213 (45), 212 (22), 194 (12), 185 (17), 173 (55), 167 (47), 166 (100), 141 (43), 139 (39), 138 (24), 127 (46), 113 (24), 95 (60), 93 (44), 81 (34), 79 (28), 67 (45), 55 (24). - $\text{C}_{13}\text{H}_{22}\text{O}_5$ (258.3): calcd. C 60.45, H 8.59; found C 60.68, H 8.40.

(2*R,5*S**)-Diethyl 2-[(*S**)-2-Hydroxy-1-methylethyl]-5-methylcyclopentane-1,1-dicarboxylate (19c) and (2*R**,5*R**)-Diethyl 2-[(*S**)-2-Hydroxy-1-methylethyl]-5-methylcyclopentane-1,1-dicarboxylate (20c):** Yield: 106 mg (86%) [R_f (2/1) = 0.38]. - IR (neat): $\tilde{\nu}$ = 3533 cm^{-1} , 2976, 2966, 2939, 2902, 2877, 1722, 1368, 1296, 1255, 1211, 1185, 1096, 1071, 1037. - **19c:** - ^1H NMR (400 MHz): δ = 0.85 (d, J = 7.1 Hz, 3 H, CH_3CHCH_2), 1.00 (d, J = 6.8 Hz, 3 H, CH_3CHCH), 1.15 - 1.27 (m, 7 H, $\text{CH}_3\text{CH}_2\text{O}$, CH_3CHCH_2), 1.37 (m, 1 H, CHCHCH_2), 1.65 (m, 1 H, CHCH_2OH), 1.68 (br. s, 1 H, OH), 1.87 (m, 1 H, CHCHCH_2), 2.09 (m, 1 H, CH_3CHCH_2), 2.66 (dt, J = 11.6, 7.3 Hz, 1 H, CHCHCH_2OH), 2.79 (sext, J = 7.2 Hz, 1 H, CH_3CH), 3.36 (dd, J = 10.7, 6.8 Hz, 1 H, CHCH_2OH), 3.54 (dd, J = 10.7, 4.4 Hz, 1 H, CHCH_2OH), 4.09 (m, 2 H, CH_2O), 4.20 (m, 2 H, CH_2O). - ^{13}C NMR (100 MHz): δ = 13.9 (q, $\text{CH}_3\text{CH}_2\text{O}$), 14.1 (q,

CH₃CH₂O), 16.7 (q, CH₃CHCH), 18.6 (q, CH₃CHCH₂), 27.3 (t, CHCHCH₂), 32.3 (t, CH₃CHCH₂), 37.4 (d, CHCH₂OH), 41.0 (d, CH₃CHC), 47.6 (d, CHCHCH₂OH), 60.8 (t, CH₃CH₂O), 60.9 (t, CH₃CH₂O), 66.4 (t, CH₂OH), 66.5 (s, CCO₂), 171.5 (s, C=O), 171.9 (s, C=O). - **20c**: - ¹H NMR (400 MHz): δ = 0.97 (d, *J* = 6.8 Hz, 3 H, CH₃CHC), 1.07 (d, *J* = 6.9 Hz, 3 H, CH₃CHCH₂OH), 1.24 (m, 6 H, CH₃CH₂O), 1.63 (m, 3 H, CHCH₂OH, CH₃CHCH₂, CHCHCH₂), 2.01 (m, 3 H, CH₃CHCH₂, CHCHCH₂, OH), 2.27 (m, 1 H, CH₃CH), 2.57 (m, 1 H, CHCHCH₂OH), 3.42 (m, 1 H, CH₂OH), 3.58 (m, 1 H, CH₂OH), 4.15 (m, 4 H, CH₂O). - ¹³C NMR (100 MHz): δ = 14.0 (q, CH₃CH₂O), 15.2 (q, CH₃CHCH), 16.4 (q, CH₃CHC), 27.6 (t, CHCHCH₂), 30.8 (t, CH₃CHCH₂), 38.4 (d, CHCH₂OH), 45.1 (d, CH₃CHC), 50.0 (d, CHCHCH₂OH), 60.4 (t, CH₃CH₂O), 61.0 (t, CH₃CH₂O), 65.8 (t, CH₂OH), 66.8 (s, CCO₂), 169.8 (s, C=O), 172.5 (s, C=O). - MS *m/z* (%): 261 (2), 255 (17), 241 (10), 227 (13), 213 (4), 195 (30), 185 (100), 181 (28), 167 (167), 153 (21), 141 (29), 125 (18), 121 (27), 107 (28), 81 (43). - C₁₅H₂₆O₅ (286.4) calcd. C 62.91, H 9.15; found C 63.33, H 9.08.

Lactonization of 19a-c (General Procedure): Alcohols **19a-c** (0.2 mmol) were placed in a NMR-tube filled with CDCl₃. *p*-TsOH (3 mg, 0.015 mmol) was added and the reaction was followed by ¹H NMR. After completion (ca. 30 min), a few drops of Na₂CO₃ solution were added. The solvent was evaporated *in vacuo* and the product was purified immediately by column chromatography (30/1 gradient to 10/1) to give lactones **21a-c** as colorless oils as single diastereomers. In a competition experiment, it was shown that a prior separation of alcohols **19c** and **20c** was actually not necessary, since **20c** did not react under the same conditions.

(±)-**Ethyl 4-Methyl-1-oxohexahydrocyclopenta[*c*]pyran-7a-carboxylate (21a)**: Yield: 45 mg (quant.) [*R*_f (2/1) = 0.26]. - IR (neat): $\tilde{\nu}$ = 2965 cm⁻¹, 1743, 1728, 1252, 1159, 1129, 1029. - ¹H NMR (400 MHz): δ = 0.91 (d, *J* = 6.7 Hz, 3 H, CHCH₃), 1.24 (t, *J* = 7.1 Hz, 3 H, OCH₂CH₃), 1.45 (m, 1 H, CHCH₂CH₂), 1.62 (m, 1 H, CH₂CH₂CCO₂), 1.76 (m, 2 H, CHCH₂CH₂), 1.89 (ddd, *J* = 13.4, 10.0, 6.7 Hz, 1 H, CH₂CCO₂), 2.48 (m, 2 H, CHCHCH₂O), 2.59 (dddd, *J* = 13.3, 7.0, 3.9, 1.4 Hz, 1 H, CH₂CCO₂), 4.11 (m, 2 H, CHCH₂O), 4.17 (q, *J* = 7.1 Hz, 1 H, OCH₂CH₃), 4.18 (q, *J* = 7.1 Hz, 1 H, OCH₂CH₃). - ¹³C NMR (100 MHz): δ = 13.8 (q, CHCH₃), 13.9 (q, OCH₂CH₃), 25.1 (t, CH₂CH₂CCO₂), 26.6 (t, CHCH₂CH₂), 29.0 (d, CHCH₃), 36.5 (t, CH₂CCO₂), 47.0 (d, CHCH₂CH₂), 59.2 (s, CCO₂), 61.8 (t, OCH₂CH₃), 70.7 (t, OCH₂CH), 171.3 (s, CO₂), 171.8 (s, CO₂). - MS *m/z* (%): 226 (7), 182 (30), 181 (24), 154 (33), 153 (100), 135 (35), 109 (60), 95 (30), 67 (57). - HRMS: C₁₂H₁₈O₄: calcd. 226.1205; found 226.1199 ± 3 ppm. - C₁₂H₁₈O₄ (226.3): calcd. C 63.70, H 8.02; found C 63.43, H 8.42.

(±)-**Ethyl 1-Oxohexahydrocyclopenta[*c*]pyran-7a-carboxylate (21b)**: Yield: 42 mg (quant.) [*R*_f (2/1) = 0.25]. - IR (neat): $\tilde{\nu}$ = 2962 cm⁻¹, 1744, 1728, 1253, 1241, 1158, 1095. - ¹H NMR (200 MHz):

δ = 1.21 (t, J = 7.1 Hz, 3 H, OCH_2CH_3), 1.34 (m, 1 H), 1.43 – 1.72 (m, 3 H), 1.90 – 2.24 (m, 3 H), 2.35 (m, 1 H), 2.71 (m, 1 H), 4.14 (q, J = 7.1 Hz, 1 H, OCH_2CH_3), 4.15 (q, J = 7.1 Hz, 1 H, OCH_2CH_3), 4.23 (m, 2 H, OCH_2CH_2). - ^{13}C NMR (50 MHz): δ = 13.9 (q, OCH_2CH_3), 24.8 (t), 29.0 (t), 33.6 (t), 36.0 (t), 42.5 (d, CHCH_2), 59.8 (s, CCO_2), 62.0 (t, OCH_2CH_3), 67.2 (t, OCH_2CH_2), 171.4 (s, CO_2), 172.1 (s, CO_2). - MS m/z (%): 212 (1) [M^+], 168 (13), 167 (15), 140 (66), 139 (53), 112 (38), 111 (25), 95 (100), 94 (27), 93 (31), 79 (32), 67 (76), 55 (25), 53 (27). - HRMS: $\text{C}_{11}\text{H}_{16}\text{O}_4$: calcd. 212.1049; found 212.1043 $\pm$ 2 ppm.

($\pm$)-Ethyl 4,7-Dimethyl-1-oxohexahydrocyclopenta[*c*]pyran-7a-carboxylate (**21c**): Yield: 44 mg (89%) [R_f (5/1) = 0.30]. - IR (neat): $\tilde{\nu}$ = 2965 cm^{-1} , 2942, 2906, 2880, 1744, 1727, 1461, 1377, 1253, 1233, 1204, 1173, 1122, 1069, 1039, 1013. - ^1H NMR (400 MHz): δ = 0.91 (d, J = 7.1 Hz, 3 H, CH_3CHCH), 1.12 (d, J = 6.9 Hz, 3 H, CH_3CHC), 1.24 (t, J = 7.1 Hz, 3 H, $\text{CH}_3\text{CH}_2\text{O}$), 1.34 – 1.54 (m, 2 H, CH_3CHCH_2 , $\text{CH}_3\text{CHCHCH}_2$), 1.80 (m, 2 H, CH_3CHCH_2 , $\text{CH}_3\text{CHCHCH}_2$), 2.32 (m, 1 H, $\text{CH}_3\text{CHCH}_2\text{O}$), 2.47 (m, 1 H, CH_3CHC), 2.95 (dt, J = 11.0, 7.1 Hz, 1 H, CCHCHCH_3), 4.08 (d, J = 5.3 Hz, 2 H, CHCH_2O), 4.18 (dq, J = 7.1, 1.2 Hz, 2 H, $\text{CH}_3\text{CH}_2\text{O}$). - ^{13}C NMR (100 MHz): δ = 13.7 (q, CH_3CHCH_2), 14.1 (q, $\text{CH}_3\text{CH}_2\text{O}$), 15.5 (q, CH_3CHC), 26.3 (t, $\text{CCHCH}_2\text{CH}_2$), 29.5 (d, CHCH_2O), 33.2 (t, CCHCH_2), 44.0 (d, CCHCH_3), 47.2 (d, CHCHCH_3), 61.3 (s, CCO_2), 61.7 (t, $\text{CH}_3\text{CH}_2\text{O}$), 71.8 (t, CHCH_2O), 171.0 (s, C=O), 171.8 (s, C=O). - MS m/z (%): 240 (1), 199 (1), 198 (7), 197 (9), 185 (69), 167 (69), 153 (18), 139 (25), 123 (32), 107 (23), 93 (23), 81 (100), 79 (47), 67 (29), 67 (29), 53 (25), 41 (73). - $\text{C}_{13}\text{H}_{20}\text{O}_4$ (240.29) calcd. C 64.98, H 8.39; found C 65.10, H 8.40.

Dihydronepetalactone 11c and nor-Methyl Dihydronepetalactone 11a (General Procedure): A carefully N_2 -flushed mixture of 0.10 mmol lactones **21a** or **21c**, 7 mg (0.17 mmol) of anhydrous LiCl, 0.006 mL (0.33 mmol) of water, and 0.59 mL of DMSO was heated to 185°C for three hours. After cooling to room temperature, 3 mL sat'd NaHCO_3 solution and 3 mL of water were added. The aqueous layer was extracted three times with ether, washed with brine and dried over Na_2SO_4 . The solvent was evaporated under reduced pressure. The crude product was passed through a small silica gel column (pentane/ether 1/1) to give **11a,c** as colorless sole products.

($\pm$)-7-nor-Methyldihydronepetalactone **11a**: Yield: 11 mg (73%) of **11a** and 3 mg (13%) of recovered **20a**. - ^1H NMR (400 MHz): δ = 0.93 (d, J = 7.0 Hz, 3 H, CHCH_3), 1.36 – 1.55 (m, 2 H, $\text{CH}_3\text{CHCHCH}_2\text{CH}_2$), 1.74 – 1.84 (m, 3 H, $\text{CH}_2\text{CH}_2\text{CH}_2$), 2.20 (m, 1 H, CH_2CHCO_2), 2.31 (m, 1 H, CHCH_3), 2.42 (m, 1 H, CH_3CHCH), 2.95 (m, 1 H, CHCO_2), 4.10 (m, 2 H, OCH_2). - ^{13}C NMR (100 MHz): δ = 13.6 (q, CHCH_3), 25.8 (t, $\text{CH}_2\text{CH}_2\text{CHCO}_2$), 26.8 (t, $\text{CH}_3\text{CHCHCH}_2$), 30.3 (d, CHCH_3), 31.8 (t, CH_2CHCO_2), 40.6 (d, CHCHCH_3), 43.2 (d, CHCO_2), 70.5 (t, OCH_2CH), 175.1 (s, CO_2). - GC/MS m/z (%): 154 (9) [M^+], 113 (18), 95 (24), 81 (44), 68 (41), 67 (100), 41 (29).

(±)-**Dihydronepetalactone 11c**: Yield: 17 mg (97%) [R_f (5/1) = 0.29]. - IR (neat): $\tilde{\nu}$ = 2958 cm^{-1} , 2928, 2872, 1732, 1457, 1379, 1246, 1206, 1166, 1118, 1091, 1058. - ^1H NMR (400 MHz): δ = 0.86 (d, J = 7.0 Hz, 3 H, $\text{CH}_3\text{CHCH}_2\text{O}$), 1.10 - 1.22 (m, 1 H, CH_3CHCH_2), 1.16 (d, J = 6.4 Hz, 3 H, $\text{CH}_3\text{CHCHC=O}$), 1.40 (dddd, J = 18.7, 12.6, 10.5, 6.1 Hz, 1 H, $\text{CH}_2\text{CHCHCH}_2\text{O}$), 1.71 (m, 1 H, $\text{CH}_2\text{CHCHCH}_2\text{O}$), 1.87 (m, 1 H, CH_3CHCH_2), 1.97 (m, 1 H, $\text{CH}_3\text{CHCHC=O}$), 2.20 (m, 1 H, CHCH_2O), 2.39 (dd, J = 10.7, 9.4 Hz, 1 H, CHC=O), 2.49 (m, 1 H, $\text{CH}_2\text{CHCHCH}_3$), 3.99 (ddd, J = 11.0, 3.9, 1.5 Hz, 1 H, CH_2O), 4.05 (dd, J = 10.9, 10.4 Hz, 1 H, CH_2O). - ^{13}C NMR (100 MHz): δ = 13.1 (q, $\text{CH}_3\text{CHCH}_2\text{O}$), 19.3 (q, $\text{CH}_3\text{CHCHC=O}$), 26.3 (t, $\text{CH}_2\text{CHCHCH}_2\text{O}$), 31.0 (d, CHCH_2O), 35.0 (t, MeCHCH_2), 40.5 (d, CH_2CHCHMe), 41.5 (d, MeCHCHC=O), 50.5 (d, CHC=O), 69.9 (t, CH_2O), 174.3 (s, C=O). - GC/MS m/z (%): 168 (17), 153 (43), 139 (5), 126 (28), 113 (58), 109 (12), 100 (5), 95 (34), 81 (100), 67 (67), 55 (25), (39). - HRMS: $\text{C}_{10}\text{H}_{16}\text{O}_2$: calcd. 168.1150; found 168.1145 $\pm$ 2 ppm.

GC trace of **11c**: