Supporting Information

Efficient Rhodium Catalyzed Asymmetric Hydrogenation for the Synthesis of a New Class of *N*-Substituted β -Amino Acid Derivatives

Qian Dai, Weiran Yang and Xumu Zhang*

* Corresponding author. Email: xumu@chem.psu.edu

General procedure for preparation of 3-phenylamino-but-2-enoic acid methyl ester **1a (analogously **1a-1d**, **1f-1i**).** A mixture of methyl acetoacetate (1.16 g, 10 mmol), aniline (0.93 g, 10 mmol) and acetic acid (60.1 mg, 1mmol) was placed in a ultrasound bath Branson 1210 for 3h. At the end of the reaction, 5 mL of ethanol was added. The solution was dried with Na₂SO₄, filtered and concentrated with reduced pressure. The residue was purified by passing through a silica gel column to give the pure product as a white solid 1.35 g (70 %).

¹H NMR (300 MHz, CDCl₃) δ 10.38 (br s, 1H), 7.27-7.36 (m, 2H), 7.08-7.19 (m, 3H), 4.71 (s, 1H), 3.69 (s, 3H), 2.00 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 170.7, 159.1, 139.2, 129.0, 125.0, 124.4, 85.5, 50.2, 20.3.

3-Phenylamino-but-2-enoic acid ethyl ester (1b**).** This compound was obtained from ethyl acetoacetate by the same method used for **1a** as a yellow oil (60%). ¹H NMR (300 MHz, CDCl₃) δ 10.44 (br s, 1H), 7.30-7.35 (m, 2H), 7.08-7.18 (m, 3H), 4.72 (s, 1H), 4.17 (q, *J* = 7.1 Hz, 2H), 2.00 (s, 3H), 1.30 (t, *J* = 7.1 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 170.2, 158.7, 139.2, 128.9, 124.7, 124.1, 85.9, 58.5, 20.1, 14.4.

3-Phenylamino-pent-2-enoic acid ethyl ester (1c). This compound was obtained from 3-oxo-pentanoic acid ethyl ester by the same method used for **1a** as a yellow oil (58%). ^1H NMR (400 MHz, CDCl_3) δ 10.42 (br s, 1H), 7.30-7.34 (m, 2H), 7.09-7.18 (m, 3H), 4.75 (s, 1H), 4.16 (q, $J = 7.1$ Hz, 2H), 2.33 (q, $J = 7.5$ Hz, 2H), 1.30 (t, $J = 7.2$ Hz, 3H), 1.04 (t, $J = 7.5$ Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 171.2, 165.1, 139.7, 129.5, 125.6, 125.5, 84.5, 59.2, 25.9, 15.0, 12.8; MS (ESI) m/e 220 (MH^+); HRMS Calcd. for $\text{C}_{13}\text{H}_{18}\text{NO}_2$: 220.1338, found : 220.1336.

6-Methyl-3-phenylamino-hept-2-enoic acid ethyl ester (1d). This compound was obtained from 6-Methyl-3-oxo-heptanoic acid ethyl ester by the same method used for **1a** as a yellow oil (55%). ^1H NMR (300 MHz, CDCl_3) δ 10.34 (br s, 1H), 7.31-7.36 (m, 2H), 7.09-7.20 (m, 3H), 4.74 (s, 1H), 4.16 (q, $J = 7.1$ Hz, 2H), 2.28-2.33 (m, 2H), 1.48 (sept, $J = 6.6$ Hz, 1H), 1.28-1.37 (m, 5H), 0.77 (d, $J = 6.5$ Hz, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ 170.6, 163.8, 139.1, 129.0, 125.1, 125.0, 84.8, 58.6, 37.1, 30.1, 27.6, 22.1, 14.5; MS (ESI) m/e 262 (MH^+); HRMS Calcd. for $\text{C}_{16}\text{H}_{24}\text{NO}_2$: 262.1799, found : 262.1807.

3-(4-Fluoro-phenylamino)-but-2-enoic acid ethyl ester (1f). This compound was obtained from ethyl acetoacetate and 4-fluoro aniline by the same method used for **1a** as a yellow oil (68%). ^1H NMR (300 MHz, CDCl_3) δ 10.24 (br s, 1H), 6.97-7.07 (m, 4H), 4.68 (s, 1H) 4.14 (q, $J = 7.1$ Hz, 2H), 1.91 (s, 3H), 1.27 (t, $J = 7.1$ Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 170.4, 160.2 (d, $J = 244.9$ Hz), 159.0, 135.2 (d, $J = 2.9$ Hz), 126.6 (d, $J = 8.2$ Hz), 115.7 (d, $J = 22.5$ Hz), 85.8, 58.7, 20.0, 14.5.

3-(3-Bromo-phenylamino)-but-2-enoic acid ethyl ester (1g). This compound was obtained from ethyl acetoacetate and 3-bromo aniline by the same method used for **1a** as a colorless oil (59%). ^1H NMR (300 MHz, CDCl_3) δ 10.38 (br s, 1H), 7.21-7.24 (m, 2H), 7.14 (t, $J = 7.9$ Hz, 1H), 6.98 (d, $J = 8.3$ Hz, 1H), 4.71 (s, 1H), 4.12 (q, $J = 7.1$ Hz, 2H), 1.99 (s, 3H), 1.26 (t, $J = 7.1$ Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 170.0, 157.7, 140.7, 130.1, 127.3, 126.5, 122.4, 122.2, 87.4, 58.7, 20.1, 14.4; MS (ESI) m/e 284 (MH^+); HRMS Calcd. for $\text{C}_{12}\text{H}_{15}\text{NO}_2\text{Br}$: 284.0286, found : 284.0286.

3-(4-Methoxy-phenylamino)-but-2-enoic acid ethyl ester (1h). This compound was obtained from ethyl acetoacetate and *p*-anisidine by the same method used for **1a** as a yellow solid (70%). ^1H NMR (300 MHz, CDCl_3) δ 10.16 (br s, 1H), 7.01-7.04 (d, $J = 8.5$ Hz, 2H), 6.84-6.86 (d, $J = 8.6$ Hz, 2H), 4.65 (s, 1H), 4.14 (q, $J = 7.1$ Hz, 2H), 3.79 (s, 3H), 1.88 (s, 3H), 1.28 (t, $J = 7.1$ Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 170.4, 159.9, 157.4, 132.0, 126.7, 114.1, 84.6, 58.5, 55.3, 20.0, 14.5.

3-(3-Methoxy-phenylamino)-but-2-enoic acid ethyl ester (1i). This compound was obtained from ethyl acetoacetate and *m*-anisidine by the same method used for **1a** as a yellow oil (63%). ^1H NMR (300 MHz, CDCl_3) δ 10.41 (br s, 1H), 7.20 (t, $J = 8.1$ Hz, 1H), 6.61-6.70 (m, 3H), 4.69 (s, 1H), 4.14 (q, $J = 7.1$ Hz, 2H), 3.77 (s, 3H), 2.01 (s, 3H), 1.28

(t, $J = 7.1$ Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 170.2, 160.0, 158.6, 140.4, 129.5, 116.3, 110.1, 109.7, 86.1, 58.6, 55.0, 20.2, 14.4.

General procedure for preparation of 3-phenyl-3-phenylamino-acrylic acid ethyl ester **2a (analogously **1e**, **2b-2e**).** A mixture of ethyl benzoylacetate (1.92 g, 10 mmol), aniline (0.93g, 10 mmol) and *p*-toluenesulfonic acid monohydrate (0.19 g, 1mmol) was dissolved in 10 mL of ethanol and refluxed overnight. After the reaction mixture was cooled to room temperature, the solution was dried with Na_2SO_4 and concentrated under reduced pressure. The residue was purified by passing through a silica gel column to give the final product as a pale yellow oil (1.36g, 51%).

^1H NMR (300 MHz, CDCl_3) δ 10.65 (br s, 1H), 7.60-7.70 (m, 5H), 7.42 (t, $J = 7.4$ Hz, 2H), 7.16-7.30 (m, 1H), 7.00 (d, $J = 7.5$ Hz, 2H), 5.34 (s, 1H), 4.55 (q, $J = 7.1$ Hz, 2H), 1.66 (t, $J = 7.1$ Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 170.1, 159.0, 140.3, 135.9, 129.4, 128.6, 128.4, 128.2, 122.9, 122.2, 91.1, 59.3, 14.5.

4,4,4-Trifluoro-3-phenylamino-but-2-enoic acid ethyl ester (1e**).** This compound was obtained from ethyl 4,4,4-trifluoroacetoacetate and aniline by the same method used for **2a** as a colorless oil (45%). ^1H NMR (300 MHz, CDCl_3) δ 9.90 (br s, 1H), 7.32-7.37 (m, 2H), 7.20-7.27 (m, 3H), 5.38 (s, 1H), 4.22 (q, $J = 7.1$ Hz, 2H), 1.31 (t, $J = 7.1$ Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 169.6, 147.0 (q, $J = 31.5$ Hz), 138.3, 129.2, 128.9, 126.6, 126.0, 88.4 (d, $J = 5.3$ Hz), 60.0, 14.2.

3-Phenylamino-3-*p*-tolyl-acrylic acid methyl ester (2b). This compound was obtained from methyl 4-methyl-benzoylacetate and aniline by the same method used for **2a** as a pale yellow solid (60%). ^1H NMR (360 MHz, CDCl_3) δ 10.33 (br s, 1H), 7.27-7.29 (m, 2H), 7.11-7.15 (m, 4H), 6.93-6.98 (m, 1H), 6.72-6.75 (m, 2H), 5.05 (s, 1H), 3.78 (s, 3H), 2.37 (s, 3H); ^{13}C NMR (90 MHz, CDCl_3) δ 170.4, 159.3, 140.5, 139.6, 132.9, 129.1, 128.6, 128.1, 122.9, 122.2, 90.3, 50.6, 21.3; MS (ESI) m/e 268 (MH^+); HRMS Calcd. for $\text{C}_{17}\text{H}_{18}\text{NO}_2$: 268.1338, found : 268.1340.

3-(4-Fluoro-phenyl)-3-phenylamino-acrylic acid methyl ester (2c). This compound was obtained from methyl 4-fluoro-benzoylacetate and aniline by the same method used for **2a** as a white solid (58%). ^1H NMR (360 MHz, CDCl_3) δ 10.30 (br s, 1H), 7.33-7.36 (m, 2H), 7.09-7.13 (m, 2H), 6.94- 7.00 (m, 3H), 6.68-6.70 (m, 2H), 5.01 (s, 1H), 3.76 (s, 3H); ^{13}C NMR (90 MHz, CDCl_3) δ 170.2, 163.2 (d, $J = 249.5$ Hz), 157.9, 140.1, 131.8 (d, $J = 3.1$ Hz), 130.0 (d, $J = 8.4$ Hz), 128.6, 123.1, 122.2 (d, $J = 6.6$ Hz), 115.4 (d, $J = 21.4$ Hz), 90.7, 50.6; MS (ESI) m/e 272 (MH^+); HRMS Calcd. for $\text{C}_{16}\text{H}_{15}\text{NO}_2\text{F}$: 272.1087, found : 272.1073.

3-(2-Methoxy-phenyl)-3-phenylamino-acrylic acid methyl ester (2d). This compound was obtained from methyl 2-methoxy-benzoylacetate and aniline by the same method used for **2a** as a white solid (67%). ^1H NMR (300 MHz, CDCl_3) δ 10.52 (br s, 1H), 7.33-7.37 (m, 2H), 6.91-7.08 (m, 4H), 6.66-6.73 (m, 3H), 4.86 (s, 1H), 3.74 (s, 3H), 3.40 (s,

3H); ^{13}C NMR (75 MHz, CDCl_3) δ 170.5, 157.6, 155.9, 140.3, 130.8, 130.0, 128.1, 124.9, 123.1, 121.3, 120.7, 110.9, 88.4, 55.0, 50.4; MS (ESI) m/e 284 (MH^+); HRMS Calcd. for $\text{C}_{17}\text{H}_{18}\text{NO}_3$: 284.1287, found : 284.1283.

3-Phenylamino-3-*o*-tolyl-acrylic acid methyl ester (2e). This compound was obtained from methyl 2-methyl-benzoylacetate and aniline by the same method used for **2a** as a white solid (62%). ^1H NMR (360 MHz, CDCl_3) δ 10.59 (br s, 1H), 7.12-7.31 (m, 3H), 6.95-7.03 (m, 3H), 6.77-6.86 (m, 1H), 6.52-6.55 (m, 2H), 4.73 (s, 1H), 3.68 (s, 3H), 2.07 (s, 3H); ^{13}C NMR (90 MHz, CDCl_3) δ 170.5, 159.4, 139.7, 135.7, 135.4, 130.3, 129.1, 128.7, 128.6, 125.9, 123.0, 120.6, 88.9, 50.5, 19.3; MS (ESI) m/e 268 (MH^+); HRMS Calcd. for $\text{C}_{17}\text{H}_{18}\text{NO}_2$: 268.1338, found : 268.1341.

General Asymmetric Hydrogenation Procedure.

In a glovebox, 0.002 mmol of catalyst and 0.2 mmol of substrates were dissolved in 1 mL of TFE (2,2,2-trifluoroethanol). This solution was then transferred into an autoclave. The hydrogenation was performed at 50 °C (for **1**) and 80 °C (for **2**) under 6 atm of H_2 for 18 to 24 h. After carefully releasing the hydrogen, the reaction mixture was directly passed through a short silica gel plug. The enantiomeric excess was determined by Chiral HPLC and the conversion was determined by crude NMR spectrum.

3-Phenylamino-butylricacid methyl ester (4a).

^1H NMR (360 MHz, CDCl_3) δ 7.17-7.27 (m, 2H), 6.72 (t, J = 7.3 Hz, 1H), 6.64 (d, J = 7.7 Hz, 2H), 3.95-3.99 (m, 2H), 3.69 (s, 3H), 2.66 (dd, J = 5.2, 15.0 Hz, 1H), 2.44 (dd, J = 6.9, 15.1 Hz, 1H), 1.29 (d, J = 6.4 Hz, 3H); ^{13}C NMR (90 MHz, CDCl_3) δ 172.2, 146.7, 129.4, 117.7, 113.6, 51.6, 46.0, 40.7, 20.6; $[\alpha]_D^{20}$ = -5.5 (c = 1, CHCl_3) for 90.9 % ee; Chiralpak OD-H, hex:ipa = 99:1, 1mL/min, t_1 = 21.5 min, t_2 = 27.0 min.

3-Phenylamino-butyracetic acid ethyl ester (4b).

^1H NMR (300 MHz, CDCl_3) δ 7.18-7.23 (m, 2H), 6.64-6.76 (m, 3H), 4.16 (q, $J = 7.1$ Hz, 2H), 3.94-4.01 (m, 1H), 3.81 (br s, 1H), 2.65 (dd, $J = 5.3, 15.0$ Hz, 1H), 2.43 (dd, $J = 6.9, 15.0$ Hz, 1H), 1.26-1.34 (m, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ 172.3, 147.3, 129.8, 118.1, 114.0, 60.9, 46.4, 41.5, 21.1, 14.7; $[\alpha]_D^{20} = -2.2$ ($c = 1$, CHCl_3) for 94.8 % ee; Chiralpak OD-H, hex:ipa = 99:1, 1mL/min, $t_1 = 15.3$ min, $t_2 = 18.1$ min.

3-Phenylamino-pentanoic acid ethyl ester (4c).

^1H NMR (300 MHz, CDCl_3) δ 7.15-7.20 (m, 2H), 6.62-6.72 (m, 3H), 4.12 (q, $J = 7.2$ Hz, 2H), 3.74-3.78 (m, 1H), 2.56 (dd, $J = 5.8, 15.0$ Hz, 1H), 2.48 (dd, $J = 6.2, 15.0$ Hz, 1H), 1.60-1.66 (m, 2H), 1.24 (t, $J = 7.1$ Hz, 3H), 0.99 (t, $J = 7.4$ Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 172.1, 147.3, 129.4, 117.5, 113.5, 60.5, 51.8, 39.0, 27.8, 14.2, 10.5; MS (ESI) m/e 222 (MH^+); HRMS Calcd. for $\text{C}_{13}\text{H}_{20}\text{NO}_2$: 222.1494, found : 222.1498; $[\alpha]_D^{20} = 7.9$ ($c = 1$, CHCl_3) for 94.9 % ee; Chiralpak OD-H, hex:ipa = 99:1, 1mL/min, $t_1 = 12.8$ min, $t_2 = 14.4$ min.

6-Methyl-3-phenylamino-heptanoic acid ethyl ester (4d).

^1H NMR (300 MHz, CDCl_3) δ 7.16-7.27 (m, 2H), 6.63-6.73 (m, 3H), 4.13 (q, $J = 7.1$ Hz, 2H), 3.77 (m, 2H), 2.51-2.62 (m, 2H), 1.54-1.61 (m, 3H), 1.23-1.37 (m, 5H), 0.88-0.91 (m, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ 172.0, 147.2, 129.3, 117.3, 113.3, 60.4, 50.6, 39.3, 35.2, 32.8, 28.0, 22.6, 22.5, 14.2; MS (ESI) m/e 264 (MH^+); HRMS Calcd. for

C₁₆H₂₆NO₂ : 264.1964, found : 264.1957; [α]_D²⁰ = 2.7 (*c* = 1, CHCl₃) for 89.7 % ee;

Chiralpak OD-H, hex:ipa = 99:1, 1mL/min, *t*₁ = 10.5 min, *t*₂ = 12.5 min.

4,4,4-Trifluoro-3-phenylamino-butyric acid ethyl ester (4e).

¹H NMR (300 MHz, CDCl₃) δ 7.25 (t, *J* = 7.9 Hz, 2H), 6.85 (t, *J* = 7.0 Hz, 1H), 6.77 (d, *J* = 8.2 Hz, 2H), 4.46-4.64 (m, 1H), 4.17 (q, *J* = 7.1 Hz, 2H), 3.95-4.00 (m, 1H), 2.87 (dd, *J* = 4.5, 15.6 Hz, 1H), 2.66 (dd, *J* = 8.8, 15.6 Hz, 1H), 1.24 (t, *J* = 7.1 Hz, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 169.9, 146.2, 129.8, 119.8, 114.4, 61.8, 53.8 (q, *J* = 30.4 Hz), 35.5 (d, *J* = 1.7 Hz), 14.4; [α]_D²⁰ = 1.4 (*c* = 1, CHCl₃) for 78.9 % ee; Chiralpak OJ-H, hex:ipa = 99:1, 1mL/min, *t*₁ = 31.2 min, *t*₂ = 40.7 min.

3-(4-Fluoro-phenylamino)-butyric acid ethyl ester (4f).

¹H NMR (300 MHz, CDCl₃) δ 6.86-6.93 (m, 2H), 6.55-6.61 (m, 2H), 4.17 (q, *J* = 7.2 Hz, 2H), 3.84-3.92 (m, 1H), 3.69 (br s, 1H), 2.59 (dd, *J* = 5.4, 15.0 Hz, 1H), 2.42 (dd, *J* = 6.6, 15.0 Hz, 1H), 1.24-1.34 (m, 6H); ¹³C NMR (75 MHz, CDCl₃) δ 171.8, 155.9 (d, *J* = 235.3 Hz), 143.1, 115.7 (d, *J* = 22.2 Hz), 114.7 (d, *J* = 7.4 Hz), 60.5, 46.8, 40.8, 20.5, 14.2; MS (ESI) *m/e* 226 (MH⁺); HRMS Calcd. for C₁₂H₁₇NO₂F : 226.1243, found : 226.1248; [α]_D²⁰ = -2.3 (*c* = 1, CHCl₃) for 96.3 % ee; Chiralpak OJ-H, hex:ipa = 99:1, 1mL/min, *t*₁ = 30.1min, *t*₂ = 36.1 min.

3-(3-Bromo-phenylamino)-butyric acid ethyl ester (4g).

^1H NMR (300 MHz, CDCl_3) δ 7.02 (t, J = 8.0 Hz, 1H), 6.76-6.82 (m, 2H), 6.52-6.55 (m, 1H), 4.15 (q, J = 7.1 Hz, 2H), 3.88-3.94 (m, 2H), 2.59 (dd, J = 5.3, 15.1 Hz, 1H), 2.45 (dd, J = 6.4, 15.1 Hz, 1H), 1.24-1.33 (m, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ 171.5, 148.1, 130.6, 123.3, 120.2, 115.9, 112.1, 60.6, 45.8, 40.8, 20.4, 14.2; MS (ESI) m/e 286 (MH^+); HRMS Calcd. for $\text{C}_{12}\text{H}_{17}\text{NO}_2$: 286.0443, found : 286.0424; $[\alpha]_D^{20}$ = -1.6 (c = 0.9, CHCl_3) for 96.1 % ee; Chiralpak OJ-H, hex:ipa = 99:1, 1mL/min, t_1 = 33.4 min, t_2 = 44.9 min.

3-(4-Methoxy-phenylamino)-butyric acid ethyl ester (4h).

^1H NMR (300 MHz, CDCl_3) δ 6.76-6.81 (m, 2H), 6.59-6.64 (m, 2H), 4.14 (q, J = 7.1 Hz, 2H), 3.79-3.90 (m, 1H), 3.74 (s, 3H), 3.41 (br s, 1H), 2.60 (dd, J = 5.3, 15.0 Hz, 1H), 2.39 (dd, J = 6.8, 14.9 Hz, 1H), 1.23-1.28 (m, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ 171.9, 152.3, 140.9, 115.3, 114.8, 60.3, 55.6, 47.1, 40.9, 20.5, 14.1; $[\alpha]_D^{20}$ = -4.0 (c = 0.5, MeOH) for 93.8 % ee, and the absolute configuration was determined to be *R* by comparison with the reported data¹; Chiralpak OJ-H, hex:ipa = 99:1, 1mL/min, t_1 = 80.0 min, t_2 = 89.9 min.

3-(3-Methoxy-phenylamino)-butyric acid ethyl ester (4i).

^1H NMR (300 MHz, CDCl_3) δ 7.09 (t, J = 8.1 Hz, 1H), 6.19-6.30 (m, 3H), 4.15 (q, J = 7.1 Hz, 2H), 3.79-3.94 (m, 2H), 3.78 (s, 3H), 2.64 (dd, J = 5.1, 15.0 Hz, 1H), 2.43 (dd, J = 6.9, 15.0 Hz, 1H), 1.24-1.29 (m, 6H); ^{13}C NMR (75 MHz, CDCl_3) δ 171.7, 160.8, 148.2, 130.0, 106.5, 102.7, 99.4, 60.4, 55.0, 45.9, 40.9, 20.5, 14.2; MS (ESI) m/e 238 (MH^+); HRMS Calcd. for $\text{C}_{13}\text{H}_{20}\text{NO}_3$: 238.1443, found : 238.1428; $[\alpha]_D^{20}$ = -3.1 (c = 1, CHCl_3) for 95.7 % ee; Chiralpak OJ-H, hex:ipa = 99:1, 1mL/min, t_1 = 28.9 min, t_2 = 35.1 min.

3-Phenyl-3-phenylamino-propionic acid ethyl ester (5a).

^1H NMR (300 MHz, CDCl_3) δ 7.29-7.44 (m, 5H), 7.11-7.17 (m, 2H), 6.59-6.71 (m, 3H), 4.87 (t, J = 6.6 Hz, 1H), 4.62 (br s, 1H), 4.15 (q, J = 7.1 Hz, 2H), 2.83-2.90 (m, 2H), 1.23 (t, J = 7.1 Hz, 3H); ^{13}C NMR (75 MHz, CDCl_3) δ 171.6, 147.2, 142.6, 129.6, 129.2, 127.9, 126.7, 118.2, 114.1, 61.3, 55.4, 43.4, 14.6; $[\alpha]_D^{20}$ = -1.3 (c = 1, CHCl_3) for 92.3 % ee; Chiralpak OJ-H, hex:ipa = 90:10, 1mL/min, t_1 = 42.0 min, t_2 = 47.5 min.

3-Phenylamino-3-*p*-tolyl-propionic acid methyl ester (5b).

^1H NMR (360 MHz, CDCl_3) δ 7.28-7.31 (m, 2H), 7.12-7.18 (m, 4H), 6.60-6.73 (m, 3H), 4.85 (t, J = 6.7 Hz, 1H), 4.57 (br s, 1H), 3.68 (s, 3H), 2.84 (d, J = 6.5 Hz, 2H), 2.35 (s, 3H); ^{13}C NMR (90 MHz, CDCl_3) δ 171.6, 146.7, 139.1, 137.0, 129.4, 129.1, 126.1, 117.7, 113.6, 54.6, 51.8, 42.6, 21.0; MS (ESI) m/e 270 (MH^+); HRMS Calcd. for $\text{C}_{17}\text{H}_{20}\text{NO}_2$: 270.1494, found : 270.1493; $[\alpha]_D^{20}$ = -16.9 (c = 0.5, CHCl_3) for 91.1 % ee; Chiralpak OD-H, hex:ipa = 95:5, 1mL/min, t_1 = 10.3 min, t_2 = 13.3 min.

3-(4-Fluoro-phenyl)-3-phenylamino-propionic acid methyl ester (5c).

^1H NMR (300 MHz, CDCl_3) δ 7.34-7.38 (m, 2H), 6.99-7.15 (m, 4H), 6.67-6.73 (m, 1H), 6.54-6.57 (m, 2H), 4.83 (t, J = 6.6 Hz, 2H), 4.57 (br s, 1H), 3.67 (s, 3H), 2.81 (d, J = 6.6 Hz, 2H); ^{13}C NMR (75 MHz, CDCl_3) δ 171.4, 146.5, 137.8, 129.2, 127.7 (d, J = 8.1 Hz), 118.0, 115.6 (d, J = 21.2 Hz), 113.7, 54.2, 51.9, 42.6; MS (ESI) m/e 274 (MH^+); HRMS

Calcd. for C₁₆H₁₇NO₂F : 274.1243, found : 274.1240; $[\alpha]_D^{20} = 12.2$ ($c = 0.5$, CHCl₃) for 95.3 % ee; Chiralpak OD-H, hex:ipa = 95:5, 1mL/min, $t_1 = 11.6$ min, $t_2 = 18.4$ min.

3-(2-Methoxy-phenyl)-3-phenylamino-propionic acid methyl ester (5d).

¹H NMR (300 MHz, CDCl₃) δ 7.28-7.33 (m, 2H), 7.09-7.15 (m, 2H), 6.89-6.92 (m, 2H), 6.58-6.67 (m, 3H), 5.13-5.18 (m, 1H), 4.74 (br s, 1H), 3.93 (s, 3H), 3.65 (s, 3H), 2.94 (dd, $J = 5.1, 14.9$ Hz, 1H), 2.83 (dd, $J = 7.9, 14.9$ Hz, 1H); ¹³C NMR (75 MHz, CDCl₃) δ 172.1, 156.8, 146.9, 129.2, 129.1, 128.4, 127.6, 120.7, 117.5, 113.7, 110.5, 55.3, 51.7, 50.6, 40.2; MS (ESI) m/e 286 (MH⁺); HRMS Calcd. for C₁₇H₂₀NO₃ : 286.1443, found : 286.1444; $[\alpha]_D^{20} = -1.7$ ($c = 0.5$, CHCl₃) for 89.7 % ee; Chiralpak AS, hex:ipa = 99:1, 1mL/min, $t_1 = 14.8$ min, $t_2 = 20.4$ min.

3-Phenylamino-3-*o*-tolyl-propionic acid methyl ester (5e).

¹H NMR (360 MHz, CDCl₃) δ 7.39-7.42 (m, 1H), 7.09-7.19 (m, 5H), 6.68 (t, $J = 7.3$ Hz, 1H), 6.52 (d, $J = 7.7$ Hz, 2H), 5.02-5.06 (m, 1H), 4.50 (br s, 1H), 3.67 (s, 3H), 2.74-2.83 (m, 2H), 2.48 (s, 3H); ¹³C NMR (75 MHz, CDCl₃) δ 172.2, 147.1, 140.2, 135.4, 131.3, 129.6, 127.8, 127.1, 125.5, 118.2, 113.9, 52.4, 51.9, 41.4, 19.5; MS (ESI) m/e 270 (MH⁺); HRMS Calcd. for C₁₇H₂₀NO₂ : 270.1494, found : 270.1494; $[\alpha]_D^{20} = 2.7$ ($c = 0.5$, CHCl₃) for 79.3 % ee; Chiralpak OD-H, hex:ipa = 95:5, 1mL/min, $t_1 = 11.1$ min, $t_2 = 15.6$ min.

Reference

1. Park, S.; Lee, S. *J. Chem. Res. Synop.* **2001**, *11*, 498.